

The Risk of Rheumatoid Arthritis among Patients with Inflammatory Bowel Disease: A Systematic Review and Meta-analysis

Yi Chen

Xiangya Hospital Central South University

Lang Chen

Central South University Third Xiangya Hospital

Changsheng Xing

Houston Methodist Research Institute

Guangtong Deng

Xiangya Hospital Central South University

Furong Zeng

Xiangya Hospital Central South University

Tingting Xie

Xiangya Hospital Central South University

Lei Gu

Xiangya Hospital Central South University

Hui Xiang Yang (✉ yang_hx430@csu.edu.cn)

Xiangya Hospital Central South University <https://orcid.org/0000-0001-5467-2927>

Research article

Keywords: inflammatory bowel disease, Crohn disease, ulcerative colitis, rheumatoid arthritis, meta-analysis

Posted Date: January 24th, 2020

DOI: <https://doi.org/10.21203/rs.2.21860/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on June 17th, 2020. See the published version at <https://doi.org/10.1186/s12876-020-01339-3>.

Abstract

Background: Studies have suggested that patients with inflammatory bowel disease (IBD) have an increased risk of rheumatoid arthritis (RA). However, the available data on this association are inconsistent. This meta-analysis aimed to determine the association between IBD and the risk of RA.

Methods: Observational studies investigating the RA risk among patients with IBD (Crohn disease (CD) and/or ulcerative colitis (UC)) were searched in PubMed, Embase, and Web of Science from the date of inception to December 2019. The methodological quality of the included studies was assessed using the Newcastle-Ottawa Scale. Relative risks (RRs) and corresponding 95% confidential intervals (CIs) were pooled with a random-effects model. Heterogeneity was evaluated using I² statistics while publication bias was determined using Begg's and Egger's tests. Subgroup and sensitivity analyses were performed.

Results: A total of three cohort studies, three cross-sectional studies, and two case-control study were included in the meta-analyses. Compared to the non-IBD control or general population, there was a significantly higher risk of RA among patients with IBD (RR=2.59; 95% CI: 1.93–3.48). Moreover, both CD (RR=3.14; 95% CI: 2.46–4.01) and UC (RR=2.29; 95% CI: 1.76–2.97) were associated with a significantly increased risk of RA. However, heterogeneity was substantial across studies and the subgroup analyses failed to identify the potential source of heterogeneity.

Conclusions: Patients with IBD have a greater risk of developing RA. Rheumatologists should be consulted when patients with IBD present with undifferentiated joint complaints. However, more prospective cohort studies are needed to validate these results.

Introduction

Inflammatory bowel disease (IBD), which comprises Crohn disease (CD) and ulcerative colitis (UC), is a chronic non-specific inflammatory condition of the gastrointestinal tract (GI) [1]. While CD is characterized by discontinuous and transmural inflammation, which can affect all parts of the gastrointestinal tract, UC causes a continuous and superficial inflammation of the large intestine [1]. Besides the pathological changes and discomfort in the GI tract, up to 50% of the patients with IBD experience extra-GI manifestations, the most common being arthritis [2–4]. The majority of IBD-related arthritis cases parallel IBD activity and are self-limited and non-erosive [5].

Rheumatoid arthritis (RA) is a chronic synovitis-based systemic disease of unknown etiology, which is characterized by symmetrical invasive inflammation of multiple joints of the body [6]. Without early and adequate treatment, deformity and dysfunction of the affected joints due to severe bone destruction and absorption will occur in the late stage [7]. However, the signs and symptoms of RA may be quite similar to those of IBD-related arthritis, especially at the early stage of RA. Therefore, it may be difficult to distinguish them in the clinical setting, which may delay the diagnosis and treatment of RA. Moreover, as both IBD and RA are progressive and disabling diseases, the co-occurrence of these two conditions,

therefore, severely increases the disease burden and compromises the quality of life and prognosis of these patients [8].

In recent years, an increasing number of studies have noted that RA tended to cluster with IBD [9–14] and that patients with IBD are more likely to develop RA [15–18]. In contrast, a nationwide population-based case-control study indicated there was a decreased risk of RA in patients with IBD, although the decrease was not statistically significant [19]. Moreover, there is controversy regarding the risk of RA in patients with IBD when the association between RA and CD and between RA and UC is analyzed separately [11]. However, a systematic evaluation and quantitative synthesis of studies on the association between IBD and RA is still not available. Evaluating the current evidence on the association between them will facilitate our understanding of both conditions and provide a reference for better healthcare practices. Therefore, we aimed to conduct a meta-analysis to systematically analyze the association between IBD and the risk of RA; we discussed the underlying mechanisms as well. To the best of our knowledge, this is the first systematic review and meta-analysis focusing on the association between IBD with RA.

Methods

This study was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) [20] and Meta-analysis of Observational Studies in Epidemiology (MOOSE) guidelines [21].

Search strategy

Related publications were comprehensively searched in online databases (including PubMed, Embase, and Web of Science) from the inception date to December 2019 by two independent reviewers (Y.C. and L.C.). Terms used in the search included a combination of the following terms and their corresponding synonyms: “inflammatory bowel disease”, “Crohn disease”, “ulcerative colitis”, and “rheumatoid arthritis”. Moreover, the references of relevant reviews and included studies were screened manually to prevent omission. The detailed search strategy is shown in supplementary Table 1 (Additional file 1).

Study selection

We included studies meeting the following inclusion criteria: 1) written in English; 2) cross-sectional, case-control, or cohort studies which reported the risk estimates of RA among patients with IBD (CD and/or UC), or presented data to calculate these effect sizes; 3) containing a control group consisting of individuals without IBD or the general population. If different publications utilized duplicated data, the most comprehensive and newest one was included in this study. If different publications recruited overlapping population from the same database, we included the one with the best methodological quality in the statistical synthesis. Cohort study was considered as the best quality design, followed by case-control and cross-sectional study. The quality of studies with the same design was ranked by the quality assessment scale mentioned below. We excluded any reviews, expert opinions, case reports, and case series.

All the retrieved publications were exported to EndNote for selection. After removal of duplicated studies, the titles and abstracts of unique publications were screened to discard ineligible studies. Next, the full text of the remaining articles was checked for their eligibility according to the selection criteria. The entire selection process was independently performed by two authors (Y.C and L.C) and any disagreement was resolved by discussion and a third reviewer (G.T.D.) was consulted when necessary.

Data extraction

The following information was extracted from eligible studies: last name of the first author, year of publication, country of the population studied, study design, source of case and control groups, definition criteria for IBD and RA, age and gender of the population, sample size of each group, and effect sizes of the association between IBD and RA. Data were independently extracted and cross-checked by two investigators (Y.C. and L.C.).

Quality assessment

The methodological quality of included case-control and cohort studies on the association between IBD and the risk of RA was assessed using the Newcastle-Ottawa Scale [22], which included 3 domains: selection (4 items, 1 point for each item), comparability (1 item with 2 points), and ascertainment of exposure or outcome (3 items, 1 point for each item). A modification of the Newcastle-Ottawa Scale as described by Herzog et al [23] was used for the assessment of cross-sectional studies. Studies that scored ≥ 6 points were considered to be of good quality. The quality assessment was performed independently by 2 investigators (Y.C. and L.C), with disagreements resolved by discussion or consulting another investigator (G.T.D.).

Statistical analysis

Stata software version 12.0 (StataCorp, College Station, TX) was employed to conduct all the meta-analyses. Odds ratio, hazards ratio, and incident rate ratio were assumed to be similar estimates of relative risk (RR) given that both IBD and RA are relatively rare diseases [24]. If more than one risk estimate was presented in a study, the most adjusted one was used for the meta-analysis. All summary estimates were calculated with a DerSimonian-Laird random-effects model [25] due to the anticipated high level of study heterogeneity. Heterogeneity among studies was assessed by means of the I^2 statistics, in which $I^2 > 50\%$ was regarded as a marker of statistically significant heterogeneity. Additionally, subgroup analyses were conducted to better assess between-study variability and explore the potential impact factors. Sensitivity analysis was performed by excluding one study at a time and pooling the prevalence and risk estimates for the remaining studies, to evaluate the stability of the summary estimates. Publication bias was assessed with Egger's test [26] and Begg's test [27]. $P < 0.05$ was considered statistically significant in all cases, except for the publication bias test, in which the threshold value was set at 0.10.

Results

Study selection and characteristics

The electronic literature search identified 20,124 publications. The titles and abstracts of 16,109 unique studies were screened using the selection criteria, leaving 35 papers. The full texts of these articles were carefully reviewed. Finally, 11 studies meeting the selection criteria were selected for further qualitative analysis, as shown in Fig. 1.

The characteristics of these studies are outlined in Tables 1 and 2. The studies were published between 2007 and 2019 and conducted in the USA [10, 11, 13, 17, 18], Korea [9, 14, 15], Denmark [12, 16], and Finland [19]. Three publications [15–17] were retrospective cohort studies, two [18, 19] were case-control studies and the other six [9–14] were cross-sectional studies. Six studies [9, 12, 14–16] were based on nationwide population-based databases, four [10, 11, 13, 17] were based on large administrative databases and the other one [18] was based on single-center biobank. Two cross-sectional studies [9, 14] and one cohort study [15] recruited an overlapping population from the Korean National Health Insurance Database; we only included the cohort study in the statistical analysis. Likewise, since two studies [12, 16] recruited overlapping samples from the Danish National Patient Database, we only included the cohort study in the meta-analysis. The cross-sectional study conducted by Cohen et al [13] utilized two independent datasets which were marked as Cohen-2008-1 and Cohen-2008-2, respectively. Dataset Cohen-2008-1 was only included in the meta-analysis of risk of RA in patients with CD or UC alone, as it shared an overlapping sample with the cohort study conducted by Aletaha et al [17] when investigated the risk of RA in patients with IBD.

Table 1
Characteristics of the included studies-Part I

Study	Country	Design	Sample source	Definition criteria	
				IBD	RA
Aletaha-2019	USA	Retrospective Cohort	MarketScan Commercial Claims and Encounters database	≥ 2 claims at least 30 days apart with ICD-9 codes (555.x, 556.x)	≥ 2 claims at least 30 days apart with ICD-9 codes (714, 714.0, 714.8, 714.89, 714.9)
Bae-2016†	Korea	Cross-sectional	National Health Insurance Database	≥ 5 health contacts due to UC/ CD as the principal diagnosis with ICD-9 codes (K50, K51)	≥ 5 health contacts due to RA as the principal diagnosis with ICD-9 codes (M05, M06)
Burisch-2019	Denmark	Retrospective Cohort	National Patient Registry	≥ 2 diagnoses with ICD-8 codes (563.01, 563.19) or ICD-10 codes (K50, K51)	≥ 1 diagnosis with ICD-10 codes (M05, M06.0, M06.8, M06.9)
Cohen-2008-1	USA	Cross-Sectional	IMS Health Integrated Administration Claims Database	≥ 1 claim with ICD-9 codes (555.x, 556.x)	≥ 1 claim with ICD-9 codes (714.x)
Cohen-2008-2	USA	Cross-Sectional	MarketScan Commercial Claims and Encounters Database	≥ 1 claim with ICD-9 codes (555.x, 556.x)	≥ 1 claim with ICD-9 codes (714.x)
Halling-2017†	Denmark	Cross-Sectional	National Patient Registry	ICD-10 codes (K50, K51)	ICD-10 codes

CD, Crohn Disease; IBD, Inflammatory Bowel Diseases; ICD-8, International Classification of Disease 8th revision; ICD-9, International Classification of Disease 9th revision; ICD-10, International Classification of Disease 10th revision; RA, Rheumatoid Arthritis; UC, Ulcerative Colitis.

† These cross-sectional studies were not included in the statistical meta-analysis as the populations in these studies overlapped with those in the other cohort studies.

Study	Country	Design	Sample source	Definition criteria	
				IBD	RA
Kappelman-2011	USA	Cross-Sectional	PharMetrics Patient-Centric Database	≥ 3 health contacts with associated ICD-9 codes (555.x, 556.x), or ≥ 1 claim with associated medication history	ICD-9 codes
Park-2019	Korea	Retrospective cohort	National Health Insurance Database	ICD-10 codes (K50, K51) and registration codes (V130, V131) for the rare/intractable disease patient-support program	ICD-10 codes and registration codes for the rare/intractable disease patient-support program
Puolakka-2014	Finland	Case-control	National Registry of the Social Insurance Institution	ICD-10 codes (K50, K51)	ICD-10 codes (M05, M06)
Vanessa-2019	USA	Case-control	Single-center biobank	Questionnaire	≥ 2 diagnoses at least 30 days apart with ICD-9 codes (714.0, 714.9) plus use of a disease modifying antirheumatic drug
Weng-2007	USA	Cross-sectional	Northern California region of the Kaiser Permanente Medical Care Plan	≥ 2 diagnoses with ICD-9 codes (555.x, 556.x)	≥ 2 diagnoses with ICD-9 codes (714.0, 714.3x)

CD, Crohn Disease; IBD, Inflammatory Bowel Diseases; ICD-8, International Classification of Disease 8th revision; ICD-9, International Classification of Disease 9th revision; ICD-10, International Classification of Disease 10th revision; RA, Rheumatoid Arthritis; UC, Ulcerative Colitis.

† These cross-sectional studies were not included in the statistical meta-analysis as the populations in these studies overlapped with those in the other cohort studies.

Study	Country	Design	Sample source	Definition criteria	
				IBD	RA
Yang-2018†	Korea	Cross-Sectional	National Health Insurance Database	ICD-10 codes (K50, K51) and registration codes (V130, V131) for the rare/intractable disease patient-support program	ICD-10 codes (M05, M06)
<p>CD, Crohn Disease; IBD, Inflammatory Bowel Diseases; ICD-8, International Classification of Disease 8th revision; ICD-9, International Classification of Disease 9th revision; ICD-10, International Classification of Disease 10th revision; RA, Rheumatoid Arthritis; UC, Ulcerative Colitis.</p>					
<p>† These cross-sectional studies were not included in the statistical meta-analysis as the populations in these studies overlapped with those in the other cohort studies.</p>					

Table 2
Characteristics of the included studies-Part II

Study	Case type	Sample (size (n)/Age range(years)/Male (%))		Adjusted RR (95%CI)	Covariates adjustment	NOS score
		Cases	Controls			
Aletaha-2019	IBD	68,535/18–64/NM	42,371,769/18–64/NM	3.5 (3.3–3.8)	Age, sex, region, and insurance	7
Bae-2016	IBD	40,843/All/60.2	122,529/All/60.2	3.41 (2.61–4.44)	Age, sex, and insurance	7
	CD	12,646/All/69.1		3.74 (2.47–5.65)		
	UC	28,197/All/59.2		3.31 (2.49–4.40)		
Burisch-2019	IBD	11,777/≥18/NM	1 for 5 matched† /≥18/NM	2.11 (1.66–2.67)	Age, sex	8
	CD	NM/≥18/NM		3.59 (2.32–5.54)		
	UC	NM/≥18/NM		1.67 (1.21–2.30)		
Cohen-2008-1	IBD	18,603/≥18/44.3	66,969/≥18/44.3	2.72 (2.43–3.04)	Age, sex, region, and insurance	7
	CD	9,267/≥18/43.0	37,068/≥18/43.0	2.76 (2.40–3.18)		
	UC	11,220/≥18/46.0	44,880/≥18/46.0	2.76 (2.40–3.18)		

BMI, Body Mass Index; CCI, Charlson Comorbidity Index; CD, Crohn Disease; IBD, Inflammatory Bowel Diseases; RA, Rheumatoid Arthritis; RR, Relative Risk; UC, Ulcerative Colitis.

†. The sample size of CD and UC was not mentioned; however, each patient was paired with 5 age- and sex-matched 5 individuals from the general population.

Study	Case type	Sample (size (n)/Age range(years)/Male (%))		Adjusted RR (95%CI)	Covariates adjustment	NOS score
		Cases	Controls			
Cohen-2008-2	IBD	16,139/ ≥18/45.5	64,556/ ≥18/45.5	2.05 (1.84–2.29)	Age, sex, region, and insurance	7
	CD	7,404/ ≥18/42.7	29,604/ ≥18/42.7	2.37 (2.02–2.77)		
	UC	10,104/ ≥18/45.7	40,416/ ≥18/45.7	2.16 (1.88–2.49)		
Halling-2017	IBD	47,325/All/46	92,839/All/46	1.8 (1.5–2.0)	Age, sex, and region	8
	CD	13,343/All/42	26,172/All/42	2.1 (1.6–2.8)		
	UC	31,066/All/47	60,951/All/47	1.6 (1.3–1.9)		
Kappelman-2011	IBD	1,242/≤20/55	3,353/≤20/54	9.6 (3.9–23.8)	Age, sex, and region	8
	CD	737/≤20/56	1,997/≤20/56	15.7 (4.6–53.7)		
	UC	488/≤20/53	1,310/≤20/52	3.6 (0.8–16.1)		
Park-2019	IBD	35,581/ ≤80/61.6	142,324/ ≤80/61.6	4.24 (3.25–5.52)	Age, sex	8
	CD	11,803/ ≤80/70.9	47,212/ ≤80/70.9	5.59 (3.32–9.41)		

BMI, Body Mass Index; CCI, Charlson Comorbidity Index; CD, Crohn Disease; IBD, Inflammatory Bowel Diseases; RA, Rheumatoid Arthritis; RR, Relative Risk; UC, Ulcerative Colitis.

†. The sample size of CD and UC was not mentioned; however, each patient was paired with 5 age- and sex-matched 5 individuals from the general population.

Study	Case type	Sample (size (n)/Age range(years)/Male (%))		Adjusted RR (95%CI)	Covariates adjustment	NOS score
		Cases	Controls			
	UC	23737/ ≤80/56.9	94,948/ ≤80/56.9	3.90 (2.86– 5.32)		
Puolakka-2014	RA	7209/≥17/32	NM/NM/NM	0.74 (0.51– 1.03)	Age, sex	7
Vanessa-2019	RA	821/≥18/27	2455/≥18/27	3.82 (1.77– 8.41)	Age, sex, region, race, BMI, education, smoking, CCI	6
Weng-2007	IBD	12,601/ ≤89/47.5	50,404/ ≤89/47.5	1.9 (1.5– 2.3)	Age, sex, insurance, and smoking	9
	CD	4,021/ ≤89/44.1	1 for 4 matched/ ≤89/44.1	2.5 (1.8– 3.5)		
	UC	7,525/ ≤89/49.3	1 for 4 matched/ ≤89/49.3	1.5 (1.2– 2.0)		
Yang-2018	IBD	43,281/19– 75/70.1	1,127,261/19– 75/48.4	1.60 (1.52– 1.68)	No	6
	CD	13,925/19– 75/63.5		1.47 (1.34– 1.61)		
	UC	29,356/19– 75/57.6		1.67 (1.57– 1.77)		
BMI, Body Mass Index; CCI, Charlson Comorbidity Index; CD, Crohn Disease; IBD, Inflammatory Bowel Diseases; RA, Rheumatoid Arthritis; RR, Relative Risk; UC, Ulcerative Colitis.						
†. The sample size of CD and UC was not mentioned; however, each patient was paired with 5 age- and sex-matched 5 individuals from the general population.						

Study quality

A summary of the methodological quality scores of the included studies is shown in Table 2 and the detailed information is presented in supplementary Table 2 (Additional file 2). With respect to the risk of RA among patients with IBD, 10 studies with 11 datasets showed “good quality”, with a median score of 7 (range: 6–9).

Risk of RA in patients with IBD

The combined evidence from eight studies (three cohort studies, two case-control studies, and three cross-sectional studies) showed a significant increased risk of RA among patients with IBD (RR = 2.59, 95% CI: 1.93–3.48, $I^2 = 94.2\%$; Fig. 2). In addition, the pooled risk estimates of six datasets from five studies showed that the corresponding risks were both significantly increased in patients with CD (RR = 3.14, 95% CI: 2.46–4.01, $I^2 = 74.9\%$; Fig. 3), and UC (RR = 2.29, 95% CI: 1.76–2.97, $I^2 = 84.7\%$; Fig. 4).

Subgroup analyses stratified by study design, age of the targeted population, and year of publication were further performed. Results showed that the risk of RA among patients with IBD was significantly increased in all subgroups except for the group in the case-control studies (RR = 1.62, 95% CI: 0.32–8.07, $I^2 = 92.9\%$; Table 3). The corresponding risk in CD remained significantly increased in all subgroups. The association between UC and the risk of RA was less significant in the pediatric group (RR = 3.60, 95% CI: 0.80–16.10) and in the group comprised of both children and adults (RR = 2.41, 95% CI: 0.94–6.14, $I^2 = 95.4\%$). However, we failed to identify the source of heterogeneity based on these factors.

Table 3
Subgroup analysis on the risk of RA in IBD.

Subgroup analysis	No. of Studies	Pooled estimates (random-effects model)		
		Relative risk (95% CI)	P value	I ² statistics
The risk of RA in IBD	8	2.59 (1.93–3.48)	0.000	94.2%
Study design				
Cross-sectional	3	2.84 (1.85–4.37)	0.000	88.1%
Case-control	2	1.62 (0.32–8.07)	0.557	92.9%
Retrospective cohort	3	3.16 (2.26–4.41)	0.000	89.4%
Age				
Children	1	9.60 (3.90–23.80)	0.000	-
Adult	5	2.20 (1.53–3.17)	0.000	95.6%
Children and adult	2	2.83 (1.29–6.21)	0.010	95.3%
Publication year				
Recent 5 years	5	2.42 (1.45–4.02)	0.001	95.5%
Over 5 years	3	2.84 (1.85–4.37)	0.000	88.1%
The risk of RA in CD	6	3.14 (2.46–4.01)	0.000	74.9%
Study design				
Cross-sectional	4	2.71 (2.14–3.42)	0.000	70.9%
Retrospective cohort	2	4.37 (2.84–6.73)	0.000	38.8%
Age				
Children	1	15.70 (4.60–53.70)	0.000	-
Adult	3	9.59 (3.86–23.83)	0.000	51.0%
Children and adult	2	3.64 (1.66–8.00)	0.001	84.6%
Publication year				
Recent 5 years	2	4.37 (2.84–6.73)	0.000	38.8%
Over 5 years	4	3.17 (2.05–4.90)	0.000	75.1%
The risk of RA in UC	5	2.29 (1.76–2.97)		84.7%

CD, Crohn Disease; IBD, Inflammatory Bowel Disease; RA, Rheumatoid Arthritis; UC, Ulcerative Colitis.

Subgroup analysis	No. of Studies	Pooled estimates (random-effects model)		
		Relative risk (95% CI)	P value	I ² statistics
Study design				
Cross-sectional	4	2.15 (1.62–2.86)	0.000	83.6%
Retrospective cohort	2	2.55 (1.11–5.87)	0.000	92.8%
Age				
Children	1	3.60 (0.80–16.10)	0.094	-
Adult	3	2.22 (1.74–2.84)	0.000	81.4%
Children and adult	2	2.41 (0.94–6.14)	0.066	95.4%
Publication year				
Recent 5 years	2	2.55 (1.11–5.87)	0.027	92.8%
Over 5 years	4	2.15 (1.62–2.86)	0.000	83.6%
CD, Crohn Disease; IBD, Inflammatory Bowel Disease; RA, Rheumatoid Arthritis; UC, Ulcerative Colitis.				

Publication bias and Sensitivity analysis

No significant publication bias was found in the meta-analysis of the risk of RA among patients with IBD or UC. However, significant publication bias was detected in the meta-analysis of the association between CD and the risk of RA (Begg's test: $P = 0.060$; Egger's test: $P = 0.048$; Supplementary Fig. 1, Additional file 3).

The results of sensitivity analysis showed that the risk estimates of RA among patients with IBD, and with CD or UC alone, were stable (Supplementary Table 3, Additional file 4).

Discussion

Previous studies have proposed a link between IBD and RA. However, whether there is an increased risk of RA among patients with IBD remains controversial. In this study, we systematically reviewed the published literature and analyzed the association between IBD and the risk of RA. Our results showed that there was a significantly increased risk of RA in patients with IBD (RR = 2.59, 95% CI: 1.93–3.48). Moreover, the increase in risk remained significant when CD and UC were separately analyzed (CD: RR = 3.14, 95% CI 2.46–4.01; UC: RR = 2.29, 95% CI 1.76–2.97).

Although there was a similar trend in the correlation between IBD and RA among almost all the datasets included in this study, the effect sizes varied greatly among studies and considerable heterogeneity was detected. We performed a subgroup analysis based on the study design, age of the recruited participants,

and year of publication, but failed to identify the source of heterogeneity. The heterogeneity may be partly explained by the different studied populations, disease definition criteria, and covariate adjustment. However, these factors could not be effectively stratified due to limited datasets. To generate more conservative results, the random-effects model was applied in all scenarios. In cross-sectional studies, the IBD and RA were measured during the same timeframe and therefore provide weaker evidence than longitudinal studies for the causal relationship between IBD and RA. However, this issue did not affect the findings, since in the subgroup analysis stratified by study design, the correlations between IBD and the increased risk of RA were consistently significant in both cross-sectional and cohort studies.

There are a few potential mechanisms that may contribute to the association between IBD and RA. Genetic susceptibility plays a vital role in the pathogenesis of both IBD and RA and several susceptibility genes which code key modulators of immune and inflammatory response are shared by IBD and RA. It has been reported that polymorphisms in the IRF5 locus can confer susceptibility to UC, CD, and RA [28–30]. Meanwhile, studies have determined that IL2/IL21 and TNFRSF14 loci are common risk loci for RA and UC [31–34]. Likewise, the risk loci shared by CD and RA have been identified in two genome-wide association studies [35, 36]. It has been suggested that a chronically imbalanced mucosal immune response to gut microbiota in the genetically susceptible patients may play an important role in the pathogenesis of IBD [37–40]. A few studies indicated that there was a link between gut microbiota and RA as well. It has been reported that *Bacteroides* was decreased in amount in both CD and RA patients [41]. Additionally, studies utilizing animal models demonstrated that RA could be rescued by eliminating the bacteria residing in the intestine, while treating sterile mice with a specific intestine-residing bacterium, segmented filamentous bacteria, led to the development of RA [42].

RA is a chronic and progressive disease; early diagnosis and intervention are essential for better prognosis. The management strategy of RA has focused on the early identification of high-risk patients [43, 44]. However, it might be difficult for clinicians to distinguish arthritis related to IBD and early RA due to the obscure signs and symptoms. Collaboration between gastroenterologists and rheumatologists and close follow-up are suggested for the management of IBD patients presenting with undifferentiated arthritis.

The results presented in this meta-analysis must be interpreted with caution due to the several limitations of the study. First, our results were based on cross-sectional, case-control, and retrospective cohort studies; no prospective cohort study was available. Second, only a few studies were included in our meta-analyses. However, this could be partly compensated for by the large number of patients as all the studies included were based on nationwide or large-scale databases. Third, while smoking is the most important environmental risk factor for RA, only 2 studies [10, 18] included in this meta-analysis adjusted for the smoking status between the IBD group and the non-IBD control group. Fourth, there was considerable heterogeneity in our meta-analysis and we failed to identify the source of heterogeneity in the subgroup and sensitivity analyses. Fifth, significant publication bias was observed in the meta-analysis of the association between CD and risk of RA. Sixth, only the studies from USA, Denmark,

Finland, and Korea analyzed the risk of RA in patients with IBD, which may compromise the generalizability of the results for the populations of other countries.

Conclusion

Our study results indicate there is a significant association between IBD and the risk of RA. The findings of this meta-analysis suggest that rheumatologist consultation may be indicated when patients with IBD present with undifferentiated joint complaints. However, more prospective cohort studies are needed to validate these results.

Abbreviations

CD, Crohn Disease; CI, Confidence Interval; GI, Gastrointestinal Tract; IBD, Inflammatory Bowel Disease; I^2 , I-squared; MOOSE, Meta-analysis of Observational Studies in Epidemiology; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; RA, Rheumatoid Arthritis; RR, Relative Risk; UC, Ulcerative Colitis

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Acknowledgements

Not applicable.

Authors' contribution

In this study, Huixiang Yang supervised the whole project; Yi Chen and Huixiang Yang designed the study; Yi Chen, Lang Chen and Guangtong Deng were involved in article screening, data extraction, and data analyses; Yi Chen, Lang Chen and Changsheng Xing wrote the manuscript; and Guangtong Deng, Furong Zeng, Lei Gu, and Tingting Xie assisted in data extraction and analyses.

Funding

The authors received no financial support for this study.

Conflicts of interest

The authors declare that they have no conflict of interest.

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article and its additional files.

References

1. Baumgart DC, Sandborn WJ. Inflammatory bowel disease: clinical aspects and established and evolving therapies. *Lancet* 2007; 369(9573):1641-1657.
2. Ott C, Scholmerich J. Extraintestinal manifestations and complications in IBD. *Nat Rev Gastroenterol Hepatol* 2013; 10(10):585-595.
3. Olpin JD, Sjoberg BP, Stilwill SE, Jensen LE, Rezvani M, Shaaban AM. Beyond the Bowel: Extraintestinal Manifestations of Inflammatory Bowel Disease. *Radiographics* 2017; 37(4):1135-1160.
4. Harbord M, Annese V, Vavricka SR, et al. The First European Evidence-based Consensus on Extraintestinal Manifestations in Inflammatory Bowel Disease. *J Crohns Colitis* 2016; 10(3):239-254.
5. Orchard TR, Wordsworth BP, Jewell DP. Peripheral arthropathies in inflammatory bowel disease: their articular distribution and natural history. *Gut* 1998; 42(3):387-391.
6. Gibofsky A. Epidemiology, pathophysiology, and diagnosis of rheumatoid arthritis: A Synopsis. *Am J Manag Care* 2014; 20(7 Suppl):S128-135.
7. McInnes IB, Schett G. Pathogenetic insights from the treatment of rheumatoid arthritis. *Lancet Gastroenterol Hepatol* 2017; 389(10086):2328-2337.
8. Argollo M, Gilardi D, Peyrin-Biroulet C, Chabot JF, Peyrin-Biroulet L, Danese S. Comorbidities in inflammatory bowel disease: a call for action. *Lancet Gastroenterol Hepatol* 2019; 4(8):643-654.
9. Yang BR, Choi NK, Kim MS, et al. Prevalence of extraintestinal manifestations in Korean inflammatory bowel disease patients. *PloS one* 2018; 13(7):e0200363.
10. Weng X, Liu L, Barcellos LF, Allison JE, Herrinton LJ. Clustering of inflammatory bowel disease with immune mediated diseases among members of a northern california-managed care organization. *Am J Gastroenterol* 2007; 102(7):1429-1435.
11. Kappelman MD, Galanko JA, Porter CQ, Sandler RS. Association of paediatric inflammatory bowel disease with other immune-mediated diseases. *Arch Dis Child* 2011; 96(11):1042-1046.
12. Halling ML, Kjeldsen J, Knudsen T, Nielsen J, Hansen LK. Patients with inflammatory bowel disease have increased risk of autoimmune and inflammatory diseases. *World J Gastroenterol* 2017; 23(33):6137-6146.
13. Cohen R, Robinson D, Jr., Paramore C, Fraeman K, Renahan K, Bala M. Autoimmune disease concomitance among inflammatory bowel disease patients in the United States, 2001-2002. *Inflamm Bowel Dis* 2008; 14(6):738-743.

14. Bae JM, Choo JY, Kim KJ, Park KS. Association of inflammatory bowel disease with ankylosing spondylitis and rheumatoid arthritis: A nationwide population-based study. *Mod Rheumatol* 2017; 27(3):435-440.
15. Park SW, Kim TJ, Lee JY, et al. Comorbid immune-mediated diseases in inflammatory bowel disease: a nation-wide population-based study. *Aliment Pharmacol Ther* 2019; 49(2):165-172.
16. Burisch J, Jess T, Egeberg A. Incidence of Immune-Mediated Inflammatory Diseases Among Patients With Inflammatory Bowel Diseases in Denmark. *Clin Gastroenterol Hepatol*. <https://doi.org/10.1016/j.cgh.2019.03.040>.
17. Aletaha D, Epstein AJ, Skup M, Zueger P, Garg V, Panaccione R. Risk of Developing Additional Immune-Mediated Manifestations: A Retrospective Matched Cohort Study. *Adv Ther* 2019; 36(7):1672-1683.
18. Kronzer VL, Crowson CS, Sparks JA, Myasoedova E, Davis JM, 3rd. Comorbidities As Risk Factors for Rheumatoid Arthritis and Their Accrual After Diagnosis. *Mayo Clin Proc* 2019; 94(12):2488-2498.
19. Puolakka K, Rantalaiho V, Kautiainen H, Farkkila M. The concomitance of inflammatory bowel disease with incident rheumatoid arthritis and ankylosing spondylitis in Finland [Abstract]. *Ann Rheum Dis* 2014; 73:403-404.
20. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009; 62(10):1006-1012.
21. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *Jama* 2000; 283(15):2008-2012.
22. Wells GA, Shea D, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Available at: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp. Accessed 25 Jul 2019.
23. Herzog R, Alvarez-Pasquin MJ, Diaz C, Del Barrio JL, Estrada JM, Gil A. Are healthcare workers' intentions to vaccinate related to their knowledge, beliefs and attitudes? A systematic review. *BMC Public Health* 2013; 13:154.
24. Greenland S. Quantitative methods in the review of epidemiologic literature. *Epidemiol Rev* 1987; 9:1-30.
25. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7(3):177-188.
26. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; 315(7109):629-634.
27. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994; 50(4):1088-1101.
28. Chua KH, Lian LH, Khor WC, et al. Association between genetic polymorphisms in interferon regulatory factor 5 (IRF5) gene and Malaysian patients with Crohn's disease. *J Dig Dis* 2015; 16(4):205-216.

29. Li P, Lv H, Yang H, Qian JM. IRF5, but not TLR4, DEFB1, or VDR, is associated with the risk of ulcerative colitis in a Han Chinese population. *Scand J Gastroenterol* 2013; 48(10):1145-1151.
30. Dawidowicz K, Allanore Y, Guedj M, et al. The interferon regulatory factor 5 gene confers susceptibility to rheumatoid arthritis and influences its erosive phenotype. *Ann Rheum Dis* 2011; 70(1):117-121.
31. Festen EA, Goyette P, Scott R, et al. Genetic variants in the region harbouring IL2/IL21 associated with ulcerative colitis. *Gut* 2009; 58(6):799-804.
32. Zhernakova A, Alizadeh BZ, Bevova M, et al. Novel association in chromosome 4q27 region with rheumatoid arthritis and confirmation of type 1 diabetes point to a general risk locus for autoimmune diseases. *Am J Hum Genet* 2007; 81(6):1284-1288.
33. Ye BD, Choi H, Hong M, et al. Identification of Ten Additional Susceptibility Loci for Ulcerative Colitis Through Immunochip Analysis in Koreans. *Inflamm Bowel Dis* 2016; 22(1):13-19.
34. Raychaudhuri S, Remmers EF, Lee AT, et al. Common variants at CD40 and other loci confer risk of rheumatoid arthritis. *Nat Genet* 2008; 40(10):1216-1223.
35. Barrett JC, Hansoul S, Nicolae DL, et al. Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's disease. *Nat Genet* 2008; 40(8):955-962.
36. Stahl EA, Raychaudhuri S, Remmers EF, et al. Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. *Nat Genet* 2010; 42(6):508-514.
37. Xavier RJ, Podolsky DK. Unravelling the pathogenesis of inflammatory bowel disease. *Nature* 2007; 448(7152):427-434.
38. Strober W, Fuss I, Mannon P. The fundamental basis of inflammatory bowel disease. *J Clin Invest* 2007; 117(3):514-521.
39. Garrett WS, Lord GM, Punit S, et al. Communicable ulcerative colitis induced by T-bet deficiency in the innate immune system. *Cell* 2007; 131(1):33-45.
40. Ni J, Wu GD, Albenberg L, Tomov VT. Gut microbiota and IBD: causation or correlation? *Nat Rev Gastroenterol Hepatol* 2017; 14(10):573-584.
41. Yeoh N, Burton JP, Suppiah P, Reid G, Stebbings S. The role of the microbiome in rheumatic diseases. *Curr Rheumatol Rep* 2013; 15(3):314.
42. Wu HJ, Ivanov II, Darce J, et al. Gut-residing segmented filamentous bacteria drive autoimmune arthritis via T helper 17 cells. *Immunity* 2010; 32(6):815-827.
43. Allen PB, Olivera P, Emery P, et al. Review article: moving towards common therapeutic goals in Crohn's disease and rheumatoid arthritis. *Aliment Pharmacol Ther* 2017; 45(8):1058-1072.
44. Hitchon CA, Peschken CA, Shaikh S, El-Gabalawy HS. Early undifferentiated arthritis. *Rheum Dis Clin North Am* 2005; 31(4):605-626.

Additional Files

Additional file 1: Supplementary Table 1. Search strategies.

Additional file 2: Supplementary Table 2. Quality assessment the of included studies.

Additional file 3. Supplementary Fig. 1. Publication bias assessed by Begg's test and Egger's test.

Additional file 4. Supplementary Table 3. Sensitivity analysis.

Figures




Figure 1

Flowchart of the literature selection process.




Figure 2

Forest plots on the risk of rheumatoid arthritis among patients with inflammatory bowel disease.




Figure 3

Forest plots on the risk of rheumatoid arthritis among patients with Crohn disease.




Figure 4

Forest plots on the risk of rheumatoid arthritis among patients with ulcerative colitis.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)
- [Additionalfile2.docx](#)
- [Additionalfile3.docx](#)
- [Additionalfile4.docx](#)