

Ketone Body Rescued Seizure Behavior of LRP1 Deficiency by Modulating Glutamate Transport

Jin-Ming Zhang

The Second Clinical Medicine School of Guangzhou Medical University, Second Affiliated Hospital of Guangzhou Medical University

Ming-Jie Chen

The Third Medicine School of Guangzhou Medical University

Jiong-Hui He

The Third Medicine School of Guangzhou Medical University

Ya-Ping Li

The Second Clinical Medicine School of Guangzhou Medical University, Second Affiliated Hospital of Guangzhou Medical University

Zhi-Cai Li

The First Clinical Medicine School of Guangzhou Medical University

Zi-Jing Ye

The Second Clinical Medicine School of Guangzhou Medical University, Second Affiliated Hospital of Guangzhou Medical University

Yong-Hui Bao

Guangzhou Medical University

Bing-Jun Huang

Guangzhou Medical University

Wen-Jie Zhang

Guangzhou Medical University

Ping Kwan

University of Sydney

Yu-Ling Mao

The Third Affiliated Hospital of Guangzhou Medical University

Jing-da Qiao (✉ Joaquinjd@163.com)

The Second Clinical Medicine School of Guangzhou Medical University, Second Affiliated Hospital of Guangzhou Medical University

Research Article

Keywords: LRP1, Epilepsy, Ketone body, Glutamate transport, Drosophila

Posted Date: April 6th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1244600/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

LRP1, the low-density lipoprotein receptor 1, would be a novel candidate gene of Epilepsy according to our bioinformatic results and the animal study. In this study, we explored the role of LRP1 in Epilepsy and whether Beta-hydroxybutyrate, the principal ketone body of the ketogenic diet, can treat epilepsy caused by LRP1 deficiency. UAS/GAL4 system was used to establish different genotype models. Flies were given Standard, High-sucrose, and ketone body food randomly. The bang-sensitive test was performed on flies and seizure-like behavior was assessed. In morphologic experiments, we found that LRP1 deficiency caused partial loss of the ellipsoidal body and partial destruction of the fan-shaped body. Whole-body and glia LRP1 defect flies had a higher seizure rate compared to the control group. Ketone body decreased the seizure rate in behavior test in all LRP1 defect flies, compared to Standard and High sucrose diet. Overexpression of glutamate transporter gene *Eaat1* could mimic the ketone body effect on LRP1 deficiency flies. This study demonstrated that LRP1 defect globally or in astrocytes or neurons could induce epilepsy. The ketone body efficaciously rescued epilepsy caused by LRP1 knockdown. The results support screening for LRP1 mutations as discriminating conduct for individuals who require clinical attention and further clarify the mechanism of the ketogenic diet in Epilepsy, which could help Epilepsy patients making a precise treatment case by case.

1. Introduction

Epilepsy is a common neurological disorder affecting all age groups and is related to seizures, which are abnormal discharge of brain neurons. It is one of the transient dysfunctions of the brain that can occur in sudden and can be recurring (Fisher, van Emde Boas et al. 2005). Recent studies have shown that abnormal brain energy metabolism is closely related to the onset of epilepsy (Greene, Todorova et al. 2003, Tumiene, Ferreira et al. 2022). Myoclonic epilepsy with ragged-red fibers (MERRF), affecting complex I of the electron transport chain, is one of the examples concerning mitochondrial DNA (mtDNA) mutations that result in epileptic phenotypes (Silvestri, Moraes et al. 1992). Glucose is the predominant energy source in the brain (Pardridge 1983). The imbalance of glucose can lead to seizures (Singh and Strobos 1980, Greene, Todorova et al. 2003, Huang, Hsieh et al. 2005, Robinson, Krishnakumar et al. 2009). However, the mechanisms underlying glucose regulation and altered neuronal excitability remain poorly understood (Schwechter, Veliskova et al. 2003).

LRP1, the low-density lipoprotein receptor 1, has more than 40 different ligands and participates in many physiological processes. In the central nervous system (CNS), LRP1 is ubiquitously expressed and serves as a critical transport receptor as well as a modulator of several distinct signaling pathways in neurons (May, Rohlmann et al. 2004), astrocytes (Wyss-Coray, Loike et al. 2003), and microglia. It has been demonstrated that LRP1 is involved in nerve excitability. In neurons, the deletion of LRP1 caused hyperactivity, dystonia (May, Rohlmann et al. 2004). In astrocytes, early astrocyte dysfunction caused by LRP1 deletion is an important factor leading to abnormal excitability and morphologic changes of epileptic seizures during brain development in mice (Romeo, Mourabit et al. 2021). In embryonic cortical radial glia stem cells located in the telencephalon, the deletion of LRP1 caused a severe epileptic

phenotype (May, Rohlmann et al. 2004). However, the mechanism of LRP1 induced epilepsy remains unknown.

The ketogenic diet (KD) is a high-fat, calorie-restricted diet used to treat childhood epilepsies that do not respond to available drugs. Despite its clinical use for nearly 100 years, how the KD controls seizures remains unknown (DeVivo, Leckie et al. 1978, Bough, Wetherington et al. 2006, D'Andrea Meira, Romao et al. 2019). As the beta-hydroxybutyrate is the major bioactive metabolite of KD (Kraeuter, Guest et al. 2020, Ferrere, Tidjani Alou et al. 2021), in this study, we will explore the role of LRP1 in Epilepsy and the potential mechanism of KD in epilepsy treatment by using beta-hydroxybutyrate. Our results highlighted that the knockdown of LRP1 can cause epilepsy and abnormal brain structure, and we also found that LRP1 induced epilepsy is associated with glutamate imbalance and KD can rescue LRP1 induced epilepsy.

2. Methods And Materials

2.1 Data sources

To explore potential novel epilepsy-related genes by bio-information program., we investigated the gene expression profiles acquired by high-throughput sequencing of GSE139914 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE139914>) and GSE134697 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE134697>) from the GEO database (<https://www.ncbi.nlm.nih.gov/geo/>) after searching for keywords related to epilepsy. Then the gene expression matrix data of the epilepsy group and the control group were imported into GraphPad Prism (version 7.0.0) for t-tests. $P < 0.05$ was used to indicate a significant difference.

2.2 Fly stocks

Drosophila melanogaster (Fruit fly) is a classical animal model for epilepsy studies (Parker, Howlett et al. 2011). It had been used to investigate the pathogenic genes for epilepsy and other neuro-development disorders (Chung, Mao et al. 2020, Li, Wang et al. 2021). Recently, we also used *Drosophila* as a model to discover the novel epilepsy gene-UNC13B (Wang, Qiao et al. 2021). Thus, we used *Drosophila* as a model here to study the role of LRP1 in epilepsy.

The flies were fed standard cornmeal and maintained in the incubator at 25°C and 60–70% humidity on a 12:12-h light/dark cycle. *UAS-LRP1-RNAi* (THU2483/FBst0029548), *UAS-Scn1a-RNAi* (para-RNAi, positive control), and *UAS-Eaat1-RNAi* flies were donated by Tsing Hua Fly Center (Tsinghua University, Beijing, China). The double balancer line was purchased from Bloomington Fly Stock Center (Bloomington, IN, USA). The Gal4 driver line *tub-Gal4*, *elav-Gal4*, *repo-Gal4* were a gift from Prof. LIU Ji-Yong (Guangzhou Medical University, Guangzhou, China), and the *UAS-mCD8::GFP* line was a gift from Prof. KE Ya (The Chinese University of Hong Kong, Hong Kong). Canton-S was used as the WT line in this study. The *tub-Gal4*, *elav-Gal4*, *repo-Gal4* line was crossed with *LRP1-RNAi* to establish global *LRP1*

knockdown flies (*tub-Gal4 > LRP1-RNAi*), glial LRP1 knockdown flies (*repo-Gal4 > LRP1-RNAi*), and neuronal LRP1 knockdown flies (*elva-Gal4 > LRP1-RNAi*) (Ni, Zhou et al. 2011).

2.3 Seizure behavior test

Stress sensitivity or bang sensitivity (BS) experiments to assess stress-induced seizures were conducted as originally described by Ganetzky and Wu (Ganetzky and Wu 1982) with some modifications that have been described in detail previously (Fogle, Hertzler et al. 2016). The bang-sensitive (BS) test was conducted on flies 3–5 days after eclosion and seizure-like behavior was assessed (Parker, Howlett et al. 2011, Echeveste and Gros 2015). Flies were anesthetized with CO₂ and were transferred to new clean food vials one day before testing. About two to five flies were collected in empty plastic fly vials (2–5 flies/vial). Flies were allowed to recover for at least 30 min and mechanically stimulated with a vortex mixer (VWR, Radnor, PA, USA) at maximum speed for 10 second. The time for each fly to recover, which defined as standing, was scored, and a mean value from the scoring was taken for each vial. One vial was calculated as one N number in each group.

2.4 Three types of Diet

Standard medium: 7.12g Cornmeal, 1.68g of yeast powder, 0.56g AGAR, 8.5g sucrose, and 100ml D-D water, was blended and heated to the appropriate temperature (about 100 °C). After that, 0.47ml propionic acid was added.

High-sucrose medium: 7.12g cornmeal, 1.68g of yeast powder, 0.56g AGAR, 30g sucrose, and 100ml D-D water, was blended and heated and 0.47ml propionic acid was added.

Ketone Body medium: 3mM 0.04g beta-HB was added to the standard medium.

2.5 Morphology

Morphologic alteration of LRP1 defect in the brain was detected using GFP expression flies. The LRP1 knockdown and control flies labeled with membrane GFP, *tub-Gal4 > UAS-mCD8::GFP/UAS-Unc13b-RNAi*, and *tub-Gal4 > UAS-mCD8::GFP*, were generated by UAS-mCD8::GFP and double balancer flies. The fly brain was dissected, fixed, and permeabilized following the previous description (Wang, Qiao et al. 2021). Samples were observed using a confocal microscope (SP8; Zeiss, Jena, Germany) and analyzed with ImageJ software (National Institutes of Health, Bethesda, MD, USA).

2.6 Statistical analysis

All quantitative data are presented as mean \pm S.D. The Student's t-test was used to compare 2 independent or paired samples. One-way ANOVA was used to compare multiple samples, and Tukey's posthoc test was used to evaluate differences between two groups. Statistical analyses were performed with GraphPad Prism 7.00 and SPSS 20. The cutoff value for statistical significance is $P < 0.05$.

3. Results

3.1 LRP1 was Low Expression in Epilepsy Patients

GSE139914 contained 6 epilepsy samples and 41 controls from the human Brodmann Area 38. Samples of GSE134697 were from Neocortex and consisted of 17 epilepsy samples and 2 controls. With $P < 0.05$ as the threshold, we found that LRP1 expression was statistically significantly lower in the Epilepsy group than that in the control group. (Fig. 1).

3.2 Global Knockdown of LRP1 Induced Seizure Behavior

To further confirm the role of LRP1 in Epilepsy, we examined BS seizure-like behavior in *tub-Gal4 > LRP1-RNAi* flies. The *tub-Gal4 > LRP1-RNAi* flies exhibited the typical seizure-like behavior (Perkins 2006). About 30.72% of *tub-Gal4 > LRP1-RNAi* flies showed obvious seizure-like behavior, which was higher than the rate in *LRP1-RNAi* control flies (30.72%±13.28% [n = 10] vs 4.16%±7.20% [n = 7]; **P = 0.003). The *tub-Gal4 > Scn1a-RNAi* positive control flies had a higher rate of seizures than *tub-Gal4 > LRP1-RNAi* flies (48.92%±18.85% [n = 5] vs 30.72%±13.28% [n = 10]; *P = 0.05) (Figure. 2B).

3.3 LRP1 deficiency induced partial loss of ellipsoidal body

We examined possible morphologic changes in LRP1 knockdown flies based on GFP expression. The confocal images showed that LRP1 knockdown did not affect the main brain structures including the ellipsoidal body and fan sharp body (Figure. 3A–C), except for the partial loss of the ellipsoidal body (Figure. 3D) and partial destruction of the fan-shaped body (Figure. 3E, 3F).

3.4 Ketone body can Reduce Seizure-Like Behavior in LRP1 Defect Flies

To further explore the effects of the ketogenic diet and high glucose diet on the incidence of epilepsy in *tub-Gal4 > LRP1-RNAi* LRP1 knockdown flies and WT flies (*Canton-S*), we performed a bang-sensitive (BS) test to different diet application groups. About 30.72% of *tub-Gal4 > LRP1-RNAi* flies fed with Standard food showed obvious seizure-like behavior, which was higher than the rate in *tub-Gal4 > LRP1-RNAi* flies fed with Ketone body (beta-hydroxybutyrate, BHB) food (30.72%±13.28% [n = 10] vs 2.08%±5.89% [n = 8]; $P < 0.001$) (Figure. 2C). The *Canton-s* flies fed with high sucrose food also had a higher rate of seizures than those fed with Standard food (24.44%±10.99% [n = 6] vs 3.07%±5.63% [n = 7]; $P = 0.02$). However, there is no difference between the two groups of *tub-Gal4 > LRP1-RNAi* flies fed with the Standard food and high sucrose food (30.72%±13.28% [n = 10] vs 23.94%±16.90% [n = 11]; $P = 0.75$) (Figure. 2C).

3.5 Glial LRP1 Defect is Sufficient to Induce Seizure Behavior and can be Rescued by Ketone body

It is well known that glial LRP1 performs in regulating energy homeostasis, so we speculate that the expression of LRP1 in glia is closely related to epilepsy. Therefore, we established *repo-Gal4 > LRP1-RNAi* LRP1 flies and *elav-Gal4 > LRP1-RNAi* flies. In the same way, we analyzed the BS percentage of the three

models. We found that about 17.94% of *repo-Gal4 > LRP1-RNAi* flies showed obvious seizure-like behavior, which was higher than the rate in *elav-Gal4 > LRP1-RNAi* LRP1 knockdown flies (17.94% \pm 13.19% [n = 22] vs 4.60% \pm 8.46% [n = 21]; ***P = 0.001) (Figure. 4A). At the same time, the *tub-Gal4 > LRP1-RNAi* flies had a higher rate of seizures than *repo-Gal4 > LRP1-RNAi* flies (30.72% \pm 13.28% [n = 10] vs 17.94% \pm 13.19% [n = 22]; P = 0.02). These results indicate that the incidence of epileptic behavior in *Drosophila* by *repo-Gal4 > LRP1-RNAi* LRP1 knockdown can be further increased, which further suggests that not only is epilepsy caused by astrocytes closely related to LRP1 but also there are other ways causing epilepsy induced by LRP1 knockdown, suggesting a new research direction for us. To our surprise, contrasting with *tub-Gal4 > LRP1-RNAi* flies fed with high sucrose diet previously mentioned, the *elav-Gal4 > LRP1-RNAi* flies fed with high sucrose food had a higher rate of seizures than that feeding standard food (30.75% \pm 15.93% [n = 5] vs 4.60% \pm 8.46% [n = 21]; ****P < 0.001), which is worth further discussion. (Figure. 4B). The same as *tub-Gal4 > LRP1-RNAi* flies fed with KD, the *repo-Gal4 > LRP1-RNAi* flies fed with standard food also had a higher rate of seizures than that feeding BHB (17.94% \pm 13.19% [n = 22] vs 3.57% \pm 7.10% [n = 14]; **P = 0.004). (Figure. 4C).

3.6 Ketone body Rescued LRP1 Defect by Modulating Glutamate transporter

A previous study had shown that LRP1 defect may cause seizure by affecting glutamate transmission. Glutamate transporter is also important to synaptic glutamate transmission. Thus, we hypothesized that loss of function of LRP1 can cause seizure via modulating the glutamate transporters. To investigate this hypothesis, we knock down glutamate transporter, EAAT1, in various tissues. Global (*tub > Eaat1-RNAi*) or glial (*repo > Eaat1-RNAi*) knockdown of EAAT1 could induce seizure behavior (Figure. 5A). Glial specific knockdown of EAAT1 could induce a serious seizure and this phenotype could not be rescued by ketone body (BHB) (39.25% \pm 18.9%, n = 4 v.s. 27.13% \pm 13.16%, n = 4, p = 0.33). Interestingly, overexpression of EAAT1 in LRP1 knockdown flies could partially rescue (29.69% \pm 9.35%, n = 9 v.s. 9.06% \pm 5.44%, n = 7, ***p = 0.0001) (Figure. 5B). It suggested that downregulated Eaat1 was the mechanism of LRP1 defect causing seizure and the ketogenic diet could partially rescue it by up-regulating EAAT1.

4. Discussion

Epilepsy is a chronic brain disease that affects approximately 65 million people worldwide (Devinsky, Vezzani et al. 2018). Therefore, it is important to understand the underlying mechanisms for epilepsy, which can improve our management to this disease. In our present study, the bioinformatics results indicated that LRP1 would be a potential epilepsy gene. Animal experiments confirmed that LRP1 defects could induce seizures and abnormal brain structure. Additionally, we found that KD is a specific treatment with excellent outcomes, through the modulation of glutamate transporters, for epilepsy induced by LRP1 deficiency.

LRP1 plays a role in neural excitability. Here we used bioinformatic and fly models to further confirm its important role in Epilepsy. Our flies' data indicated that LRP1 defect could induce seizure behavior and

abnormal brain structure, which is consistent with the previous mouse study (Bres, Safina et al. 2020).

It has been reported that LRP1 is closely related to glucose homeostasis. Neuronal LRP1 deficiency impairs insulin signal transduction and downregulates GLUT3 expression in neurons, leading to reduced glucose uptake (Liu, Hu et al. 2015). In astrocytes, LRP1 mediates the molecular interaction between insulin-like growth factor 1 receptor (IGF1-R) and GLUT1 (Hernandez-Garzon, Fernandez et al. 2016). Glucose imbalance leads to seizures, and LRP1 plays an important role in maintaining glucose homeostasis. Thus, we use high sucrose and KD to feed the flies. Our data demonstrated that high sucrose diet can induce epilepsy in wild type flies, consistent with previous studies that high glucose concentrations provoked seizure in the adult rat model (Schwechter, Veliskova et al. 2003). However, to our surprise, LRP1 deficiency in global cells did not show a higher seizure rate in high sucrose feeding. This suggested the presence of another mechanism that LRP1 deficiency could result in further glucose accumulation, which meant that great seizure susceptibility could not be further aggravated by high sucrose. Further study required to demonstrate this hypothesis.

In this study, we found that the Epilepsy caused from LRP1 deficiency can be rescued by ketone body. The ketone body triggers a systemic shift from glucose metabolism toward the metabolism of fatty acids yielding ketone bodies, such as acetoacetate and BHB as substrates for energy (Ulamek-Kozioł, Pluta et al. 2016). Ketones, unlike glucose, are not likely deliver an immediate and large amount of energy necessary to initiate or sustain seizure activity (Greene, Todorova et al. 2001). But in our case of LRP1 deficiency, the BHB should not act via the glucose metabolism since high sucrose feeding did not induce seizure in LRP1 knockdown flies (Fig. 2C). Instead, BHB can rescue seizure in LRP1 deficiency by modulating synaptic glutamate cycling. Our behavior data showed that overexpression of glutamate transport gene EAAT1 could reduce the seizure rate of LRP1 deficiency flies, and BHB could not decrease seizure rate in EAAT1 knockdown flies. This set of data suggested that glutamate transporter would be one of the targets of KD for epilepsy. Additionally, in a mouse study, LRP1 deficiency result in NMDA receptor decrease [26]. These evidences suggested that KD may rescue epilepsy in LRP1 deficiency by modulating synaptic glutamate cycling.

In summary, we found that LRP1 can be a potential novel epilepsy gene. Screening for LRP1 mutations can identify individuals who require clinical attention for possibly frequent daily seizures. And we firstly found that BHB showed a rescue effect in the LRP1 deficiency model by modulating glutamate transport, which may help to understand the KD mechanism and establish precise treatment toward epilepsy case by case.

Declarations

Acknowledgments

This study was supported by Guangzhou Medical University (No. 2019A023). RNAi fly lines were donated from Tsing-Hua Fly Stock Center.

Funding

This work was supported by Guangzhou Medical University (No. 2019A023) and Guangzhou Postdoc Startup Fund (Jing-Da Qiao).

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

Study design: JM Zhang and JD Qiao. Bioinformatic experiment: YP Li. Manuscript writing: MJ Chen, JH He, P Kwan, YL Mao and JD Qiao. Experiment performance and data analysis: JM Zhang, MJ Chen, JH He, ZC Li, YP Li, ZJ Ye, YH Bao, BJ Huang, WJ Zhang, YL Mao and JD Qiao.

Data availability

Raw data were generated at Institute of Neuroscience, The Second Affiliated Hospital of Guangzhou Medical University. Derived data supporting the findings of this study are available from the corresponding author on request.

Ethics approval

No ethical approval is required in this study.

Consent to participate

Not applicable

Consent to publish

Not applicable

References

1. Bough, K. J., J. Wetherington, B. Hassel, J. F. Pare, J. W. Gawryluk, J. G. Greene, R. Shaw, Y. Smith, J. D. Geiger and R. J. Dingledine (2006). "Mitochondrial biogenesis in the anticonvulsant mechanism of the ketogenic diet." *Ann Neurol* 60(2): 223–235.
2. Bres, E. E., D. Safina, J. Muller, P. Bedner, H. Yang, X. Helluy, O. Shchyglo, S. Jansen, M. D. Mark, A. Esser, C. Steinhauser, S. Herlitze, C. U. Pietrzik, S. Sirko, D. Manahan-Vaughan and A. Faissner (2020). "Lipoprotein receptor loss in forebrain radial glia results in neurological deficits and severe seizures." *Glia* 68(12): 2517–2549.
3. Chung, H. L., X. Mao, H. Wang, Y. J. Park, P. C. Marcogliese, J. A. Rosenfeld, L. C. Burrage, P. F. Liu, D. R. Murdock, S. Yamamoto, M. F. Wangler, H. T. Chao, H. Y. Long, L. Feng, C. A. Bacino, H. J. Bellen and

- B. Xiao (2020). "De Novo Variants in CDK19 Are Associated with a Syndrome Involving Intellectual Disability and Epileptic Encephalopathy." *American Journal of Human Genetics* 106(5): 717–725.
4. D'Andrea Meira, I., T. T. Romao, H. J. Pires do Prado, L. T. Kruger, M. E. P. Pires and P. O. da Conceicao (2019). "Ketogenic Diet and Epilepsy: What We Know So Far." *Front Neurosci* 13: 5.
 5. Devinsky, O., A. Vezzani, T. J. O'Brien, N. Jette, I. E. Scheffer, M. de Curtis and P. Perucca (2018). "Epilepsy." *Nat Rev Dis Primers* 4: 18024.
 6. DeVivo, D. C., M. P. Leckie, J. S. Ferrendelli and D. B. McDougal, Jr. (1978). "Chronic ketosis and cerebral metabolism." *Ann Neurol* 3(4): 331–337.
 7. Echeveste, R. and C. Gros (2015). "Two-trace model for spike-timing-dependent synaptic plasticity." *Neural Comput* 27(3): 672–698.
 8. Ferrere, G., M. Tidjani Alou, P. Liu, A. G. Goubet, M. Fidelle, O. Kepp, S. Durand, V. Iebba, A. Fluckiger, R. Daillere, C. Thelemaque, C. Grajeda-Iglesias, C. Alves Costa Silva, F. Aprahamian, D. Lefevre, L. Zhao, B. Ryffel, E. Colomba, M. Arnedos, D. Drubay, C. Rauber, D. Raoult, F. Asnicar, T. Spector, N. Segata, L. Derosa, G. Kroemer and L. Zitvogel (2021). "Ketogenic diet and ketone bodies enhance the anticancer effects of PD-1 blockade." *JCI Insight* 6(2).
 9. Fisher, R. S., W. van Emde Boas, W. Blume, C. Elger, P. Genton, P. Lee and J. Engel, Jr. (2005). "Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE)." *Epilepsia* 46(4): 470–472.
 10. Fogle, K. J., J. I. Hertzler, J. H. Shon and M. J. Palladino (2016). "The ATP-sensitive K channel is seizure protective and required for effective dietary therapy in a model of mitochondrial encephalomyopathy." *J Neurogenet* 30(3–4): 247–258.
 11. Ganetzky, B. and C. F. Wu (1982). "Indirect Suppression Involving Behavioral Mutants with Altered Nerve Excitability in *DROSOPHILA MELANOGASTER*." *Genetics* 100(4): 597–614.
 12. Greene, A. E., M. T. Todorova, R. McGowan and T. N. Seyfried (2001). "Caloric restriction inhibits seizure susceptibility in epileptic EL mice by reducing blood glucose." *Epilepsia* 42(11): 1371–1378.
 13. Greene, A. E., M. T. Todorova and T. N. Seyfried (2003). "Perspectives on the metabolic management of epilepsy through dietary reduction of glucose and elevation of ketone bodies." *J Neurochem* 86(3): 529–537.
 14. Hernandez-Garzon, E., A. M. Fernandez, A. Perez-Alvarez, L. Genis, P. Bascunana, R. Fernandez de la Rosa, M. Delgado, M. Angel Pozo, E. Moreno, P. J. McCormick, A. Santi, A. Trueba-Saiz, C. Garcia-Caceres, M. H. Tschop, A. Araque, E. D. Martin and I. Torres Aleman (2016). "The insulin-like growth factor I receptor regulates glucose transport by astrocytes." *Glia* 64(11): 1962–1971.
 15. Huang, C. W., Y. J. Hsieh, M. C. Pai, J. J. Tsai and C. C. Huang (2005). "Nonketotic hyperglycemia-related epilepsy partialis continua with ictal unilateral parietal hyperperfusion." *Epilepsia* 46(11): 1843–1844.
 16. Kraeuter, A. K., P. C. Guest and Z. Sarnyai (2020). "Protocol for the Use of the Ketogenic Diet in Preclinical and Clinical Practice." *Methods Mol Biol* 2138: 83–98.

17. Li, D., Q. Wang, N. N. Gong, A. Kurolap, H. B. Feldman, N. Boy, M. Brugger, K. Grand, K. McWalter, M. J. G. Sacoto, E. Wakeling, J. Hurst, M. E. March, E. J. Bhoj, M. J. M. Nowaczyk, C. Gonzaga-Jauregui, M. Mathew, A. Dava-Wala, A. Siemon, D. Bartholomew, Y. Huang, H. Lee, J. A. Martinez-Agosto, E. M. C. Schwaibold, T. Brunet, D. Choukair, L. S. Pais, S. M. White, J. Christodoulou, D. Brown, K. Lindstrom, T. Grebe, D. Tiosano, M. S. Kayser, T. Y. Tan, M. A. Deardorff, Y. Q. Song and H. Hakonarson (2021). "Pathogenic variants in SMARCA5, a chromatin remodeler, cause a range of syndromic neurodevelopmental features." *Science Advances* 7(20).
18. Liu, C. C., J. Hu, C. W. Tsai, M. Yue, H. L. Melrose, T. Kanekiyo and G. Bu (2015). "Neuronal LRP1 regulates glucose metabolism and insulin signaling in the brain." *J Neurosci* 35(14): 5851–5859.
19. May, P., A. Rohlmann, H. H. Bock, K. Zurhove, J. D. Marth, E. D. Schomburg, J. L. Noebels, U. Beffert, J. D. Sweatt, E. J. Weeber and J. Herz (2004). "Neuronal LRP1 functionally associates with postsynaptic proteins and is required for normal motor function in mice." *Mol Cell Biol* 24(20): 8872–8883.
20. Ni, J. Q., R. Zhou, B. Czech, L. P. Liu, L. Holderbaum, D. Yang-Zhou, H. S. Shim, R. Tao, D. Handler, P. Karpowicz, R. Binari, M. Booker, J. Brennecke, L. A. Perkins, G. J. Hannon and N. Perrimon (2011). "A genome-scale shRNA resource for transgenic RNAi in *Drosophila*." *Nat Methods* 8(5): 405–407.
21. Pardridge, W. M. (1983). "Brain metabolism: a perspective from the blood-brain barrier." *Physiol Rev* 63(4): 1481–1535.
22. Parker, L., I. C. Howlett, Z. M. Rusan and M. A. Tanouye (2011). "Seizure and epilepsy: studies of seizure disorders in *Drosophila*." *Int Rev Neurobiol* 99: 1–21.
23. Perkins, K. L. (2006). "Cell-attached voltage-clamp and current-clamp recording and stimulation techniques in brain slices." *J Neurosci Methods* 154(1–2): 1–18.
24. Robinson, R., A. Krishnakumar, C. J. C. Paulose and m. neurobiology (2009). "Enhanced dopamine D1 and D2 receptor gene expression in the hippocampus of hypoglycaemic and diabetic rats." 29(3): 365–372.
25. Romeo, R., D. B. E. Mourabit, A. Scheller, M. D. Mark and A. Faissner (2021). "Low-Density Lipoprotein Receptor-Related Protein 1 (LRP1) as a Novel Regulator of Early Astroglial Differentiation." *Frontiers in Cellular Neuroscience* 15.
26. Schwechter, E. M., J. Veliskova and L. Velisek (2003). "Correlation between extracellular glucose and seizure susceptibility in adult rats." *Ann Neurol* 53(1): 91–101.
27. Silvestri, G., C. T. Moraes, S. Shanske, S. J. Oh and S. DiMauro (1992). "A new mtDNA mutation in the tRNA(Lys) gene associated with myoclonic epilepsy and ragged-red fibers (MERRF)." *Am J Hum Genet* 51(6): 1213–1217.
28. Singh, B. M. and R. J. Strobos (1980). "Epilepsia partialis continua associated with nonketotic hyperglycemia: clinical and biochemical profile of 21 patients." *Ann Neurol* 8(2): 155–160.
29. Tumiene, B., C. R. Ferreira and C. D. M. van Karnebeek (2022). "2022 Overview of Metabolic Epilepsies." *Genes* 13(3): 508.

30. Ulamek-Koziol, M., R. Pluta, A. Bogucka-Kocka and S. J. Czuczwar (2016). "To treat or not to treat drug-refractory epilepsy by the ketogenic diet? That is the question." *Ann Agric Environ Med* 23(4): 533–536.
31. Wang, J., J. D. Qiao, X. R. Liu, D. T. Liu, Y. H. Chen, Y. Wu, Y. Sun, J. Yu, R. N. Ren, Z. Mei, Y. X. Liu, Y. W. Shi, M. Jiang, S. M. Lin, N. He, B. Li, W. J. Bian, B. M. Li, Y. H. Yi, T. Su, H. K. Liu, W. Y. Gu, W. P. Liao and P. China Epilepsy Gene (2021). "UNC13B variants associated with partial epilepsy with favourable outcome." *Brain*.
32. Wyss-Coray, T., J. D. Loike, T. C. Brionne, E. Lu, R. Anankov, F. Yan, S. C. Silverstein and J. Husemann (2003). "Adult mouse astrocytes degrade amyloid-beta in vitro and in situ." *Nat Med* 9(4): 453–457.

Figures





Figure 1

LRP1 in Epilepsy patient and normal person. (A) There is no difference between normal person (Control) and epilepsy patients in GSE139914 database. (B) The LRP1 level of epilepsy patient is lower than that of normal person in GSE14697 database.

A



B



C




Figure 2

Knockdown of LRP1 induces seizure-like behavior in *Drosophila*. (A) Seizure behavior in flies. (B) Seizures occurred at a higher rate in LRP1 knockdown flies (*tub-Gal4>LRP1-RNAi*) than in WT flies (*Canton-S*) and other control groups. The *tub-Gal4>Scn1a-RNAi* positive controls had a higher rate of seizures than the *tub-Gal4>LRP1-RNAi* group. (C) Ketone body diet largely inhibited the seizure in LRP1 global knockdown flies.




Figure 3

LRP1 defect induced partial loss of ellipsoidal body (A-C) Serial images of the brain of WT flies from anterior to posterior showing the structure of Ellipsoidal body (A), fan shaped body (surf) (B), fan shaped body (deep) (C). (D-F) Serial images of the brain of a LRP1 knockdown fly from anterior to posterior showing the structure of Ellipsoidal body (D), fan shaped body (surf) (E), fan shaped body (deep) (F). Scale bar 50 μ m.




Figure 4

Glial knockdown of LRP1 sufficiently induced Seizure behavior. (A) Seizures occurred at a higher rate in *tub-Gal4>LRP1-RNAi* flies than in *repo-Gal4>LRP1-RNAi* flies. The *repo-Gal4>LRP1-RNAi* flies had a higher rate of seizures than of the *elva-Gal4>LRP1-RNAi* group. (B) The pan-neural LRP1 knockdown (*elav-Gal4>LRP1-RNAi*) flies fed with high sucrose food had a higher rate of seizures than that feeding standard food. Ketone body diet completely inhibited the seizure of those flies. (C) Seizures occurred at a higher rate in glial knockdown (*repo-Gal4>LRP1-RNAi*) flies fed with standard food than that feeding for ketone body diet.




Figure 5

Ketone body diet rescued LRP1 deficiency by downregulating glutamate transmission. (A) Knockdown of glial glutamate transport (*repo>Eaat1-RNAi*) induce serious seizure behavior, and ketone body diet could not inhibit the seizure rate in glial glutamate transporter knockdown flies. (B) Over-expression of Eaat1 could inhibit seizure caused by LRP1 deficiency. ST is short for Standard diet; KB is short for Ketone body diet.