Asymmetric Three-Component Olefin Dicarbofunctionalization Enabled by Photoredox and Copper Dual Catalysis

Wen-Jing Xiao (✉ wxiao@mail.ccnu.edu.cn)

Central China Normal University

Peng-Zi Wang

Central China Normal University

Yuan Gao

Central China Normal University

Xiao-Die Huan

Central China Normal University

Jun Chen

Central China Normal University

Jia-Rong Chen

Central China Normal University

Article

Keywords: copper dual catalysis, photoredox, dicarbofunctionalization

Posted Date: December 15th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-124676/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Communications on March 22nd, 2021. See the published version at <https://doi.org/10.1038/s41467-021-22127-x>.

Abstract

An intermolecular, enantioselective three-component radical vicinal dicarbofunctionalization reaction of olefins enabled by merger of radical addition and cross-coupling using photoredox and copper dual catalysis is presented here. Key to the success of this protocol relies on chemoselective addition of acyl and cyanoalkyl radicals, generated in situ from the redox-active oxime esters by a photo-catalytic N-centered iminyl radical-triggered C-C bond cleavage event, onto the alkenes to form new carbon radicals. Single electron metalation of such newly formed carbon radicals to TMSCN-derived $L1Cu(II)(CN)_2$ complex leads to asymmetric cross-coupling. This three-component process proceeds under mild conditions, and tolerates a diverse range of functionalities and synthetic handles, leading to valuable optically active β -cyano ketones and alkyldinitriles, respectively, in a highly enantioselective manner (> 60 examples, up to 97% ee).

Introduction

Alkenes are arguably one of the most privileged and versatile chemicals in organic synthesis because a diverse range of functional groups could be readily introduced across the $C = C$ π system by many classic vicinal difunctionalization reactions¹⁻⁴. In this context, one of the most investigated and fundamental transformations is the intermolecular three-component alkene vicinal dicarbofunctionalization (DCF) reaction, which allows installation of two different carbon fragments. While these reactions have been extensively explored in ionic chemistry, radical-mediated, especially the enantioselective versions of such reaction class remain largely unexplored⁵⁻¹⁰. Given the unique reactivity modes of radical species, the development of radical-mediated alkene vicinal DCF reactions would enable a wider variety of functional groups and synthetic handles to be incorporated, and hence produce valuable target molecules¹¹⁻¹⁵. While particularly promising, controlling the stereoselectivity in radical alkene vicinal dicarbofunctionalization reactions is a long-standing and fundamental challenge due to the intrinsically high reactivity and instability of radical intermediates¹⁶⁻¹⁸. In recent years, owing to their unique single-electron transfer ability and good coordination with chiral ligands, chiral copper catalysis opened a new and robust platform for development of asymmetric radical-mediated alkene vicinal DCF reactions (Fig. 1a)¹⁹⁻²¹. For example, the Liu group disclosed that carbon-centered radicals, which are formed in situ upon addition of CF_3 and (fluoro)alkyl²²⁻²⁷ or aryl radicals²⁸ to alkenes or enamides, could couple with TMSCN, boronic acid, or alkyne-derived chiral copper(II) complexes, leading to alkene vicinal DCF products with excellent enantiomeric excess. Recently, Zhang²⁹ and Liu³⁰⁻³² reported elegant examples of copper-catalyzed highly enantioselective radical alkene 1,2-dicarbofunctionalization with diaryliodonium salts and alkyl halides as radical sources, respectively. Despite the broad synthetic applicability of these methods, however, the intrinsic redox potential window of copper catalysts, which is critical to the generation of radicals, still results in significant limitations on the scope of radical precursors.

In recent years, visible-light photoredox catalysis has emerged as a powerful tool for organic chemists to develop many elusive radical-mediated chemical transformations with high levels of functional group tolerance^{33–36}. This activation mode also provides a promising approach for development of radical multicomponent reactions (in some cases with excellent stereoselectivity)^{37–42}. For example, the Studer group recently developed an asymmetric three-component cascade reaction of quinolines or pyridines with enamides using α -bromo carbonyl compounds as radical precursors under photoredox and phosphoric acid dual catalysis⁴³. Using this strategy, a range of chiral γ -amino-acid derivatives could be achieved with high chemo-, regio-, and enantioselectivity. Recently, our group^{44–47} and others^{48–54} introduced readily accessible redox-active oxime derivatives as precursors to generate iminyl radicals under single-electron transfer (SET) reduction or oxidation conditions (Fig. 1b). The resultant iminyl radicals further triggered smooth formation of sp^3 cyanoalkyl and sp^2 acyl^{55–57} radicals via a C-C bond cleavage process. Despite extensive synthetic utility of these carbon radicals, to our knowledge, their engagement in asymmetric multicomponent alkene vicinal DCF reactions is still challenging and unexplored^{58–59}. As a result, we aimed at developing a catalytic asymmetric three-component radical vicinal DCF reaction of alkenes with oximes and TMSCN (Fig. 1b). This protocol would provide an efficient and general approach for preparation of valuable optically active β -cyano ketones and alkyldinitriles^{60–62}.

Results

Optimization of reaction conditions. Our optimization began by reacting 2-vinylnaphthalene **1a** with oxime ester **3a** and TMSCN in a ratio of 1:3:3 in DMA under the dual photoredox and copper catalysis (Table 1). After some experimentation⁶³, we found that the target three-component reaction indeed occurred to give a moderate yield of desired product **6aa** with 88% ee, when using photocatalyst *fac*-Ir(ppy)₃ (1 mol%) and a combination of Cu(CH₃CN)₄PF₆ (0.5 mol%) and Box-type ligand **L1** (0.6 mol%) under irradiation of purple LEDs (Table 1, entry 1). However, several competing processes were also involved. For example, in addition to **6aa**, a significant amount of side products **sp-1**, **sp-2**, and **sp-3** were also detected, which might result from acyl radical **3a-II**-mediated two-component cross-coupling with **1a** and TMSCN, or its own dimerization. A brief screening of typical solvents such as DMF, CH₃CN, and THF showed that DMA was still the best of choice in terms of reaction efficiency (Table 1, entry 1 vs 2–5). Notably, when the catalyst loading of *fac*-Ir(ppy)₃ was decreased to 0.8 mol%, a cleaner reaction was observed, and a 64% yield of **6aa** was obtained without effect on the enantioselectivity (Table 1, entry 6). An extensive survey of other commonly used copper salts and chiral ligands established that the combination of Cu(CH₃CN)₄PF₆ and Box-type ligand **L1** were superior to others (Table 1, entries 6–8). Further optimization studies with respect to the loading of copper salt and concentration confirmed that a combination of 1.5 mol% of Cu(CH₃CN)₄PF₆ and 2.25 mol% of ligand **L1** with 0.8 mol% of *fac*-Ir(ppy)₃ at a concentration of 0.04 M gave the best results, with **6aa** being isolated with 74% yield and 90% ee (Table 1, entry 11). It was postulated that rational tuning of catalyst loading can help regulate the concentration of the reactive radical species, thus suppressing the competing side reaction pathways. As

expected, a series of control experimental results established that each component (light, photocatalyst, and copper salt) is critical to this asymmetric alkene vicinal dicarbofunctionalization reaction⁶³.

Substrate scope. With optimized reaction conditions established, we first investigated the substrate scope of alkenes by reacting with **3a** and TMSCN (Fig. 2). Notably, most of these alkenes are inexpensive and commercially available feedstock chemicals. As shown in Fig. 2A, aside from **1a**, simple neutral styrene **1b** and a range of styrene derivatives **1c-j** with electron-donating (e.g., Me, ^tBu, Ph) or electron-withdrawing (e.g., F, Cl, Br, OAc, Bpin) functional groups at the *para*-position of the aromatic ring are well tolerated, furnishing the corresponding products **6ba-ja** in 64–79% yields with 86–90% ee. Notably, halide substituents, F, Cl, and Br, as well as Bpin moiety remained intact after the reaction, thereby facilitating further modifications at their positions (e.g., products **6fa-ha** and **6ja**). A 1.0 mmol scale reaction of **1b** also proceeded smoothly to give comparable results (**6ba**, 74% yield, 90% ee), demonstrating the scalability of this process. Moreover, the reactions with alkenes **1k-o** bearing common substituents such as methoxy, methyl, fluoro, and bromo at the *meta*- or *ortho*-positions also worked well; and the expected products **6ka-oa** were isolated in 61–81% yields with 89–90% ee. 2-Vinylnaphthalene **1p** having a methoxy group and heterocycle-containing alkenes **1q-s** all proved to be suitable coupling partners, leading to **6pa-sa** with good yields and 86–93% ee. Remarkably, this protocol can also be successfully extended to biologically relevant molecule and pharmaceutical-derived styrene analogues (Fig. 2B). For instance, estrone, febuxostat-, and simple amino acid-derived alkenes **1t-v** reacted well to give the desired acylcyanation products **6ta**, **6ua**, and **6va** with good stereoselectivity, respectively. As a result, our protocol should be of potential use for late-stage structural modification of drug and complex compounds. Unfortunately, the current catalytic system is not applicable to simple unactivated or electron-deficient alkenes.

Then, we continued to evaluate the generality of this asymmetric three-component reaction by using a representative set of oxime esters, which can be easily prepared in two steps from the relevant ketone precursors. As shown in Fig. 3A, a range of aryl ketone-derived oxime esters **3b-g** with electronically diverse functional groups (e.g., Me, OMe, ^tBu, F, Cl, or Br) at the *para*-position of the phenyl ring reacted well with **1b** and TMSCN. And the expected alkene acylcyanation products **6bb-bg** were obtained with good yields (70–86%) and excellent enantiomeric excess (83–92% ee). As shown in the cases of **3h-k**, the change of the substitution pattern and steric hindrance of their phenyl ring has deleterious effect on the reaction efficiency or enantioselectivity, with the corresponding products **6bh-bk** being obtained with 67–85% yields and 89–92% ee. Single crystals of product **6bj** were obtained, and the absolute stereochemistry was determined to be *S* by X-ray crystallographic analysis⁶⁴, and all other coupling products were tentatively assigned by analogy with **6bj**. Remarkably, oxime esters **3m-p** derived from aliphatic ketone with various lengths of alkyl chains also reacted well with 2-vinylnaphthalene **1a** and TMSCN (Fig. 3B). The relative products **6am-ap** were isolated with 61–70% yields and 86–90% ee.

Encouraged by these results, we further attempted to extend the current dual photoredox and copper catalysis strategy to the asymmetric three-component vicinal DCF reaction of cycloketone-derived oxime esters, alkenes, and TMSCN (Fig. 4). Minor modification of the reaction conditions identified that a

combination of organic photocatalyst Ph-PTZ (1.25 mol%) and $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.5 mol%)/ligand **L1** (0.6 mol%) enabled the desired reaction to proceed smoothly under irradiation of 2×3 W purple LEDs at room temperature⁶³. This process also exhibited broad substrate scope and good functional tolerance with respect to both alkenes and oxime esters. As shown in Fig. 4A, a wide variety of commercially available styrenes containing neutral (**2a**), alkyl (**2b-c**), electron-withdrawing (**2e-h**), or aryl (**2i-j**) groups at the *para*-position of the phenyl ring could react well with oxime ester **4a**. The corresponding dinitrile products **7aa-ja** were obtained in 51–75% yields with 83–96% ee. The absolute stereochemistry of **7ja** was also confirmed to be *S*-configuration by X-ray diffraction⁶⁴. Again, as shown in the reactions of alkenes **2 k-n**, variation of the substitution pattern and steric hindrance of the aromatic ring could be well tolerated, leading to formation of products **7 ka-na** with 51–78% yields and 82–97% ee. Moreover, the reactions of 2-vinylnaphthalene **2o** and substrates containing heterocycle-fused ring (**2p**) or heteroaryl groups (e.g., **2q**, **2r**) all proceeded well to give products **7oa-ra** with moderate to good yields and excellent enantioselectivity (84–90% ee). Notably, styrenes (e.g., **2 s** and **2t**) derived from dihydroartemisinin and gibberellic acid could also participate in the reaction with good stereoselectivity, suggesting that the method can potentially be used in the late-stage modification of pharmaceutically relevant compounds.

Finally, we turned our attention to study the substrate scope of cyclobutanone oxime esters by reacting with styrene **2i** and TMSCN (Fig. 4B). Both oxetan-3-one and 1-Cbz-3-azetidinone derived oxime esters **4b** and **4c** reacted well to afford **7ib** and **7ic** in moderate to good yields with excellent enantioselectivity (87–89% ee). Mono-substituted oxime ester **4d** could participate in the reaction smoothly to deliver product **7id** as a mixture of diastereomers with good yield and high enantioselectivity. Note that sterically demanding oxime esters **4e-g** also proved to be compatible with the reaction, giving the expected products **7ie-ig** in good yields with 85–92% ee.

Synthetic applications. To showcase the potential synthetic utility of this asymmetric method in the construction of valuable skeletons, we performed diverse further transformations with the chiral β -cyano ketones and alkyldinitriles (Fig. 5)^{65–66}. For example, the cyano group of **6ba** and **7ia** could be easily converted to amide group by Pd-catalyzed hydrolysis using stoichiometric acetaldoxime in refluxing aqueous EtOH, giving the corresponding products **8** and **9** with good yields, respectively (Fig. 5a). Moreover, treatment of **7ia** with $\text{NiCl}_2/\text{NaBH}_4$ and Boc_2O in MeOH allowed efficient sequential reduction and protection of both cyano groups, with aliphatic chiral amine **10** being obtained with 66% yield and 90% ee (Fig. 5b). The synthesis of chiral ester **11** can also be achieved by treatment of **7ia** with alcoholysis. Notably, no notable loss of optical purity was detected in these manipulations.

Mechanistic studies. To gain some insight into the mechanism, we carried out several control experiments by using the substrates **1b**, **3a**, and TMSCN (Fig. 6). The target three-component reaction was completely inhibited, when stoichiometric radical scavenger 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) was introduced (Fig. 6a). Instead, the relevant TEMPO-adduct **12** was obtained in 76% yield, suggesting the possible involvement of acyl radical **3a-I** in this process. Moreover, the reaction of radical clock substrate **13** having a cyclopropyl moiety also proceeded smoothly to give ring-opening product **14**

in 64% yield with good stereoselectivity (Fig. 6b). These results indicated the intermediacy of radical species **13-A** and **13-B**, as well as the radical property of the reaction. Similar control experimental results were also observed in the case of cycloketone oxime ester-based asymmetric three-component reaction⁶³.

On the basis of these mechanistic studies and related literature^{19–21, 44–47}, we proposed a dual photoredox and copper-catalyzed mechanism for the present asymmetric three-component reaction as depicted in Fig. 7. The reaction starts with SET reduction of redox-active oxime esters **3a** and **4a** by the excited state photocatalyst to give iminyl radicals **3a-I** and **4a-I**, with release of carboxylic anion (RCO_2^-). Then, **3a-I** and **4a-I** undergoes C-C bond β -cleavage to form acyl and cyanoalkyl radicals **3a-II** and **4a-II**. Further facile trap of these carbon radicals by styrene derivatives **1** and **2** forms relatively more stable benzylic radicals **III**. On the other hand, the initially formed carboxylic anion (RCO_2^-) could also facilitate the ligand exchange between **L1**/copper(I) complex and TMSCN to form **L1Cu(I)CN** species. Such **L1Cu(I)CN** complex can further be oxidized by the oxidizing photocatalyst (PC^{*+}) via a SET process, and undergoes another ligand exchange with TMSCN to form **L1Cu(II)(CN)₂** complex, regenerating ground-state photocatalyst to close the photoredox catalysis cycle. Finally, **L1Cu(II)(CN)₂** traps the prochiral benzylic radical **III** to form a chiral high-valent Cu(III) complex **IV**, which undergoes reductive elimination to afford the coupled product **6** or **7**, with regeneration of **L1Cu(I)CN** species to complete the copper catalysis cycle. It should be noted that an alternative process involving direct cyano transfer from the **L1Cu(II)(CN)₂** complex to the benzylic radical **III** through an outer-sphere pathway is also possible. Notably, the whole process is redox-neutral and does not need any external stoichiometric oxidants or reductants.

Discussion

In summary, we have developed an intermolecular, highly enantioselective three-component radical vicinal DCF reaction of alkenes for the first time using oxime esters and TMSCN by dual photoredox and copper catalysis. Key to the success of this protocol relies on chemoselective addition of acyl and cyanoalkyl radicals, generated in situ from the redox-active oxime esters by a photocatalytic N-centered iminyl radical-triggered C-C bond cleavage event, onto the alkenes to form new carbon radicals. Single electron metalation of such carbon radicals to TMSCN-derived **L1Cu(II)(CN)₂** complex leads to asymmetric cross-coupling. This three-component reaction proceeds under mild conditions, and demonstrates broad substrate scope and high functional group tolerance, providing a general approach to optically active β -cyano ketones and alkylidinitriles. Many exciting extensions of this strategy to other radical precursors and nucleophiles can be envisaged; current investigations into these subjects are ongoing in our laboratory.

Methods

General procedure for the synthesis of products 6. In a flame-dried 10 mL Schlenk tube equipped with a magnetic stirrer bar was charged sequentially with $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.56 mg, 0.0015 mmol) and chiral ligand **L1** (0.80 mg, 0.00225 mmol), followed by the addition of DMA (2.5 mL). Then the mixture was stirred at room temperature for 30 min. To the resulting mixture were added **3** (0.30 mmol), **1** (0.10 mmol), *fac*- $\text{Ir}(\text{ppy})_3$ (0.53 mg, 0.0008 mmol). Then, the resulting mixture was degassed (3 times) under argon atmosphere. After that, TMSCN (0.3 mmol) was added into the mixture. At last, the mixture was stirred at a distance of ~ 1 cm from a 2×3 W purple LEDs at room temperature for 24 h until the reaction was completed, as monitored by TLC analysis. The reaction mixture was quenched with water (10 mL), diluted with EtOAc (3×10 mL), washed with NaCl (aq.) and dried over with anhydrous Na_2SO_4 . After filtration and concentration, the residue was purified by silica gel chromatography afford the final products.

General procedure for the synthesis of products 7. In a flame-dried 10 mL Schlenk tube equipped with a magnetic stirrer bar was charged sequentially with $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.001 mmol), chiral ligand **L1** (0.0012 mmol) and organo-photocatalyst Ph-PTZ (0.0025 mmol), followed by the addition of DMA (4 mL). Then the mixture was stirred at room temperature for 30 min. To the resulting mixture were added **2** (0.20 mmol) and **4** (0.60 mmol). Then, the resulting mixture was degassed (3 times) under argon atmosphere. After that, TMSCN (0.60 mmol) was added into the mixture. At last, the mixture was stirred at a distance of ~ 1 cm from a 2×3 W purple LEDs at room temperature 6 h until the reaction was completed, as monitored by TLC analysis. The reaction mixture was diluted with water (10 mL). The mixture was firstly extracted with EtOAc (3×10 mL), then washed with NaHCO_3 (aq.) (15 mL), and finally washed with NaCl (aq.), dried over with anhydrous Na_2SO_4 . After filtration and concentration, the residue was purified by silica gel chromatography afford the final products. Full experimental details and characterization of new compounds can be found in the Supplementary Methods.

Declarations

Data availability

The authors declare that the main data supporting the findings of this study, including experimental procedures and compound characterization, are available within the article and its Supplementary Information files. X-ray structural data of compounds **6bj** and **7ja** are available free of charge from the Cambridge Crystallographic Data Center under the deposition number CCDC 2047031 (**6bj**) and 2047032 (**7ja**). These data can be obtained free of charge from The Cambridge Crystallographic Data Center via www.ccdc.cam.ac.uk/data_request/cif.

Author contributions

P.-Z.W., Y. G., J. C. and X.-D.H. are responsible for the plan and implementation of the experimental work. J.-R.C. and W.-J.X. supervised the project and wrote the manuscript. All authors discussed the results and commented on the manuscript.

Competing interests

The authors declare no competing interests.

Acknowledgements

We are grateful to the financial support from the National Natural Science Foundation of China (No. 21971081, 91856119, 21772053, 21820102003, and 91956201), and the Program of Introducing Talents of Discipline to Universities of China (111 Program, B17019).

References

1. Kolb, H. C., VanNieuwenhze, M. S. & Sharpless, K. B. Catalytic asymmetric dihydroxylation. *Chem. Rev.* **94**, 2483–2547 (1994).
2. McDonald, R. I., Liu, G. & Stahl, S. S. Palladium(II)-catalyzed alkene functionalization via nucleopalladation: stereochemical pathways and enantioselective catalytic applications. *Chem. Rev.* **111**, 2981–3019 (2011).
3. Cresswell, A. J., Eey, S. T. & Denmark, S. E. Catalytic, stereoselective dihalogenation of alkenes: challenges and opportunities. *Angew. Chem. Int. Ed.* **54**, 15642–15682 (2015).
4. Luo, Y.-C., Xu, C. & Zhang, X. Nickel-catalyzed dicarbofunctionalization of alkenes. *Chin. J. Chem.* **38**, 1371–1394 (2020).
5. Tang, S., Liu, K., Liu, C. & Lei, A. Olefinic C-H functionalization through radical alkenylation. *Chem. Soc. Rev.* **44**, 1070–1082 (2015).
6. Liu, Q., Dong, X., Li, J., Xiao, J., Dong, Y. & Liu, H. Recent advances on palladium radical involved reactions. *ACS Catal.* **5**, 6111–6137 (2015).
7. Crossley, S. W. M., Obradors, C., Martinez, R. M. & Shenvi, R. A. Mn-, Fe-, and Co-catalyzed radical hydrofunctionalizations of olefins. *Chem. Rev.* **116**, 8912–9000 (2016).
8. Liu, L., Ward, R. M. & Schomaker, J. M. Mechanistic aspects and synthetic applications of radical additions to allenes. *Chem. Rev.* **119**, 12422–12490 (2019).
9. Bao, X., Li, J., Jiang, W. & Huo, C. Radical-mediated difunctionalization of styrenes. *Synthesis* **51**, 4507–4530 (2019).
10. Jiang, H. & Studer, A. Intermolecular radical carboamination of alkenes. *Chem. Soc. Rev.* **49**, 1790–1811 (2020).
11. Qin, T., Cornella, J., Li, C., Malins, L. R., Edwards, J. T., Kawamura, S., Maxwell, B. D., Eastgate, M. D. & Baran, P. S. A general alkyl-alkyl cross-coupling enabled by redox-active esters and alkylzinc reagents. *Science* **352**, 801–805 (2016).
12. Kischkewitz, M., Okamoto, K., Muck-Lichtenfeld, C. & Studer, A. Radical-polar crossover reactions of vinylboron ate complexes. *Science* **355**, 936–938 (2017).

13. Garcia-Dominguez A., Li, Z. & Nevado, C. Nickel-catalyzed reductive dicarbofunctionalization of alkenes. *J. Am. Chem. Soc.* **139**, 6835–6838 (2017).
14. Liu, L., Lee W., Youshaw, C. R., Yuan, M., Geherty, M. B., Zavalij, P. Y. & Gutierrez, O. Fe-catalyzed three-component dicarbofunctionalization of unactivated alkenes with alkyl halides and Grignard reagents. *Chem. Sci.* **11**, 8301–8305 (2020).
15. Campbell, M. W., Compton, J. S., Kelly, C. B. & Molander, G. A. Three-component olefin dicarbofunctionalization enabled by nickel/photoredox dual catalysis. *J. Am. Chem. Soc.* **141**, 20069–20078 (2019).
16. Encyclopedia of Radicals in Chemistry, Biology and Materials. Chatgililoglu, C., Studer, A., Eds.; Chechester, (2012).
17. Sibi, M. P., Manyem, S. & Zimmerman, J. Enantioselective radical processes. *Chem. Rev.* **103**, 3263–3296 (2003).
18. Lipp, A., Badir, S. O. & Molander, G. Stereoinduction in metallaphotoredox catalysis. *Angew. Chem. Int. Ed.* 10.1002/anie.202007668 (2020).
19. Wang, F., Chen, P. & Liu, G. Copper-catalyzed radical relay for asymmetric radical transformations. *Acc. Chem. Res.* **51**, 2036–2046 (2018).
20. Li, Z.-L., Fang, G.-C., Gu, Q.-S. & Liu, X.-Y. Recent advances in copper-catalyzed radical-involved asymmetric 1,2-difunctionalization of alkenes. *Chem. Soc. Rev.* **49**, 32–48 (2020).
21. Gu Q.-S., Li, Z.-L. & Liu, X.-Y. Copper(I)-catalyzed asymmetric reactions involving radicals. *Acc. Chem. Res.* **53**, 170–181 (2020).
22. Wang, F., Wang, D., Wan, X., Wu, L., Chen, P. & Liu, G. Enantioselective copper-catalyzed intermolecular cyanotrifluoromethylation of alkenes via radical process. *J. Am. Chem. Soc.* **138**, 15547–15550 (2016).
23. Zhou, S., Zhang, G., Fu, L., Chen, P., Li, Y. & Liu, G. Copper-catalyzed asymmetric cyanation of alkenes via carbonyl-assisted coupling of alkyl-substituted carbon-centered radicals. *Org. Lett.* **22**, 6299–6303 (2020).
24. Zhang, G., Zhou, S., Fu, L., Chen, P., Li, Y., Zou, J. & Liu, G. Asymmetric coupling of carbon-centered radicals adjacent to nitrogen: copper-catalyzed cyanation and etherification of enamides. *Angew. Chem. Int. Ed.* **59**, 20439–20444 (2020).
25. Wu, L., Wang, F., Wan, X., Wang, D., Chen, P. & Liu, G. Asymmetric Cu-catalyzed intermolecular trifluoromethylarylation of styrenes: enantioselective arylation of benzylic radicals. *J. Am. Chem. Soc.* **139**, 2904–2907 (2017).
26. Wu, L., Wang, F., Chen, P. & Liu, G. Enantioselective construction of quaternary all-carbon centers via copper-catalyzed arylation of tertiary carbon-centered radicals. *J. Am. Chem. Soc.* **141**, 1887–1892 (2019).
27. Fu, L., Zhou, S., Wan, X., Chen, P. & Liu, G. Enantioselective trifluoromethylalkynylation of alkenes via copper-catalyzed radical relay. *J. Am. Chem. Soc.* **140**, 10965–10969 (2018).

28. Zhuang, W., Chen, P. & Liu, G. Enantioselective arylcyanation of styrenes via copper-catalyzed radical relay. *Chin. J. Chem.* 10.1002/cjoc.202000494 (2020).
29. Lei, G., Zhang, H., Chen, B., Xu, M. & Zhang, G., Copper-catalyzed enantioselective arylalkynylation of alkenes. *Chem. Sci.* **11**, 1623–1628 (2020).
30. Lin, J.-S., Li, T.-T., Liu, J.-R., Jiao, G.-Y., Gu, Q. S., Cheng, J.-T., Guo, Y.-L., Hong, X. & Liu, X.-Y. Cu/chiral phosphoric acid-catalyzed asymmetric three-component radical-initiated 1,2-dicarbofunctionalization of alkenes. *J. Am. Chem. Soc.* **141**, 1074–1083 (2019).
31. Dong, X.-Y., Cheng, J.-T., Zhang, Y.-F., Li, Z.-L., Zhan, T.-Y., Chen, J.-J., Wang, F.-L., Yang, N.-Y., Ye, L., Gu, Q.-S. & Liu, X.-Y. Copper-catalyzed asymmetric radical 1,2-carboalkynylation of alkenes with alkyl halides and terminal alkynes. *J. Am. Chem. Soc.* **142**, 9501–9509 (2020).
32. Dong, X.-Y., Zhan, T.-Y., Jiang, S.-P., Liu, X.-D., Ye, L., Li, Z.-L., Gu, Q.-S. & Liu, X.-Y. Copper-catalyzed asymmetric coupling of allenyl radicals with terminal alkynes to access tetrasubstituted allenes. *Angew. Chem. Int. Ed.* 10.1002/anie.202013022 (2020).
33. Photochemically Generated Intermediates in Synthesis (Eds.: A. Albini, M. Fagnoni), Wiley, Hoboken, (2013).
34. Prier, C. K., Rankic, D. A. & MacMillan, D. W. C. Visible light photoredox catalysis with transition metal complexes: applications in organic synthesis. *Chem. Rev.* **113**, 5322–5363 (2013).
35. Marzo, L., Pagire, S. K., Reiser, O. & König, B. Visible-light photocatalysis: does it make a difference in organic synthesis? *Angew. Chem. Int. Ed.* **57**, 10034–10072 (2018).
36. Crisenza, G. E. M., Mazzarella, D. & Melchiorre, P. Synthetic methods driven by the photoactivity of electron donor-acceptor complexes. *J. Am. Chem. Soc.* **142**, 5461–5476 (2020).
37. Garbarino, S., Ravelli, D., Protti, S. & Basso, A. Photoinduced multicomponent reactions. *Angew. Chem. Int. Ed.* **55**, 15476–15484 (2016).
38. Courant, T. & Masson, G. Recent progress in visible-light photoredox-catalyzed intermolecular 1,2-difunctionalization of double bonds via an ATRA-type mechanism. *J. Org. Chem.* **81**, 6945–6952 (2016).
39. Pantaine, L., Bour, C. & Masson, G. *Photochemistry*, **46**, 395–431 (2019).
40. Silvi, M. & Melchiorre, P. Enhancing the potential of enantioselective organocatalysis with light. *Nature* **554**, 41–49 (2018).
41. Zhang, H.-H., Chen, H., Zhu, C. & Yu, S. A review of enantioselective dual transition metal/photoredox catalysis. *Sci. China Chem.* **63**, 637–647 (2020).
42. Rigotti, T. & Alemán, J. Visible light photocatalysis – from racemic to asymmetric activation strategies. *Chem. Commun.* **56**, 11169–11190 (2020).
43. Zheng, D. & Studer, A. Asymmetric synthesis of heterocyclic gamma-amino-acid and diamine derivatives by three-component radical cascade reactions. *Angew. Chem. Int. Ed.* **58**, 15803–15807 (2019).

44. Yu, X. Y., Zhao, Q. Q., Chen, J., Xiao, W. J. & Chen, J. R. When light meets nitrogen-centered radicals: from reagents to catalysts. *Acc. Chem. Res.* **53**, 1066–1083 (2020).
45. Yu, X.-Y., Chen, J.-R., Wang, P.-Z., Yang, M.-N., Liang, D. & Xiao, W.-J. A visible-light-driven iminyl radical-mediated C-C single bond cleavage/radical addition cascade of oxime esters. *Angew. Chem. Int. Ed.* **57**, 738–743 (2018).
46. Yu, X. Y., Zhao, Q. Q., Chen, J., Chen, J. R. & Xiao, W. J. Copper-catalyzed radical cross-coupling of redox-active oxime esters, styrenes, and boronic acids. *Angew. Chem. Int. Ed.* **57**, 15505–15509 (2018).
47. Chen, J., Wang, P.-Z., Lu, B., Liang, D., Yu, X.-Y., Xiao, W.-J. & Chen, J.-R. Enantioselective radical ring-opening cyanation of oxime esters by dual photoredox and copper catalysis. *Org. Lett.* **21**, 9763–9768 (2019).
48. Xiao, T., Huang, H., Anand, D. & Zhou, L. Iminyl-radical-triggered C–C bond cleavage of cycloketone oxime derivatives: generation of distal cyano-substituted alkyl radicals and their functionalization. *Synthesis* **52**, 1585–1601 (2020).
49. Song, C., Shen, X., Yu, F., He, Y.-P. & Yu, S. Generation and application of iminyl radicals from oxime derivatives enabled by visible light photoredox catalysis. *Chin. J. Org. Chem.* 10.6023/cjoc202004008 (2020).
50. Zhao, B. & Shi, Z. Copper-catalyzed intermolecular Heck-like coupling of cyclobutanone oximes initiated by selective C – C bond cleavage. *Angew. Chem. Int. Ed.* **56**, 12727–12731 (2017).
51. Dauncey, E. M., Morcillo, S. P., Douglas, J. J., Sheikh, N. S. & Leonori, D. Photoinduced remote functionalisations by iminyl radical promoted C-C and C-H bond cleavage cascades. *Angew. Chem. Int. Ed.* **57**, 744–748 (2018).
52. Fan, X., Lei, T., Chen, B., Tung, C. H. & Wu, L.-Z. Photocatalytic C-C bond activation of oxime ester for acyl radical generation and application. *Org. Lett.* **21**, 4153–4158 (2019).
53. Cheng, Y.-Y., Lei, T., Su, L., Fan, X., Chen, B., Tung, C.-H. & Wu, L.-Z. Visible light irradiation of acyl oxime esters and styrenes efficiently constructs β -carbonyl imides by a scission and four-component reassembly process. *Org. Lett.* **21**, 8789–8794 (2019).
54. Wang, T., Wang, Y.-N., Wang, R., Zhang, B.-C., Yang, C., Li, Y.-L. & Wang, X.-S. Enantioselective cyanation via radical-mediated C-C single bond cleavage for synthesis of chiral dinitriles. *Nat. Commun.* **10**, 5373 (2019).
55. Ruan, L., Chen, C., Zhang, X. & Sun, J. Recent advances on the photo-induced reactions of acyl radical. *Chin. J. Org. Chem.* **38**, 3155–3164 (2018).
56. Banerjee, A., Lei, Z. & Ngai, M. Y. Acyl radical chemistry via visible-light photoredox catalysis. *Synthesis* **51**, 303–333 (2019).
57. Raviola, C., Protti, S., Ravelli, D. & Fagnoni, M. Photogenerated acyl/alkoxycarbonyl/carbamoyl radicals for sustainable synthesis. *Green Chem.* **21**, 748–764 (2019).

58. Chu, X.-Q., Ge, D., Shen, Z.-L. & Loh, T.-P. Recent advances in radical-initiated C(sp³)-H bond oxidative functionalization of alkyl nitriles. *ACS Catal.* **8**, 258–271 (2017).
59. Wu, W.-B., Yu, J.-S. & Zhou, J. Catalytic enantioselective cyanation: recent advances and perspectives. *ACS Catal.* **10**, 7668–7690 (2020).
60. Wang, Y., Du, Y. & Huang, N. A survey of the role of nitrile groups in protein-ligand interactions. *Future Med. Chem.* **10**, 2713–2728 (2018).
61. Lopez, R. & Palomo, C. Cyanoalkylation: alkyl nitriles in catalytic C-C bond-forming reactions. *Angew. Chem. Int. Ed.* **54**, 13170–13184 (2015).
62. Hayrapetyan, D. & Khalimon, A. Y. Catalytic nitrile hydroboration: a route to N,N-diborylamines and uses thereof. *Chem. Asian. J.* **15**, 2575–2587 (2020).
63. See supporting information for more details.
64. CCDC 2047031 (**6bj**) and 2047032 (**7ja**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via .
65. Yang, S., Wang, L., Zhang, H., Liu, C., Zhang, L., Wang, X., Zhang, G., Li, Y. & Zhang, Q. Copper-catalyzed asymmetric aminocyanation of arylcyclopropanes for synthesis of γ -amino nitriles. *ACS Catal.* **9**, 716–721 (2019).
66. Long, J., Yu, R., Gao, J. & Fang, X. Access to 1,3-dinitriles by enantioselective auto-tandem catalysis: merging allylic cyanation with asymmetric hydrocyanation. *Angew. Chem. Int. Ed.* **59**, 6785–6789 (2020).

Tables

Table 1 | Optimization of the reaction conditions.

Entry	x / y / z [mol%]	Solvent	Yield [%] ^a	ee [%] ^b
1	0.5 / 0.6 / 1.0	DMA (0.05 M)	41	88
2	0.5 / 0.6 / 1.0	DMF (0.05 M)	39	89
3	0.5 / 0.6 / 1.0	CH ₃ CN (0.05 M)	6	90
4	0.5 / 0.6 / 1.0	THF (0.05 M)	15	90
5	0.5 / 0.6 / 1.0	CH ₂ Cl ₂ (0.05 M)	9	90
6	0.5 / 0.6 / 0.8	DMA (0.05 M)	64	88
7 ^c	0.5 / 0.6 / 0.8	DMA (0.05 M)	51	85
8 ^d	0.5 / 0.6 / 0.8	DMA (0.05 M)	69	81
9 ^e	0.5 / 0.6 / 0.8	DMA (0.05 M)	16	68
10	1.5 / 2.25 / 0.8	DMA (0.05 M)	78	90
11	1.5 / 2.25 / 0.8	DMA (0.04 M)	88 (74)	90

Reaction conditions: **1a** (0.1 mmol), **3a** (0.3 mmol), TMSCN (0.3 mmol), Cu(CH₃CN)₄PF₆ (x mol%), ligand **L1** (y mol%), *fac*-Ir(ppy)₃ (z mol%), solvent (2.0-2.5 mL), 2x3 W purple LEDs, at room temperature.

^aYields were determined by GC analysis, with isolated yield in parentheses.

^bDetermined by HPLC analysis on a chiral stationary phase.

^cWith CuCl.

^dWith CuI.

^eWith Cu(OTf)₂.

ppy = 2-phenylpyridine. DMA = N,N-dimethylacetamide. DMF = N,N-dimethylformamide.

Figures

A Cu-catalyzed, three-component radical alkene vicinal dicarbofunctionalization

-Previous work

B First asymmetric, three-component alkene DCF with oxime esters and TMSCN

-This work

Figure 1

Catalytic asymmetric three-component radical dicarbofunctionalization reactions of alkenes. a) Cu-catalyzed, three-component radical alkene vicinal dicarbofunctionalization. b) First asymmetric, three-component alkene DCF with oxime esters and TMSCN.

A Cu-catalyzed, three-component radical alkene vicinal dicarbofunctionalization

-Previous work

B First asymmetric, three-component alkene DCF with oxime esters and TMSCN

-This work

Figure 1

Catalytic asymmetric three-component radical dicarbofunctionalization reactions of alkenes. a) Cu-catalyzed, three-component radical alkene vicinal dicarbofunctionalization. b) First asymmetric, three-component alkene DCF with oxime esters and TMSCN.

(A) scope of alkenes

(B) biologically relevant alkenes

Figure 2

Scope of the alkenes in asymmetric three-component alkene acylcyanation. Reaction conditions: **1** (0.1 mmol), **3a** (0.3 mmol), TMSCN (0.3 mmol), Cu(CH₃CN)₄PF₆ (1.5 mol%), ligand L1 (2.25 mol%), fac-Ir(ppy)₃ (0.8 mol%), DMA (2.5 mL), 2x3 W purple LEDs, at room temperature. Isolated yields were reported. The ee and d.r. values were determined by HPLC analysis on a chiral stationary phase. ^a1.0 mmol scale reaction, 24 h.

(A) scope of alkenes

(B) biologically relevant alkenes

Figure 2

Scope of the alkenes in asymmetric three-component alkene acylcyanation. Reaction conditions: **1** (0.1 mmol), **3a** (0.3 mmol), TMSCN (0.3 mmol), Cu(CH₃CN)₄PF₆ (1.5 mol%), ligand L1 (2.25 mol%), fac-Ir(ppy)₃ (0.8 mol%), DMA (2.5 mL), 2x3 W purple LEDs, at room temperature. Isolated yields were reported. The ee and d.r. values were determined by HPLC analysis on a chiral stationary phase. ^a1.0 mmol scale reaction, 24 h.

(A) with alkene **1b**

(B) with alkene **1a** (R = 2-naphthyl)

Figure 3

Scope of the oxime esters in asymmetric three-component alkene acylcyanation. Reaction conditions: **1** (0.1 mmol), **3** (0.3 mmol), TMSCN (0.3 mmol), Cu(CH₃CN)₄PF₆ (1.5 mol%), ligand L1 (2.25 mol%), fac-Ir(ppy)₃ (0.8 mol%), DMA (2.5 mL), 2x3 W purple LEDs, at room temperature. Isolated yields were reported. The ee and d.r. values were determined by HPLC analysis on a chiral stationary phase.

(A) with alkene **1b**

6bb, R = Me, 73%, 83% ee
6bc, R = OMe, 75%, 86% ee
6bd, R = *t*-Bu, 81%, 89% ee
6be, R = F, 77%, 91% ee
6bf, R = Cl, 86%, 91% ee
6bg, R = Br, 70%, 86% ee

(B) with alkene **1a** (R = 2-naphthyl)

Figure 3

Scope of the oxime esters in asymmetric three-component alkene acylcyanation. Reaction conditions: **1** (0.1 mmol), **3** (0.3 mmol), TMSCN (0.3 mmol), Cu(CH₃CN)₄PF₆ (1.5 mol%), ligand **L1** (2.25 mol%), fac-Ir(ppy)₃ (0.8 mol%), DMA (2.5 mL), 2x3 W purple LEDs, at room temperature. Isolated yields were reported. The ee and d.r. values were determined by HPLC analysis on a chiral stationary phase.

Figure 4

Scope of the alkenes and cycloketone oxime esters in asymmetric three-component alkene cyanoalkylcyanation reaction. Reaction conditions: **2** (0.2 mmol), **4** (0.6 mmol), TMS-CN (0.6 mmol), $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.5 mol%), ligand **L1** (0.6 mol%), Ph-PTZ (1.25 mol%), DMA (4.0 mL), 2x3 W purple LEDs, at room temperature. Isolated yields were reported. The ee and d.r. values were determined by HPLC analysis on a chiral stationary phase.

Figure 4

Scope of the alkenes and cycloketone oxime esters in asymmetric three-component alkene cyanoalkylcyanation reaction. Reaction conditions: **2** (0.2 mmol), **4** (0.6 mmol), TMS-CN (0.6 mmol), $\text{Cu}(\text{CH}_3\text{CN})_4\text{PF}_6$ (0.5 mol%), ligand **L1** (0.6 mol%), Ph-PTZ (1.25 mol%), DMA (4.0 mL), 2x3 W purple LEDs, at room temperature. Isolated yields were reported. The ee and d.r. values were determined by HPLC analysis on a chiral stationary phase.

Figure 5

Synthetic applications.

Figure 5

Synthetic applications.

a) radical trapping experiment

b) radical clock experiment

Figure 6

Mechanistic studies.

a) radical trapping experiment

b) radical clock experiment

Figure 6

Mechanistic studies.

Figure 7

Proposed reaction mechanism.

Figure 7

Proposed reaction mechanism.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupportingInformationFinal20201209.pdf](#)
- [SupportingInformationFinal20201209.pdf](#)
- [6bjcheckcifReport.pdf](#)
- [6bjcheckcifReport.pdf](#)
- [7jcheckcifReprot.pdf](#)
- [7jcheckcifReprot.pdf](#)
- [cifof6bj.cif](#)

- [cifof6bj.cif](#)
- [cifof7ja.cif](#)
- [cifof7ja.cif](#)