An Assessment of Current Concussion Identification and Diagnosis Methods in Sports Settings. A Systematic Review

Ed Daly

Galway-Mayo-Institute of Technology

Alan J Pearce

La Trobe University

Emma Finnegan

Galway-Mayo-Institute of Technology

Ciara Cooney

Galway-Mayo-Institute of Technology

Maria McDonagh

Galway-Mayo-Institute of Technology

Grainne Scully

Galway-Mayo-Institute of Technology

Michael McCann

Letterkenny Institute of Technology

Ronan Doherty

Letterkenny Institute of Technology

Adam White

Oxford Brookes University

Simon Phelan

Oxford Brookes University

Nathan Howarth

Oxford Brookes University

Lisa Ryan (✉ lisa.ryan@gmit.ie)

Galway Mayo Institute of Technology <https://orcid.org/0000-0002-5505-7130>

Research Article

Keywords: Concussion, Head Injury Assessment, Screening tools, Assessment Protocols, Sport Injury Risk, SCAT

Posted Date: February 21st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1256804/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Concussion in sport is an ongoing global concern. The head injury assessment (HIA) by the field of play is acknowledged as the first step in recognising and identifying concussion. While previous systematic literature reviews have evaluated the sensitivity of sideline screening tools and assessment protocols, no systematic review has evaluated the research designs and assessments used in a field setting.

This systematic review investigated existing screening and diagnostic tools used in research as part of the HIA protocol to identify concussion that are currently used in professional, semi-professional and amateur (club) sports settings.

Methods

A systematic searching of relevant databases was undertaken for peer-reviewed literature between 2015 and 2020.

Results

Twenty-five studies met the inclusion criteria. Studies were of moderate to good quality, reporting a variety of designs. The majority of studies were undertaken in professional/elite environments with medical doctors and allied health practitioners (e.g., physical therapists) involved in 88% of concussion assessments. While gender was reported in 24 of the 25 studies, the majority of participants were male (77%). There was also a variety of concussion assessments (n=20) with the sports concussion assessment tool (SCAT) used in less than half of the included studies.

Conclusion

The majority of studies investigating concussion HIAs are focused on professional/elite sport. With concussion an issue at all levels of sport, future research should be directed at non-elite sport. Further, for research purposes, the SCAT assessment should also be used more widely to allow for consistency across studies.

Key Points

- Head injury assessments are vital as part of the diagnostic pathway for suspected concussion at all levels of sport.
- This systematic review found that in studies between 2015 and 2020, the majority were focused on elite sports and male participants.
- Studies also utilized a disparate array of assessments making comparison difficult.
- Future studies should aim to focus on amateur (club) sports, include more female participants, and as a minimum include the sports concussion assessment tool (SCAT), along with other assessments to allow for consistency across studies.

Background

Despite consensus on the clinical definition of concussion in sport, its immediate and accurate recognition in a clinical setting / pitch side setting remains a challenge to implement beyond the elite level of sport. Currently, only a licensed medical practitioner is entitled to diagnose a concussion [1]. However, even at the elite level, clinical decision-making is guided on subjective athlete self-report with observation of symptoms and severity [1]. Further, determination of when an athlete is appropriately ready to return to sport is limited also by subjective symptom scores and imperfect clinical and neuropsychological testing [2]. Despite these challenges, the consensus remains a medical decision based upon resolution of symptoms, as reported by the athlete, as well as completing the return to sport guideline protocols [1].

It is important to accurately diagnose a concussion and have the athlete complete a full recovery. It is well described that sustaining a concussion increases the likelihood of incurring a subsequent injury [3, 4]. While epidemiological studies only describe the risk, observational studies have suggested that increased risk occurs as complete recovery may not have been possible affecting neuromuscular function [5–7]. It has been suggested that experiencing numerous concussions, or exposure to repetitive head trauma, could be associated with long-term consequences such as persistent post concussive symptoms [8] or increasing risk of neurodegenerative disorders [9, 10].

Given these risks, it is important that athletes suspected of concussion following an impact are assessed and identified via the removal of participants for further evaluation and, if diagnosed with concussion, consequently engaged in a graduated return to play protocol [1]. At the elite levels of competitions, the use of head injury assessment (HIA) tools, such as the Sports Concussion Assessment Tool (SCAT) have been evolving since the first consensus statement published in 2001 [11]. While these assessment tools do not replace the medical decision, their use certainly assists the medical practitioner with clinical decision making as well as guidance for return to play clinical decisions for athletes' post-concussion. While previous systematic literature reviews have evaluated the sensitivity of sideline screening tools and assessment protocols [12–14]. To date no systematic review has evaluated the overall efficacy of these tools and protocols in a field setting. Moreover, it is unknown if HIAs are utilized in non-elite environments, such as amateur club competitions. This systematic review and qualitative analysis aimed to investigate the efficacy of current off-field or 'sideline' recognition of suspected concussions. The primary objective was to investigate existing screening and diagnostic tools that are used in identifying concussion, or head injury assessment protocols that are currently used in professional, semi-professional and amateur (club) sports settings.

Methods

Study design

The review protocol was prospectively registered in the PROSPERO database for systematic reviews (protocol ID: CRD42021214339) and complies with Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines [15]. The review question and inclusion/exclusion criteria and search terms are presented in Table 1.

Identification of evidence

An electronic search was conducted on 01 July 2020. The following electronic databases were queried: PsychInfo (OVID), PubMed, Science Direct, SPORTDiscus (EBSCOhost), Web of Science, and the Cumulative Index to Nursing and Allied Health Literature (CINAHL) (EBSCOhost). The search terms included: (sport related concussion OR SRC OR mTBI OR mild traumatic brain injury) AND (diagnosis OR treatment) AND (sport). Articles met inclusion (eligibility criteria) based on the following priori inclusion criteria: (i) participants were involved in professional/elite, semi-professional/sub-elite or amateur (club) sport at the time of injury; (ii) individuals aged 18 years or greater; (iii) concussion diagnosis assessment was administered acutely (here, we define acute as ≤ 7 days of injury); (iv) peer-reviewed journal articles published since 2015. Articles prior to 2015 would not be reflective of concussion recognition strategies as outlined in the most recent Concussion Consensus Statement (2016). However, part of our *a-priori* inclusion criteria was modified. For the purpose of this review we allowed for the inclusion of athletes younger than age 18 years (e.g., studies with an age range of 5 – 23 years). These studies were considered important as a majority of athletes in these included studies were 18 years or older. Studies in which all participants were under 18 years of age were not included. Exclusionary criteria applied included: (i) nonathlete (e.g., participants involved in military duties at the time of injury or a motor vehicle accident-related concussion, healthy adults) and mixed (athlete plus non-athlete) populations were excluded to ensure a focus on sport-related concussion; (ii) the investigation did not perform concussion testing pitch side defined in Table 1 (as any testing away from the immediate sporting environment, for example, rink side, trackside, locker room, medical room, touch line and so on for the purpose of this research); (iii) the type of publication was a conference abstract (without full data set), narrative review, editorial or letter to editor, student theses (e.g., honours/masters/PhD) or book chapter.

Selection of evidence and data extraction

Following compilation of online search results, record titles and abstracts were screened by four authors (CC, ED, MMD, AJP), full text articles were reviewed by five authors (CC, ED, MMD, AJP, LR). The reference lists from review articles were assessed for pertinent studies that may have been overlooked. Data extraction was performed independently by two authors (GS and ED) and data reviewed by a third author (CC) for consistency and accuracy. In cases of disagreement at any stage, consultations with other authors (CC, ED, MMD, AJP, LR) were planned and disagreement resolved by joint discussion and consensus. Figure 1 illustrates the literature review process using PRISMA flow chart [15].

Data synthesis, statistical analyses and assessment of overall quality of evidence

References were managed in EndNote (Clarivate Analytics, Berkeley California, USA), extracted data were collated in Excel 2013. The overall quality of evidence for each outcome was assessed using the Downs and Black [16] checklist for measuring study quality by three authors (AP, ED, LR). The Downs and Black [16] checklist was developed to assess both randomised and non-randomised research designs. The checklist contains 27 'yes'-or-'no' questions across five sections including (i) study quality (10 items); (ii) external validity (3 items); (iii) study bias (6 items); (iv) confounding and selection bias (6 items); (v) power of the study (1 item). Higher total scores for this

checklist reflect increased study quality and confidence in conclusions. However, as some questions were not pertinent to this review, a modified form of the checklist was used for a maximum score of 22. Due to the heterogeneity of the studies a meta-analysis could not be performed, and a qualitative analysis of studies was instead conducted.

Table 1
PICO-model and Medline search strategy in accordance with PRISMA statement

Primary review question/aim	
What are the current side-line screening methods used to establish the diagnosis of acute concussion or suspected concussion across sports in an adult population?	
Inclusion criteria	
Population	Athletes aged 18 years or greater, involved in amateur, semi-professional or professional sport and sustaining a suspected concussive injury. For the purpose of this review, we allowed for the inclusion of athletes younger than 18 years (e.g., studies with an age range of 5 – 23 years). These studies were considered important as the cohort included were over 18 years of age.
Intervention	Any side-line* screening assessment used to detect suspected concussion following sports-related head impact event in the acute phase of injury. For this review the acute phase of injury will be defined as minutes after the event up to and including 7 days post event. These will include (but may not be limited to) reported (i). Concussion 2, (ii) mTBI, (iii) Cervical neck injury.
Outcomes	Acute concussion diagnosis methods.
Study design	Published research, retrospective data analysis, cross sectional study design, parallel studies, prospective, observational, systematic reviews where data meeting the PICO can be extracted. Abstracts (with data) will be included initially. Research published from 2015 onwards.
Exclusion criteria	
Population	Not related to sport, animal studies or studies in which all participants were under 18 years of age.
Intervention	Non-side-line testing, testing conducted >7 days post event.
Outcomes	Concussion/suspected concussion not examined side-line or diagnosis assessment requiring advanced medical training/technology or referral to secondary care for diagnosis to be made.
Study design	Case reports, editorials, commentary's, review articles (in the case of systematic reviews if relevant data cannot be extracted or does not meet PICO), consensus statements, position stands and non-English publications. Research published in 2014 and prior.
Search terms	
† ((sport related concussion OR SRC OR mTBI OR mild traumatic brain injury) AND (diagnosis OR treatment)) AND (sport)	
*Side-line' is used generally to denote testing away from the immediate sporting environment, for example rink side, track side, locker room, medical room, touch line and so on. †MeSH terms were exploded to include more specific terms; MeSH terms were translated into the appropriate subject headings for other databases. Keywords were the same for each database searched.	

Results

Study selection

A total of 7679 citations were screened for eligibility, with the full text of 258 articles retrieved for detailed evaluation. Twenty-five studies met the inclusion criteria following full text review. Figure 1 presents the PRISMA flow diagram [15] of identification, screening, eligibility and inclusion criteria for the literature review of side-line diagnosis of concussion. Table 2 presents the studies meeting the inclusion criteria.

Characteristics of included studies

Twenty-five studies (Table 2) met the review inclusion criteria and reported interpretable data on side-line assessments for the diagnosis of concussion. Characteristics of the included studies are summarised in Table 2. Studies consisted of prospective (n = 12), pilot (n = 2), cross sectional (n = 2), prospective observational (n = 1), prospective longitudinal (n = 1), cohort (n = 1), observational (n = 1), descriptive (n = 1), prospective cross sectional (n = 1), pilot case study (n = 1), case control (n = 1) study and retrospective (n=1). Gender was reported in 24 of the 25 studies with a total population of 7126 participants (5484 males and 1642 females). Nineteen studies examined baseline to post concussion testing. The remaining six studies examined post-concussion exclusively. All studies assessed concussion

within the acute diagnosis phase (≤ 7 days). A definition of acute concussion diagnosis was defined in the included studies within the first 3 to 5 days post-concussion [17], ≤ 48 h after concussion for SCAT3 [18], 24–48 hours for BESS, SAC testing, subacute measures of neurocognitive impairment (i.e., ImPACT at 5–7 days and 10–14 days postinjury) [19], 24 h [20] and 72 h post-injury [21]. Included studies were conducted between 2015 to 2020 in a range of countries, including the USA, United Kingdom, Finland, Australia, Canada, South Africa and New Zealand.

Test Categories

Included studies (Table 2) used a battery of tests in the diagnosis of side-line concussion (i.e. 'side-line' refers to pitch-side, rink side, changing rooms, or an assessment area immediately available that is not a clinical or hospital setting), that fell into three main testing categories (i) cognitive, (ii) observational and (iii) visual. Fifty-six percent of studies employed cognitive tests, 8% observational, a further 8% visual and 28% used a combination of the three test categories. Eighteen studies used one test category, six studies used two test categories and one study reported the use of tests within all three categories.

Cognitive tests were most commonly employed ($n=14$) with a combination of cognitive and observational tests used in a further four studies [19, 22–24]. Two studies used a combination of cognitive and visual [25, 26]. Of the 14 studies that recorded the use of cognitive tests, seven studies used one cognitive test, six studies used two cognitive tests and one study used three cognitive tests. Of the studies that recorded the use of observational testing only, one study used one test while the second study used three testing methods. The two studies that used visual tests exclusively, one test method per study were recorded. The remaining seven studies [19, 22–24] used a combination of cognitive and observational test methods, cognitive and visual test methods [25, 26] or all three test categories [21].

Tests and screening methods used

A total of 20 different concussion diagnosis tests employed across all studies, differing in frequency. Tests included King-Devick (KD; 10 studies), Sports Concussion Assessment Tool (SCAT) version 3 (SCAT3; 9 studies), Balance Error Scoring System (BESS; 5 studies), Standardised Assessment of Concussion (SAC; 5 studies), tandem gait (TG; 2 studies), immediate post-concussion assessment and cognitive testing (ImPACT; 4 studies), video assessment (VA; 1 study), SCAT version 2 (SCAT2; 2 studies) and vestibular and oculomotor screening (VOMS; 2 studies). The following 11 tests were used once within the range of 16 studies (Brief Symptom Inventory-18, Gait Initiation, Mobile Universal Lexicon Evaluation System, Pitch Side Concussion Assessment, BTrackS Balance Test, Post Concussion Symptom Scale, Pitch Side Concussion Assessment (version 2), CogSport, modified Balance Error Scoring System, Motor Cognitive Test battery, SCAT5). SCAT3 was used on its own in three studies [17, 27, 28]. Three studies used SCAT3 and KD testing in combination with each other [29–31]. Three Studies used KD, individually [25, 32, 33]. One study used TG testing along with a combination of cognitive tests [22]. VOMS was used along with a set of cognitive/cognitive and observational tests [21, 25].

Administration of tests

Tests were administered by medical and non-medical personnel. Eighty-eight percent of participants were tested for concussion by doctors, clinicians, orthopaedic support, neurologists, or the assistance of certified athletic trainers or physiotherapists. Twelve percent of studies used team members, trained volunteers or the study research coordinator to conduct side-line concussion testing for scientific purposes only.

Level of sport

As the included studies are focused on diagnosis techniques for side-line concussion, there were a range of sports used throughout. For example, sports included rugby (union and league), American Football, ice hockey, baseball, soccer, Australian rules football, basketball, volleyball and wrestling. The sports activities can be divided into different levels of participation. Consisting of (1) professional/elite, semi-professional and (2) amateur/club, community level sports, 56% of participants fell into the professional/elite or semi-professional levels of sport, 36% were categorised as amateur/club or community sports. Another 4% were classified as a mixed level cohort of participations who were professional/elite, semi professional and amateur level athletes within a study. The remaining 4% did not state the level of sports played by those participants, however it is important to note the study did not report the level of sports played and testing were carried out within multidisciplinary concussion centre.

Most significant symptoms

Two studies employing the SCAT3 reported symptom frequency. Fatigue or low energy, along with neck pain [27] and pressure in head, headache and "don't feel right" [28] were reported most common post injury symptoms compared to baseline. One study [34] observed

“slow to get up” a total of 2240 times on 223 different occasions. Signs of clutching was reported 212 times (58.7%) in concussion assessment. Other concussion diagnosis signs reported within this study included unresponsiveness (n = 52), gait ataxia (n = 102), vacant stare (n = 98), and a post-impact seizure (n = 4).

Test scores and gender

Two of the 25 studies examined the differences in tests scores by gender. Results were significantly different when compared to each other. Both of these studies used SCAT3 as part (Kyle Harrold, Hasanaj [29]) or all (Downey, Hutchison [17]) of the testing protocols. It's clear the results are not similar when compared to each other and require further investigation. For example, Downey et al. [17] reported that while using SCAT3, male participants reported significantly more symptoms ($p = 0.012$), of greater severity ($p = 0.025$); and performed significantly worse on the SAC compared to females ($p = 0.012$). While the study by Kyle Harrold et al. [29] reported that using SCAT3, women reporting more total symptoms ($p = 0.001$, linear regression, accounting for age) and had higher symptom severity scores ($p = 0.006$).

Table 2
Overall study characteristics and modified Downs and Black [16] checklist (maximum score 22).

Author (Year)	Study Type	Assessment	Sample size (Concussed/Control)	Sex M/F	Mean Age (Range) Years	Sport	Test Post-Concussion	Score (max 22)
Broglia et al. [21]	PCS	BESS, SAC, SCAT5, ImPACT, VOMS, BSI-18	N = 1458	919/539	19.0 (N/A)	NCAA (sport unspecified)	3 times in 72hours (h): 0–1.25 h (side-line), 1.25–24 h (post-event), 24–72 h (clinic)	17
Buckley et al. [39]	Cohort	GI	N = 84 (42/42)	40/44	19.2 (N/A)	NCAA (sport unspecified)	24 hours	16
Downey et al. [17]	PCS	SCAT3	N = 45 (23/22)	19/26	20.0 (N/A)	Football, rugby, ice hockey, soccer, lacrosse, basketball, volleyball, field hockey, baseball, wrestling	3 to 5 days (acute), 3 weeks (post-acute)	14
Fallon et al. [26]	PCS	MULES, SCAT3	N = 681 (17)	422/259	17 (6-37)	Ice hockey, soccer, football	Side-line	13
Fuller et al. [27]	Cross Sectional	SCAT3	N = 639 (24)	All male	27.4 (N/A)	Rugby	After game of injury	14
Fuller et al. [40]	Pilot	PSCA	N = 165	All male	N/A	Rugby	Side-line (Time frame not stated)	17
Fuller et al. [33]	PCS	KD	N = 261	All male	27.6 (N/A)	Rugby	48 hours	18
Galetta et al. [22]	PCS	KD, SAC, TG	N = 332	192/140	11.0 (5-23)	Ice hockey, lacrosse	Side-line /rink-side. (Time frame not stated)	15
Gardner et al. [34]	Observational	VA	N = 400	All male	N/A	Rugby league	N/A	
Goble et al. [41]	Cross-Sectional	BBT	N = 25	11/14	20.7 (N/A)	College athletes- (unspecified)	48 hours	9
Graves et al. [23]	PCS	SOT, BESS	N = 15 (15)	All male	18.9 (N/A)	Football	1-14 days	15
Hanninen et al. [28]	PCS	SCAT3	N = 283 (27)	All male	27.0 (N/A)	Ice hockey	24 hours	15
Harrold et al. [29]	PCS	KD, SCAT3	N = 426	177/249	35.0 (N/A)	Sport, other	N/A	16

KD: King Devick test, GI: Gait Initiation, SOT: Sensory Organization, TG: Tandem Gait, GT: Gait Termination MULES: Mobile Universal Lexicon Evaluation Systems, PSCA: Pitch-Side Concussion Assessment Version 1, PSCA2: Pitch-Side Concussion Assessment Version 2, MotCoTe: Motor Cognitive Test Battery, VOMS: Vestibular/Ocular Motor Screening, SCAT2: Sport Concussion Assessment Tool Version 2, SCAT3: Sport Concussion Assessment Tool Version 3, SCAT5: Sport Concussion Assessment Tool Version 5, ImPACT: Immediate Post-Concussion and Cognitive Testing, SAC, standardized assessment of concussion; PCSS: Post-Concussion Symptom Scale, VA: Video Assessment, BESS: Balance Error Scoring System, mBESS: modified Balance Error Scoring System, MotCoTe: Motor Cognitive Test battery, PCS, prospective cohort study; RCS, retrospective cohort study

Author (Year)	Study Type	Assessment	Sample size (Concussed/Control)	Sex M/F	Mean Age (Range) Years	Sport	Test Post-Concussion	Score (max 22)
Hecimovich et al. [32]	PCS	KD	N = 22 (7/15)	All male	19.6 (N/A)	Australian football	10-20 min post-game	14
King et al. [30]	PCS Observational	KD, SCAT3	N = 104 (52/52)	All male	23.7 (N/A)	Rugby	Days 3, 7, 14, and 21	15
Leong et al. [42]	PCS	KD, SCAT2	N = 127 (11)	119/8	19.5 (N/A)	Football, basketball	N/A	17
Marinides et al. [24]	RCS	KD, PCS, SAC, BESS, ImPACT	N = 221 (30)	150/71	N/A	Football, lacrosse, soccer	87mins	14
Merritt et al. [43]	PCS	PCSS, ImPACT	N = 846 (86)	637/209	19.9 (N/A)	Football, basketball, ice hockey, soccer, lacrosse, wrestling, other	2, 7, and 14 days post injury.	18
Molloy et al. [44]	Descriptive	KD, PSCA2, CogSport	N = 176 (19/33)	All male	23.8 (N/A)	Rugby	48 hours	18
Oldham et al. [18]	Prospective, longitudinal	TG, BESS, mBESS	N = 76 (38/38)	All male	N/A	NCAA student-athletes	<48 hours	16
Putukian et al. [45]	Prospective cross-sectional	SCAT2	N = 263 (85/178)	184/79	20.3 (N/A)	Football, rugby, volleyball, football, crew	0.52 ± 1.18 days	13
Russell-Giller et al. [25]	Pilot	KD, VOMS	N = 71	N/A	14.0 (N/A)	Sports (unspecified), other	1 - 5 days	17
Seidman et al. [31]	PCS	KD, SCAT3	N = 337 (9/328)	All male	15.0 (N/A)	American Football	72 hours	12
Sufrinko et al. [19]	Case-control	ImPACT, SAC, BESS	N = 125(125)	85/40	16.8 (N/A)	Football, soccer, volleyball, basketball, wrestling, ice hockey, softball,	SAC, BESS: 24 to 48 hours, ImPACT: 5 to 7 and 10 to 14	13
Vartiainen et al. [20]	Pilot	SCAT3, MotCoTe	N= 16 (9/7)	All male	23.4 (N/A)	Ice hockey	36 hours	15
KD: King Devick test, GI: Gait Initiation, SOT: Sensory Organization, TG: Tandem Gait, GT: Gait Termination MULES: Mobile Universal Lexicon Evaluation Systems, PSCA: Pitch-Side Concussion Assessment Version 1, PSCA2: Pitch-Side Concussion Assessment Version 2, MotCoTe: Motor Cognitive Test Battery, VOMS: Vestibular/Ocular Motor Screening, SCAT2: Sport Concussion Assessment Tool Version 2, SCAT3: Sport Concussion Assessment Tool Version 3, SCAT5: Sport Concussion Assessment Tool Version 5, ImPACT: Immediate Post-Concussion and Cognitive Testing, SAC, standardized assessment of concussion; PCSS: Post-Concussion Symptom Scale, VA: Video Assessment, BESS: Balance Error Scoring System, mBESS: modified Balance Error Scoring System, MotCoTe: Motor Cognitive Test battery, PCS, prospective cohort study; RCS, retrospective cohort study								

Discussion

This systematic literature review is an analysis of current side-line assessments for efficacy of diagnosis of concussion in adults participating in professional, semi-professional and amateur sports. A definition of 'pitch side' in this review included side-line, rink side, changing rooms, or an assessment area immediately available that is not a clinical or hospital setting. The focus for this review utilised papers from 2015 onwards to align with the most recent consensus statement of concussion in sport (October 2016), as the authors deemed any study carried out prior to that year (2015) would be outdated. The main findings showed studies overall were of moderate to good quality and a variety of cognitive, observational and visual tests were utilised pitch side by mostly medical and allied health

personnel (e.g. physical therapists) to assess acute concussion in adults. However, the review also found that the majority of studies have investigated mostly professional/elite or semi-professional cohorts, aligning with these studies likely having access to medical and allied health staff to undertake these assessments. Conversely the limitation in non-medically trained researchers operating assessments are likely to be contributing to the paucity of studies not being undertaken at non-elite/club levels of participation where research is urgently required.

Key findings from the qualitative review showed a large range of different assessments used to quantify concussion. Interestingly, we found that less than half of the studies employed the SCAT assessment (either SCAT2, 3 or 5 for this systematic review). We found this observation surprising given that it has been suggested that the SCAT is the most widely accepted and deployable sport concussion assessment and screening tool currently available [12]. We appreciate that the use of SCAT in research is not a mandatory requirement, and that the objective of studies would be to test the efficacy of other modalities and cohorts, such as in a laboratory environment. However, in field studies such as those included in this review, that to reduce disparity in findings, that the SCAT assessment is used consistently across future studies.

The review also found that 25% of studies used a multi-modality approach to assess concussion by combining two or more testing batteries. While it may be argued that the SCAT does incorporate a multi-modality approach [1], there are elements that the SCAT assessment does not measure, such as the oculo-motor system [12]. Indeed, a previous study where the SCAT and an oculomotor test such as the King-Devick test have been implemented together, results showed a 100% sensitivity in diagnosing athletes suspected of concussion [35]. For clinically focused studies and application to clinical-practice, future work should incorporate two or more rapid and non-invasive pitch-side assessments for the diagnosis of concussion. This has been recently suggested in a systematic review by [36].

An ongoing concern, particularly for concussion in non-elite/community club-based sports, is the paucity of suitably qualified people who are allowed to administer the SCAT HIA. This is reflected in the current review where the majority of studies used SCAT at professional/elite levels of sport, where access to a medical practitioner was possible. As the consensus statement strictly asserts that only a medical practitioner can administer the SCAT as part of the clinical diagnosis [1], this may limit opportunities for suitably qualified scientists who are technically proficient at the SCAT but cannot provide a result, limiting research. Conversely other assessments such as the VOMS can be delivered by allied health professionals and the KD can be delivered by anyone, increasing the ability to undertake research, but without the use of the SCAT despite assertions that the SCAT is the most widely agreed upon assessment tool for concussion [12]. As studies have argued that when used in isolation assessments such as the KD or VOMS may not be sensitive enough alone to detect concussion [33, 37], we suggest from this systematic review that the SCAT should be incorporated with other testing modalities for data collection purposes, with any clinical diagnosis made by the associated team doctor outside of the study scope.

Interestingly, despite consensus on the use of video identification of concussion [38], only one study in this systematic review utilised video. A reason for this was because the video was used for post-event confirmation rather than used for confirmation of concussion at time of incident. With network media covering professional/elite events, video is easily accessible at matches. However, although multiple and multi-angled cameras are not available at amateur/community club levels, many sub-elite competitions will now incorporate a fixed camera supplied by the clubs themselves or via the local league for streaming or replay on the league's webpage. While not optimal in terms of clarity, it may assist the detection of concussion and we suggest that future studies involve video to confirm concussion events to improve study quality. Similarly, we found no studies employing impact sensors in a surveillance capacity. The use of impact sensors (attachments behind the ear or embedded within a gumshield/mouth guard) will assist video confirmation of suspected concussion events, however with no studies to support this hypothesis, we conclude there is no evidence to support their use at present for the detection of concussion.

Limitations of this review include research involving lab-based clinical assessments. However, the focus of this review was on published research that could assist at the lower levels of sport in community or amateur settings (i.e., in settings where the presence of medical professionals' pitch side may be limited or indeed non-existent). In these instances, the general approach would be to remove the participants from the sports activity where there is any indication that a concussion has occurred. These initial sideline tests may be substantiated at a later time using advanced diagnostic techniques by a medical professional. The studies included in this review utilised well established testing methods which offered some form of sideline tests for suspected concussions and have highlighted the necessity for a multimodal concussion assessment tool for the initial identification and assessment of concussion. This review highlighted the need for multi-modality concussion testing and that there is a clear disparity in research focusing on professional/elite levels and the lack of studies in amateur/club levels.

Conclusions

Recognising suspected concussion in sports participants is most effectively realized by using multimodal test protocols that are guided (via primary or secondary confirmation) by medical experts. Based on this review, the KD and SCAT appear to be the most robust tool for the primary assessment of concussion currently. Using additional tests such as VOMS from an observational perspective and balance testing such as BESS show promise in conjunction with cognitive testing. The addition of concurrent video review could potentially offer a promising approach to improve identification and evaluation of significant head impact events, and a multi-modality-based concussion evaluation process appears to be important to detect delayed-onset SRC, however current evidence does not support the use of impact sensor systems for real-time concussion screening. As shown in a recent systematic review [36] there is an urgent need to conduct research, using multi-modality assessment methods, but focusing on non-elite levels where concussion injuries occur regularly but a lack of resources and education preclude effective assessment and management.

Abbreviations

KD: King Devick test, GI: Gait Initiation, SOT: Sensory Organization, TG: Tandem Gait, GT: Gait Termination MULES: Mobile Universal Lexicon Evaluation Systems, PSCA: Pitch-Side Concussion Assessment Version 1, PSCA2: Pitch-Side Concussion Assessment Version 2, MotCoTe: Motor Cognitive Test Battery, VOMS: Vestibular/Ocular Motor Screening, SCAT2: Sport Concussion Assessment Tool Version 2, SCAT3: Sport Concussion Assessment Tool Version 3, SCAT5: Sport Concussion Assessment Tool Version 5, ImPACT: Immediate Post-Concussion and Cognitive Testing, SAC, standardized assessment of concussion; PCSS: Post-Concussion Symptom Scale, VA: Video Assessment, BESS: Balance Error Scoring System, mBESS: modified Balance Error Scoring System, MotCoTe: Motor Cognitive Test battery, PCS, prospective cohort study; RCS, retrospective cohort study

Declarations

Funding:

This study was funded by a grant from the Erasmus+ strategic partnerships program (2019-1-IE01-KA202-051555). The funder had no role in the study design, data collection, analysis or interpretation or the manuscript preparation.

Competing Interests:

AJP currently receives partial research salary funding from Sports Health Check charity (Australia) and Erasmus+ strategic partnerships program (2019-1-IE01-KA202-051555). AJP has previously received partial research funding from the Australian Football League, Impact Technologies Inc., and Samsung Corporation, and has provided expert reports in concussion legal proceedings. No other author has any declaration of interest.

Ethics approval and Consent to participate:

Not applicable

Consent for publication:

Not applicable

Availability of Data and Material:

Available upon request

Code availability:

Not applicable

Authors' contributions:

All authors have read and approved the manuscript.

ED and AJP are joint first authors and contributed to the searching, extraction, analysis and writing of the publication.

CC, EF, MMcd, GS, AW, MMcc, RD, SP & NH contributed to the initial search phases and first and second pass, all authors contributed and approved the final publication.

LR secured the funding for the work, informed the PICO, helped to develop the search strategy, contributed to extraction and analysis and publication preparation.

Acknowledgements:

The authors would like to acknowledge the contribution of ICHIRF (International Concussion and Head Injury Research Foundation) to the compilation of this Systematic Literature Review.

References

1. McCrory P, Meeuwisse W, Dvořák J, Aubry M, Bailes J, Broglio S, et al. Consensus statement on concussion in sport-the 5(th) international conference on concussion in sport held in Berlin, October 2016. *Br J Sports Med*. 2017 Jun;51(11):838-47.
2. Kamins J, Bigler E, Covassin T, Henry L, Kemp S, Leddy JJ, et al. What is the physiological time to recovery after concussion? A systematic review. *Br J Sports Med*. 2017 Jun;51(12):935-40.
3. Nordström A, Nordström P, Ekstrand J. Sports-related concussion increases the risk of subsequent injury by about 50% in elite male football players. *Br J Sports Med*. 2014 Oct;48(19):1447-50.
4. McPherson AL, Nagai T, Webster KE, Hewett TE. Musculoskeletal Injury Risk After Sport-Related Concussion: A Systematic Review and Meta-analysis. *Am J Sports Med*. 2019 Jun;47(7):1754-62.
5. Howell DR, Lynall RC, Buckley TA, Herman DC. Neuromuscular Control Deficits and the Risk of Subsequent Injury after a Concussion: A Scoping Review. *Sports Med*. 2018 May;48(5):1097-115.
6. Wilkerson GB, Grooms DR, Acocello SN. Neuromechanical Considerations for Postconcussion Musculoskeletal Injury Risk Management. *Curr Sports Med Rep*. 2017 Nov/Dec;16(6):419-27.
7. Scott E, Kidgell DJ, Frazer AK, Pearce AJ. The Neurophysiological Responses of Concussive Impacts: A Systematic Review and Meta-Analysis of Transcranial Magnetic Stimulation Studies. *Front Hum Neurosci*. 2020;14:306.
8. Pearce AJ, Tommerdahl M, King DA. Neurophysiological abnormalities in individuals with persistent post-concussion symptoms. *Neuroscience*. 2019 Jun 1;408:272-81.
9. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. 2020 Aug 8;396(10248):413-46.
10. Mez J, Alosco ML, Daneshvar DH, Salte N, Baucom Z, Abdolmohammadi B, et al. Validity of the 2014 traumatic encephalopathy syndrome criteria for CTE pathology. *Alzheimers Dement*. 2021 Apr 7.
11. Aubry M, Cantu R, Dvorak J, Graf-Baumann T, Johnston K, Kelly J, et al. Summary and agreement statement of the First International Conference on Concussion in Sport, Vienna 2001. Recommendations for the improvement of safety and health of athletes who may suffer concussive injuries. *Br J Sports Med*. 2002 Feb;36(1):6-10.
12. Echemendia RJ, Broglio SP, Davis GA, Guskiewicz KM, Hayden KA, Leddy JJ, et al. What tests and measures should be added to the SCAT3 and related tests to improve their reliability, sensitivity and/or specificity in sideline concussion diagnosis? A systematic review. *Br J Sports Med*. 2017 Jun;51(11):895-901.
13. Swart J, Bigard X, Fladischer T, Palfreeman R, Riepenhof H, Jones N, et al. Harrogate consensus agreement Cycling specific Sport Related Concussion. *Sports Medicine and Health Science*. 2021.
14. Patricios J, Fuller GW, Ellenbogen R, Herring S, Kutcher JS, Loosemore M, et al. What are the critical elements of sideline screening that can be used to establish the diagnosis of concussion? A systematic review. *British journal of sports medicine*. 2017;51(11):888-94.
15. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71.
16. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health*. 1998 Jun;52(6):377-84.
17. Downey RI, Hutchison MG, Comper P. Determining sensitivity and specificity of the Sport Concussion Assessment Tool 3 (SCAT3) components in university athletes. *Brain Inj*. 2018;32(11):1345-52.
18. Oldham JR, Difabio MS, Kaminski TW, Dewolf RM, Howell DR, Buckley TA. Efficacy of Tandem Gait to Identify Impaired Postural Control after Concussion. *Medicine & Science in Sports & Exercise*. 2018;50(6):1162-8.

19. Sufrinko A, McAllister-Deitrick J, Womble M, Kontos A. Do Sideline Concussion Assessments Predict Subsequent Neurocognitive Impairment After Sport-Related Concussion? *J Athl Train*. 2017 Jul;52(7):676-81.
20. Vartiainen MV, Holm A, Lukander J, Lukander K, Koskinen S, Bornstein R, et al. A novel approach to sports concussion assessment: Computerized multilimb reaction times and balance control testing. *J Clin Exp Neuropsychol*. 2016;38(3):293-307.
21. Broglio SP, Harezlak J, Katz B, Zhao S, McAllister T, McCrea M. Acute Sport Concussion Assessment Optimization: A Prospective Assessment from the CARE Consortium. *Sports Med*. 2019 Dec;49(12):1977-87.
22. Galetta KM, Morganroth J, Moehringer N, Mueller B, Hasanaj L, Webb N, et al. Adding Vision to Concussion Testing: A Prospective Study of Sideline Testing in Youth and Collegiate Athletes. *J Neuroophthalmol*. 2015 Sep;35(3):235-41.
23. Graves BS. UNIVERSITY FOOTBALL PLAYERS, POSTURAL STABILITY, AND CONCUSSIONS. *Journal of Strength & Conditioning Research*. 2016;30(2):579-83.
24. Marinides Z, Galetta KM, Andrews CN, Wilson JA, Herman DC, Robinson CD, et al. Vision testing is additive to the sideline assessment of sports-related concussion. *Neurol Clin Pract*. 2015;5(1):25-34.
25. Russell-Giller S, Toto D, Heitzman M, Naematullah M, Shumko J. Correlating the King-Devick Test With Vestibular/Ocular Motor Screening in Adolescent Patients With Concussion: A Pilot Study. *Sports Health*. 2018 Jul/Aug;10(4):334-9.
26. Fallon S, Akhand O, Hernandez C, Galetta MS, Hasanaj L, Martone J, et al. MULES on the sidelines: A vision-based assessment tool for sports-related concussion. *Journal of the Neurological Sciences*. 2019 2019/07/15;402:52-6.
27. Fuller CW, Fuller GW, Kemp SP, Raftery M. Evaluation of World Rugby's concussion management process: results from Rugby World Cup 2015. *Br J Sports Med*. 2017 Jan;51(1):64-9.
28. Hänninen T, Parkkari J, Tuominen M, Öhman J, Howell DR, Iverson GL, et al. Sport Concussion Assessment Tool: Interpreting day-of-injury scores in professional ice hockey players. *Journal of Science & Medicine in Sport*. 2018;21(8):794-9.
29. Kyle Harrold G, Hasanaj L, Moehringer N, Zhang I, Nolan R, Serrano L, et al. Rapid sideline performance meets outpatient clinic: Results from a multidisciplinary concussion center registry. *J Neurol Sci*. 2017 Aug 15;379:312-7.
30. King D, Gissane C, Hume PA, Flaws M. The King-Devick test was useful in management of concussion in amateur rugby union and rugby league in New Zealand. *J Neurol Sci*. 2015 Apr 15;351(1-2):58-64.
31. Seidman DH, Burlingame J, Yousif LR, Donahue XP, Krier J, Rayes LJ, et al. Evaluation of the King-Devick test as a concussion screening tool in high school football players. *J Neurol Sci*. 2015 Sep 15;356(1-2):97-101.
32. Hecimovich M, King D, Dempsey AR, Murphy M. The King-Devick test is a valid and reliable tool for assessing sport-related concussion in Australian football: A prospective cohort study. *J Sci Med Sport*. 2018 Oct;21(10):1004-7.
33. Fuller GW, Cross MJ, Stokes KA, Kemp SPT. King-Devick concussion test performs poorly as a screening tool in elite rugby union players: a prospective cohort study of two screening tests versus a clinical reference standard. *British Journal of Sports Medicine*. 2019;53(24):1526-32.
34. Gardner AJ, Howell DR, Levi CR, Iverson GL. Evidence of Concussion Signs in National Rugby League Match Play: a Video Review and Validation Study. *Sports Medicine-Open*. 2017;3.
35. Ventura RE, Jancuska JM, Balcer LJ, Galetta SL. Diagnostic tests for concussion: is vision part of the puzzle? *J Neuroophthalmol*. 2015 Mar;35(1):73-81.
36. Harris SA, Dempsey AR, Mackie K, King D, Hecimovich M, Murphy MC. Do Sideline Tests of Vestibular and Oculomotor Function Accurately Diagnose Sports-Related Concussion in Adults? A Systematic Review and Meta-analysis. *Am J Sports Med*. 2021 Aug 25:3635465211027946.
37. Yorke AM, Smith L, Babcock M, Alsalaheen B. Validity and Reliability of the Vestibular/Ocular Motor Screening and Associations With Common Concussion Screening Tools. *Sports Health*. 2017 Mar/Apr;9(2):174-80.
38. Davis GA, Makdissi M, Bloomfield P, Clifton P, Echemendia RJ, Falvey É C, et al. International consensus definitions of video signs of concussion in professional sports. *Br J Sports Med*. 2019 Oct;53(20):1264-7.
39. Buckley TA, Oldham JR, Munkasy BA, Evans KM. Decreased Anticipatory Postural Adjustments During Gait Initiation Acutely Postconcussion. *Archives of Physical Medicine and Rehabilitation*. 2017 2017/10/01;98(10):1962-8.
40. Fuller GW, Kemp SP, Decq P. The International Rugby Board (IRB) Pitch Side Concussion Assessment trial: a pilot test accuracy study. *Br J Sports Med*. 2015 Apr;49(8):529-35.
41. Goble DJ, Manyak KA, Abdenour TE, Rauh MJ, Baweja HS. AN INITIAL EVALUATION OF THE BTRACKS BALANCE PLATE AND SPORTS BALANCE SOFTWARE FOR CONCUSSION DIAGNOSIS. *International Journal of Sports Physical Therapy*. 2016;11(2):149-55.

42. Leong DF, Balcer LJ, Galetta SL, Evans G, Gimre M, Watt D. The King-Devick test for sideline concussion screening in collegiate football. *J Optom.* 2015 Apr-Jun;8(2):131-9.
43. Merritt VC, Meyer JE, Arnett PA. A novel approach to classifying postconcussion symptoms: The application of a new framework to the Post-Concussion Symptom Scale. *J Clin Exp Neuropsychol.* 2015;37(7):764-75.
44. Molloy JH, Murphy I, Gissane C. The King-Devick (K-D) test and concussion diagnosis in semi-professional rugby union players. *J Sci Med Sport.* 2017 Aug;20(8):708-11.
45. Putukian M, Echemendia R, Dettwiler-Danspeckgruber A, Duliba T, Bruce J, Furtado JL, et al. Prospective clinical assessment using Sideline Concussion Assessment Tool-2 testing in the evaluation of sport-related concussion in college athletes. *Clin J Sport Med.* 2015 Jan;25(1):36-42.

Figures

Figure 1

Preferred Reporting Items for Systematic Reviews and Meta-analysis flow diagram of the studies included in the review [15].

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [PRISMA2020checklistBMCED.docx](#)