Modeling of CO2 Emission Statistics in Turkey by Fuzzy Time Series Analysis

Fatih Cemrek (✉ fcemrek@ogu.edu.tr)

Eskişehir Osmangazi University, Faculty of Science and Letters, Department of Statistics

<https://orcid.org/0000-0002-6528-7159>

Research Article

Keywords: Time Series Analysis, Fuzzy Time Series Analysis, CO2 emission, RMSE, Chen Models, Gustafson-Kessel clustering algorithm

Posted Date: January 17th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1261965/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Modeling of CO₂ emission statistics in Turkey by fuzzy time series analysis

Fatih ÇEMREK

Eskişehir Osmangazi University, Faculty of Science and Letters, Department of Statistics,
Meşelik Kampüsü, Büyükdere Mah. Prof. Dr. Nabi AVCI Bulvarı No: 4, 26040,
Odunpazarı/ESKİŞEHİR/TURKEY

fcemrek@ogu.edu.tr +905336305751

ORCID ID: <https://orcid.org/0000-0002-6528-7159>

Modeling of CO₂ emission statistics in Turkey by fuzzy time series analysis

Abstract

The process of determining the values which a time series will receive in the future is a very important concept. The fuzzy time series method has been widely used in recent years as it is more convenient to process data in small samples which are incomplete and/or ambiguous, and it does not contain any assumptions for time series. In this study, fuzzy time series analysis was used to predict CO₂ emission values for Turkey. For this purpose, time series (annual) for total greenhouse gas emissions by sectors (CO₂ equivalent) between 1990 and 2016 were analyzed. The main goal of this study is to model greenhouse gas emission statistics in Turkey with fuzzy time series analysis. The RMSE value was taken into consideration to determine the most suitable model among the analysis performed.

Key Words: Time Series Analysis, Fuzzy Time Series Analysis, CO₂ emission, RMSE, Chen Models, Gustafson-Kessel clustering algorithm

1. Introduction

Global warming and climate change are among the most important problems in recent years. Global warming is caused by global economy, energy consumption, and by gases with greenhouse effect such as carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), Hydro flora carbon (HFC), Fluorocarbons (PFC), sulfur hexafluor (SF₆). Carbon dioxide (CO₂) is one of the most important greenhouse gases that cause climate change and global warming (Ayvaz et al 2017; Mohsen 2019).

Greenhouse gases are released from both natural and human sources. The major greenhouse gases which exist in the atmosphere are water vapor (H₂O), carbon dioxide (CO₂), nitrogen oxide (N₂O), methane (CH₄) and ozone (O₃) gases. The most important greenhouse gases which are not naturally present in the atmosphere and which occur as a result of human activities are hydrofluorocarbons (HFCs), perfluorocarbons (PFCs) and sulfur hexafluoride (SF₆) gases. The amount of greenhouse gases has been increasing rapidly in recent years with economic activity and the use of fossil fuels. Consequently, this causes global warming (Kuşkaya and Gençglu 2017). With the atmosphere's heat retention feature, the seas and oceans do not freeze. This heating and heat retention feature of the atmosphere is called **the greenhouse effect**.

Emission factors are provided as carbon dioxide (CO₂) equivalent (CO₂e). Emissions of greenhouse gases other than CO₂ (CH₄, N₂O and CFC, HCFC) are calculated separately and converted into CO₂ equivalent. During this

conversion, the release amounts of each greenhouse gas are multiplied by the global warming potential of that gas ((Tatar and Ozer 2018).

Carbon dioxide (CO₂): its share in greenhouse gases is 82%. CO₂, whose amount did not change in the atmosphere for millions of years, has increased by 31% since the start of the Industrial Revolution. According to the calculations, the annual increase of CO₂ gas in the atmosphere in the 20 years prior to 1990 was 0.4%, while the amount of increase in the following years ranged from 0.2% to 0.8%. The biggest source of CO₂ released into the atmosphere is fossil fuels, which are used a lot (Akın 2006).

The Turkish Statistical Institute (TÜİK, 2018) announced that the total greenhouse gas emissions in Turkey in 2016 were 496.1 million tons of carbon dioxide equivalents. In this period, energy-derived emissions as CO₂ equivalent received the largest share of total emissions (72.8%). This was followed by industrial operations and crop use (12.6%), agricultural activities (11.4%), and waste (3.3%), respectively. Total greenhouse gas emissions as CO₂ equivalent increased by 135.4% compared to 1990. CO₂ equivalent emissions per capita were 3.8 tonnes per capita in 1990, while in 2016 it was 6.3 tonnes per capita (Dikmen 2019).

2. Literature Review

2.1. Examining classical ARIMA and CO₂ studies,

Abdullah and Pauzi (2015) examined the methods used to estimate CO₂ emissions. The articles published in international journals between 2003 and 2013 were analyzed to determine which methods should be applied and which factors have been regularly investigated.

Ozceylan (2018) estimated CO₂ emissions with the help of particle optimization (PSO) and artificial bee colony (ABC) techniques (models were used as linear, exponential and quadratic) and by means of the socio-economic indicators in Turkey (energy consumption, population, GDP and number of motor vehicles). The data used are from 1980-2008 and the predictions have been made until 2030.

Ayvaz et al. (2017) used different discrete grey models (DGM) to estimate energy-related CO₂ emissions in Turkey, Europe and the Eurasian region. The data used covers the years between 1965 and 2014. With this data, CO₂ emissions from 2015-2030 have been estimated.

Liu et al. (2017) handled the problem of carbon emission based on carbon emission time series data and chaos theory to make the relationships between the data clearer. The BP neural network model was used to estimate carbon emissions.

Appiah et al. (2018) exploited a two-layer forward feed neural network model with Tangent activation function that occurs with hidden neurons where neurons are used as linear output in their work. In the study, a nonlinear least squares algorithm such as LM (Levenberg-Marquardt) was applied to estimate emission for the selected emerging economies.

Garip and Oktay (2018) estimated Turkey's CO₂ emissions using random forest and support vector machines methods from popular machine learning methods. In the study, data from 1965-2003 were used for education and estimates for 2004-2014 were obtained. According to the results of the study, the Support Vector Machine yielded more successful results than the random forest method.

Li et al. (2019), in their study, used metabolic grey model (MGM), adapted exponential curve model (MECM), autoregressive integrated moving average model (ARIMA), and neural network model BP (BP) to estimate the metabolic energy demand of Central Africa.

Sutthichaimethee et al. (2019) made use of a second-degree autoregressive structural equality model (second-degree autoregressive SEM).

Dorogoi and Mokhtar (2019) implemented trend analysis and double exponential smoothing method using the data between 1967 and 2014 to estimate energy consumption in some sectors (industry, agriculture, transport and households - in general-commercial) which have significant relationship with the greenhouse gas emissions in Iran.

Oyehan et al. (2017) applied trend analysis to CO₂ data for the years 1980-2008 for Persian Gulf Countries (Bahrain, Iran, Iraq, Qatar, Saudi Arabia, Kuwait and the United Arab Emirates).

Maleki et al. (2018) employed autoregressive integrated moving average (ARIMA) and Sini Network autoregressive (NNAR) techniques to the time series on water characteristics of water treatment plants. As a result of the study, it was determined that, compared to ARIMA, NNAR provided better predictive success for CO₂ in terms of R².

2.2. On reviewing the studies using Fuzzy ARIMA

In their study, Abd Rahman et al. (2013) envisioned a monthly air pollution index (API) for 10 years with the data obtained from industrial and residential monitoring stations in Malaysia. In the study, ARIMA and Fuzzy Time Series (Chen's method and Yu's method) were used to predict API measurements. Comparison was made by

obtaining RMSE values for the three proposed models. It was determined that the neural network model (ANN) provided better results.

Karaaslan and Gezen (2017) attempted to predict the total energy demand for Turkey and determine the amount of unused energy and the distribution of this demand among sectors. The study used annual data from 1990 to 2012 and estimated energy demand until 2023 using a fuzzy grey regression model.

Mahla et al. (2019) used the ARIMA model to predict emissions from biogas.

Atsalakis et al. (2015) examined hourly data ($n=8760$) for January 1, 2009-December 31-, 2009 period using an integrated neuro-fuzzy controller (PATSON) technique. The prediction system is based on Adaptive Neural Fuzzy Systems (ANFIS).

2.3. When Fuzzy and CO₂ studies are inspected,

Tavan M (2019) introduced a new hybrid modelling to predict carbon dioxide emissions in order to make the correct decision to reduce air pollution in Iran. The paper, CO₂ emissions in Iran in the period of 1980-2014 was predicted using three models of Auto-Regressive Distributed Lag (ARDL), Fuzzy Linear Regression (FLR) and hybrid model based on the combination of ARDL and FLR models, and then the prediction accuracy of the models is compared.

Examining the literature, it is observed that studies on greenhouse gas emissions and classical ARIMA, and studies on Fuzzy ARIMA and other application areas exist. However, the aim of this study is to model greenhouse gas emission statistics in Turkey with fuzzy time series analysis. For this purpose, greenhouse gas emission was modeled using four different Fuzzy ARIMA models, and their performance was evaluated.

This paper consists of four parts. The first section is the introductory section, and the second part contains fuzzy time series models. The third section is the application section. The fourth part is the concluding part.

3. Fuzzy Time Series

The fuzzy time series method has been widely used in recent years as it is more convenient to process data in small samples which are incomplete and/or ambiguous, and it does not contain any assumptions for time series. There is a fundamental difference between fuzzy time series and traditional time series. The values in fuzzy time series are fuzzy clusters, whereas the values used in traditional time series are real numbers. A fuzzy set is a cluster with fuzzy boundaries.

The first definition of the fuzzy time series was proposed by Song and Chissom (1993a, 1993 b), and the method was developed by Chen (1996). Later, other studies were conducted on prediction problems using fuzzy time series concept (Chen 1996, 2000; Chen and Hwang 2000; Hwang et al. 1998; Song and Chissom 1994; Song and Leland 1996; Song et al. 1995; Sullivan and Woodall 1994).

Song and Chissom (1993a, 1993b, 1994) suggested the model of first-degree time-invariant fuzzy time series, and to predict an annual time series, they proposed first-degree fuzzy time-varying time series.

Sullivan and Woodall (1994) examined the first-degree time-varying time series model proposed by Song and Chissom and the first-degree time-invariant fuzzy time-series model. They have compared the models to each other.

Song et al (1995) presented a new fuzzy time series model for a fuzzy number observation. Chen (1996) developed a new prediction method using fuzzy time series. Hwang et al (1998) proposed a method which eliminates the prediction problems with the use of fuzzy time series. Chen and Hwang (2000) developed a method for temperature estimation using fuzzy time series.

Chen (2000) developed a new method for the prediction problem using high-degree fuzzy time series.

3.1. Chen (1996) model: This method consists of 6 steps defined as follows.

Step 1: Determination of universal sets and subintervals.

In this step, the universal set is determined according to the smallest and largest values of the time series. The defined universal set is divided into a predetermined number of subintervals. If the smallest value of the time series is $Dmin$ and the largest value is $Dmax$, the universal set is defined as follows.

$$U=[Dmin-D1,Dmax+D2] \quad (1)$$

Here, $D1$ and $D2$ are two small randomly chosen numbers.

Step 2: Determination of fuzzy sets: fuzzy sets defined in peer (1) are determined.

Step 3: Observations are fuzzy

Subinterval for the observation of each classical time series is determined. The fuzzy set with the largest membership value of this subinterval gives the fuzzy value of the classical time series observation.

Step 4: Determination of fuzzy relations

Determination of the fuzzy logical relationships can be explained with an example. Let the elements of (t) , a BZS with five observations, be as follows. $A1,1,A2,A2,A3$.

In this case, fuzzy relations are defined as follows.

$$A1 \rightarrow A1 \quad A1 \rightarrow A2 \quad A2 \rightarrow A2 \quad A2 \rightarrow A3$$

Fuzzy relations are grouped as follows.

$$A1 \rightarrow A1,2 \quad A2 \rightarrow A2, A3$$

Step 5: Predictions are obtained.

There are 3 different situations in the process of obtaining predictions.

Case 1, A_j is the predictive value for time t when there is only one fuzzy logical relationship as $A_i \rightarrow A_j$ in the fuzzy relations sequence.

Case 2, If $A_i \rightarrow A_j, s, Al$, the predictive value is equal to A_j, As, Al .

Case 3, If $A_i \rightarrow \emptyset$, the predictive value equals A_i .

Step 6: Obtained predictive values are defuzzified.

‘Centralization-defuzzification method’ is used to obtain the results. There are 3 different situations in defuzzification process.

Case 1, if the predictive value is equal to A_j , defuzzified predictive value is the cluster center c_j of A_j fuzzy set.

Case 2, if the predictive value is $A_j,,l$, defuzzified predictive value is calculated in the form of $(c_j+cs+cl)3'$ as arithmetic average of the cluster centers of the fuzzy sets A_j,As,Al .

Case 3, if A_i is equal to the empty set ($\emptyset$), then the defuzzified predictive value is cluster center c_i of A_i fuzzy set (Eğrioğlu et al 2011).

3.2. Chen (2002) Model: General Definitions

Let $U=\{u_1, u_2, u_3, \dots, u_n\}$ be the universal cluster. The elements of U have intervals. These intervals are obtained by breaking down the universal set according to a previously determined fixed interval length. Fuzzy sets of U are defined as:

$$A=A(u_1)/u_1+(u_2)/u_2 +\dots+f(u_n)/u_n \quad (2)$$

f_A : is the membership function of A function. $f_A : U \rightarrow [0,1]$ and (u_i) denote the membership degree of the UI in A [31].

Definition 1:

Let $X(t)$ be a real-valued subset with $(t=..., 0,1,2,...)$. $F(t)$ is the sum including the fuzzy set defined as $f_i(t)$ ($i = 1,2,...$) and is defined as a fuzzy time series to $F(t)$.

$F(t)$ time series obtained after determining A_i fuzzy set corresponding to appropriate sub-intervals of the X time series and to observation of each time series is called “**fuzzy time series**”. $F(t)$ is also a function of time. In other words, since the universal set can take different values at different times, $F(t)$ can also take different values at different times.

In definition 1;

1. $F(t)$ is a function of time.
2. $F(t)$ can be thought of as a verbal variable, taking verbal values where all the values it will take can be represented by fuzzy sets.
3. $f_i(t)$ ($i = 1,2,...$) are possible verbal values of $F(t)$. $f_i(t)$ ($i = 1,2,...$) is represented by fuzzy sets.

Definition 2: including $F(T)$ fuzzy time series, for any time $F(T) = F(T-1)$, and $F(T)$ has only finite elements. Thus, $F(t)$ is called time-invariant fuzzy time series.

Definition 3:

Let $F(t)$ be a fuzzy time series. If $F(t), F(t-1), F(t-2),...$ causes $F(t-n)$, then the fuzzy logical relationship is expressed as follows.

$$F(t-n), ..., F(t-2), F(t-1) \rightarrow F(t) \quad (3)$$

and it is called n . degreed fuzzy time series predictive model.

Chen (2002) method consists of the following stages.

Step 1: Determination of universal sets and subintervals.

In this step, the universal set (U) is determined with the smallest (min) and the largest value (max) of the time series.

$$U = [D_{min} - D_1, D_{max} + D_2] \quad (4)$$

Here, D_1 and D_2 are two randomly chosen positive numbers.

A predetermined number of subintervals of u_i are defined so that this universal set is $U=\{u_1, u_2, u_3, \dots, u_n\}$.

Step 2: Determination of fuzzy sets

The fuzzy $A_1, A_2, \dots, A_k$ sets are defined in the universal set of U .

$$A_1 = a_{11}/u_1 + a_{12}/u_2 + \dots + a_{1m}/u_m$$

$$A_2 = a_{21}/u_1 + a_{22}/u_2 + \dots + a_{2m}/u_m \quad (5)$$

.

.

.

$$A_k = a_{k1}/u_1 + a_{k2}/u_2 + \dots + a_{km}/u_m$$

Here, $a_{ij} \in [0,1]$ ve $1 \leq i \leq k$ $1 \leq j \leq m$. The values of a_{ij} in the A_i fuzzy set.

Step 3: Observations are defuzzified.

It is matched with the fuzzy set with the largest membership value for the subinterval set determined for each time series data. Thus, the time series is defuzzified.

Step 4: Determination of fuzzy relations

The determination of the fuzzy logical relationships can be explained with an example., let the elements of (t) , a BZS with five observations, be as follows. $A1,1,A2,A2,A3$. In this case, fuzzy relations are determined as follows.

$A1 \rightarrow A1$ $A1 \rightarrow A2$ $A2 \rightarrow A2$ $A2 \rightarrow A3$ Fuzzy relations are grouped as follows. $A1 \rightarrow A1,2$ $A2 \rightarrow A2, A3$

Fuzzy predictions are obtained. Using the artificial neural network model obtained in the previous step, the outputs of the artificial neural network are calculated. The calculated outputs are the numbers of fuzzy sets which the predictions belong to. Accordingly, fuzzy predictions are derived from the outputs of the artificial neural network model obtained in the previous step.

Step 5: Calculating predictions.

The following principles are taken into account at this stage.

Case 1: If fuzzified observations for year $i \dots$ from $K(k \geq 2)$ k^{th} level are $A_{ik}, A_{i(k-1)}, \dots$, and A_{il} , and there is a fuzzy logic present in k^{th} level fuzzy logic relations, then the groups are defined as follows.

$$A_{ik}, A_{i(k-1)}, \dots, A_{il} \rightarrow A_j$$

$A_{ik}, A_{i(k-1)}, \dots, A_{il}$ ve A_j are fuzzy sets, and they are in m_j interval which is the midpoint of u_j and u_j , which is the largest membership of A_j , and predictive value for year i is m_j .

Case 2 If observations fuzzified from the k th level ($k \geq 2$) for the year are $A_{ik}, A_{i(k-1)}, \dots$, and A_{il} , if there is a fuzzy logic relationship in the k th level fuzzy logic relations, the groups are defined as follows.

$$A_{ik}, A_{i(k-1)}, \dots, A_{il} \rightarrow A_{j1}$$

$$A_{ik}, A_{i(k-1)}, \dots, A_{il} \rightarrow A_{j2}$$

.

.

.

$$A_{ik}, A_{i(k-1)}, \dots, A_{il} \rightarrow A_{jp}$$

$A_{ik}, A_{i(k-1)}, \dots, A_{il}, A_{j1}, A_{j2}, \dots, A_{jp}$ are fuzzy sets. It can be seen that there is an uncertainty to predict the record of year i (fuzzy observation for year i can be A_{j1}, A_{j2} or A_{jp}). In this case, there should be high-grade fuzzy observations for year i . Therefore, there will be no uncertainty for the prediction of the year i . To eliminate this uncertainty, suppose there is an integer value of m ($m \geq k$). For the year i , the m^{th} degree fuzzified observation values are $A_{im}, A_{i(m-1)}, \dots, A_{i1}$, and the fuzzy logic relationships in m^{th} degree fuzzy logic relation groups are defined as follows:

$$A_{im}, A_{i(m-1)}, \dots, A_{i1} \rightarrow A_j$$

$A_{im}, A_{i(m-1)}, \dots, A_{i1}$ ve A_j are fuzzy sets.

The largest membership value of A_j is u_j , and it occurs in m_j interval which is the midpoint of $u_2, \dots, u_j$. Consequently, the predictive value of year i is m_j .

Case 3: Let the observations fuzzified from k^{th} degree for year i be $A_{ik}, A_{i(k-1)}, \dots$, and A_{il} . If the right side of the fuzzy logic relation is empty, the logical relation groups of k^{th} degree are defined as follows.

$$A_{ik}, A_{i(k-1)}, \dots, A_{il} \rightarrow \#$$

Here, $A_{ik}, A_{i(k-1)}, \dots$ and A_{il} are fuzzy sets, and the largest membership value of $A_{ik}, A_{i(k-1)}, \dots$ and A_{il} is $u_{ik}, u_{i(k-1)}, \dots$; and it occurs in m_{ik} , the midpoint of $m_{ik}, m_{i(k-1)}, \dots, i$ and m_{il} interval. Thus, the predicted value for year i is calculated as follows [31].

$$\frac{1 \cdot m_{ik} + 2 \cdot m_{i(k-1)} + \dots + k \cdot m_{il}}{1 + 2 + \dots + k} \quad (6)$$

Step 6. Defuzzification process for fuzzy predictions

Defuzzification process is applied to fuzzy predictions. During defuzzification, centralization method is used. If fuzzy prevision is A_j , defuzzified prediction is the midpoint value of u_j which is the interval with the highest membership value for this fuzzy set.

3.3. Fuzzy time series models based on fuzzy clustering

Gustafson ve Kessel (1979)

Gustafson and Kessel (1979) proposed an adapted version of the fuzzy clustering algorithm. This algorithm uses the Mahalanobis distance instead of the Euclidean distance. The distance of Mahalanobis forms ellipse-shaped clusters .

Mahalanobis distance($d_{ikA_i}^2$), the distance norm of the GK algorithm, is defined as

$$d_{ikA_i}^2 = (Z_k - V_i)^T A_i (Z_k - V_i) \quad (7)$$

The purpose function of the GK clustering algorithm is expressed in equality (8).

$$J(X, U, V, A) = \sum_{i=1}^N \sum_{k=1}^c (u_{ik})^m d_{ikA_i}^2 \quad (8)$$

The steps of the Gustafson-Kessel clustering algorithm are as follows:

Step 1: Initial values, cluster number (c), fuzzification index (m), number of iterations (i), membership degrees matrix (u), cluster center values (v) and stop criteria (ε) are determined. Here, it is defined as ($\varepsilon > 0$), ($1 < c < N$), ($m > 1$).

Step 2: Fuzzy cluster centers are calculated.

$$v_k = \frac{\sum_{i=1}^N u_{ik}^m x_i}{\sum_{i=1}^N u_{ik}^m}, \quad 1 \leq k \leq c \quad (9)$$

Step 3: The Mahalanobis distance is calculated for each element.

Step 4: Fuzzy covariance matrices (F_i) are calculated for each set separately.

$$F_i = \frac{\sum_{k=1}^N (d_{ik})^m (Z_k - V_i)(Z_k - V_i)^T}{\sum_{k=1}^N (d_{ik})^m} \quad (10)$$

Step 5: The membership matrices are updated.

$$u_{ik} = \left[\sum_{j=1}^c \left(d_{ikA_k} / d_{ijA_k} \right)^{2/(m-1)} \right]^{-1} \quad (11)$$

Step 6: The process is terminated when $\|V_t - V_{t-1}\| < \varepsilon$ condition is met. If not, return to step 2 [Gustafson and Kessel 1979; Balasko et. al. 2005, 2009).

4. Analysis Of Greenhouse Gas Emission Statistics in Turkey With Fuzzy Time Series Analysis

Fuzzy logic applications, unlike statistical methods, may include methods which can be analyzed with a small number of data which do not require any assumptions. Fuzzy time series is also a method that can be applied in

cases where the number of samples is small without requiring any assumption on the time series. The most important step in the implementation of the fuzzy time series models in the literature is that the fuzzy equivalent of the classical time series can be obtained, that is they can be fuzzified. The main purpose of the application is to model the greenhouse gas emission statistics in Turkey with fuzzy time series analysis. Expressing which model is the most appropriate in the analysis performed will be evaluated by considering the RMSE value.

In the implementation phase of the study, total greenhouse gas emissions (CO₂ equivalent) in Turkey between 1990 and 2016 will be evaluated using fuzzy time series analysis. For this purpose, data (Table 1) will be analyzed using Chen (1996) method, Chen (2002) method, and fuzzy time series models based on fuzzy clustering, which are fuzzy time series based on fuzzy C-average (FCMFTS) and fuzzy time series based on Gustafson - Kessel clustering (GKFTS).

Table 1: Total greenhouse gas emissions by sector in Turkey (CO₂ equivalent)

Year	Greenhouse gas emission (million tons)	Year	Greenhouse gas emission (million tons)	Year	Greenhouse gas emission (million tons)
1990	210,7	1999	272,1	2008	387,9
1991	218,7	2000	293,5	2009	395,9
1992	224,7	2001	274,4	2010	402,6
1993	233,4	2002	280,8	2011	431,4
1994	227,6	2003	300,3	2012	445,6
1995	242,2	2004	311,2	2013	439,0
1996	261,2	2005	332,7	2014	451,8
1997	272,6	2006	356,8	2015	469,9
1998	274,5	2007	390,5	2016	496,1

(Source: TURKISH STATISTICAL INSTITUTE 13th 2018)

4.1. Chen 1996 model Results: Implementing prediction algorithm of Chen 1996 fuzzy time series model, fuzzy time series model prediction analysis was performed for CO₂ emission value in Turkey. After the related analysis, RMSE value for CO₂ emissions of Chen (1996) model was obtained as 11.61.

Table 2: Chen (1996) Model Results

Year	Greenhouse gas emission (million tons)	Chen (1996)
2012	445,6	445,6
2013	439	439
2014	451,8	451,8
2015	469,9	469,9
2016	496,1	496,1
RMSE		11,61

4.2. Chen 2002 model Results:

Chen (2002) model, one of fuzzy time series analysis methods, was applied for CO₂ emission value in Turkey. The Chen (2002) model is the high-order fuzzy time series model. Analyzing the Chen 2002 model for the related data, the values in Table 3 were obtained. In the analysis of time series, Chen (2002) model was tested from 2nd degree to 12th degree model by changing the number of fuzzy sets between 3 and 10.

The optimal model degree and number of fuzzy sets were determined according to the RMSE criteria calculated for validity set by taking the time series training set between 1990 and 2006 and observations validity set between 2007 and 2011. The performance of the method was observed using as a test set between 2012-2016 in the last 5 observations. The best results were obtained when the model rating was chosen as 2 and the number of fuzzy sets as 5. For CO₂ emissions in Turkey, RMSE value obtained by Chen 2002 method was obtained as 24.86.

Table 3: Chen (2002) Model Results

Year	Greenhouse gas emission	Chen (2002)
	(milyon Ton)	Order=2, number of cluster=5
Test Data	2012	445,6
	2013	439
	2014	451,8
	2015	469,9
	2016	496,1
RMSE		24,86

4.3. Fuzzy time series model based on fuzzy sets:

Fuzzy time series analysis based on fuzzy clustering uses fuzzy cluster analysis methods in the step of converting into fuzzy time series. In this study, analysis was performed using fuzzy time series based on fuzzy c-averages and fuzzy time series models based on Gustafson-Kessel clustering. In both methods, the number of clusters was taken as 5. As a result of the analysis, the RMSE value of the FCMTS (Fuzzy-Cluster Means Time Series) method was obtained as 30.42, and the RMSE value of the GKFTS method was 25.85. The related results are given in Table 3.

Table 4: Fuzzy Time Series Model Based on Fuzzy Sets Results :

	Year	Greenhouse gas emission	FCMTS	GKFTS
		(million tons)		
Test Data	2012	445.6	417,4968	430,0034
	2013	439	417,4968	430,0034
	2014	451,8	417,4968	430,0034
	2015	469,9	417,4968	430,0034
	2016	496,1	417,4968	430,0034
		RMSE	30,42	25,85

5. Conclusions

In this study, greenhouse gas emissions statistics in Turkey was modeled with fuzzy time series analysis. Greenhouse gas emission was modeled using four different Fuzzy ARIMA models, and the performances of the models were evaluated. Expressing which model is the most appropriate in the analyses performed was put forward considering the RMSE value. As a result, modeling the greenhouse gas emission statistics with fuzzy time series analysis as the main goal of the study was achieved, and the applicability of four different fuzzy time series models proposed to the related data set was revealed. Comparing RMSE values, it was seen that Chen 1996 model had the smallest value.

Declarations

Ethical approval: This paper does not contain any studies with human participants or animals performed by any of the authors.

Funding: The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

Conflict of interest: The authors declare that they have no conflict of interest.

Author's Contributions: FC contributed to designed the model, collected dataset, performed the analysis, validated the results, written and reviewed the manuscript.

Data availability: All data analyzed during this study are included in Tables 1.

References

- Abd Rahman NH, Lee MH, Talib ML, Suhartono S (2013) Forecasting of air pollution index with artificial neural network. *Jurnal Teknologi (Sciences and Engineering)*. 63(2):59–64. <https://doi.org/10.11113/jt.v63.1913>
- Abdullah L, Pauzi HM (2015) Methods in forecasting carbon dioxide emissions: A decade review”, *Jurnal Teknologi*, 75(1):67-82. <https://doi.org/10.11113/jt.v75.2603>
- Akın G (2006) Küresel ısınma, nedenleri ve sonuçları. *Ankara Üniversitesi Dil ve Tarih-Coğrafya Fakültesi Dergisi*. 46(2):29-43.
- Appiah K, Du J, Appah R, Quacoe D (2018) Prediction of potential carbon dioxide emissions of selected emerging economies using artificial neural network. *Journal of Environmental Science and Engineering A*, 7: 321-335. DOI:10.17265/2162-5298/2018.08.003
- Atsalakis G, Frantzis D, Zopounidis C (2015) Energy’s exports forecasting by a neuro-fuzzy controller. *Energy Systems*.6(2):249-267. DOI 10.1007/s12667-015-0140-1
- Ayvaz B, Kusakçı AO, Temur GT (2017) Energy-related CO2 emission forecast for Turkey and Europe and Eurasia A discrete grey model approach..*Grey Systems: Theory and Application*. 7(3):437-4532. DOI:10.1080/15567036.2014.978086.
- Balasko B, Abonyi J, Feil B (2005) Fuzzy clustering and data analysis toolbox. Department of Process Engineering University of Veszprem.
- Chen SM (1996) Forecasting enrollments based on fuzzy time-series. *Fuzzy Sets and Systems*. 81(3):311-319. [https://doi.org/10.1016/0165-0114\(95\)00220-0](https://doi.org/10.1016/0165-0114(95)00220-0)
- Chen SM (2000) Using high-order fuzzy time series for handling forecasting problems. *Proceedings of the 11th National Conference on Information Management, Kaohsiung, Taiwan, Republic of China, 2000*.
- Chen SM, Hwang JR (2000) Temperature predication using fuzzy time series. *IEEE Transactions on Systems, Man, and Cybernetics, Part B: Cybernetics* 30(2):263-275. DOI: 10.1109/3477.836375
- Chen SM (2002) Forecasting enrollments based on highorder fuzzy time series. *Cybernetics and Systems: An International Journal*, 33(1): 1-16.<https://doi.org/10.1080/019697202753306479>.
- Dikmen AÇ (2019) Türkiye’de güneş ve rüzgar enerjisinden elektrik üretiminin sera gazı emisyonları ve çevre maliyetinin azalmasına katkıları. *Turkish Studies*.14(2):275-293. <https://doi.org/10.7827/TurkishStudies.15149>

- Doroodi M, Mokhtar A (2019) Comparison of time series approaches for prediction of energy consumption focusing on greenhouse gases emission in Iran. *International Journal of Energy Sector Management*, 13(3):486-499. DOI 10.1108/IJESM-08-2018-0002
- Garip E, Oktay AB (2018) CO2 emisyonunun makine öğrenmesi metotları ile tahmin edilmesi. Conference: 2018 International Conference on Artificial Intelligence and Data Processing (IDAP), 262-266, 28-30 September 2018, Malatya, Turkey.
- Gustafson D, Kessel W (1979) Fuzzy clustering with a fuzzy covariance matrix. *Proc. 18th IEEE CDC Conference*, San Diego, USA, 761-766, December 1979.
- Hwang JR, Chen SM, Lee CH (1998) Handling forecasting problems using fuzzy time series. *Fuzzy Sets and Systems*. 100 (2):217- 228. [https://doi.org/10.1016/S0165-0114\(97\)00121-8](https://doi.org/10.1016/S0165-0114(97)00121-8)
- Karaaslan A, Gezen M (2017) Forecasting of Turkey's sectoral energy demand by using fuzzy grey regression model. *International Journal of Energy Economics and Policy*. 7(1): 67-77.
- Kuşkaya S, Gençoğlu P (2017) OECD ülkelerinin 1995-2015 yılları itibarıyla sera gazı salınımları açısından karşılaştırılması: istatistiksel bir analiz. *International Journal of Disciplines Economics and Administrative Sciences Studies*. 3(3):177-188.
- Liu Y, Tian Y, Chen M. (2017) Research on the prediction of carbon emission based on the chaos theory and neural network. *International Journal of Bioautomation*. 21(4), Special Issue, 339-348.
- Maleki A, Nasseri S, Aminabad MS, Hadi M (2018) Comparison of ARIMA and NNAR models for forecasting water treatment plant's influent characteristics. *KSCE Journal of Civil Engineering*. 22(9):3233-3245. DOI:10.1007/s12205-018-1195-z.
- Oyehan TA, Tawabini BS (2017) Forecasting CO2 emissions in the Persian Gulf States. *Global Journal of Environmental. Science and Management*. 3(1): 1-10, Winter 2017. 10.22034/GJESM.2017.03.01.001
- Özceylan E (2016) Forecasting CO2 Emission of Turkey: Swarm intelligence approaches. *International Journal of Global Warming*, 9(3): 337-61.
- Song Q, Chissom BS (1993a) Forecasting enrollments with fuzzy time series, Part I. *Fuzzy Sets and Systems*. 54(1):1-9. DOI: 10.1016/0165-0114(93)90355-L
- Song Q, Chissom BS (1993b) Fuzzy time series and its models", *Fuzzy Sets and Systems*, 54(3): 269-277. [https://doi.org/10.1016/0165-0114\(93\)90372-0](https://doi.org/10.1016/0165-0114(93)90372-0)
- Song Q, Chissom.BS(1994) Forecasting enrollments with fuzzy time series, Part II.. *Fuzzy Sets and Systems*, 62(1):1- 8. DOI: 10.1016/0165-0114(94)90067-1

- Song Q, Leland.RP (1996) Adaptive learning defuzzification techniques and applications. *Fuzzy Sets and Systems*. 81(3):321-329. DOI:10.1016/0165-0114(95)00219-7
- Song Q, Leland RP, Chissom BS (1995) A new fuzzy time-series model of fuzzy number observations. *Fuzzy Sets and Systems*, 73(3):341- 348. DOI: 10.1016/0165-0114(94)00315-X
- Sullivan J, Woodall WH (1994). A Comparison of fuzzy forecasting and markov modeling. *Fuzzy Sets and Systems* 64(3):279- 293. DOI:10.1016/0165-0114(94)90152-X
- Sutthichaimethee P, Apinyar C.,Suyaprom S (2019) A forecasting model for economic growth and co2 emission based on industry 4.0 political policy under the government power: adapting a second-order autoregressive-sem. *Journal of Open Innovatin Technolgy Market and Complexity*. 5(3): 69-89. doi:10.3390/joitmc5030069
- Tatar V., Özer MB (2018) Sera gazi emisyonlarının iklim değışikliğı üzerindeki etkileri: Türkiye’de mevcut durum analizi. *Journal of Social and Humanities Sciences Research*.5(30):3993-3999. DOI:10.26450/jshsr.805.
- Tavan M (2019) Estimation and forecast of carbon dioxide emission in Iran: Introducing A New Hybrid Modelling. *Journal of Economics and Management*, 36 (2) 144-167. <https://doi.org/10.22367/jem.2019.36.08>
- Wang L, Zhan L.,Rongrong L (2019) Prediction of the energy demand trend in middle africa a comparison of MGM, MECM, ARIMA and BP models. *Sustainability*.11(8):2436. <https://doi.org/10.3390/su11082436>