Patch Selection by Bumble Bees Navigating Discontinuous Landscapes

Fabiana Fragoso

Oak Ridge Institute for Science and Education

Qi Jiang

University of Wisconsin–Madison

Murray Clayton

University of Wisconsin–Madison

Johanne Brunet (✉ Johanne.Brunet@usda.gov)

USDA-Agricultural Research Service

Research Article

Keywords: Bumble bee, discontinuous landscapes, isolation distance, probability model of bee 34 movement, patch attractiveness, patch selection, patch size, resource assessment

Posted Date: December 17th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-126254/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on April 26th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-88394-2>.

Patch selection by bumble bees navigating discontinuous landscapes

Fabiana P. Fragoso¹, Qi Jiang^{2,4}, Murray K. Clayton² and Johanne Brunet^{3*}

¹ Agricultural Research Service Research Participation Program, Oak Ridge Institute for Science and Education, 455 Science Drive, Madison, WI 53711, USA

² Department of Statistics, University of Wisconsin - Madison, 1300 University Ave, Madison, WI 53706, USA

³ United States Department of Agriculture, Agricultural Research Service, Vegetable Crop Research Unit, 455 Science Drive, Madison, WI 53711, USA

⁴ Present address: Goldman Sachs, 200 West Street, New York, NY 10282, USA

*Correspondence to Johanne Brunet (johanne.brunet@usda.gov)

Abstract

Pollen and nectar resources are unevenly distributed over space and bees must make routing decisions when navigating patchy resources. Determining the patch selection process used by bees is crucial to understanding bee foraging over discontinuous landscapes. To elucidate this process, we developed four distinct probability models of bee movement where the size and the distance to the patch determined the attractiveness of a patch. A field experiment with a center patch and four peripheral patches of two distinct sizes and distances from the center was set up in two configurations. Empirical transition probabilities from the center to each peripheral patch were obtained at two sites and two years. The best model was identified by comparing observed and predicted transition probabilities, where predicted values were obtained by incorporating the spatial dimensions of the field experiment into each model's mathematical expression. Bumble bees used both patch size and isolation distance when selecting a patch and could assess the total amount of resources available in a patch. Bumble bees prefer large, nearby patches. This information will facilitate the development of a predictive framework to the study of bee movement and of models that predict the movement of genetically engineered pollen in bee-pollinated crops.

Keywords: Bumble bee, discontinuous landscapes, isolation distance, probability model of bee movement, patch attractiveness, patch selection, patch size, resource assessment

Introduction

Pollen and nectar resources are unevenly distributed over time and space and bees must make many decisions when foraging over the landscape¹. They must select which patch to visit and choose individual plants, inflorescences and flowers within each patch. Their decision process must remain flexible because resource availability varies both spatially and temporally. Bees often prefer to visit nearby plants and plants that offer constant rewards^{2,3}. They use various visual and olfactory cues to select plants⁴⁻⁶ and they also learn to associate floral traits and reward⁷. Bees can switch their preferred flower type, for example a flower of certain color or size, based on reward availability⁸⁻¹⁰. In addition, bees prefer inflorescences with larger floral displays^{4,8,11,12}. They can detect the number of flowers with pollen or nectar on an inflorescence and thus assess resource availability^{8,13}. As bees navigate the landscape to obtain food, their decisions of which plants and inflorescences to visit are strongly influenced by resource availability^{8,13}.

While the process used by bees to select plants, inflorescences and flowers has received a lot of attention, less is known about how bees select patches over the landscape. Bees demonstrate patch fidelity¹⁴⁻¹⁶ and the total number of flowers they visit in a patch during a foraging bout (residence) is affected by patch size¹⁷. Previous research of bumble bee foraging at a spatial scale has been guided by the concept of trapline foraging, where bees optimize their travel route over time¹⁸⁻²⁰. Bumble bees do not simply move to nearest neighbor flowers but exhibit a tradeoff between minimizing travel distance and favoring more rewarding locations²¹⁻²³. However, most experiments on trapline foraging have been done either in the greenhouse or cages using a few (<10) artificial flowers or in the wild using a few feeding stations. In general, all flowers or feeding stations present equal reward to the bees. In a typical experiment, bees

used the shortest route developed in only 21.4% of all foraging bouts and started using this route after an average of 26.6 foraging bouts²³. Under these conditions, bees perform 65-80 potential foraging bouts per day and traplining can develop over a matter of hours¹⁸. Under natural field conditions, however, bees visit patches with many plants and each plant offers a large number of flowers with variable reward. In addition, individual bumble bees in the field make an average of 5.9 foraging trips daily and a foraging trip tends to last over 55 minutes^{14,24}. As a reference point, in a patch of 169 plants, bumble bees were observed visiting, on average, 53.9 flowers in the patch over 3.9 minutes²⁵. Bees are therefore likely to visit many patches, plants and flowers during a foraging trip and would need many days to establish a trapline route, assuming it develops after an average of 27 foraging bouts. Moreover, they would need to remember the location of patches, various plants and numerous flowers over many days. While bees learn to quickly optimize a foraging route under controlled conditions, with few flowers or feeding stations presenting similar reward, it is less clear how the concept of trapline foraging can explain bee foraging within and among patches of plants in the field, where each patch contains many plants, each with many inflorescences and flowers offering variable reward. Given these limitations, feeding stations and artificial flowers used in traplining experiments have been equated to patches over the landscape²⁶. However, this interpretation assumes that the patch selection process is similar to the process used by bees to select individual flowers. Such an assumption must be tested and it is therefore crucial to examine how bees select patches in a landscape with patchily distributed resources.

To fill this critical gap in our knowledge of bee foraging behavior, we developed four models of patch attractiveness that differed in how the probability of a bee moving to a patch was influenced by the size of a patch (resources available) and the isolation distance between

patches (Fig. 1a). Resource availability influences bee choice of individual plants and inflorescences^{8,13} and minimizing traveling distances over a foraging trip is central to trapline foraging although reward also plays a role^{20,22,23,26}. The models also differed in whether bees could estimate resources fully or only partially. Many animals use encounter rate with preys as a measure of resource availability and cannot estimate total resources available²⁷. The ability to estimate or not estimate resource availability represents an important assumption of optimal foraging models²⁸.

To determine which model best described the process by which bumble bees select patches, we set up a field experiment consisting of a center patch and four peripheral patches of two distinct sizes and located at two distances from the center patch (Fig. 1b). The experiment was set up in two configurations with similar size patches on the same or opposite side of the center patch (Fig. 2a, b). Bumble bee hives were placed along one edge of the center patch (Fig. 2c) and empirical transition probabilities from the center to each peripheral patch were obtained over two years. Predicted transition probabilities from the center to each of the four peripheral patches were obtained for each of the four models by incorporating the spatial dimensions of a field experiment into the mathematical expression for each model. The best model was selected by comparing observed transition probabilities to the probabilities predicted by each of the four models. By answering how distance and resource availability affect patch selection, and whether bumble bees can estimate resources fully or partially, this study advances our knowledge of bee foraging behavior over patchily distributed resources and contributes to linking behavioral ecology to landscape ecology^{29,30}. In addition, by elucidating the rules bees used to select patches, this study will facilitate the development of models to predict the movement of genetically engineered pollen in bee-pollinated crops³¹.

Results

Predicted values from each model

The first model predicted bees would preferentially move to the large near (LN) patch, followed by small near (SN), large far (LF) and small far (SF) patches (Fig. 1c). A similar trend was predicted for Model II, with smaller probabilities of moving to LN and SN patches and greater probabilities of movement to SF and LF patches, relative to Model I. Model III predicted a greater probability of movement to smaller relative to larger patches. Lastly, Model IV predicted an equal probability of movement to any of the four peripheral patches (Fig. 1c).

Observed bee transitions

In total, we recorded 224 transitions from the center patch to one of the four peripheral patches. In 2017, we observed 52 transitions at the site with ‘opposite side’ configuration (Fig. 2a) and 67 transitions at the site with the “same side” configuration (Fig. 2b). In 2018, these numbers were, respectively, 54 and 51 transitions. Observed transition probabilities from the center patch to each of the four peripheral patches (LN, SN, LF and SF) were calculated from these transition data over both years, per year and separately for each site and year (Fig. 3; Table 1).

Selection of the best model

Model I best explained the observed transition data in all cases examined (Table 2). This means bees are more likely to move to large near patches and can estimate the total amount of resources in a patch. For site M600, Model II could not be rejected in 2017 and Model III in 2018 (Table 2). As discussed in the Methods section for selecting the best model, an outlier was detected for

M600 in 2018, and floral density was uneven among peripheral patches at that site that year, with greater density in the small near patch. The fact that a second model, besides Model I, could not be rejected at site M600 each year, suggests an impact of patch configuration on bee movement. However, because Model I consistently explained the observed transition patterns (Table 2), we conclude that bumble bees use information on both distance and total resources available when selecting patches.

Discussion

Bumble bees consider both patch size and isolation distance when selecting a patch and prefer larger and nearby patches. Visiting large patches provides more resources to bumble bees and flying to the nearest patch minimizes flight energy expenditure. These decisions are likely to optimize the bees net energy intake as optimal foraging theory would predict^{28,32,33}. Unlike many animals that use encounter rate with preys as a measure of patch quality and cannot estimate the total resources available²⁷, bumble bees possess such an ability. They can estimate the total amount of resources available in the patch. Determining the patch selection process used by bumble bees improves our understanding of bee movement and helps translate the study of such movement into a predictive framework³². For example, an individual based model describing movement in a continuous landscape, an approach used by Levey and collaborators^{34,35} to examine the effect of corridors on seed dispersal in the eastern blue bird (*Sialia sialis*), can be used to describe bee movement within patches²⁵ and be combined with the patch selection process to describe and predict bee movement in different landscapes³¹. There is major interest in modelling the potential spread of pollen from genetically engineered crops into conventional crops³⁶ and into populations of cross-compatible weedy or wild relatives³⁷. The basic

understanding of bee movement patterns provided by this study will facilitate the development of such models for species pollinated by bees foraging in discontinuous landscapes. The impending release of new varieties produced via gene editing technologies³⁸ for various insect-pollinated crops and the new USDA's plant biotechnology regulations 'SECURE' only heighten the criticality of this already imperative issue.

The question remains how bees can estimate distances and the level of resources in a patch. Bees use optic flow – or the retinal image motion experienced during travel – to estimate distance^{39,40}. They rely on path integration to keep track of homeward directions by using a sun and polarized-light compass when assessing direction of travel^{41,42} and they can also make use of prominent landmarks to pinpoint their destinations⁴³. All these abilities are presumably being employed by bees when estimating distances between patches. Little is known about how bees evaluate total resources available in a patch. The local discontinuity in apparent motion across boundaries allows for flying bees to detect a figure raised from the ground⁴⁴. Pattern and color discrimination, as well as shape detection⁴⁵ would allow bees to evaluate the patch they are heading to. Various visual and potentially olfactory cues⁴⁻⁶ are likely being computed by bees to determine the attractiveness of the patch. More work is needed to determine how bees estimate the total amount of resources in a patch but our results indicate that they do not passively visit large patches because they are more apparent, as has been previously suggested⁴⁶.

Landscape configuration, similarly to spatial configuration at smaller scales, can affect bee behavior. Bumble bees have more difficulty making decisions with some spatial configurations than others^{18,47} and our findings indicate that landscape configuration may also influence bee behavior. Bees exhibited stronger responses with the opposite side configuration, where only Model I was a good fit to the data. However, the fact that Model 1 represented a

good fit to the data for both field configurations suggest that the patterns of transitions observed are not the results of environmental factors such as wind direction⁴⁸. In addition, we observed variation in the response of bees as some bees did not move to the larger, nearby patch. However, we did not mark individual bees in this experiment and could not follow their decision process over time and thus with experience. A bee may in fact learn the optimal decision with learning and experience⁴⁹ and some bees may learn faster than others reflecting inter-individual differences in how bees handle cognitive tasks and make decisions^{50,51}. Future patch selection experiments, marking individual bees in a similar experimental design, would help further clarify the role of learning and experience and individual cognitive differences in the decision-making process of bumble bees selecting patches in a discontinuous landscape.

This study highlights various similarities between the decision-making process bees used to select patches and the one they use to choose inflorescences. When bumble bees visit inflorescences, they prefer inflorescences with larger floral displays^{4,11,12} and base their decision on the number of pollen- or nectar-producing flowers^{8,13}. They can estimate the total resources available on inflorescences. Similarly, when visiting patches of plants, bumble bees can estimate the total amount of resources available and prefer patches with more resources. Therefore, bumble bees use a similar criterion, total resources available, when selecting the next inflorescence or the next patch they move to. Moreover, when foraging on inflorescences, bumble bees visit more flowers on larger inflorescences but visit a smaller proportion of the flowers^{12,52 12,52}. Bumble bees exhibit a similar behavior when foraging on patches of plants where they visit more flowers but a smaller proportion of the flowers in larger patches¹⁷. With respect to distance, bumble bees prefer nearby patches and seem to prefer nearby plants² although they do not necessarily select nearest neighboring flowers when establishing an optimal

trapline²³. More work is needed to determine the impact of distance and the combined impact of distance and resource availability on inflorescence selection. However, currently available data highlight similarities between the decision-making process of bumble bees when selecting inflorescences and patches of plants.

Conclusions

Bees consider both distance and patch size and can estimate total resources when selecting patches over the landscape. Moreover, bumble bees follow many similar rules when moving between patches and when selecting inflorescences. This study expands our understanding of bee foraging over patchy resources and discontinuous landscapes and contributes to a more predictive framework to the study of bee movement. For example, results of this study can facilitate the development of models of the spread of genetically modified pollen in bee-pollinated crops.

Methods

Models of patch selection

We contrast four distinct probability models of bee movement between patches which differ in theories of patch attractiveness. In particular, we assume that the probability of a bee moving to one patch – when given the choice of moving to one of several patches – can be described in terms of the size of and the distance to the patch, and that these two factors determine the attractiveness of the patch (Fig. 1a). For mathematical simplicity (and tractability) we assume

that a given patch can be represented as a disc of radius R whose center is a distance d from the pollinator. Moreover, we assume that d is much larger than R .

Our first two models are the most complex and borrow notions from physics in their construction. In *Model I*, we assume that each point in the patch emits an attractant, whose effect is inversely proportional to the square of the distance from the point to the bee. Suppose that (x, y) is a point lying on the disc, a distance r from the disc center. Then it can be shown that the total attractiveness is given by

$$\iint_P 1/(x^2 + y^2 + d^2) \, dx dy$$

where P is the set of points (x, y) on the disc. By converting to polar coordinates and solving the resultant integral, we see that the total attractiveness is given by

$$2\pi(\ln\left(1 + \left(\frac{R}{d}\right)^2\right))$$

Because d is assumed to be much larger than R , Taylor's expansion gives the total attractiveness to be proportional to $2\pi(\frac{R}{d})^2$, or equivalently, the disc Area divided by d^2 . This model incorporates both patch size and isolation distance. It illustrates a case where a pollinator can integrate the area of the patch and determine the total amount of resources available in the patch.

In *Model II*, we assume that the pollinator views the patch (*i.e.* disc) edge-on, and thus only perceives its diameter, and not its area. In that case, the total attractiveness of the patch can be approximated by $\arctan(\frac{R}{d})$ which, by a Taylor series approximation, can be shown to be proportional to the diameter over distance or $2R/d$. In this model, a pollinator can estimate distance but the probability of choosing a patch is a function of its diameter rather than its area, and therefore the pollinator only obtains partial information about the total resources available.

The next two models are simpler in construction. In *Model III*, we assume that the attractiveness of a patch depends solely on its distance, with the attractiveness being inversely proportional to the squared distance between the pollinator and the center of the patch. This is the closest approximation to a nearest neighbor model except that we are using a more flexible probabilistic approach than the pure nearest neighbor model. Finally, in *Model IV*, we assume that all patches are equally attractive to pollinators, without regard to their size or distance. This model represents a purely random patch choice model.

Experimental design

We set up an experiment at different sites at the West Madison Agricultural Research Station, WI, USA. The experiment consisted of five patches of alfalfa, a center and two peripheral 9.14m x 9.14m patches containing 100 plants (small patches) and two peripheral 13.7m x 13.7m patches with 225 plants (large patches). Plants were originally grown from seeds in the greenhouse and transplanted 90 cm apart from each other within each patch. One of the large and one of the small patches were placed 9.14m diagonally from the center patch (near) while the other large and small patches were placed 18.3m away (far). This arrangement ensured the presence of four types of peripheral patches: large near (LN), small near (SN), large far (LF) and small far (SF) (Fig. 1b). The spatial configuration of the peripheral patches differed between some sites, with the “opposite side” configuration having same size peripheral patches on opposite sides of the center patch (Fig. 2a) while the “same side” configuration had the two large or two small patches on the same side of the center patch (Fig. 2b). The type of configuration depended on whether similar-size peripheral patches were situated on opposite sides or on the same side of the center patch. The use of different configurations permitted generalization of the

results because it minimized potential bias resulting from environmental factors. Tall fescue grass was grown for weed control among patches and among plants within patches and the grass was mowed periodically. At each site, the five alfalfa patches for the experiment were set up within a larger alfalfa hay field which was harvested at least three times throughout the summer prior to flowering. Each year, beehives of the common eastern bumble bee *Bombus impatiens* Cresson, an effective pollinator of alfalfa⁵³, were set up along the center patch at each site at least one week prior to the beginning of bee observations (Fig. 2c). Observations took place throughout the day, when bees were active.

Predictions from each of the four models

As noted above, we assume in the mathematical construction that patches are discs to facilitate the necessary calculus. To create a landscape configuration involving patches of different sizes and isolation distances and generate predicted patterns of movement to be tested empirically, we use square patches because these are easier to implement in the field. The mathematically derived attractiveness of a patch above is therefore an approximation to the true attractiveness in the field experiment.

Predictions for the probability of a bee transitioning from the center patch to each of the four conventional patches were made by assigning the dimensions of the field experiment to the variables in the respective equation for each of the four models. Consider our landscape configuration as given in Figure 2. The areas of the large patches are $(13.7\text{ m})^2$ and the areas of the small patches are $(9.1\text{ m})^2$; the near and far distances between patches are 9.1 m and 18.3 m, respectively, and the distances from the corners of the patches to their centers are 6.5 m for small patches and 9.7 m for large patches. This allows us to calculate the total distance from the edge

of the center patch to the center of each peripheral patch. For the patches abbreviated as LN, SN, LF, and SF for large near, small near, large far and small far, these distances are: 18.8, 15.6, 28.0, and 24.8m, respectively. Thus, for Model I, the probabilities of movement to the four patches LN, SN, LF, and SF are proportional to $187.69/18.8^2$, $82.81/15.6^2$, $187.69/28.0^2$, and $82.81/24.8^2$, respectively. We therefore calculate those four quantities and multiply by a proportionality constant to ensure that they add to one (because they represent probabilities). Similarly, for the different attractiveness models, we obtained the probabilities of movement to each of the four peripheral patches (Fig. 1c).

Observed bee transitions

We collected bee transition data over two summers and at two sites each summer. Transition data were collected throughout the day, when bees were active, during the weeks of alfalfa peak bloom. During periods of bee observations, at least two trained observers were positioned on opposite sides of the center patch and followed bumblebees foraging in the patch, especially along the edges. A transition was recorded when a bumblebee was observed to leave the center patch and moved to a peripheral patch and foraged on flowers in that new patch. During the summer of 2017, transition data were collected from site B400 with the ‘opposite side configuration’ and site M600 with the ‘same side’ configuration. In summer 2018, transition data were collected from site M600 and from a new site, M1000, with the opposite side configuration. Site B400 did not have enough plants that survived the winter. Peak flowering at the two sites each year were separated by approximately two weeks by mowing the plants at different times in late spring-early summer. Transition data were collected in July for site M600 in 2017 and site M1000 in 2018 and during August for site B400 in 2017 and site M600 in 2018. Observed

transition probabilities from the center patch to each of the four peripheral patches were calculated over both years, per year and separately for each site and year.

Selecting the best model

To identify the model that best explained the empirical data, observed transition probabilities were compared to the predictions derived for each of the four models using chi-square goodness of fit tests. For these tests, probability values of $P \leq 0.05$ indicate that a model is not a good fit to the empirical data. We contrasted observed and predicted transition probabilities over all empirical data (combining sites and years), per year and for each site within each year. All analyses were performed using R 3.5.1⁵⁴.

In 2018, patch SN at site M600 had higher floral density relative to the other patches and transition counts were also high. Before performing further analyses for that site and year we first examined whether that patch should be considered an outlier. For the outlier test, a Poisson (generalized linear) regression model was fit whereby we assumed that the transition counts follow a Poisson distribution, and that the log mean of the Poisson distribution could be modeled by a linear combination of “distance” (near or far), “size” (small or large), “year”, “site”, and an indicator variable coding for the observation in question⁵⁵. The resulting P-value corresponding to the indicator variable was 0.0016. Multiplying that P-value by 4, we had the Bonferroni corrected P-value of 0.0064 for testing for an outlier for the M600/2018 site. Given the outlier indicator was statistically significant under these conditions, and given that this patch had a different floral density, we concluded that the high number ($n=26$) of transitions for the small near patch for site M600 in 2018 could reasonably be considered an outlier. Therefore, to

perform the test for site M600 in year 2018 the outlier, patch SN, was excluded and the remaining transition probabilities were scaled up to ensure that they still added to 1.

Data availability

The data supporting the findings of this study are available within this paper.

References

1. Becher, M. A. *et al.* BEESCOUT: A model of bee scouting behaviour and a software tool for characterizing nectar/pollen landscapes for BEEHAVE. *Ecol. Modell.* **340**, 126–133 (2016).
2. Marden, J. H. & Waddington, K. D. Floral choices by honeybees in relation to the relative distances to flowers. *Physiol. Entomol.* **6**, 431–435 (1981).
3. Waddington, K. D., Allen, T. & Heinrich, B. Floral preferences of bumblebees (*Bombus edwardsii*) in relation to intermittent versus continuous rewards. *Anim. Behav.* **29**, 779–784 (1981).
4. Bauer, A. A., Clayton, M. K. & Brunet, J. Floral traits influencing plant attractiveness to three bee species: consequences for plant reproductive success. *Am. J. Bot.* **104**, 1–10 (2017).
5. Bradshaw, H. D. & Schemske, D. W. Allele substitution at a flower colour locus produces a pollinator shift in monkeyflowers. *Nature* **426**, 176–178 (2003).
6. Klahre, U. *et al.* Pollinator choice in petunia depends on two major genetic loci for floral scent production. *Curr. Biol.* **21**, 730–739 (2011).
7. Muth, F., Papaj, D. R. & Leonard, A. S. Bees remember flowers for more than one reason: Pollen mediates associative learning. *Anim. Behav.* **111**, 93–100 (2016).
8. Brunet, J., Thairu, M. W., Henss, J. M., Link, R. I. & Kluever, J. A. The effects of flower, floral display, and reward sizes on bumblebee foraging behavior when pollen is the reward and plants are dichogamous. *Int. J. Plant Sci.* **176**, 811–819 (2015).
9. Nicholls, E. & De Ibarra, N. H. Bees associate colour cues with differences in pollen rewards. *J. Exp. Biol.* **217**, 2783–2788 (2014).
10. Thairu, M. W. & Brunet, J. The role of pollinators in maintaining variation in flower

- 361 colour in the Rocky Mountain columbine, *Aquilegia coerulea*. *Ann. Bot.* **115**, 971–979
362 (2015).
- 363 11. Ishii, H. S. Floral display size influences subsequent plant choice by bumble bees. *Funct.*
364 *Ecol.* **20**, 233–238 (2006).
- 365 12. Mitchell, R. J., Karron, J. D., Holmquist, K. G. & Bell, J. M. The influence of *Mimulus*
366 *ringens* floral display size on pollinator visitation patterns. *Funct. Ecol.* **18**, 116–124
367 (2004).
- 368 13. Makino, T. T. & Sakai, S. Experience changes pollinator responses to floral display size:
369 from size-based to reward-based foraging. *Funct. Ecol.* **21**, 854–863 (2007).
- 370 14. Osborne, J. L. *et al.* A landscape-scale study of bumble bee foraging range and constancy,
371 using harmonic radar. *J. Appl. Ecol.* **36**, 519–533 (1999).
- 372 15. Osborne, J. L. & Williams, I. H. Site constancy of bumble bees in an experimentally
373 patchy habitat. *Agric. Ecosyst. Environ.* **83**, 129–141 (2001).
- 374 16. Saville, N. M., Dramstad, W. E., Fry, G. L. A. & Corbet, S. A. Bumblebee movement in a
375 fragmented agricultural landscape. *Agric. Ecosyst. Environ.* **61**, 145–154 (1997).
- 376 17. Cresswell, J. E. & Osborne, J. L. The effect of patch size and separation on bumblebee
377 foraging in oilseed rape: implications for gene flow. *J. Appl. Ecol.* **41**, 539–546 (2004).
- 378 18. Ohashi, K. & Thomson, J. D. Trapline foraging by pollinators: its ontogeny, economics
379 and possible consequences for plants. *Ann. Bot.* **103**, 1365–1378 (2009).
- 380 19. Saleh, N. & Chittka, L. Traplining in bumblebees (*Bombus impatiens*): A foraging
381 strategy's ontogeny and the importance of spatial reference memory in short-range
382 foraging. *Oecologia* **151**, 719–730 (2007).
- 383 20. Woodgate, J. L., Makinson, J. C., Lim, K. S., Reynolds, A. M. & Chittka, L. Continuous
384 radar tracking illustrates the development of multi-destination routes of bumblebees. *Sci.*
385 *Rep.* **7**, 17323 (2017).
- 386 21. Lihoreau, M., Chittka, L. & Raine, N. E. Trade-off between travel distance and
387 prioritization of high-reward sites in traplining bumblebees. *Funct. Ecol.* **25**, 1284–1292
388 (2011).
- 389 22. Lihoreau, M., Chittka, L. & Raine, N. E. Travel optimization by foraging bumblebees
390 through readjustments of traplines after discovery of new feeding locations. *Am. Nat.* **176**,
391 744–757 (2010).
- 392 23. Lihoreau, M., Chittka, L., Le Comber, S. C. & Raine, N. E. Bees do not use nearest-
393 neighbour rules for optimization of multi-location routes. *Biol. Lett.* **8**, 13–16 (2012).
- 394 24. Minahan, D. F. & Brunet, J. Strong interspecific differences in foraging activity observed
395 between honey bees and bumble bees using miniaturized radio frequency identification

(RFID). *Front. Ecol. Evol.* **6**, (2018).

25. Brunet, J., Zhao, Y. & Clayton, M. K. Linking the foraging behavior of three bee species to pollen dispersal and gene flow. *PLoS One* **14**, e0212561 (2019).

26. Reynolds, A. M., Lihoreau, M. & Chittka, L. A simple iterative model accurately captures complex trapline formation by bumblebees across spatial scales and flower arrangements. *PLoS Comput. Biol.* **9**, e1002938 (2013).

27. Marschall, E. A., Chesson, P. L. & Stein, R. A. Foraging in a patchy environment: prey-encounter rate and residence time distributions. *Anim. Behav.* **37**, 444–454 (1989).

28. Pyke, G. H. Optimal foraging theory : a critical review. *Ann. Rev. Ecol. Syst.* **15**, 523–575 (1984).

29. Rands, S. A. Landscape fragmentation and pollinator movement within agricultural environments : a modelling framework for exploring foraging and movement ecology. *PeerJ* **e269**, (2014).

30. Lima, S. L. & Zollner, P. A. Towards a behavioral ecology of ecological landscapes. *Trends Ecol. Evol.* **11**, 131–135 (1996).

31. Brunet, J. A conceptual framework that links pollinator foraging behavior to gene flow. *Proceedings for the 2018 WASGA Winter Seed Conference* 63–67 (2018).

32. Pyke, G. H. Understanding movements of organisms: It’s time to abandon the Lévy foraging hypothesis. *Methods Ecol. Evol.* **6**, 1–16 (2015).

33. Macarthur, R. H. & Pianka, E. R. On optimal use of a patchy environment. *Am. Nat.* **100**, 603–609 (1966).

34. Levey, D. J., Bolker, B. M., Tewksbury, J. J., Sargent, S. & Haddad, N. M. Effects of Landscape Corridors on Seed Dispersal by Birds. *Science*. **309**, (2005).

35. Levey, D. J., Tewksbury, J. J. & Bolker, B. M. Modelling long-distance seed dispersal in heterogeneous landscapes. *J. Ecol.* **96**, 599–608 (2008).

36. Smith, K. & Spangenberg, G. Considerations for managing agricultural co-existence between transgenic and non-transgenic cultivars of outcrossing perennial forage plants in dairy pastures. *Agronomy* **6**, 59–68 (2016).

37. Ellstrand, N. C. *et al.* Introgression of crop alleles into wild or weedy populations. *Annu. Rev. Ecol. Evol. Syst.* **44**, 325–345 (2013).

38. Gupta, R. M. & Musunuru, K. Expanding the genetic editing tool kit: ZFNs, TALENs, and CRISPR-Cas9. *J. Clin. Invest.* **124**, 4154–4161 (2014).

39. Esch, H. E. & Burns, J. E. Distance estimation by foraging honeybees. *J. Exp. Biol.* **199**, 155–162 (1996).

- 430 40. Srinivasan, M. V., Zhang, S., Altwein, M. & Tautz, J. Honeybee navigation: nature and
431 calibration of the 'odometer'. *Science*. **287**, 851–853 (2000).
- 432 41. Collett, M. & Collett, T. S. How do insects use path integration for their navigation? *Biol.*
433 *Cybern.* **83**, 245–259 (2000).
- 434 42. Collett, M., Chittka, L. & Collett, T. S. Spatial memory in insect navigation. *Curr. Biol.*
435 **23**, R789–800 (2013).
- 436 43. Chittka, L., Geiger, K. & Kunze, J. The influences of landmarks on distance estimation of
437 honey bees. *Anim. Behav.* **50**, 23–31 (1995).
- 438 44. Srinivasan, M. V., Lehrer, M. & Horridge, G. A. Visual figure-ground discrimination in
439 the honeybee: the role of motion parallax at boundaries. *Proc. R. Soc. B Biol. Sci.* **238**,
440 331–350 (1990).
- 441 45. Lehrer, M. Looking all around: honeybees use different cues in different eye regions. *J.*
442 *Exp. Biol.* **201**, 3275–3292 (1998).
- 443 46. Goulson, D. Foraging strategies of insects for gathering nectar and pollen, and
444 implications for plant ecology and evolution. *Perspect. Plant Ecol. Evol. Syst.* **2**, 185–209
445 (1999).
- 446 47. Ohashi, K., Thomson, J. D. & D'Souza, D. Trapline foraging by bumble bees: IV.
447 Optimization of route geometry in the absence of competition. *Behav. Ecol.* **18**, 1–11
448 (2007).
- 449 48. Comba, L. Patch use by bumblebees (hymenoptera apidae): Temperature, wind, flower
450 density and traplining. *Ethol. Ecol. Evol.* **11**, 243–264 (1999).
- 451 49. Ohashi, K., Leslie, A. & Thomson, J. D. Trapline foraging by bumble bees: V. Effects of
452 experience and priority on competitive performance. *Behav. Ecol.* **19**, 936–948 (2008).
- 453 50. Klein, S., Pasquaretta, C., Barron, A. B., Devaud, J. M. & Lihoreau, M. Inter-individual
454 variability in the foraging behaviour of traplining bumblebees. *Sci. Rep.* **7**, 4561 (2017).
- 455 51. Chittka, L. Bee cognition. *Curr. Biol.* **27**, R1049–R1053 (2017).
- 456 52. Ohashi, K. & Yahara, T. Visit larger displays but probe proportionally fewer flowers:
457 Counterintuitive behaviour of nectar-collecting bumble bees achieves an ideal free
458 distribution. *Funct. Ecol.* **16**, 492–503 (2002).
- 459 53. Brunet, J. & Stewart, C. M. Impact of bee species and plant density on alfalfa pollination
460 and potential for gene flow. *Psyche A J. Entomol.* **2010**, 1–7 (2010).
- 461 54. R Core Team. *R: A language and environment for statistical computing*. (R Foundation
462 for Statistical Computing, 2019).
- 463 55. Weisberg, S. *Applied Linear Regression*. (John Wiley & Sons, 2013).

Acknowledgments

Various undergraduate and graduate students and technicians helped gather field data. Ethan Fox performed the Chi-square tests. This project was supported by a Biotechnology Risk Assessment Grant Program, competitive grant no. 2018-33522-28707 from the USDA National Institute of Food and Agriculture to J. Brunet and by funds from the USDA Agricultural Research Service to J. Brunet. F. P Fragoso was supported in part by an appointment to the Agricultural Research Service (ARS) Research Participation Program administered by the Oak Ridge Institute for Science and Education (ORISE) through an interagency agreement between the U.S. Department of Energy (DOE) and the U.S. Department of Agriculture (USDA). ORISE is managed by ORAU under DOE contract number DE-SC0014664. All opinions expressed in this paper are the author's and do not necessarily reflect the policies and views of USDA, ARS, DOE, or ORAU/ORISE.

Author Contributions

J. B. and M.K.C. conceived the study and designed the experiments. Q.J. and M.K.C. developed the patch attractiveness models and generated predictions. J.B. and F.P.F. supervised field data collection. J.B., M.K.C and F.P.F analyzed the data. F.P.F. and J.B. wrote the original draft and M.K.C provided comments. J. B. supervised the entire project.

Competing interests

The authors declare no competing interests.

487

488

489 **Figure 1.** Models of patch attractiveness and derived predictions. a. The parameters of the four
 490 models of patch attractiveness. b. Parameters of the four models of patch attractiveness
 491 illustrated within the experimental design. c. Predicted probabilities of transitions from the center
 492 patch to each of the four peripheral patches derived for each of the four models.

493

494

495

496

505

506 **Figure 2.** Configurations of the experiment and bee transitions. A center patch surrounded by
 507 four peripheral patches of two sizes and at two distances from the center were set up at different
 508 sites in two configurations, a. an “opposite side” or b. a “same side” configuration, referring to
 509 the location of patches of similar sizes relative to the center patch. c. Bumblebee hives were
 510 positioned facing east along the center patch (dark grey) and bee transitions from the center to
 511 the peripheral patches (light grey) were recorded over two summers.

512

513

514

515

521

528

529

530

531

532

533

534

535

536

537

Figure 3. Observed and predicted transition probabilities. For both years combined (all), for 2017, and for 2018, observed probabilities of moving to large near (LN), small near (SN), large far (LF), and small far (SF) peripheral patches together with probabilities predicted by each of the four models of patch attractiveness (I, II, III and IV). A non-significant (ns) model indicates a good fit to the data. A * refers to a model with a probability $P < 0.05$. Model I, where bees estimate distance and total resources available, best fit the experimental data.

Table 1. Observed transition probabilities for each site each year. Transition probabilities from the center patch to peripheral patches of two sizes, located at two distances from the center patch with LN= large near, SN = small near, LF= large far and SF = small far. N is the total number of observed transitions for the site and year.

year	site	N	LN	SN	LF	SF
2017	M600	67	0.463	0.269	0.134	0.134
2017	B400	52	0.481	0.308	0.115	0.096
2018	M1000	54	0.500	0.148	0.222	0.130
2018	M600	51	0.333	0.510	0.039	0.118

Table 2. Selection of the best model. Observed transition probabilities were compared against predicted values for each of the four models of patch attractiveness (defined in the text). The Chi-square value and associated probability, P value, are presented here. A model with a probability > 0.05 (bolded) indicates a good fit to the empirical data.

	Model I		Model II		Model III		Model IV	
	χ^2	P	χ^2	P	χ^2	P	χ^2	P
Both years	5.85	0.12	21.47	< 0.0001	29.54	< 0.0001	65.17	< 0.0001
2017	3.39	0.33	13.65	0.003	20.59	0.0001	39.42	< 0.0001
2018	3.13	0.37	8.58	0.035	9.69	0.02	26.69	< 0.0001
2017 M600	1.79	0.62	6.16	0.104	11.17	0.01	19.38	0.0002
2017 B400	2.24	0.52	7.98	0.046	9.89	0.02	20.46	0.0001
2018 M600	4.98	0.083	6.89	0.032	4.23	0.120	14.48	0.0007
2018 M1000	4.24	0.236	7.75	0.051	21.88	< 0.0001	19.03	0.0002

Figures

Figure 1

Models of patch attractiveness and derived predictions. a. The parameters of the four models of patch attractiveness. b. Parameters of the four models of patch attractiveness illustrated within the experimental design. c. Predicted probabilities of transitions from the center patch to each of the four peripheral patches derived for each of the four models.

Figure 2

Configurations of the experiment and bee transitions. A center patch surrounded by four peripheral patches of two sizes and at two distances from the center were set up at different sites in two configurations, a. an “opposite side” or b. a “same side” configuration, referring to the location of patches of similar sizes relative to the center patch. c. Bumblebee hives were positioned facing east along the center patch (dark grey) and bee transitions from the center to the peripheral patches (light grey) were recorded over two summers.

Figure 3

Observed and predicted transition probabilities. For both years combined (all), for 2017, and for 2018, observed probabilities of moving to large near (LN), small near (SN), large far (LF), and small far (SF) peripheral patches together with probabilities predicted by each of the four models of patch attractiveness (I, II, III and IV). A non-significant (ns) model indicates a good fit to the data. A * refers to a model with a probability $P < 0.05$. Model I, where bees estimate distance and total resources available, best fit the experimental data.