Re-Irradiation for Recurrent Brain Tumors: A Retrospective Study from a Tertiary Hospital in Saudi Arabia

HAFIZ ASIF IQBAL (✉ asifzunu@gmail.com)

King Faisal Specialist Hospital and Research Center <https://orcid.org/0000-0001-7082-5700>

Rolina K Alwassia

King Abdulaziz University Hospital

Zaheeda Mulla

King Faisal Specialist Hospital and Research Center

Ahmed Abdelkhalek Hussein

King Faisal Specialist Hospital and Research Center

Hane Mohammad Muamenah

King Faisal Specialist Hospital and Research Center

Research Article

Keywords: Re-irradiation, primary brain tumor, glioblastoma, glioma

Posted Date: January 24th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1265731/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective

To analyze the post-re-RT progression-free survival (PFS) and incidence of radio-necrosis (BRN) in patients with recurrent primary brain tumors and to explore the associated factors.

Method

A retrospective cohort study that included 15 pediatric and adult patients with primary brain tumors who were treated with re-RT between 2011 and 2020. The study endpoints included the post-re-RT PFS, which were analyzed using Kaplan-Meier survival analysis, and the incidence of radio-necrosis. Baseline demographic and clinical data, primary radiation therapy (RT1) parameters and outcomes, and re-RT parameters and outcomes, were analyzed as factors for the two outcomes.

Result

Of the 15 participants, 7 had glioblastoma and 5 had anaplastic ependymoma. The mean interval from first RT to re-RT was 24 months (range=2 – 60 months). The mean total cumulative dose after re-RT as per EQD2 (equivalent dose in 2 Gy) fractions was 101.97 Gy (max 135.6 Gy). The total mean (max) cumulative doses for organs at risk as per EQD2 after re-RT were 54.05 (92.93) Gy for brain stem, 41.19 (87.94) Gy for optic chiasma, and 28.79 (77.18) Gy and 28.6 (88.71) Gy for left and right optic nerves respectively. Disease progression occurred in 10/15 patients, and the median PFS was 4 months (95%CI=0 – 9.1). Although not statistically significant, PFS was likely to be prolonged in case of low-grade tumors, longer RT1 – re-RT time. Radiation necrosis occurred in 2 patients.

Conclusion

The expected clinical benefits against the adverse effects should be contemplated for re-irradiation in primary brain tumors.

Introduction

Primary brain tumors constitute a heterogenous group of benign and malignant tumors that develop in the brain structures including parenchyma and annex tissues. They are characterized by a high mortality and poor functional outcome, resulting into a substantial disease burden, and high propensity for recurrence.^{1–3} The worldwide incidence of primary brain cancers was estimated as 10.82 per 100 000 person-years, with a significant age-disparity depending on the tumor type and histology.⁴

Recent progress in molecular biology, imaging, surgery and radiation therapy has enabled advanced understanding and management of primary brain tumors, especially gliomas and primary central nervous system lymphomas that represent the most common pathological types. Nevertheless, the standard treatment remains relatively aggressive, including surgical resection followed by adjuvant radiotherapy (RT), using either conformal intensity-modulated radiation therapy (IMRT) or stereotactic radiotherapy (SRS), and in some cases adjuvant chemotherapy (CT).^{3,5-8} Furthermore, in refractory or relapsing cases a re-irradiation (re-RT) and or re-resection can be proposed, entailing concerns regarding the risk of radio-necrosis that may occur several months to several years following re-RT.⁹⁻¹³

Currently, there is no standard of care, notably dose regimens, regarding re-RT of brain tumors and the prospective data addressing this approach are scarce⁹. The safety of re-RT approach is limited by the capacity of brain recovery after radiation, which depends on the initial biologically effective dose (BED) as well as on the time interval between the primary radiation (RT1) and re-RT (RT1–re-RT interval).¹¹ Consequently, a low RT1 dose and an increased RT1–re-RT interval are supposedly favorable for a re-RT indication¹¹; however, some data have shown no correlation between RT1–re-RT interval and brain tolerance to re-RT.^{9,10} On the other hand, recent advances in RT technique, such as IMRT and SRS, allowed reduction of the treatment volume, thereby sparing normal tissue and reducing the proportional risk of radiation toxicity.^{9,10,14}

In our center, which is a referral center in radiation oncology, a selected number of patients with recurrent primary brain tumor have benefitted from re-RT over the last years. This study aims at analyzing the safety and efficacy of re-RT among this cohort of patients, by estimating the incidence risk of brain radio-necrosis (BRN) and the progression-free survival (PFS), post re-RT, and analyzing the associated treatment parameters, notably the cumulative radiation doses for brain and organs at risk. Such data would provide a more nuanced insight into the clinical benefit of re-RT for brain tumors and enable determine eventual irradiation dose thresholds and toxicity profiles, which may give direction for future randomized trials. Additionally, data from this study would supply further systematic reviews and meta-analyses.

Methods

Design & Setting

A retrospective cohort study was conducted at the Radiotherapy Unit of XXX in XXX, XXX, between 2011 and 2020. The study received ethical approval of the institutional review board of XXX.

Participants

Fifteen patients were identified with primary brain tumors who were treated with re-RT during the study period, either with 3-D conformal radiotherapy, IMRT or SRS. Both pediatric and adult patients were included.

Data collection

A pre-formatted Excel sheet was used to collect the following data: 1) baseline demographic and clinical data including age, gender, and pathology type, grade, and side; 2) RT1 parameters including radiation technique, total radiation dose, dose per fraction, number of fractions, duration, and maximal dose for optic nerves, optic chiasm and brainstem; 3) RT1 outcomes including time from RT1 to disease progression and re-resection; 4) re-RT parameters including RT1–re-RT interval, technique, total radiation dose, dose per fraction, number of fractions, duration, and maximal dose for optic nerves, optic chiasm and brainstem, in addition to the cumulative radiation doses for the same structures; 5) re-RT outcomes including progression and time from end of re-RT to progression, and BRN. Alpha beta ratio of 3 (for late responding organs) was used for re-RT composite dose calculations. Patients were assessed clinically for neuropathy in follow-up clinic.

Statistical methods

Statistical analysis was performed with the Statistical Package for Social Sciences version 21.0 for Windows (SPSS Inc., Chicago, IL, USA). Categorical variables are presented as frequency and percentage, while numerical variables are presented as mean $\pm$ standard deviation (SD) or median (range) as applicable. Kaplan-Meier survival function was used to plot the survival curve for the progression-free time after re-RT, and to estimate the mean and median PFS. Further, factors associated with PFS were analyzed using Kaplan-Meier survival with calculation of the Log-rank; a Log-rank <0.05 was considered to reject the null hypothesis.

Results

Demographics and first radiation therapy parameters

Fifteen patients with brain re-irradiation were eligible and were included in the study, age range was 11 – 75 years and 8 of them were male. Pathology showed 7 cases of glioblastoma and 5 cases of anaplastic ependymoma, and 13 patients were grade III or IV. The first RT used VMAT in 12 patients, the mean duration was 42 days, and the median number of fractions was 30. The median maximal equivalent dose in 2 Gy (EQD2) received to brainstem was 37.05 Gy. Disease progression occurred after a mean time of 29 months and 7/15 underwent re-resection (Table 1).

Table 1
Patients' demographics data and first radiation therapy parameters

Parameter	Category	Frequency	Percentage
<i>Demographic and clinical data</i>			
Age (years)	Mean, SD (range=11, 75)	33.73	19.11
Gender	Male	8	53.3
	Female	7	46.7
Radiation side	Right hemisphere	9	60.0
	Left hemisphere	4	26.7
	Midline	2	13.3
Lobe	Frontal	6	40.0
	Parietal	4	26.7
	Occipital	1	6.7
	Temporal	1	6.7
	Other (central or diffuse)	3	20.0
Pathology	Glioblastoma	7	46.7
	Anaplastic Ependymoma	5	33.3
	Astrocytoma	1	6.7
	Ependymoma	1	6.7
	Meningeal melanomatous	1	6.7
Grade	Grade II	1	6.7
	Grade III	6	40.0
	Grade IV	7	46.7
	Missing	1	6.7
<i>Treatment parameters</i>			
Radiation technique	3-D	3	20.0
	VMAT	12	80.0
Values are frequency and percentage, except where otherwise specified.			

Parameter	Category	Frequency	Percentage
Total Radiation dose (Gy) / Fractions	35/10#	1	6.7
	54/30#	1	6.7
	56/28#	1	6.7
	59.4/33#	4	26.7
	60/30#	8	53.3
Dose per fraction (Gy)	1.8	6	40.0
	2	8	53.3
	3.5	1	6.7
No. fractions	Median, range	30	10, 33
Radiation therapy duration (days)	Mean, SD	42	11.13
	Median, range	42	11, 64
Maximal dose (Gy)			
Left optic nerve	Median, range	10.6	0.4 - 51.8
Right optic nerve	Median, range	7.0	0.4 - 54.7
Optic chiasm	Median, range	37.6	0.6 - 55.8
Brainstem	Median, range	50.6	1.0 - 57.6
<i>Outcomes</i>			
Time from end of radiation to progression (months)	Mean, SD	29.0	15.11
	Median, range	30.0	11 - 47
Re-resection	No	8	53.3
	Yes	7	46.7
Time from end of first radiation to re-resection (months)	Median, range	30.0	11 - 56
Values are frequency and percentage, except where otherwise specified.			

Re-irradiation parameters

The mean interval from first RT to re-RT was 24 months (range=2 – 60). The mean total cumulative dose after re-RT as per EQD2 was 101.97 Gy (max 135.6 Gy). The total mean (max) cumulative doses for organs at risk as per EQD2 after re-RT were 54.05 (92.93) Gy for brain stem, 41.19 (87.94) Gy for optic chiasma, and 28.79 (77.18) Gy and 28.6 (88.71) Gy for left and right optic nerves respectively. The

median total re-radiation dose was 35 Gy delivered over a median 10 fractions and over a mean period of 18.7 days. The brainstem received a median maximal dose EQD2 of 17 Gy, resulting in a median cumulative dose of 54.7 Gy (Table 2). No evidence of neuropathy was noted clinically.

Table 2
Re-irradiation parameters

Parameter	Category/(unit)	Mean	SD	Median	Range	N	%
<i>Treatment parameters</i>							
RT1 –re-RT interval	(months)	24.0	16.0	20.0	2, 60		
Radiation technique	3-D					4	26.7
	VMAT					9	60.0
	SRT					2	13.3
Total radiation dose	(Gy)	32,43	13.09	35	10.00, 54.00		
Dose per fraction	(N)	4,38	4.57	3	1.80, 18.00		
No. fractions	(N)			10	1, 30		
RT duration	(days)	18.7	15.5	13.0	1.0, 50.0		
Maximal doses							
Left optic nerve	(Gy)			4.0	0.2, 38.2		
Right optic nerve	(Gy)			2.0	0.3, 37.6		
Optic chiasma	(Gy)			6.0	0.5, 36.8		
Brainstem	(Gy)			10.0	0.5, 38.0		
Cumulative doses[§]							
Total EQD2	(Gy)	101.97	15.68	102.75	66.00, 135.00		
Total BED	(Gy)	168.04	26.73	163.34	110.00, 226.00		
Max. left optic nerve	(Gy)	28.79	26.79	22.22	0.96, 77.18		
Max. Right optic nerve	(Gy)	28.62	29.57	18.90	1.05, 88.71		
Max. Optic chiasma	(Gy)	41.20	30.17	49.41	1.82, 87.94		

RT1: First radiotherapy; re-RT: Re-irradiation; EQD2: equivalent dose in 2 Gy fractions; BED: Biologically Effective Dose.

[§] Cumulative radiation dose = dose for 1st radiotherapy + dose of re-irradiation

Parameter	Category/(unit)	Mean	SD	Median	Range	N	%
Max. Brainstem	(Gy)	54.06	32.96	60.24	2.40, 92.93		
Chemotherapy	No					9	60.0
	Yes					6	40.0
<i>Outcomes</i>							
Progression	No					5	33.3
	Yes					10	66.7
Time from end of RT to progression	(months)			4.5	0.0, 13.0		
Radiation necrosis	No					13	86.7
	Yes					2	13.3
RT1: First radiotherapy; re-RT: Re-irradiation; EQD2: equivalent dose in 2 Gy fractions; BED: Biologically Effective Dose.							
§ Cumulative radiation dose = dose for 1st radiotherapy + dose of re-irradiation							

Outcomes – Progression and radiation necrosis

Subsequent to re-RT, disease progression occurred in 10/15 patients after a median follow up time of 4.5 months (range = 0 – 13). The mean and median PFS were 5.13 months (95%CI=2.51 – 7.76) and 4 months (95%CI=0 – 9.1) respectively (Table 2, Figure 1).

Radiation necrosis occurred in 2 patients, giving an incidence of 13.3% (95%CI = 1.7% – 40.5%) (Table 2). It was confirmed on MRI spectroscopy. The characteristics of the two patients who developed brain radiation necrosis are presented in Table 3. Both were female, aged 11 and 46 years old. Intervals between first RT and re-RT were 20 and 12 months. Both patients had disease progression.

Table 3
Characteristics of the two radiation necrosis cases

Parameter	Case 1	Case 2
<i>Demographic and clinical data</i>		
Age (years)	46	11
Gender	Female	Female
Radiation side	Left	Right
Lobe	Frontal	Frontal
Pathology	Glioblastoma	Anaplastic ependymoma
Grade	IV	III
<i>First radiotherapy parameters</i>		
Radiation technique	VMAT	VMAT
Total Radiation dose (Gy)	60	59.4
Dose per fraction	2	1.8
No. fractions	30	33
Radiation therapy duration (days)	55	46
Maximal dose (Gy)		
Left optic nerve	2.6	42.9
Right optic nerve	2.0	53.1
Optic chiasma	4.5	53.5
Brainstem	4.7	53.8
Re-resection	Yes	No
Time to progression (months)	17	10
<i>Re-irradiation parameters</i>		
RT1-RE-RT interval (months)	20	12
Radiation technique	VMAT	3-D
Total Radiation dose (Gy)	27,5	30
Dose per fraction	55	3

RT1: First radiotherapy; re-RT: Re-irradiation; PFS: progression-free survival; EQD2: equivalent dose in 2 Gy fractions; BED: Biologically Effective Dose.

Parameter	Case 1	Case 2
No. fractions	5	10
Radiation therapy duration (days)	8	11
Maximal dose (Gy)		
Left optic nerve	0.4	30.0
Right optic nerve	0.3	30.0
Optic chiasma	0.5	30.0
Brainstem	0.5	30.0
<i>Cumulative doses (Gy)</i>		
Total EQD2	91.58	93.02
Total BED	152.64	155.04
Max. left optic nerve	70.92	77.18
Max. Right optic nerve	88.71	86.98
Max. Optic chiasma	87.94	87.36
Max. Brainstem	88.7	87.65
<i>Outcome</i>		
Progression	Yes	Yes
Death	No	Yes
PFS (months)	4	2
RT1: First radiotherapy; re-RT: Re-irradiation; PFS: progression-free survival; EQD2: equivalent dose in 2 Gy fractions; BED: Biologically Effective Dose.		

Factors associated with progression-free survival

The Kaplan-Meier survival analysis showed no statistically significant factor associated with PFS (Log-rank>0.05); however, some observations are worth noting. PFS was relatively longer among patients with grade II disease (9.0 vs <6 months). Regarding treatment parameters, a longer PFS was associated with a longer RT1 – re-RT interval, a longer re-RT duration. The two patients who developed brain necrosis -one had glioblastoma and the other anaplastic ependymoma- had a relatively shorter PFS (mean=3.0 vs 5.6 months, p=0.342) compared to their counterparts respectively (Table 4).

Table 4
Factors associated with progression-free survival (Kaplan-Meier survival analysis)

Parameter	Category	Mean	95%CI	Log-Rank
<i>Demographic and clinical data</i>				
Age (years)	≤33	5.5	1.4	9.6
	>33	4.8	1.1	8.5
Gender	Male	5.9	1.8	9.9
	Female	4.1	1.1	7.1
Radiation side	Rt hemisphere	3.3	0.5	6.2
	Lt hemisphere	8.7	3.6	13.8
	Midline	9.0	9.0	9.0
Grade	Grade II	9.0	9.0	9.0
	Grade III	5.8	0.2	11.3
	Grade IV	4.0	0.1	7.9
<i>Treatment parameters</i>				
RT1 to re-RT interval (months)	≤20	3.7	0.4	7.0
	>20	7.1	3.1	11.1
re-RT duration (days)	≤13	4.5	1.7	7.3
	>13	7.0	0.0	14.1
Cumulative total equivalent dose (Gy)	≤102	18.20	3.12	33.29
	>102	7.03	2.10	11.96
Chemotherapy	No	3.7	0.5	6.9
	Yes	7.2	3.1	11.3
<i>Outcomes</i>				
Radiation necrosis	No	5.6	2.5	8.7
	Yes	3.0	1.0	5.0
Kaplan-Meier survival analysis; event= progression or death after re-radiation				
RT1: First radiotherapy; re-RT: Re-irradiation.				

Discussion

Summary of findings

Over the 10-year period of the study, only 15 patients underwent re-RT for primary brain tumor in our institution. With respect of the size limitation of this cohort, findings support that recurrent or treatment-resistant, high-grade glioblastomas and anaplastic ependymomas represent the most common indications of re-RT in primary brain tumors. These tumors are characterized with early progression after the first treatment, leading to shortened intervals between the first RT and re-RT in majority patients. Regarding safety in our institute, findings suggest that re-RT is associated with an incidence of brain radio-necrosis of 13.3%, depending on radiation dose and pathology.

Indications of brain re-irradiation

The first published data regarding re-RT of the cranium and CNS date back to nearly one century, with the works of Beclere and Levy reporting cases of single or twice re-RT.^{15,16} In a historical series of 16 patients with brain tumors, mainly gliomas, published in 1926, 2 of the cohort patients were reirradiated after 6 and 12 months of the first RT respectively.¹⁷ Relapsed brain tumors constitute the most common indication of fractionated re-RT, which remains the only treatment option in majority of these patients. Several pathological types of primary brain tumors have been reported in re-RT cohorts, notably glioblastomas, anaplastic gliomas, medulloblastomas, ependymomas and meningiomas.^{9,18} In the pediatric population, recurrent medulloblastoma/primitive neuroectodermal tumors, germinoma and non-germinomatous germ cell tumors, and all-grade gliomas were reported as indications for re-RT, which is often adjunctive to re-resection.^{19–21} Both in adults and children, recurrence in primary brain tumors is often observed in high-grade tumors or atypical ones.^{21–23} This is consistent with the present study's cohort including 40.0% and 46.7% of grade III and IV tumors, respectively. Consequently, the primary therapeutic plan for such patients with high recurrence profile should be adapted at two levels: first, to reduce the risk of recurrence or delay its occurrence, by performing a more aggressive surgery to minimize the residual disease;

The indication of re-RT is not limited to primary brain tumors, brain metastasis was early considered among the eventual indications of brain re-RT.²⁴ However, unlike primary brain tumors, the metastatic indications often require whole brain radiotherapy due to the frequent presence of multiple, diffused lesions to be treated, which may result in a higher risk of brain atrophy or failure.^{25,26} Nonetheless, this indication was out of the scope of the present study.

Efficacy and safety balance of brain re-irradiation

It is common knowledge that the growing use of re-RT in both primary and metastatic brain tumors is associated with improved overall survival especially with the advent of high-precision RT techniques. However, the expected clinical efficacy should be assessed cautiously and weighed against the risk of radiotoxicity on the healthy structures notably the brain parenchyma and organs at risk such as the optic chiasm and optic nerves. Such caution balance combined with the scarcity of clinical data result in lack of evidence-based consensus and persisting controversies regarding the dosimetry and treatment

regimens of such patients.^{27,28} Further, the growing number of single-center reports is inconsistent with respect of the efficacy and safety profiles.

Findings from the present study suggest a low-efficacy profile for re-RT given the high incidence of disease progression (66.7%) occurring early, i.e., with a median time of 4.5 months from the end of re-RT. On the other hand, the safety profile of re-RT in the present cohort was relatively high, as only 2 patients developed brain radio-necrosis representing an incidence risk of 13.3%; however, the severity of the necrotic lesions in the two concerned patients and their respective functional outcomes were not reported. Further, given the small size of the present cohort, the incidence of radio-necrosis can statistically be inferred with a large confidence interval of 1.7% – 40.5%, which is conclusive regarding the safety profile.

By comparison, a retrospective cohort by Stiefel et al. analyzed the outcomes of 76 patients with recurrent brain tumors, of whom 34 (44.7%) were primary and the others metastatic tumors. Outcomes in patients with primary tumors showed a median overall survival of approximately 14 months (range=21 – 42 months) after re-RT, while the median time to local recurrence was not reached due to high proportion of censoring and was less than 6 months in majority of non-censored patients. Regarding radio-toxicity, authors did not carry out separate analysis for the primary tumors group; instead, they reported an overall incidence of radio-necrosis of ~12%, including 5.3% in the acute phase (<12 weeks post-re-RT). Furthermore, low-grade toxicity events such as edema, headache and fatigue, were observed in 74% of the patients.²³ A comparable incidence of radio-necrosis was in our present study; however, we did not report the low-grade toxicity events.

A clinical trial by Møller et al. randomized 31 candidates for brain re-RT for recurrent high-grade glioma, of whom 81% had glioblastoma, into 4 treatment groups, each a specific sequential regimen. Of the 4 groups, 3 had a planning target volume (PTV)<100cm³ including Group 1 (3.5 Gy x 10); Group 2 (3.5 Gy x 10 + 7Gy boost); and Group 3 (5.9 Gy x 5); whereas the 4th group (39 Gy x 10) had a PTV 100–300cm³. The median PFS in the total study population was 2.8 months and the median overall survival was 7 months. However, due to high censoring and early mortality, only 7 patients reached a PFS of 2.5 months, 5 of them received the first regimen (3.5 Gy x 10, PTV<100 cm³) and 2 received the second one (3.5Gy x 10 + 7Gy boost, PTV<100 cm³). Beside these low efficacy profiles, a low-safety profile was reported including high rate of minor early toxicity and serious late events in 3 out of 7 patients with PFS>2.5 months, including radio-necrosis and irreversible white matter changes with neurofunctional sequelae.²⁹

In pediatric patients, a 24-year retrospective study by Bouffet et al. reported a higher efficacy profile in a series 18 children who were selected for full re-RT, with or without re-resection, among a total 47 with recurrent ependymoma. Re-irradiation was associated with a 3-year survival rate of 81%, compared with only 7% among non-re-irradiated children. Additionally, re-irradiation was associated with a significantly delayed disease progression/relapse (3-year PFS=61% vs 25%) compared to the initial relapse after the first RT respectively, p=0.003. However, this high-efficacy and survival profile was achieved at the price of a significant decline in the intellectual quotient in re-irradiated children from pre- to post-re-RT times.³⁰

Improving efficacy and controlling radiotoxicity in re-irradiation

Although not statistically significant, due to small sample size, findings from the present study suggest the presence of factors that may improve the post -re-RT prognosis, notably by prolonging the PFS. Tumor characteristics that showed association trend with PFS included lower tumor grade having longer PFS than high-grade. PFS showed likelihood of positive association with RT1–re-RT interval and re-RT duration, and was likely to be increased in patients who received adjuvant chemotherapy.

Observations from animal and clinical studies support the relevance of these factors as determinant of both the efficacy and brain tolerance to re-RT.⁹ In an interesting cohort including 233 patients with recurrent grade II (40%) and III (22%) gliomas and glioblastoma (38%), Combs et al. proposed a prognostic score to determine potential candidates for re-RT based on the prognostic value of the baseline clinical and pathological characteristics. Findings demonstrated that the tumor histological type, patient age at diagnosis, and RT1–re-RT interval were the strongest predictors of survival post re-RT.³¹ The use of this score should be encouraged and further validated in different settings and cohorts, as it has a direct clinical impact on decision-making regarding the re-RT indication and could result into a consensual approach.

Further efforts are made to improve the prognosis of patients with recurrent brain tumors, while improving their quality of life. Some studies have suggested that the risk of radiotoxicity associated with re-RT can be mitigated by using proton beam therapy (PBT). A study by Mizumoto et al. (2013) analyzed the efficacy and toxicity of re-RT using conventional PBT versus conventional RT and stereotactic radiotherapy, in a cohort of 26 pediatric and adult patients with recurrent malignant brain tumors, including 15 (57.7%) glioblastomas multiform and 6 (23.1%) grade 3 gliomas. Toxicity outcomes showed 2 (7.7%) cases of radio-necrosis, which were well controlled in the second year following re-RT; beside minor acute toxicity events. Efficacy outcomes showed one- and 2-year overall survival rates (55.4% and 45.1%) and one- and 2-year local control rates (43.0% and 18.4%) respectively in the total population; and the one-year overall survival rate was higher in patients treated with PBT (75.0%).³² More recently, Scartoni et al. assessed the health-related quality of life among 33 patients with recurrent glioblastoma who underwent re-RT using PBT. Findings showed a clinically and statistically significant improvement in global health at the early post-re-RT phase, followed by progressive improvement in social functioning and motor dysfunction dimensions. However, a decline in the cognitive and emotional functioning was observed among these patients and was deemed as being non-significant. Further, the median PFS and overall survival were 5.9 and 8.7 months respectively.³³

Limitations

The present study is limited by the retrospective design and the small sample size which impact the external validity of the findings and conclusions. This highlights the urgent need for randomized trials

and prospective cohorts to provide a more accurate evaluation of the clinical benefit of re-RT in the local population.

Conclusion

Fractionated re-RT is likely to be a safe therapeutic option for recurrent primary brain tumors, with relatively low incidence of radio-necrosis. However, its efficacy may be dependent on the pathology parameters and treatment regimen, and could be confounded with factors impacting radiotoxicity notably the maximal and cumulative radiation doses on the brain parenchyma and organs at risk. Such observations highlight the relevance for an anticipative approach to determine candidates for re-RT based on predicted safety-efficacy profiles by weighing the benefits in overall and progression-free survival with the adverse effects and other relevant parameters of quality of life. There is unmet need for conducting further randomized trials and prospective cohorts to evaluate more accurately the clinical benefit of re-RT in the local population.

Declarations

Conflict of Interest:

The Authors declare that they have no competing interest.

Funding Statement

There was no funding for this work.

Acknowledgment

The authors would like to thank Dr Amine Haireche (amine.haireche@gmail.com) for his help with the statistical analysis.

Data Availability

Research data are stored in an institutional repository and will be shared upon request to the corresponding author

References

1. Jacques G, Cormac O. Central nervous system tumors. *Handbook of clinical neurology*. 2013;112:931-958.
2. Lacy J, Saadati H, Yu JB. Complications of Brain Tumors and Their Treatment. *Hematology/Oncology Clinics of North America*. 2012;26(4):779-796. doi:10.1016/j.hoc.2012.04.007

3. McFaline-Figueroa JR, Lee EQ. Brain Tumors. *The American Journal of Medicine*. 2018;131(8):874-882. doi:10.1016/j.amjmed.2017.12.039
4. de Robles P, Fiest KM, Frolkis AD, et al. The worldwide incidence and prevalence of primary brain tumors: a systematic review and meta-analysis. *Neuro-Oncology*. 2015;17(6):776-783. doi:10.1093/neuonc/nou283
5. Herholz K, Langen K-J, Schiepers C, Mountz JM. Brain Tumors. *Seminars in Nuclear Medicine*. 2012;42(6):356-370. doi:10.1053/j.semnuclmed.2012.06.001
6. Orringer DA, Golby A, Jolesz F. Neuronavigation in the surgical management of brain tumors: current and future trends. *Expert Review of Medical Devices*. 2012;9(5):491-500. doi:10.1586/erd.12.42
7. Ricard D, Idbaih A, Ducray F, Lahutte M, Hoang-Xuan K, Delattre J-Y. Primary brain tumours in adults. *The Lancet*. 2012;379(9830):1984-1996. doi:10.1016/S0140-6736(11)61346-9
8. DeAngelis LM. Chemotherapy for Brain Tumors — A New Beginning. *New England Journal of Medicine*. 2005;352(10):1036-1038. doi:10.1056/NEJMe058010
9. Nieder C, Andratschke NH, Grosu AL. Re-irradiation for recurrent primary brain tumors. *Anticancer Research*. 2016;36(10):4985-4995. doi:10.21873/anticanres.11067
10. Sminia P, Mayer R. External Beam Radiotherapy of Recurrent Glioma: Radiation Tolerance of the Human Brain. *Cancers*. 2012;4(2):379-399. doi:10.3390/cancers4020379
11. Mayer R, Sminia P. Reirradiation Tolerance of the Human Brain. *International Journal of Radiation Oncology Biology Physics*. 2008;70(5):1350-1360. doi:10.1016/j.ijrobp.2007.08.015
12. Shanker M, Chua B, Bettington C, Foote MC, Pinkham MB. Re-irradiation for recurrent high-grade gliomas: a systematic review and analysis of treatment technique with respect to survival and risk of radionecrosis. *Neuro-Oncology Practice*. 2019;6(2):144-155. doi:10.1093/nop/npy019
13. Menon D, Chelakkot PG. Reirradiation for recurrent primary central nervous system tumors: Eight-year audit from a tertiary cancer care center in South India. *Indian journal of cancer*. 2017;54(2):409.
14. Amichetti M, Amelio D. A review of the role of re-irradiation in recurrent high-grade glioma (HGG). *Cancers*. 2011;3(4):4061-4089.
15. Nieder C, Langendijk JA, Guckenberger M, Grosu AL. Preserving the legacy of reirradiation: A narrative review of historical publications. *Advances in radiation oncology*. 2017;2(2):176-182.
16. Bécélère A, Levy G. Die Strahlentherapie der Erkrankungen des Nervensystems. In: *Strahlenklinik Und Spezielle Therapeutische Methodik*. Springer; 1931:884-916.

17. Strauß O. Gehirn- und Rückenmarkstumoren. In: Gudzent F, Holthausen H, eds. *Lehrbuch Der Strahlentherapie III: Die Strahlentherapie in Der Inneren Medizin*. Urban & Schwarzenberg; 1926:433-440.
18. Chojnacka M, Skowrońska-Gardas A, Pedziwiatr K, Morawska-Kaczyńska M, Perek M, Perek D. Reirradiation of relapsed brain tumors in children. *Reports of Practical Oncology and Radiotherapy*. 2012;17(1):32-37. doi:10.1016/j.rpor.2011.10.004
19. Rao AD, Rashid AS, Chen Q, et al. Reirradiation for Recurrent Pediatric Central Nervous System Malignancies: A Multi-institutional Review. *International Journal of Radiation Oncology Biology Physics*. 2017;99(3):634-641. doi:10.1016/j.ijrobp.2017.07.026
20. Tsang DS, Laperriere NJ. Re-irradiation for Paediatric Tumours. *Clinical Oncology*. 2019;31(3):191-198. doi:10.1016/j.clon.2018.10.003
21. You SH, Lyu CJ, Kim DS, Suh CO. Second primary brain tumors following cranial irradiation for pediatric solid brain tumors. *Child's Nervous System*. 2013;29(10):1865-1870. doi:10.1007/s00381-013-2098-4
22. Combs SE, Kessel K, Habermehl D, Haberer T, Jäkel O, Debus J. Proton and carbon ion radiotherapy for primary brain tumors and tumors of the skull base. *Acta Oncologica*. 2013;52(7):1504-1509. doi:10.3109/0284186X.2013.818255
23. Stiefel I, Schröder C, Tanadini-Lang S, et al. High-dose re-irradiation of intracranial lesions – Efficacy and safety including dosimetric analysis based on accumulated EQD2Gy dose EQD calculation. *Clinical and Translational Radiation Oncology*. 2021;27:132-138. doi:10.1016/j.ctro.2021.01.011
24. Chu FCH, Hilaris BB. Value of radiation therapy in the management of intracranial metastases. *Cancer*. 1961;14(3):577-581.
25. Maranzano E, Trippa F, Casale M, et al. Reirradiation of brain metastases with radiosurgery. *Radiotherapy and Oncology*. 2012;102(2):192-197. doi:10.1016/j.radonc.2011.07.018
26. Akiba T, Kunieda E, Kogawa A, Komatsu T, Tamai Y, Ohizumi Y. Re-irradiation for metastatic brain tumors with whole-brain radiotherapy. *Japanese Journal of Clinical Oncology*. 2012;42(4):264-269. doi:10.1093/jjco/hys007
27. Nieder C. Second re-irradiation: A delicate balance between safety and efficacy. *Physica Medica*. 2019;58:155-158. doi:10.1016/j.ejmp.2019.01.004
28. Ermoian R, MacDonald S, Laack NNI, Baldini E, Breneman J. Reirradiation in Pediatric Patients With Recurrent Brain Tumors: A Last Hope, But One With Greatly Feared Consequences. *International Journal of Radiation Oncology*Biological Physics*. 2019;103(1):1-4. doi:10.1016/j.ijrobp.2018.06.036

29. Møller S, Munck af Rosenschöld P, Costa J, et al. Toxicity and efficacy of re-irradiation of high-grade glioma in a phase I dose- and volume escalation trial. *Radiotherapy and Oncology*. 2017;125(2):223-227. doi:10.1016/j.radonc.2017.09.039
30. Bouffet E, Hawkins CE, Ballourah W, et al. Survival benefit for pediatric patients with recurrent ependymoma treated with reirradiation. *International Journal of Radiation Oncology Biology Physics*. 2012;83(5):1541-1548. doi:10.1016/j.ijrobp.2011.10.039
31. Combs SE, Edler L, Rausch R, Welzel T, Wick W, Debus J. Generation and validation of a prognostic score to predict outcome after re-irradiation of recurrent glioma. *Acta Oncologica*. 2013;52(1):147-152. doi:10.3109/0284186X.2012.692882
32. Mizumoto M, Okumura T, Ishikawa E, et al. Reirradiation for recurrent malignant brain tumor with radiotherapy or proton beam therapy. *Strahlentherapie und Onkologie*. 2013;189(8):656-663. doi:10.1007/s00066-013-0390-6
33. Scartoni D, Amelio D, Palumbo P, Giacomelli I, Amichetti M. Proton therapy re-irradiation preserves health-related quality of life in large recurrent glioblastoma. *Journal of Cancer Research and Clinical Oncology*. 2020;146(6):1615-1622. doi:10.1007/s00432-020-03187-w

Figures

Figure 1

Progression-free survival after re-irradiation in patients with recurrent primary brain tumor.

Caption: Kaplan-Meier survival curve: event=progression or death; time from end of re-irradiation to progression, death or last follow up. The curve shows a median progression-free survival of 4 months.