In Vitro Screening of Antioxidant, Antimicrobial and Anticancer Activities of Cyanobacteria Found Across Odisha Coast, India

SATYABRATA DASH

Maharaja Sriram Chandra Bhanja Deo University

SABYASACHY PARIDA

Maharaja Sriram Chandra Bhanja Deo University

BIJAYANANDA SAHOO

Maharaja Sriram Chandra Bhanja Deo University

BISWAJIT RATH (✉ brath_2000@yahoo.com)

Maharaja Sriram Chandra Bhanja Deo University <https://orcid.org/0000-0002-5232-4993>

Research Article

Keywords: Antimicrobial, Anticancer, Cyanobacteria, DPPH, FRAP, GC-MS

Posted Date: January 24th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1272821/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Cyanobacteria contain a rich source of several bioprospective molecules with antibacterial, antifungal, antioxidant, and anticancer properties. The objective of the current study was to determine the antioxidant, antimicrobial and anticancer potential of methanol and acetone extracts of two cyanobacterial species (*Lyngbya majuscula* and *Lyngbya martensiana*) isolated from Odisha coast. The free radical scavenging assay of the extracts was performed against DPPH and ABTS⁺. The reducing power assay of the extracts was also determined. The maximum DPPH and ABTS radical scavenging activity is recorded on *L. majuscula* and the IC₅₀ values were 251.34±0.96 and 282.24±0.87 µg/ml respectively. Similarly, the reducing power assay was also highest in *L. majuscula* (0.762±0.0015). The total phenolic (69.18±0.79 µg gallic acid equivalents g⁻¹) and total flavonoid (38.21±0.61 µg quercetin equivalents g⁻¹) was found more in methanolic extract. The methanol extract of *L. majuscula* exhibited higher antimicrobial activity (23±0.74 mM) as compared to other test species. Further, methanol extract of *L. majuscula*, active fractions were analysed by gas chromatography-mass spectrometry (GC-MS), and bioactive compounds like 1-Monolinoleoylglycerol trimethylsilyl ether (31.182%) and n-Hexadecanoic acid (15.42%) were detected. Similarly, the cytotoxicity results exhibited decreasing cell viability of HepG2 cells in methanol extracts of *L. majuscula* in a dose-dependent manner without cytotoxicity on normal cells. These results concluded that extracts of studied cyanobacterial species exhibited appreciable antioxidant and antimicrobial activity and further the methanol extract of *L. majuscula* possess sufficient amount of bioactive compounds having antioxidant, antimicrobial and anticancer activity and thus could be a promising source for health products.

Introduction

The distinct source of antibiotics is microorganisms and the major being the marine microalgae and cyanobacteria (Blunt et al. 2009). Cyanobacteria are reported as an important source of structurally and biologically active secondary bioactive metabolites (Tan 2007; Gademann and Portmann 2008) and found to be an attractive source of antiviral, antibacterial, antioxidant and anticancer molecules (Ruiz et al. 2010). At present there has been an increasing focus on cyanobacteria as a potential source for important drugs (Dvornyk and Nevo 2003). Marine cyanobacteria have been found to be one of the ample source for unique bioactive compounds (Tan 2007; Gerwick et al. 2008; Thajuddin and Subramanian 2005). Cyanobacteria present in marine system compete actively for light and nutrient resources. Cyanobacteria have produced a wide variety of antimicrobials and allelopathic secondary metabolites to achieve a competitive edge in complex and dynamic societies (Senhorinho et al. 2015; Burja et al. 2001). Cyanobacteria are versatile organisms, according to their morphological, biochemical and metabolic properties, they are considered as attractive targets for identifying bioprospective compounds with functional utility (Falaise et al. 2016; Jena and Subudhi 2019; Barkia et al. 2019). Many Phytochemicals are present in cyanobacteria. They protect the cell constituents against destructive oxidative damage. Antioxidants present in many cyanobacteria are essential bioactive compounds that, through the defense of cells from oxidative destruction, play significant role against different diseases

and anti-ageing processes. Against numerous illnesses (e.g. chronic inflammation, cancer and coronary disorders) and aging cycles, antioxidant compounds from cyanobacteria play an important role, explaining their immense market ability in medicine, food processing and the cosmetic industry. Several experiments have reported that cyanobacteria and microalgae are found to become a profound source of secondary metabolites of antimicrobials, antioxidants and anticancer activity (Mudimu et al. 2014; Iglesias et al. 2019). Thus the present study is focused with a purpose to examine the antimicrobial, antioxidant and anticancer activity of cyanobacterial species isolated across Odisha coast.

Materials And Methods

Collection, isolation and culture conditions

Two species of Cyanobacteria (*Lyngbya majuscula* and *Lyngbya martensiana*) were collected from coastal regions of Odisha. The samples were cleaned to remove the necrotic parts and cultured in ASN III medium. The unialgal cultures were maintained under fluorescent light at an intensity of 2000-3000 lux with a photoperiod of 16hrs light and 8hrs dark at a temperature of $25 \pm 1^{\circ}\text{C}$.

Extracts preparation

The dried samples were grounded using mortar and pestle to a uniform powder. Two gram dried powder sample was dissolved in 10ml of an appropriate solvent (methanol and acetone) at room temperature for 72 hrs and the extracts were filtered with the help of Whatman No.1 filter paper. Under reduced pressure, the filtrate was evaporated and the extracts were then investigated against antioxidant, antimicrobial and anticancer activity.

Antioxidant activities

DPPH/ ABTS⁺ radical scavenging activity

The radical scavenging activity of cyanobacterial extracts is assayed by 1,1-diphenyl-2-picryl-hydrazil (DPPH) (Shah et al. 2013). One ml of DPPH (0.1 mM) was mixed with different concentration of the extracts (100, 250, 500, 750 and 1000 $\mu\text{g/ml}$) and stored for 30 minutes in dark. The absorbance was recorded using UV-visible spectrophotometer (Systronics-2202) at 517nm and ascorbic acid is used as standard. The spectrophotometric analysis of ABTS⁺ [2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonicacid) diammonium salt] radical scavenging activity was determined according to Re et al. (1999). ABTS⁺ was prepared by 5ml of ABTS (14mM) solution and $\text{K}_2\text{S}_2\text{O}_8$ (4.9mM) solution and stored in dark for 14-16 hrs. The absorbance is maintained 0.70 ± 0.020 by diluting the ABTS⁺ solution in ethanol at 734nm. The antioxidant activity was compared using ascorbic acid as a standard.

The inhibition percentage of DPPH/ABTS⁺ was obtained by means of the formula: $[(A_0 - A_1)/A_0] \times 100$, where, A_0 - control absorption, A_1 - sample absorption. The 50 percent DPPH/ABTS⁺ scavenging effect (IC_{50}) sample concentration was estimated from the DPPH/ABTS⁺ scavenging effect

percentage of graph against various concentration of sample. Lower the IC₅₀ value indicated stronger free radical scavenging activity.

Ferric reducing antioxidant power assay (FRAP)

The reducing power of the cyanobacterial extracts was determined according to the method of Oyaizu (1986). Various concentration of cyanobacterial extracts (100-1000 µg/ml) was dissolved in 2.5 ml of K₃Fe(CN)₆ (1%) and PO₄ buffer (0.2 M, pH 6.6) and stand for 25 min at 50°C followed by the addition of trichloroacetic acid (10%, 2.5ml) and centrifuged. Subsequently the supernatant (2.5ml) was added with same volume of double distilled water and FeCl₃ (0.1%, 0.5 ml). At 700 nm, the absorbance was measured. Butylated hydroxyl toluene was taken as standard.

Total Phenolic Content (TPC)

According to Slinkard et al. (1977) the total phenolic content in cyanobacterial extracts was measured by using Folin's reagent. The extract (100 µl) was mixed with Na₂CO₃ (2%, 2ml) and allowed to stand for 2 minutes. Then Folin's reagent (500 µl) was mixed and stands for 20 min in dark. The absorbance was reordered at 650 nm. The gallic acid equivalent (µg GAE/ g extract) was used to determine the total phenolic content in the extracts.

Total flavonoid contents (TFC)

The total flavonoid content was estimated by zhishen et al.(1999). One ml of cyanobacterial extract was added to NaNO₂ (5%, 500 µl) and AlCl₃.H₂O (10%, 300 µl). After 10 minutes, one ml of NaOH (1M) was mixed and distilled water was added up to 5ml make up and placed at room temperature for 30 minutes. The absorbance was measured at 510 nm. The total flavonoid in the extract was determined in the form of quercentin equivalents (µg QE /g extract).

Antimicrobial activity

Human Pathogens

Four bacterial pathogens (*Staphylococcus aureus* MTCC-96; *Bacillus subtilis* MTCC-441; *Vibrio Cholerae* MTCC-3906; *Escherichia coli* MTCC-443) and four fungal pathogens (*Candida albicans* MTCC 183; *Aspergillus niger* MTCC-1344; *Penicillium verrucosum* MTCC-1758 and *Fusarium oxysporum* MTCC-284) were obtained from IMTECH Chandigarh and maintained in Department of Biotechnology, Maharaja Sriram Chandra Bhanja Deo University, Takatpur, Mayurbhanj, Odisha. These pathogens are used for the experimental purposed.

Agar well diffusion assay

Mueller Hinton and Potato dextrose agar medium was used for evaluation of antimicrobial activity by agar well diffusion method for Bacteria and fungi respectively (Abedin and Taha 2008). The test pathogens (20 µl, 10⁶ CFU/ml) were swabbed on all the plates. The extracts were dissolved in DMSO (1 mg/ml, 50 µl) and supplied into 6mm diameter wells and kept at 37°C for 20 hours (bacteria) and 28°C

for 56 hours (fungi). The inhibited zone around the wells was measured in millimeter. Ampicillin and Clotrimazole (100g/ml) were employed as +ve controls for antibacterial and antifungal activities.

Determination of minimum inhibitory concentration (MIC)

A conventional 96-well microplate with Mueller hinton broth (MHB) was used to determine the MIC, MHB was used to developed various concentration of extracts (15.62, 31.25, 62.5, 125, 250, 500, 1000 µg/ml) using two-fold serial dilution. Fifty micro liter of pathogens (10^6 CFU/ml) was added. After 24hr (Bacteria) / 48hr (Fungus) of incubation microtiter plate reader (Bio-rad,iMark-11457) was to analyze the MIC. The minimum inhibition concentration was determined as the lowest concentration of the extracts that inhibit the visible growth of the test pathogens (Skov et al. 2019; Prasannabalaji et al. 2012).

Cytotoxicity assay

Cytotoxicity activity (*In vitro* bioassay) in human hepatocellular carcinoma cell line (HepG2) was conducted using methanol extract of *Lyngbya majuscula* by MTT assay (Mosmann1983).The cellular viability was evaluated by the reduction of the 3-(4,5-dimethylthiazole-2-yl)-2,5-diphenyltetrazoliumbromide(MTT). The cells were maintained in RPMI1640 medium. After 24h of adhesion, the media was supplemented with various concentrations of sample extracts (0.1% DMSO) and incubated for 48hours.After removal of the sample solution and washing with phosphate-buffered saline (pH 7.4) cells were exposed to 10 µL of MTT solution(0.5 mg /ml) for 4hours.MTT solution was removed after centrifugation at 2000 rpm for 10 min and the immiscible blue formazan crystals trapped in cells were dissolved and maintained in 100 µl of DMSO and the absorbance was measured at 550 nm in a microplate reader (Bio-Rad, model 3350), using wells without sample containing cells as blanks. All experiments were performed in triplicate and the concentration required for a 50% inhibition of viability (IC_{50}) was determined graphically. The effect of the samples on the proliferation of human Liver cancer cells was expressed as the % cell viability, using the following formula:

% cell viability = A_{570} of treated cells / A_{570} of control cells $\times$ 100%.

GC-MS Analysis

The chemical components of methanolic extracts of *L. majuscula* was identified by gas chromatography-mass spectrometry (GC-MS) (Swamy et al. 2017), Agilent Model 7890A-5975B [Column, DB 5ms, Agilent form (30m x 250 µm x 0.25 µm)] in the Unit of Analytical Chemistry, Triyat Scientific Co., Nagpur, Maharashtra, India. The compounds were tentatively identified by comparing their relative retention times and mass spectra to those of the NIST, WILLY library data of the GC/MS system. A relative percentage peak area was used to explore the quantification of all identified components.

Statistical analysis

For statistical purpose the experimental results were obtained in triplicate and expressed as the mean with standard deviations ($\pm$ SD).The data were analyzed by one way ANOVA, and the means were compared by the least significant difference test ($p=0.05$).

Results

Antioxidant activities

DPPH and ABTS⁺ radical scavenging assay

Antioxidant radical scavenging activities were evaluated and presented in (Table 1). The result of this assay was presented as IC₅₀ values which range from 251.34±0.96 to 309.11±0.92 µg/ml (DPPH) and 282.24±0.87 to 353.03±0.81 µg/ml (ABTS⁺). Among all the test species, methanolic extract of *L. majuscula* noted higher DPPH and ABTS⁺ radical scavenging activity (IC₅₀: 251.34± 0.96 µg/ml and 282.24±0.87 µg/ml). In contrary, lower scavenging activity was noted in acetone extract of *L. martensiana* (IC₅₀:339.11±0.92 µg/ml and 353.03±0.81 µg/ml).

Table 1
Antioxidant activity of extracts of cyanobacteria by the
DPPH and ABTS⁺ radical scavenging assays

cyanobacterial species	Methanol	Acetone
DPPH radical scavenging activity (IC ₅₀ µg/ml)		
<i>L. majuscula</i>	251.34±0.96	306.04±0.94
<i>L. martensiana</i>	257.00±0.93	339.11±0.92
ABTS ⁺ radical scavenging activity (IC ₅₀ µg/ml)		
<i>L. majuscula</i>	282.24±0.87	310.61±0.83
<i>L. martensiana</i>	326.69±0.84	353.03±0.81

Ferric Reducing Antioxidant Power assay (FRAP)

The experimental result on reducing power assay was shown in (Fig. 1A, B). Higher absorbance value suggesting higher reducing power of the sample. Methanol extract of *L. majuscula* showed highest reducing power i.e., 0.762±0.0015 µg/ml. On the other hand *L. martensiana* exhibited minimum reducing power (0.679±0.0009 µg/ml).

Total Phenolic And Flavonoid Content

The total phenolic and flavonoid content of solvent extracts (methanol and acetone) of two species of cyanobacteria was estimated in terms of Galic acid and Quercetin equivalent (µg/ml). The highest phenolic content was found in methanol (69.18±0.79 µg/ml) followed by acetone (57.14±0.75 µg/ml) extracts respectively in *L. majuscula*, whereas *L. martensiana* revealed lowest phenolic content i.e., 54.81±0.74 and 48.27±0.73 µg/ml in methanol and acetone extract respectively. Maximum flavonoid

content was observed in methanol extract of *L. majuscula* (38.21 ± 0.61 µg/ml) and minimum (22.53 ± 0.64 µg/ml) was recorded in acetone extract of *L. martensiana* (Table 2).

Table 2
Total phenolic and flavonoid contents of the cyanobacterial species

Cyanobacterial species	Methanol	Acetone
TPC (µg/ ml)		
<i>L. majuscula</i>	69.18 ± 0.79	57.14 ± 0.75
<i>L. martensiana</i>	54.81 ± 0.74	48.27 ± 0.73
TFC (µg/ml)		
<i>L. majuscula</i>	38.21 ± 0.61	30.34 ± 0.62
<i>L. martensiana</i>	29.39 ± 0.66	22.53 ± 0.64

Antimicrobial activity

The antimicrobial activity of methanol and acetone extracts of two marine cyanobacteria *L. majuscula* and *L. martensiana* showed differential zone of inhibition against bacterial and fungal pathogens. The zone of inhibition is clearly depends on the type of solvent used for extract and test bacterial and fungal pathogens. The methanol extracts of *L. majuscula* showed higher inhibition zone (23 ± 0.74 mM) against *V. cholerae* (Fig. 2A). However minimum zone of inhibition was observed in acetone extract of *L. martensiana* (10 ± 0.77 mM) against *E. coli* (Fig. 2B). Similarly maximum antifungal activity was noted in methanol extract of *L. majuscula* (21 ± 0.8 mM) (Fig. 3A), while acetone of *L. Martensiana* exhibits minimal zone of inhibition (10 ± 0.72 mM) against *C. albicans* (Fig. 3B).

Minimum inhibitory concentration (MIC)

The efficacy of antimicrobial activity against the pathogens was determined by Minimum inhibitory concentration (Table 3). The MIC for the various concentrations was within the range of 62.5 to 500 µg/ml against the tested bacterial/fungal pathogens. The highest activity was recorded in methanol extract of *L. majuscula* against *V. cholerae*, *E. coli* and *C. albicans* with MIC value (62.5 µg/ml) whereas least activity was shown in acetone extract of *L. martensiana* against all the test pathogens used. The MIC value was better against gram -ve bacteria than gram +ve bacteria.

Table 3
Minimum inhibitory concentration (MIC) of cyanobacterial species against bacterial and fungal pathogens.

Pathogens	Minimum inhibitory concentration (µg/ml)			
	<i>L. majuscula</i>		<i>L. martensiana</i>	
	Methanol	Acetone	Methanol	Acetone
<i>S. aureus</i>	125	250	500	500
<i>B. subtilis</i>	125	500	500	500
<i>V. cholerae</i>	62.5	125	250	500
<i>E. coli</i>	62.5	250	125	250
<i>A. niger</i>	125	250	250	500
<i>C. albicans</i>	62.5	125	250	500
<i>P. verrucosun</i>	250	500	125	250
<i>F. oxysporum</i>	125	250	500	500

Cytotoxic activity

Data on the cytotoxic effects of *L. majuscula* methanol extract using liver cancer cell line *In vitro* are showed in figure 3&4. The results of this study showed that the methanolic extract from *L. majuscula* can inhibit the proliferation of liver cancer cells (HepG2). Morphological changes were noted in the treated cells included the cell shrinkage resulting the cells of various shape and size as compared to normal cells and (Fig. 4). The Cytotoxic effects of various concentrations (15.62, 31.25, 62.5, 125, 250, 500, 1000µg/ml) of *L. majuscula* crude extract was investigated using MTT assay (Fig. 5). The highest percentage (88.2%) of cell viability was observed with the treatment using 15.62 µg/ml while the lowest cell viability percentage was calculated as 17.6% with the treatment using 1000µg/ml.

GC-MS analysis

GC-MS Chromatogram in methanol extract of *L. majuscula* along with molecular formula, Retention time and peak area were depicted in (Table 4). Eleven compounds were identified in methanolic extract of *L. majuscula*. The 1-Monolinoleoylglycerol trimethylsilyl ether with percentage composition of (31.182%) was the major compound followed by n-Hexadecanoic acid (15.42%), Heptadecane (13.3%), Phytol, acetate (12.47%), 3,7,11,15-Tetramethyl- 2-hexadecen-1-ol (10.37%), Hexadecanoic acid, ethylester (5.02%)

Table 4
Major chemical composition of methanol extract of *Lyngbya majuscula* using GC-MS

Sl.No	Name of the compound	Molecular Formulae	Retention time	Peak area (%)
1	2-Mercaptopropanoic acid	C ₃ H ₆ O ₂ S	4.3	0.39
2	Benzoic acid	C ₇ H ₆ O ₂	5.6	0.78
3	Undecanoic acid	C ₁₁ H ₂₂ O ₂	9.5	1.73
4	3,7,11,15-Tetramethyl- 2-hexadecen-1-ol	C ₂₀ H ₄₀ O	14	10.37
5	Phytol, acetate	C ₂₂ H ₄₂ O ₂	14	12.47
6	n-Hexadecanoic acid	C ₁₆ H ₃₂ O ₂	16	15.42
7	Hexadecanoicacid, ethylester	C ₁₈ H ₃₆ O ₂	16	5.02
8	Heptadecane	C ₁₇ H ₃₆	16.75	13.3
9	Phytol	C ₂₀ H ₄₀ O	18	1.58
10	Erucic acid	C ₂₂ H ₄₂ O ₂	25	0.87
11	1-Monolinoleoylglycerol trimethylsilyl ether	C ₂₇ H ₅₄ O ₄ Si	29	31.7

Discussion

Marine cyanobacteria are excellent producers of bioactive secondary metabolites that provide antioxidant and antimicrobial compounds (Pahlich et al. 1983). Compounds of cytotoxic, antifungal, antibacterial, antiviral and antioxidant functions have been reported in studies throughout the past decades. Previously hydrophilic and lipophilic extracts of cyanobacterial strains isolated from fresh, brackish, and marine water were tested for antibiotic efficacy against microbes (Thacker et al. 1997). In current study, four marine cyanobacteria isolated from the coastal region of Odisha were tested against various human pathogens. The findings imply that selected cyanobacterial species are potential source for antimicrobial and antioxidant compounds. Hossain et al. (2016) exhibited the antioxidant properties of four cyanobacterial species such as (*Spirulina* sp.; *Oscillatoria* sp.; *Lyngbya* sp and *Microcystis* sp.) of which the higher DPPH radical scavenging activity was expressed in *Oscillatoria* sp. and *Lyngbya* sp. which is inclined to our work. However, the DPPH assay on *Lyngbya* sp. was studied by Hossain et al. (2015) showed marginally lower scavenging activity which contradict to our findings as the test species was from a freshwater environment. Likewise, Babic et al. (2016) also reported that the *Phormidium* sp. exhibited maximum antioxidant activity followed by *Nostoc* sp. This differential response might be of varied strains and different habitats of their occurrence. The capacity to reduce indicates that the

antioxidant components are electron donors, and that the lipid peroxidation mechanism will reduce the oxidised intermediates, allowing them to function as primary and secondary antioxidants (Kokilam and Vasuki 2014). Overall, for the extraction of antioxidant compounds from cyanobacteria, methanol was the most effective solvent, this might be because methanol has a greater dielectric constant than acetone. The present study revealed that methanol extracts of *Lyngbya majuscula* marked high contents of both total phenols and total flavonoids which agree to the results reported by Sahin (2019); Patil and Kaliwal (2019) which states that higher total phenolic contents in *Scenedesmus* sp. Several workers have concluded that cyanobacterial phenolic compounds (along with flavonoids) substantially refer as their antioxidant potential (Hajimahmoodi et al. 2010; Shetty and Sibi 2015; Morowvat et al. 2019). These compounds show a broad variety of biological activities in addition to their antioxidant function, explaining their commercial applications in food and pharmaceutical industries (Machu et al. 2015; Ali and Doumandji 2017; Galasso et al. 2019). All the cyanobacteria extracts inhibited the test pathogens and were more potent against Gram-negative bacteria, for which is consistent with the findings of Caires et al. (2018) and Pramanik et al. (2011). The test species *Lyngbya majuscula* expressed the maximum inhibitory activity against both bacteria and fungi. Similar observation was reported by Sundaramanickam et al. (2015), that *Lyngbya* sp. has a significant inhibitory impact on Gram-negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*). Similarly, Sethubathi and Prabu (2010) demonstrated that solvent extracts of *Oscillatoria* sp., *Phormidium* sp., and *Lyngbya majuscula* have antimicrobial activity against human pathogens such as *S. mutants*, *S. aureus*, *P. aeruginosa*, *B. subtilis* and *K. pneumonia* which supplement to our result. Further, MIC assay result was found to be coherence to the results of earlier author showing lowest MIC against *V. cholerae* and *E. coli* among the test cyanobacterial extracts. The antibacterial activity of the cyanobacterial extracts was found to be more promising than the antifungal activity in the current investigation, which is consistent with the findings of Martins et al. (2008). GC-MS assessment of present study revealed higher Hexadecanoic acid in *L. majuscula*, similar to the Plaza 2010 results, which showed superior antimicrobial activity in the presence of different fatty acids and volatile compounds such as fucosterol, phytol, neophytadiene, palmitic, palmitoleic and oleic acids in *Synechocystis* sp. Similarly Geetha (2010) investigated the presence of diisooctyl ester, 1,2-Benzenedicarboxylic acid, 9, 12,15-Octadecatrienoic acid, Octadecanoic acid, n-Hexadecanoic acid, 9,12-ctadecadienoic acid, n-Hexanoic acid, Decanoic acid, Oleic acid, and other compounds in chlorella extracts. Inoue et al. 2005 demonstrated that the growth of *Staphylococcus aureus* is inhibited by phytol. The anticancer activity as determined in *L. majuscula* using HepG2 cells was encouraging depicting lower cell viability at higher concentration of extract which was supported the work of Yusof et al. 2010, reported anticancer screening of the microalga *Chlorella vulgaris* against cancerous cell lines HepG2 and found to prevent cell proliferation and to promote apoptosis.

Conclusions

It can be summarized that two extracts (methanol and acetone) from two test species of cyanobacteria demonstrated various levels of antioxidant and antimicrobial activity. We also observed that the phenolic content of cyanobacteria plays a significant role in antioxidants and antimicrobial activities. Eleven

compounds with various biological activities were proposed in the GC/MS study to extend our understanding of this species as a producer of biologically active substances. Our analysis indicates that the methanol extract of *Lyngbya majuscula* has substantial biological activity (antioxidant and antimicrobial), rendering this species an important topic for further studies in the food, nutraceutical and pharmaceutical sectors. Further, we found that *Lyngbya majuscula* extract, affect cell viability by inducing apoptosis in Human liver cancer cell lines *in vitro* indicating it as an anti-cancer agent. This study suggests that further work will be helpful in pharmaceutical sciences for determining cyanobacteria as potential source for therapeutic purposes.

Abbreviations

DPPH- 1,1-diphenyl-2-picryl-hydrazil, ABTS-2, 2-azinobis-(3-ethylbenzothiazoline-6)- sulfonic acid, PUFA- Poly unsaturated fatty acid, ASN III- Artificial sea nutrient, Abs- Absorbance, FRAP -Ferric reducing antioxidant power assay, BHT- Butylated hydroxyl toluene, TPC - Total phenolic content, GAE- Gallic acid equivalent, TFC- Total flavonoid contents, QE- Quercetin equivalents, MTCC- Microbial type culture collection and gene bank, MHA-Mueller-hinton agar, PDA- Potato dextrose agar, CFU- Colony forming unit, DMSO- Dimethyl sulfoxide, MIC- Minimum inhibitory concentration, MHB-Mueller hinton broth, GC-MS- Gas Chromatography-Mass Spectrometry, SD- Standard deviation, ANOVA- Analysis of variance, *P. molle*- *Phormidium molle*, *P. angustissium* – *Phormidium angustissium*, *L. majuscula*-*Lyngbya majuscula*, *L. martensiana*- *Lyngbya martensiana*, *S. aureus*- *Staphylococcus aureus*, *B. subtilis*-*Bacillus subtilis*, *V. cholerae*- *Vibrio cholerae*, *E. coli*- *Escherichia coli*, *C. albicans*-*Candida albicans*, *A. niger*- *Aspergillus niger*, *P. verrucosum*- *Penicillium verrucosum*, *F. oxysporum*- *Fusarium Oxysporum*

Declarations

Acknowledgments

The authors would like to convey their heartfelt thanks to Head, Department of Biotechnology, MSCBD University, for facilitating laboratory for smooth conduct of experiments.

Financial support

No financial support received for the work

Conflict of Interest

The authors declare that there is no conflict of interest.

References

1. Abedin RM, Taha HM (2008) Antibacterial and antifungal activity of cyanobacteria and green microalgae. Evaluation of medium components by Plackett-Burman design for antimicrobial activity

- of *Spirulina platensis*. Global Journal of Biotechnology and Biochemistry 3(1):22–31
2. Ali IH, Doumandji A (2017) Comparative Phytochemical analysis and in vitro antimicrobial activities of the cyanobacterium *Spirulina platensis* and the green alga *Chlorella pyrenoidosa*: potential application of bioactive components as an alternative to infectious diseases, vol 39. Bulletin de l'Institut Scientifique, Rabat, pp 41–49. Section Sciences de la Vie
 3. Babic O, Kovac D, Raseta M, Sibul F, Svircev Z, Simeunovic J (2016) Evaluation of antioxidant activity and phenolic profile of filamentous terrestrial cyanobacterial strains isolated from forest ecosystem. J Appl Phycol 28(4):2333–2342
 4. Blunt JW, Copp BR, Hu WP, Munro MH, Northcote PT, Prinsep MR (2009) Review Marine natural products. Nature Products 26(2):170–244
 5. Burja AM, Banaigs B, Abou-Mansour E, Burgessd JG, Wrighta PC (2001) Marine cyanobacteria Da prolific source of natural products. Tetrahedron 57:9347–9377
 6. Caires TA, da Silva AM, Vasconcelos VM, Affe HM, de Souza Neta LC, Boness HV, Sant'Anna CL, Nunes JM (2018) Biotechnological potential of *Neolyngbya* (Cyanobacteria), a new marine benthic filamentous genus from Brazil. 36:1–9Algal research
 7. Dvornyk V, Nevo E (2003) Genetic polymorphism of cyanobacteria under permanent natural stress: a lesson from the “Evolution Canyons”. Res Microbiol 154(2):79–84
 8. Falaise C, François C, Travers MA, MorgaB, Haure J, Tremblay R, Turcotte F, Pasetto P, Gastineau R, Hardivillier Y Leignel V(2016)Antimicrobial compounds from eukaryotic microalgae against human pathogens and diseases in aquaculture.Marine drugs14(9):159
 9. Gademann K, Portmann C (2008) Secondary metabolites from cyanobacteria: complex structures and powerful bioactivities. Curr Org Chem 12(4):326–341
 10. Galasso C, Gentile A, Orefice I, Ianora A, Bruno A, Noonan DM, Sansone C, Albin A, Brunet C (2019) Microalgal derivatives as potential nutraceutical and food supplements for human health: A focus on cancer prevention and interception. Nutrients 11(6):1226
 11. Geetha BV, Rajendran N, Ekambaram P(2010)Investigation of antioxidant capacity and phytochemical composition of Sun Chlorella-an in vitro study.Journal of Aquaculture Research and Development1(2)
 12. Gerwick WH, Coates RC, Engene N, Gerwick L, Grindberg RV, Jones AC, Sorrels CM (2008) Giant marine cyanobacteria produce exciting potential pharmaceuticals. Microbe-American Society for Microbiology 3(6):277
 13. Hajimahmoodi M, Faramarzi MA, Mohammadi N, Soltani N, Oveisi MR, Nafissi-Varcheh N (2010) Evaluation of antioxidant properties and total phenolic contents of some strains of microalgae. Journal of Applied Phycology22(1):43–50
 14. Hossain MF, Ratnayake RR, Meerajini K, Kumara KLW (2015) Antioxidant properties in some selected cyanobacteria isolated from fresh water bodies of Sri Lanka. Food Science & Nutrition 4(5):753–758
 15. Hossain MF, Ratnayake RR, Meerajini K, Wasantha Kumara KL (2016) Antioxidant properties in some selected cyanobacteria isolated from fresh water bodies of Sri Lanka. Food science & nutrition

16. Iglesias MJ, Soengas R, Probert I, Guilloud E, Gourvil P, Mehiri M, López Y, Cepas V, Gutiérrez-del-Río I, Redondo-Blanco S, Villar CJ (2019) NMR characterization and evaluation of antibacterial and antibiofilm activity of organic extracts from stationary phase batch cultures of five marine microalgae (*Dunaliella* sp., *D. salina*, *Chaetoceros calcitrans*, *C. gracilis* and *Tisochrysis lutea*). *Phytochemistry* 164: 192-205
17. Inoue Y, Hada T, Shiraishi A, Hirose K, Hamashima H, Kobayashi S (2005) Biphasic effects of geranylgeraniol, teprenone, and phytol on the growth of *Staphylococcus aureus*. *Antimicrob Agents Chemother* 49(5):1770–1774
18. Jena J, Subudhi E (2019) Microalgae: An untapped resource for natural antimicrobials. The role of microalgae in wastewater treatment. Springer, Singapore, pp 99–114
19. Machu L, Misurcova L, Vavra Ambrozova J, Orsavova J, Mlcek J, Sochor J, Jurikova T (2015) Phenolic content and antioxidant capacity in algal food products. *Molecules* 20(1):1118–1133
20. Martins RF, Ramos MF, Herfindal L, Sousa JA, Skærven K, Vasconcelos VM (2008) Antimicrobial and cytotoxic assessment of marine cyanobacteria-Synechocystis and Synechococcus. *Mar Drugs* 6(1):1–11
21. Morowvat MH, Goharian S, Ghasemi Y (2019) Investigation of Antioxidant Properties of Three Naturally Isolated Microalgae: Identification and Bioinformatics Evaluation of the Most Efficient Strain. *Recent Patents Biotechnol* 13(4):277–283
22. Mosmann T (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 65(1–2):55–63
23. Mudimu O, Rybalka N, Bauersachs T, Born J, Friedl T, Schulz R (2014) Biotechnological screening of microalgal and cyanobacterial strains for biogas production and antibacterial and antifungal effects. *Metabolites* 4(2):373–393
24. Oyaizu M, dietetics (1986) Studies on products of browning reaction antioxidative activities of products of browning reaction prepared from glucosamine. *The Japanese journal of nutrition and* 44(6):307–315
25. Pahlich E, Kerres R, Jäger HJ (1983) Influence of water stress on the vacuole/extra vacuole distribution of proline in protoplasts of *Nicotiana rustica*. *Plant Physiol* 72(2):590–591
26. Patil L, Kaliwal BB (2019) Microalga *Scenedesmus bajacalifornicus* BBKLP-07, a new source of bioactive compounds with in vitro pharmacological applications. *Bioprocess Biosyst Eng* 42(6):979–994
27. Plaza M, Santoyo S, Jaime L, Reina GGB, Herrero M, Señoráns FJ, Ibáñez E (2010) Screening for bioactive compounds from algae. *Journal of pharmaceutical and biomedical analysis* 51(2):450–455
28. Pramanik A, Sundararaman M, Das S, Ghosh U, Mukherjee J (2011) Isolation and characterization of Cyanobacteria possessing antimicrobial activity from the sundarbans, the World's largest tidal mangrove forest. *J Phycol* 47(4):731–743

29. Prasannabalaji N, Muralitharan G, Sivanandan RN, Kumaran S, Pugazhvendan SR (2012) Antibacterial activities of some Indian traditional plant extracts. *Asian Pacific Journal of Tropical Disease* 2:291–295
30. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med* 26(9–10):1231–1237
31. Ruiz B, Chávez A, Forero A, García-Huante Y, Romero A, Sánchez M, Rocha D, Sánchez B, Rodríguez-Sanoja R, Sánchez S, Langley E (2010) Production of microbial secondary metabolites: regulation by the carbon source. *Crit Rev Microbiol* 36(2):146–167
32. Sahin SC (2019) *Scenedesmus obliquus*: A Potential Natural Source for Cosmetic Industry. *International Journal of Secondary Metabolite* 6(2):129–136
33. Senhorinho GN, Ross GM, Scott JA (2015) Cyanobacteria and eukaryotic microalgae as potential sources of antibiotics. *Phycologia* 54(3):271–282
34. Sethubathi GVB, Prabu VA (2010) Antibacterial activity of cyanobacterial species from adirampattinam coast, southeast coast of palk bay. *Current Research Journal of Biological Sciences* 2(1):24–26
35. Shah NA, Khan MR, Ahmad B, Noureen F, Rashid U, Khan RA (2013) Investigation on flavonoid composition and anti free radical potential of *Sida cordata*. *BMC Complement Altern Med* 13(1):1–12
36. Shetty V, Sibi G (2015) Relationship between total phenolics content and antioxidant activities of microalgae under autotrophic, heterotrophic and mixotrophic growth. *Journal of Food Resource Science* 4(1):1–9
37. Skov R, Frimodt-Moller N, Blomstrom A, Espersen F (2019) Comparison of different MIC methods and establishment of zone diameter breakpoints for mecillinam, using NCCLS methodology. In Program and abstracts of the 40th Interscience Conference on Antimicrobial Agents and Chemotherapy (Toronto) Canada
38. Slinkard K, Singleton VL (1977) Total phenol analysis: automation and comparison with manual methods. *Am J Enol Viticult* 28(1):49–55
39. Sundaramanickam A, Palanivel S, Shekhar S, Kumaresan S, Balasubramanian T (2015) In vitro evaluation of Antimicrobial activity of some selected Cyanobacterial extracts against human pathogens. *International Journal of Advance Pharmacology Biology and Chemistry* 4(1):36–43
40. Swamy MK, Arumugam G, Kaur R, Ghasemzadeh A, Yusoff MM, Sinniah UR (2017) GC-MS based metabolite profiling, antioxidant and antimicrobial properties of different solvent extracts of Malaysian *Plectranthus amboinicus* leaves. *Evidence-Based Complementary and Alternative Medicine* 2017:1–11
41. Tan LT (2007) Bioactive natural products from marine cyanobacteria for drug discovery. *Phytochemistry* 68(7):954–979

42. Thacker RW, Nagle DG, Paul VJ (1997) Effects of repeated exposures to marine cyanobacterial secondary metabolites on feeding by juvenile rabbitfish and parrotfish. *Mar Ecol Prog Ser* 147:21–29
43. Thajuddin N, Subramanian G (2005) Cyanobacterial biodiversity and potential applications in biotechnology. *Current science* 47–57
44. Zhishen J, Mengcheng T, Jianming W (1999) The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem* 64(4):555–559

Figures

Figure 1

(A-B) Ferric Reducing Antioxidant Power assay in two solvent extract, Methanol(A) and Acetone(B) of two test species of cyanobacteria

Figure 2

(A-B) Antibacterial activity of *L. majuscula* & *L. martensiana* against bacterial pathogens (M-Methanol; A-Acetone; AMP- Ampicillin)

Figure 3

(A-B) Antifungal activity of *L. majuscula* & *L. martensiana* against fungal pathogens

(M-Methanol; A- Acetone; CLT- Clotrimazole)

Figure 4

Cytotoxic effect of *Lyngbya majuscula* methanol extracts against HepG2. The cells were treated with various concentrations (control; 15.62; 31.25; 62.5; 125; 250; 500 and 1000 µg/ml)

Figure 5