Using An Optimized Generative Model To Infer The Progression of Complications In Type 2 Diabetes Patients

Xiaoxia Wang

Fudan University

Yun Xiong

Fudan University

Joseph M. Plasek

Brigham and Women's Hospital

David Bates

Brigham and Women's Hospital

Chunlei Tang (✉ ctang5@partners.org)

Brigham and Women's Hospital

Research Article

Keywords: Computer Simulation, Disease Progression Model, "Diabetes Mellitus, Type 2", Probabilistic Generative Model, Electronic Health Records

Posted Date: February 4th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1274048/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Using an optimized generative model to infer the progression of complications in type 2 diabetes patients

Xiaoxia Wang¹, Yun Xiong¹, Joseph M. Plasek², David W. Bates^{2,3}, Chunlei Tang^{2,3*}

¹Shanghai Key Laboratory of Data Science, School of Computer Science, Fudan University, Shanghai, China; ²Division of General Internal Medicine and Primary Care, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA;

³Clinical and Quality Analysis, Mass General Brigham, Boston, MA, USA.

* Correspondence: C. Tang ctang5@partners.org

Division of General Internal Medicine and Primary Care

Brigham and Women's Hospital

1620 Tremont Street, BS-3

Boston, MA 02120, USA

(857) 366-7211

ABSTRACT

Background People with type 2 diabetes would progress slowly and have few symptoms before diagnosed, if any, and don't discover their condition until complications develop. Furthermore, with the development of the diabetes, patients often develop comorbidities. However, heterogeneity of progression coupled with deficiencies in electronic health records related to incomplete data, discrete events, and irregular event intervals make identification of pre-diabetes and critical points of diabetes progression challenging.

Methods We utilized longitudinal electronic health records of 9,298 patients with type 2 diabetes or prediabetes from 2005 to 2016 from a large regional healthcare delivery network in China. We optimized a generative Markov-Bayesian-based model to generate 5000 synthetic illness trajectories. The synthetic data were manually reviewed by endocrinologists.

Results We build an optimized generative progression model for type 2 diabetes using anchor information to reduce the number of parameters learning in the third layer of the model from $O(N \times W)$ to $O((N - C) \times W)$, where N is the number of clinical findings, W is the number of complications, C is the number of anchors. Based on this model, we infer the relationships between progression stages, the onset of complication categories, and the associated diagnoses during the whole progression of type 2 diabetes using electronic health records.

Conclusions The full type 2 diabetes patient trajectories generated by the chronic disease progression model can counter a lack of real-world evidence of desired longitudinal timeframe while facilitating population health management.

Keywords: Computer Simulation; Disease Progression Model; “Diabetes Mellitus, Type 2”, Probabilistic Generative Model; Electronic Health Records

Word Counts: 3,100 (233 in abstract)

I. INTRODUCTION

With total medical spending and out-of-pocket costs of patients with chronic diseases, such as diabetes, cancer, heart disease and chronic lung disease, has become increasingly expensive, modeling the entire progression path to obtain disease stages/states is one of the most important in chronic disease care [1]. It is estimated that more than 1 in 3 American adults (~88 million) and nearly 36% of the Chinese adult population have prediabetes [1, 2]. Patients with type 2 diabetes usually have few, if any, symptoms initially. Patients' persist in the prediabetes phase for years; thus, the disease is often undetected until it progresses to a chronic condition as serious complications develop. When prediabetes, patients either have insulin resistance where the body still produces insulin but is unable to effectively use insulin, or they don't produce enough insulin, leading to accumulation of glucose in the bloodstream. Diabetes affects multiple major organs and its most frequent complications include myocardial infarction, stroke, neuropathy, kidney damage, and microvascular events [3-7].

In the medical field, a large amount of data (e.g., laboratory test results, clinical findings, diagnoses, symptoms, and medication treatments) recorded in electronic health record (EHR) systems can facilitate clinical knowledge discovery. However, EHR data has inherent limitations for studying progression of type 2 diabetes from prediabetes to overt diabetes, including its static nature (e.g., family history information, genetic testing results), missing values, and irregularity (e.g., data are recorded at discrete time points with non-equal intervals). Disease progression models (DPM) require substantial domain knowledge on disease stages, vital indicators/ measurements, and insight into the target diseases epidemiology. Watabe *et al.* [8] employed a hierarchical

Bayesian framework to infer the progression level to diabetes based on oral glucose tolerance tests. It is not a true DPM due to not focusing on the slow development of diabetes over time. Marini *et al.* [9] developed a Dynamic Bayesian Network (DBN) model to simulate of development of several clinical complications of type 1 diabetes. Islam *et al.* [10] applied a machine learning pipeline to predict future development of type 2 diabetes based on finding an optimal set of risk-factors. This is not the case for our purpose because we aim to use a minimally supervised approach to generate the full trajectories of chronic diseases, which does not require either a training dataset with patient disease stages labeled or domain knowledge that specifies the indicators for stage transitions.

Type 2 diabetes is a chronic disease with a progression trajectory that can span 30+ years from pre-diabetes to severe complications and death meaning that most EHR datasets only cover a portion of the relative longitudinal trajectory. In order to cover the full longitudinal trajectory, synthetic data generated using probabilistic generative models can be a suitable proxy. Sukkar *et al.* [11] used unsupervised hidden Markov models to create a general disease progression model. Wang *et al.* [12] utilized a three-layer pipeline (Fig. 1) consisting of the Markov Jump Process, Markov Chain, and noisy-or Bayesian network to similarly create an unsupervised disease progression model that infers progression from the onset of comorbidities. In this model, a comorbidity is a disease or syndrome that co-occurs with the target disease. Comorbidities are assumed to be conditionally independent, given the state of the target disease. The bottom layer is a bipartite noisy-or Bayesian network that is used to infer the presence of the comorbidities from the observed clinical findings (e.g., ICD codes). Given a set of ICD codes, Wang *et al.* assume an observed clinical finding was "activated" by the presence of any of the comorbidities with a certain activation

Fig 1. The outline of Wang *et al.*'s model, where K is the number of disease stages, M is the number of complications, and N is the number ICD codes.

probability; it is also possible that none of the comorbidities is present and the finding was activated by an always-on hidden cause with a leak probability. This layer allows the model to deal with large amounts of clinical findings. The pros of Wang *et al.*'s model [12] are visible. Its structure is flexible enough to be well suited to the setting, especially for modeling sparse and noisy observations. The model can be used to consider either irregular patient visits or a continuous-time disease progression. The gap in the literature our study focuses on is to improve the time complexity of Wang *et al.*'s model and adapting the model design to a new population.

In this study, we aim to infer full progression trajectories of type 2 diabetes via synthetic patient generation. Firstly, to learn more efficiently the probabilistic relationships between chronic disease progression stages, complication categories, and clinical diagnoses based on the real incomplete patient records in EHR data sets, we optimize Wang *et al.*'s model [12] (see Fig. 1). Then we suit the optimized model to build a progression model for type 2 diabetes using a real EHR data set. We further demonstrate the full progression of type 2 diabetes by means of synthetic patients generated by the learned progression model and how the model can facilitate population health management.

II. METHODS

A. Dataset Settings

Data was derived from a 17-hospital-based regional healthcare delivery network managed by the local Center for Disease Control (CDC) in Shanghai, China. The data integrates real world electronic health record (EHR) data with "follow-up" data (that was generated from that tracks patient outcomes for those same patients). Data were coded using the International Classification of Diseases - Version 10 (ICD-10-CM) codes. This study was approved by Shanghai CDC's Institutional Review Board (IRB).

Our dataset consists of 9,298 real patients with confirmed type 2 diabetes over an 11-year timespan from January 2005 to January 2016, in which 43.3% (4,028/9,298) were male, 100% had been hospitalized in the facility at some point during the timespan for any cause, and a 3.9% mortality rate (367/9,298, 188 males and 179 females). We retrieved a total of 1,311 distinct ICD codes relating to these patients' comorbidities from the data. We next removed infrequent ICD-10 codes (i.e., 1,223 of 1,311) that appeared less than 30 times, leaving 88 distinct ICD-10 codes for use in the generative models. Considering that type 2 diabetes progress slowly, we integrated the patient records within one year into a time slice as an encounter. Since the model needs to calculate the interval between two adjacent encounters, we excluded patients whose total number of time slices are less than 2 from the data. Fig. 2 shows the number of positive observations in each encounter for each real patient, and demonstrates our data are very sparse. Furthermore, note that it is a big challenge to learn a full progression model for type 2 diabetes using data which time span are shorter than the common type 2 diabetes progression.

B. Model Optimization

The time efficiency of Wang *et al.*'s is largely dependent on the number of a variety of variables and time slices. More specifically, given 1,000 distinct ICD codes (i.e., N) of patients'

clinical findings relating to type 2 diabetes, 10 complications (i.e., W), and 5 disease stages (i.e., K), the number of model parameters is 10,125 calculated by the following formula.

$$(K \times K) + 2(K \times W) + (N \times W) \\ = (5 \times 5) + 2(5 \times 10) + (10 \times 1,000) = 10,125 \quad (1)$$

The idea of our optimization is to prune (i.e., reduce the number of parameters) the fully connected observable bottom layer. We consulted our medical experts to add some clinical pieces of knowledge. They believe that most ICD-10 codes only belong to one complication relating to type 2 diabetes. We referred to these pieces of medical knowledge as "anchors." A set of anchors as prior knowledge given by our medical experts is set to a known probability of obtaining a better-interpreted clustering result of the complication group. Due to no recalculation, the model can save most of the time to run and performs excellently. Also, anchors are few enough to make the model unsupervised. In our optimized model, we considered the observed data as clinical diagnoses (i.e. $F_{w,p,t}$ are ICD-10 codes), and the third layer as complications related to a group of corresponding clinical diagnosis (i.e. $C_{n,p,t}$ are complications). Fig. 2 (a) shows that the initial layer is fully connected. As shown in Fig. 2(b), given a set of anchors $\{(C_1, [F_2]), (C_2, [F_1, F_5]), (C_3, [F_3])\}$ corresponding to a set of complications $\{C_1, C_2, C_3\}$ and a set of clinical diagnoses $\{F_1, F_2, F_3, F_4, F_5\}$. The anchors help strengthen any possible links with high probability based on prior knowledge (i.e., blue lines). Using anchor $(C_2, [F_1, F_5])$, which means clinical diagnoses F_1 and F_5 should be clustered into complication C_2 , we prune all other links to F_1 and F_5 , and only links $C_2 \rightarrow F_1$ and $C_2 \rightarrow F_5$ are left. Since few anchors exist, it is impossible to determine all possible links with low probability, so that some necessary lines may remain. For example, for a clinical diagnosis F_4 , none of the anchors helped at removing the black lines, and it remains a fully connected with all three complications C_1, C_2, C_3 .

Fig 2. An illustration of before and after optimizing the observable bottom layer

In addition, we considered an alternate to improve efficiency; that is, to give an enlarged time granularity and reduce the number of time slices. We integrate the patient records within one year into a single time slice. This is a mild assumption in our case because generally type II diabetes is a common chronic disease with a slow progression.

C. Data Analysis and Statistics

As a comparison, we trained the original Wang *et al.* model on our data and compared the run time against our optimized J Med Internet Res J Med Internet Res model using a computer with 2 GeForce GTX1080 Yi 11G cards.

TABLE I. ANCHOR SETTINGS

Serial Number	Complication	Comorbidities (ICD-10 Code-Based Anchors)
1*	Diabetes	E11.9 (Diabetes without complications)
2	Acute complications	E11.0 (Diabetes with coma), E11.1 (Diabetes with ketoacidosis), K81 (Cholecystitis), J20 (Bronchitis)
3	Cardiovascular	I25 (Chronic ischemic heart disease), I10 (Hypertension)
4	Nephropathy	E11.2 (Diabetes with renal complications), N18 (Chronic nephrosis)
5	Ophthalmopathy	E11.3 (Diabetes with ophthalmic complications), H26.9 (Cataracts, unspecified)
6	Peripheral vascular	E11.5 (Diabetes with peripheral circulatory complications), I83 (Varicose vein of lower extremity)
7	Cerebrovascular	I63 (Cerebral infarction), G45 (Transient cerebral ischemic attacks)
8	Neuropathy	E11.4 (Diabetes with neurological complications), G63.2 (diabetic polyneuropathy)
9	Metabolic complications	E11.6 (Diabetes with other specified complications), E78 (Lipidemia)
10	Tumor	Z51.1 (Chemotherapy session for neoplasm), C34 (Malignant neoplasm of bronchus), C16 (Malignant neoplasm of stomach)
11	Musculoskeletal	M48 (Spondylodinia), M13 (Arthritis), M81 (Osteoporosis)
12	Autoimmune diseases	K52 (Gastroenteritis), E04 (Goiter), J45 (Asthma)

* Diabetes itself needs an anchor to deal with the case of “no complications.”

We first change the following main parameters of our optimized model to suit type 2 diabetes and learn a generative unsupervised progression model, where disease stages/states and onset of diabetes complications were hidden variables. Type 2 diabetes’ disease stages/states were specified as 5 (i.e., $K = 5$) by referencing the diabetes complications severity index (DCSI) [13]. With two medical experts’ help and literature reviewing [14], we set the number of complication categories to 12 (see Table I), namely $W = 12$. Note that $N = 88$ because there are 88 distinct ICD-10 codes after data preprocessing.

Then using the learned progression model, we infer the personal progression trajectories of real patients based on the history medical records. For comparison purposes, our medical experts helped to retrieve two additional representative cases with different development rates. We used the maximum a posteriori (MAP) inference that gives a point estimate by maximizing a posterior probability, the conventional approach in Bayesian statistics to infer progression trajectories based on existing evidence.

We next followed Wang *et al.*’s assumption to generate synthetic patient with full progression trajectories of type 2 diabetes by initializing each patient to timestamp 0 in state I, and using our optimized model to generate the patient’s subsequent stages and the complication onsets corresponding to those stages until the last stage.

III. RESULTS

A total of 35,210 encounters with 64,383 positive observations were input into our optimized model to generate 5,000 (that is a specified number) synthetic patient trajectories. We averaged over these 5,000 patient records and computed the average holding time for each state, as summarized in Fig. 3(a).

Fig. 3(a) covers a 23.9-year progression path of type 2 diabetes, which is approximately double the timespan of the available data (i.e., 11 years). Specifically, the average duration of each disease stage/state and the prevalence of each complication at different stages are computed.

The time to run the original Wang *et al.*’s model based on our data was more than ten days (each iteration needs about one hour, and each execution requires thousands of iterations). In contrast, our optimized model runs to converge within 8 days. Anchors are few enough (12 anchors) that we consider the model to be minimally supervised.

Fig. 3(b) and (c) illustrate the inferred maximum posterior probability results for the two real individuals’ personal disease progression trajectories. For individual A, the MAP inference only infers one possible complication as “complication 9: metabolic complications;” and our model also gives such a finding and further predicts that the progression will move to the next stage (i.e., stage II) in the ninth year. Individual B is the case of rapid deterioration. While having no ICD-10 codes related to diabetes directly (i.e., E11.9), our model determines the onset of diabetes because of finding some specified complications (e.g., E11.6); in clinical practice, these complications can be found after people with type 2 diabetes diagnosed. Our model assigns state II at the beginning of this progression trajectory and predicts that nephropathy (i.e., E11.2) and cardiovascular (i.e., I50.9) may show up after seven years. At that time, individual B’s transition from state II to state III is highlighted. The model also points out that ophthalmopathy (i.e., H26.9) will quickly follow developing state III in the next year. Then the progression path will subsequently move into state 4 under a life-threatening condition. Even though anemia (i.e., D64.9) belongs to peripheral vascular complications, yet there

Fig 3. A Comparison of complications of 5,000 virtual patients learned by our optimized model as well as two representative patients retrieved by our medical experts

TABLE II. SINGLE COMPLICATION STATISTICS OF TRANSITION FROM A LATER STATE TO AN EARLIER STATE GIVEN BY 5K GENERATED PATIENTS

Complication Stage	2 Acute Complications	3 Cardiovascular	4 Nephropathy	5 Ophthalmopathy	6 Peripheral vascular	7 Cerebrovascular	8 Neuropathy	9 Metabolic complications	10 Tumor	11 Musculoskeletal	12 Autoimmune diseases
Number	2,096	3,622	2,723	2,600	2,430	2,477	3,110	4,102	2,085	2,381	811
Probability	41.9%	72.4%	54.5%	52.0%	48.6%	49.5%	62.2%	82.0%	41.7%	47.6%	16.2%
Later to Earlier	$V \rightarrow IV$	1,834 87.5%	2,506 69.2%	2,617 96.1%	2,002 77.0%	2,085 85.8%	2,390 96.5%	2,746 88.3%	3,782 92.2%	2,062 98.9%	2,114 88.8%
	$V \rightarrow III$	1,352 64.5%	2,238 61.8%	2,176 79.9%	1,750 67.3%	902 37.1%	2,356 95.1%	1,567 50.4%	3,401 82.9%	1,420 68.1%	1,300 54.6%
	$V \rightarrow II$	461 22.0%	2,228 61.5%	1,822 66.9%	1,729 66.5%	423 17.4%	2,093 84.5%	1,213 39.0%	3,319 80.9%	544 26.1%	600 25.2%
	$V \rightarrow I$	161 7.7%	2,032 56.1%	634 23.3%	335 12.9%	372 15.3%	1,375 55.5%	650 20.9%	2,420 59.0%	229 11.0%	226 9.5%

All probabilities are based on stage V.

is no evidence to support it – our model indicates its activation probability is slightly small.

Fig. 3 indicates possibilities to help understand diabetes and associated complications. We wonder if any retrospective possibilities can be generated and evaluated. With illness trajectories viewed by our medical experts, we generated Table II and III to show the transition from a later state to an earlier state based on the last state (i.e., stage V). Table II is to calculate all probabilities of showing before stage V based on one single complication, and Table II is on two commonly shown complication patterns that reflects the co-occurrence of several complications (i.e., such as pattern [3, 7, 8] and pattern [4, 5, 6]). For example, complication 2 as “acute complications” occurred in state V has 2,096 virtual patients accounting for 41.9% (2,096/5,000), and is extremely low (161 virtual patients accounting for 7.7%, 161/2,096) compared with stage V. We followed our experts’ opinions to keep; for example, complication 6 as “peripheral vascular” can either occur as a single one or in with other complications such as [6] (14.6%), [4, 6] (2.8%), [5, 6] (2.5%), [4, 5, 6] (0%) in stage I (compared with

stage V). Note that relevant clinical meanings amongst the patterns are not the focus of this study. The table aims to demonstrate the information and knowledge among the entire disease progression via virtual patients, which can facilitate population health management.

IV. DISCUSSION

Our main finding was that it was feasible to optimize a minimally supervised generative model to simulate synthetic patient trajectories from EHR data focused on the progression of complications in prediabetic patients and patients diagnosed with type 2 diabetes. Our proposed model is built upon a model proposed by Wang *et al.* [12], which focused on modeling a chronic obstructive pulmonary disease (COPD) patient cohort. The model we modified, enhanced and employed for learning disease progression is scalable; it can comfortably accommodate new sources of data with clinical findings or outcomes. For instance, the lists of medications prescribed or procedures performed, the distribution over the initial disease stages/states (e.g., a function of age, gender, and family history), and patient’s

TABLE III. TWO COMMONLY COMPLICATION PATTERNS' STATISTICS OF TRANSITION FROM A LATER STATE TO AN EARLIER STATE GIVEN BY 5K GENERATED PATIENTS

Complication Pattern		[3 Cardiovascular, 7 Cerebrovascular, 8 Neuropathy]							[4 Nephropathy, 5 Ophthalmopathy, 6 Peripheral vascular]						
Stage															
Number Probability		1,132 22.6% (1,132/5,000)							717 14.3% (717/5,000)						
		[3]	[7]	[8]	[3, 7]	[3, 8]	[7, 8]	[3, 7, 8]	[4]	[5]	[6]	[4, 5]	[4, 6]	[5, 6]	[4, 5, 6]
Later															
to	V→IV	69.2	95.7	88.1	66.3	61.0	84.2	58.5	96.8	75.9	85.2	73.5	82.7	63.9	62.1
Earlier	V→III	61.7	94.3	49.7	57.9	30.4	46.7	28.4	81.3	66.4	37.2	54.3	30.7	25.2	21.3
(%)	V→II	61.2	82.4	39.2	49.6	23.2	31.9	18.5	68.9	66.1	17.4	44.2	12.3	11.2	7.9
	V→I	55.6	55.6	20.9	30.0	11.2	11.0	6.0	22.2	14.2	14.6	3.2	2.8	2.5	0

All probabilities are based on stage V.

social-behavioral history and habits (e.g., smoking, alcohol use) can be included as a supplement. With progression trajectories depicted by our optimized model, we can obtain some insights. These include, but are not limited to, what disease stages/states the patient traverses, how rapidly the disease develops, which complications can be found, and how long specified complications give the stage transition.

Previously, the idea of “synthetic patient simulations” were utilized for educational purposes (e.g., pre- and post-registration health care professional education [15]). In this way, learning of disease progression can also achieve the purpose of teaching patients or health professionals, and even healthy non-patients included. This is because; there exists no the completely GOLD criteria for partitioning type 2 diabetes stages/states, according to the American Diabetic Association guidelines. Take the DCSI (diabetes complications severity index) as an example; while having a higher citation count, the DCSI remains a reference designed by Glasheen *et al.* [13] rather than an indicator used widely in clinical practice. Our study may offer some evidence to help evaluate such indicators and then to facilitate uniform clinical guidelines. Fig. 3(a) indicates some interesting clinical insights. First, we found that metabolic complications showed up most frequently, which is throughout the entire progression of type 2 diabetes starting at the very beginning. In contrast, autoimmune diseases were the most infrequent. Second, the top 3 complications, are cardiovascular, cerebrovascular, and metabolic complications, imply that it is necessary to recommend laboratory tests regularly for prediabetic patients. Third, nephropathy and ophthalmopathy both are microvascular complications and may be useful to study in the early stages, especially in the transition from stage I to II. Last, acute complications, as well as nephropathy, peripheral vascular, and musculoskeletal complications present an increased risk after stage II. Fig. 3(a) indicates that autoimmune diseases are a relatively rare complication. This is expected as existing knowledge about type 2 diabetes; namely, developing type 2 diabetes doesn't mean that the body cannot produce insulin (such like type I diabetes, which is an autoimmune disease; the immune system attacks the pancreas, so it can't make insulin) [14]. The fact that the body is often unable to effectively use insulin to accumulate glucose in the bloodstream [3, 14]. Nevertheless, these prior research have found evidence that insulin resistance may be the result of immune system cells attacking the body's tissues rather than just a metabolic disorder, which warrants further investigation.

Although having constrained the continuous-time Markov model to allow only forward transitions, synthetic patient simulations without disease stages/states of absence can help better understand the evolution of chronic illnesses in reverse. Therefore, we can try to conduct research in population medicine through this research and take population health management as pro-active management to improve health and resolve health disparities relating to diabetes and prediabetes.

The main limitation of our study is that this work was based on a single chronic disease, type 2 diabetes, and thus our results may not generalize to other chronic conditions. In addition, our study only used one kind of data, ICD-10 codes, and therefore might limit clinical insights on population medicine which could include other sources of data, e.g., population-level data, monitoring, surveillance data, and social media data..

V. CONCLUSIONS

In this study, we employed the generative nature of Wang *et al.*'s model [12] to infer the progression of complications in type 2 diabetes patients. After adding 12 anchors based on prior domain knowledge, the model's fully connected observable bottom layer is pruned to reduce runtime significantly. These anchors only use few manual efforts so that our model is minimally supervised. Our main findings are (1) that a generative model can help solve incomplete and/or insufficient data problems to better understand the whole trajectories of lowly and long progression of chronic disease, and (2) it is feasible to facilitate population health management (e.g., prediabetes) as a statistical retrospect or prediction of synthetic patient trajectories.

DECLARATIONS

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Shanghai Putuo Supervision Institution of Health and Family Planning Commission's IRB. Informed and written consent was obtained through a regional 17-hospital-based regional healthcare delivery network by each participant according to the Declaration of Helsinki prior to beginning data collection.

CONSENT FOR PUBLICATION

Not applicable.

AVAILABILITY OF DATA AND MATERIAL

Our research data is unavailable for access because it is confidential. According to the HIPAA standard, it would be cost-prohibitive to sufficiently de-identify such a large corpus of clinical documents to remove all patient identifying data.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDING

This work was partially funded by the National Natural Science Foundation of China Projects No. U1636207, No. U1936213, and the Shanghai Science and Technology Development Fund No. 19511121204, No.19DZ1200802.

AUTHORS' CONTRIBUTIONS

CT drafted the manuscript; XW and YX designed research; YX provided funding acquisition and resources. All authors reviewed and edited the manuscript and approved the final manuscript. All the authors are accountable for the integrity of the work.

ACKNOWLEDGEMENTS

The authors thank Li Zhou for valuable comments and suggestions on the early versions.

AUTHORS INFORMATION

¹Shanghai Key Laboratory of Data Science, School of Computer Science, Fudan University, Shanghai, China; ²Division of General Internal Medicine and Primary Care, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA; ³Clinical and Quality Analysis, Mass General Brigham, Boston, MA, USA.

REFERENCES

- [1] Prediabetes - Your Chance to Prevent Type II Diabetes, "US centers for disease control and prevention," June 11, 2020. Available from: <https://www.cdc.gov/diabetes/basics/prediabetes.html#:~:text=Approximately%2088%20million%20American%20adults,%2C%20heart%20disease%2C%20and%20stroke>. [Accessed August 2020].
- [2] L. Wang, P. Gao, M. Zhang, D. Zhang, *et al.*, "Prevalence and ethnic pattern of diabetes and prediabetes in China in 2013," *Journal of the American Medical Association*, 2017, vol. 317(24): pp. 2515-2523.
- [3] V. A. Fonseca, "Defining and characterizing the progression of type II diabetes," *Diabetes Care* 32 (suppl 2), 2009, pp. S151-S156. PMID: 19875543.
- [4] S. Colagiuri, "Epidemiology of prediabetes," *Medical Clinics North America*, Mar 2011, vol. 95(2): pp. 299-307, vii. doi: 10.1016/j.mcna.2010.11.003. PMID: 21281834.
- [5] M. R. Rooney, A. M. Rawlings, J. S. Pankow, *et al.*, "Risk of progression to diabetes among older adults with prediabetes," *JAMA Internal Medicine*, 2021, vol. 181(4): pp. 511-519. doi: 10.1001/jamainternmed.2020.8774.
- [6] K. Lam, S. J. Lee. "Prediabetes—A risk factor twice removed," *JAMA Internal Medicine*. 2021, vol. 181(4): pp. 520-521. doi: 10.1001/jamainternmed.2020.8773.
- [7] R. S. DeJesus, C. R. Breitkopf, L. J. Rutten, *et al.*, "Population health management," pp. 216-223, Jun 2017.
- [8] T. Watabe, Y. Okuhara, Y. Sagara, "A hierarchical Bayesian framework to infer the progression level to diabetes based on deficient clinical data," *Computers in Biology & Medicine*, 2014, vol. 50(4): pp. 107-115.
- [9] S. Marini, E. Trifoglio, N. Barbarini, F. Sambo, *et al.*, "A Dynamic Bayesian Network model for long-term simulation of clinical complications in type 1 diabetes," *Journal of Biomedical Informatics*, 2015, vol. 57: pp. 369-376.
- [10] M. S. Islam, M. K. Qaraqe, S. B. Belhaouari, M. A. Abdul-Ghani, "Advanced techniques for predicting the future progression of type II diabetes," *IEEE Access*, 2020, vol. 8: pp. 120537-120547.
- [11] R. Sukkar, E. Katz, Y. Zhang, D. Raunig, *et al.*, "Disease progression modeling using Hidden Markov Models," *IEEE 2012 34th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC 2012)*, IEEE, San Diego CA USA, pp. 2845-2848, August 2012.
- [12] X. Wang, D. Sontag, F. Wang, "Unsupervised learning of disease progression models," *Proceedings of the 20th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining (KDD 2014)*, ACM, New York NY USA, pp. 85-94, August 2014.
- [13] W. P. Glasheen, A. Renda, Y. Dong, "Diabetes Complications Severity Index (DCSI)-update and ICD-10 translation," *Journal of Diabetes & Its Complications*, 2017, vol. 31(6): pp. 1007-1013.
- [14] J. M. Valderas, B. Starfield, B. Sibbald, *et al.*, "Defining comorbidity: implications for understanding health and health services," *The Annals of Family Medicine*, 2009, vol. 7(4): pp. 357-363.
- [15] C. O. Osborn, "Type 1 and type 2 diabetes: What's the difference?" *Healthline Media*, January 14, 2019. Available from: <https://www.healthline.com/health/difference-between-type-1-and-type-2-diabetes#symptoms> [Accessed August 2020].
- [16] A. A. Kononowicz, L. A. Woodham, S. Edelbring, *et al.*, "Virtual patient simulations in health professions education: systematic review and meta-analysis by the digital health education collaboration," *Journal of Medical Internet Research*, 2019, vol. 21(7): pp. e14676.