Weight Spectrum and Executive Function in Adolescents: The Moderating Role of Depression

Qingmin Lin

Shanghai Jiao Tong University

Yanrui Jiang

Shanghai Children's Medical Center

Xiaoning Sun

Shanghai Children's Medical Center

Yunting Zhang

Shanghai Children's Medical Center

Wenjie Shan

Shanghai Children's Medical Center

Jin Zhao

Shanghai Children's Medical Center

Xuelai Wang

Shanghai Children's Medical Center

Qi Zhu

Shanghai Children's Medical Center

Wanqi Sun

Shanghai Mental Health Center

Hui Lu

Shanghai Jiao Tong University

Fan Jiang (✉ fanjiang@shsmu.edu.cn)

Shanghai Children's Medical Center

Research Article

Keywords: Adolescents, Weight spectrum, Abdominal weight, Executive function, Depression

Posted Date: January 24th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1274226/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Overweight/obesity may impair executive function (EF). However, adolescent EF impairments across the weight spectrum were not well studied, with the moderating effect of negative emotions in the weight-EF association being limitedly investigated. We aimed to determine whether overall and abdominal weight spectrum associated with EF impairments and to identify whether negative emotions moderate the weight-EF link in adolescents. **Methods:** We applied a subsample of the SCHEDULE-A project. Adolescents (11-18 years) were recruited using a multi-stage cluster random sampling approach. Overall and abdominal weight were measured by body mass index z-score and waist-to-height ratio, respectively. We used the Behavior Rating Inventory of Executive Function (BRIEF) to evaluate adolescent EF in nature setting, and utilized the Depression Anxiety and Stress Scales (DASS-21) to assess three types of negative emotional states (i.e., depression, anxiety, and stress). **Results:** Of the 1 935 adolescents, 963 (49.8%) were male. We observed that abdominal, not overall, overweight was associated with the Global Executive Composite (GEC) impairment (OR = 1.59, 95% CI: 1.07-2.35), particularly for inhibit, emotion control, shift, working memory, and monitor domains. Furthermore, depression moderated the abdominal overweight-GEC association (P = 0.032 for interaction term), especially for emotional control, working memory, and initiate dimensions. Moreover, we also found abdominal thinness was associated with the Metacognition Index problem (OR = 1.33, 95% CI: 1.04-1.72), particularly for plan and monitor areas. **Conclusions:** Both abdominal overweight and thinness were associated with adolescent EF, and depression would be a modifiable target to improve EF in adolescents with abdominal overweight.

Background

Nearly 50 years, overweight/obesity has become a global pandemic, especially in children with a much larger rate of increase [1]. While the increasing trend of overweight/obesity prevalence has flattened in developed western countries, such increase, however, is still accelerating in most developing Asian countries [2]. Study showed that overweight/obesity rate in the Chinese children and adolescents quadrupled from 5.3% in 1995 to 20.5% in 2014 [3], and now China has the largest number of children and adolescents with excessive body weight [4]. It is worth noting that children's overweight/obesity, especially in adolescence, is more likely to persist into adulthood, not only predisposing individuals to a bulk of adverse physical consequences such as metabolic syndrome, cardiovascular disease, type 2 diabetes, and cancers [1, 5], but also leading to a high risk of cognitive impairment and lower academic achievement [6].

Executive function (EF), a sub-domain of cognitive capacity, also called cognitive or executive control, refers to a set of top-down neurocognitive mental processes against one's bottom-up instinct actions [7]. Recent work suggested that obesity may increase the risk of EF issues in children and adolescents [8]. However, the weight-EF relationship in adolescents has not been well studied. While some studies reported adolescents with overweight/obesity had poor EF performance [9], others showed null results [10, 11]. Additionally, few research has investigated the association of underweight and EF in

adolescents. Considering that most studies had a limited sample size [8], only used the body mass index (BMI) indicator (less sensitive to body fat, [12]) to assess overweight/obesity condition, and often neglected the underweight group that may present with a lower EF score [13], all of which might lead to the above-mentioned equivocal findings, and hence hinder a comprehensive understanding of the weight-EF link.

More importantly, the real situation that not all individuals with overweight/obesity will progress to EF impairment suggests that the weight-EF association is likely contingent on some factors. Negative emotions (e.g., depression) are common in children, particularly during adolescence, a period known for significant changes in the body, the brain and the mind, as well as the social environment around them [14]. Adolescents with weight concerns may tend to experience negative self-image and more maladaptive behaviors (e.g., unhealthy eating, physical inactivity, and more screen exposure) and, in turn EF problems [15], alternatively, negative emotions may serve to exacerbate the weight issue [16], which gave us a hint that negative emotions might moderate the weight-EF relationship. However, much of the current literature only focused on the solitary effect of weight status on EF without considering moderating effect of negative emotions [9-11]. Given that a growing number of adolescents suffer from weight problems, it is crucial to identify modifiable targets for the EF improvement.

To address these gaps, the current study aimed to determine whether overall and abdominal weight spectrum (including underweight and overweight) significantly associated with EF impairment and to identify whether negative emotions (e.g., depression, anxiety, and stress) moderate the weight-EF association in adolescents.

Methods

Study sample

From the study of the Shanghai Children's Health, Education and Lifestyle Evaluation-Adolescents (SCHEDULE-A), a population-based cross-sectional survey investigating risk factors of adolescent physical and mental health (11-18 years), the current study utilized a subsample of adolescents recruited from Shangrao region (December 2018), a relatively social-economic underdeveloped city located in the downstream Yangtze River in southeast China. The subsample included both overall and abdominal weight indicators, and also had a relatively high underweight rate that allowed us to examine the underweight-EF relationship. A multi-stage cluster random sampling approach (i.e., District-School-Class-Student) was used. Details were reported in **Method S**. Briefly, four districts/counties were firstly sampled according to the per capita disposal income of the Chinese residents in 2016; next, in each sampled district/county, two junior and two senior high schools were selected at random (4 districts/counties $\times$ 4 schools); then one class from each grade of the selected schools was drawn randomly; finally, all students were invited to take part in the study. The protocol was approved by the Shanghai Children's Medical Center Human Ethics Committee according to the Declaration of Helsinki (SCMCIRB-K2018103). All parents and their adolescent children provided written informed consents.

Main variables

Weight spectrum

We acquired adolescents' height, weight, and waist circumference from each school. BMI was calculated as the weight divided by the squared height, and waist-to-height ratio (WHtR) was computed as the waist circumference divided by the height. BMI value was subsequently converted into z-score using the least mean square method according to the World Health Organization growth reference. Then, we defined BMI z-score <-1 , ≥ 1 , and ≥ 2 as the overall underweight, overweight, and obesity, and defined WHtR <0.40 , ≥ 0.46 , and ≥ 0.50 as the abdominal thinness, overweight, and obesity, respectively [17, 18].

Negative emotions

We assessed three negative emotional states (i.e., depression, anxiety, and stress) using the Depression Anxiety and Stress Scales with 21 items (DASS-21), which was validated in the Chinese population with a Cronbach's alpha of 0.92 and test-retest reliability of 0.46 for the total score [19]. Each domain consists of seven items rated on a four-point Likert-type scale from 0 "did not apply to me at all" to 3 "applied to me very much or most of the time". The summed score for each domain was multiplied by 2, and higher scores indicated greater negative emotional symptoms. Adolescent who scored moderate or above with the cutoffs of ≥ 14 , ≥ 10 , and ≥ 19 were classified as experiencing depression, anxiety, and stress issues, respectively.

Executive function

We measured EF in a natural setting employing a parent form of the Behavior Rating Inventory of Executive Function (BRIEF), which can serve as a screening tool for possible executive dysfunction in children and adolescents aged 5-18 years old [20], and showed a good reliability and validity in the Chinese population [21]. Of the total 86 items, each one is rated on a three-point scale (i.e., never, sometimes and often), and parents were required to select the most suitable answer for each described behavior of their adolescents in the past 6 months. For a few individuals aged over 18 years ($n = 137$), we utilized the same version, as they lived together with their parents, and their everyday function therefore could be well-assessed by their parents [22]. We checked the raw data based on two validity indexes (i.e., negativity <5 and inconsistency <7) to reduce reporting bias according to the BRIEF manual.

The BRIEF has eight non-overlapping domains, two broader indices, and one overall index. The broad Behavior Regulation Index (BRI) includes three domains, i.e., inhibit, emotional control, and shift, which are interpreted as the ability to regulate one's own behavioral and emotional control, and to move flexibly from one action to another. The broad Metacognition Index (MI) incorporates five domains, i.e., working memory, plan, initiate, organize, and monitor, which are related to the ability to solve problem actively, and to initiate, organize, and monitor one's own actions. All eight domains form the overall index, i.e., the Global Executive Composite (GEC). T-scores were computed based on sex- and age-specific norms, with the T-score >60 and >65 being defined as potential sub-clinical and clinical EF impairments or problems.

Confounders

Parents or other main caregivers reported the socio-demographic information, including parents' educational attainment, gross household income, and adolescents' sex, birth date, and some chronic physical and mental health status. Screen exposure was measured by two widely used questions in children and adolescents [23], that is in the last month, on average, the total time he/she spent per day on 1) sitting and watching television or videos, and 2) playing games using device such as cellphone, iPad, PlayStation, etc. Each response was dichotomized, with ≥ 2 h/day indicating excessive screen time. Adolescents were also asked "At what time do you usually go to bed and get up on weekdays and weekends, respectively?" Averaged night sleep duration (i.e., time in bed in current study) was calculated by $(5 \times \text{weekdays} + 2 \times \text{weekends}) / 7$, then was classified as shorter duration by cutoffs of <9 , <8 , and <7 h for adolescents aged 12-13, 14-17, and ≥ 18 years, respectively [24]. We used a short Chinese version of the International Physical Activity Questionnaire to measure adolescent physical activity intensity, and then categorized it into low, moderate and high level [25].

Statistical analysis

Participants' characteristics were described with mean (SD) and frequency (%), and the social-demographic differences between analyzed and excluded sample were assessed by t-test and chi-squared test for continuous and categorical variables, respectively.

In considering of a non-linear association between weight and EF, we fitted a linear regression model with BMI z-score and WHtR included as both a linear and quadratic term for each EF domain. Because almost all BMI z-score quadratic terms were not statistically significant (**Fig. S1**), but WHtR quadratic terms were (**Fig. S2**), two overall weight categories (i.e., normal and overweight), and three abdominal weight categories (i.e., thinness, normal, and overweight) were used in the subsequent analyses (because of a low obesity prevalence, these adolescents therefore were included in the overweight category, **Table 1**).

We used multivariable logistic regression with standard error type of clustered robust to determine the association of weight spectrum and negative emotion with each sub-clinical EF problem (because of a low prevalence of clinical EF issues, these adolescents therefore were included in the sub-clinical category, **Table 1**). The potential moderating effect of negative emotion on the weight-EF link was tested by adding an interactive term (e.g., depression $\times$ overweight) in the model. Should the interaction term reached statistical significance, simple effect analysis was subsequently performed. We further conducted a post-hoc analysis of the associations between negative emotions and weight spectrum using multinomial logistic analysis. Finally, we performed a sensitivity analysis by limiting adolescents with WHtR values within 0.2-0.7 (removing possible outliers, **Fig. S2**), and by omitting individuals with age over 18 years (the BRIEF was developed for children and adolescents aged 5-18 years).

All analyses were performed in Stata 15.0, and P values <0.05 were considered statistical significance.

Results

Descriptive characteristics

Of the final 1 935 adolescents (**Fig. 1**), the mean age was 15.32 ± 1.79 years and 963 (49.8%) were male (**Table 1**). 26.8% of adolescents' parents obtained a high school education or more, and 41.3% of their family income were $\geq 50\,000$ RMB. Moreover, approximately 40% participants had each type of screen exposure exceeding 2 h/day, 50% were shorter sleepers, and 30% displayed low physical activity. There were no significant social-demographic differences between the analyzed and excluded individuals (**Table S1**).

Association of weight spectrum and EF problems

After adjusting for confounders, we observed that abdominal, not overall, overweight had a significant association with GEC problem (OR = 1.59, 95% CI: 1.07-2.35). However, we didn't find abdominal thinness significantly associated with GEC issue (**Table 2**).

In sub-domain analyses (**Table 2, Table S2**), the abdominal overweight significantly associated with MI problem (OR = 1.55, 95% CI: 1.21-1.98), especially in working memory and monitor. Furthermore, despite of a borderline association between abdominal overweight and BRI issue (OR = 1.59, 95% CI: 0.93-2.70), all three domains (i.e., inhibit, emotion control, and shift) reached statistical significance. Moreover, we also found that abdominal thinness obviously associated with MI problem (OR = 1.33, 95% CI: 1.04-1.72), especially in plan and monitor.

Moderating effect of negative emotions in the weight-EF link

In the interactive (**Table S3, Fig. 2A1-2A3**) and simple (**Table 3**) effect analyses, we only found depressive condition positively moderated the association of abdominal overweight and GEC problem ($P = 0.032$). That is in adolescents who suffered from depression, abdominal overweight significantly associated with GEC impairment (OR = 1.92, 95% CI: 1.26-2.93); whereas, in individuals free from that condition, no significant effect was found. Additionally, we also observed a borderline negative moderating effect of depression on the link of abdominal thinness and GEC problem ($P = 0.08$), namely in individuals without depression, abdominal thinness significantly associated with GEC impairment (OR = 1.51, 95% CI: 1.10-2.08).

In sub-domain interactive (**Table S3-S4, Fig. 2, Fig. S3**) and simple (**Table 3, Table S5**) effect analyses, similar findings were observed that depression positively moderated the link of abdominal overweight with BRI ($P = 0.016$) and MI ($P < 0.001$) problems, especially for emotional control (borderline), working memory, and initiate. That is abdominal overweight significantly associated with BRI (i.e., emotion control) and MI (i.e., working memory, and initiate) impairments only in adolescents who endorsed depression. While we observed a significant negative moderating effect of depression on the link of abdominal thinness and BRI problem ($P = 0.031$), the simple effect analysis did not reveal significant findings.

Although we observed that stress positively moderated the link of abdominal thinness and BRI issue ($P = 0.044$) (**Table S3** and **Fig. 2a3**), it didn't reach statistical significance of the moderating effect in any specific sub-domains (**Fig. S4**). We also didn't find anxiety had significant moderating effect. The post-hoc analysis also showed that only depression significantly associated with abdominal overweight (**Table S7**).

Sensitivity analysis

After limiting adolescents' WHtR value and age range, the results did not change much indicating the robustness of our findings (**Table S7-S12, Table S13-S18**).

Discussion

To our knowledge, this is the first study to examine whether negative emotion moderate the weight-EF link in adolescents, including overall and abdominal weight spectrum. Firstly, we observed abdominal overweight rather than overall overweight, significantly associated with EF problem, particularly in domains of inhibit, emotion control, shift, working memory, and monitor. Furthermore, we identified a statistically significant moderating effect of depression in the relationship of abdominal overweight and EF, especially in emotional control, working memory, and initiate domains. Moreover, we also found abdominal thinness significantly associated with plan and monitor areas.

We observed a significant association of abdominal, not overall, overweight with EF problems, particularly in the areas of inhibit, emotion control, shift, working memory, and monitor, which was consistent with two prior findings indicating that waist circumstance or WHtR was more sensitive than BMI indicator when examining the weight-EF association [26, 27]. Consistent with our findings, prior studies also suggested overweight had pronounced effect on the deficits of inhibit control, working memory, cognitive flexibility (i.e., shift in our study), as well as high-order EF (e.g., reason, problem-solve, and plan) [8]. The underlying mechanism of the overweight-EF link may be attributed to chronic low-grade inflammation. Extensive studies have documented that overweight, especially abdominal adiposity (a marker of visceral fat), can induce systemic low-grade inflammation through activating pro-inflammatory processes in adipose tissue by secreting chemokines and in gut microbiota by altering intestinal permeability and releasing endotoxins [28, 29]. Inflammation can result in decreased brain volume, reduced white matter integrity, atrophied grey matter, and lower regional blood flow, all of which likely damage EF if occurred in EF-related regions, e.g., frontal, temporal and occipital cortices [30].

Furthermore, we also found adolescent depression significantly moderated the link of abdominal overweight and EF impairment, especially in the domains of emotional control, working memory, and initiate, which was similar to one adult research that the EF difference between obesity and normal weight group were more prominent in patients with major depressive disorder compared with health controls [31]. Research showed that individuals who suffered from overweight were more vulnerable to depression due to body dissatisfaction and peer stigma than those who had normal weight [32]. Meanwhile, depressed persons were more likely to gain weight and had poor metabolic profile given

increased unhealthy diet, sleep disorder, and less physical activity [33]. A recent longitudinal study also demonstrated that higher weight predicted more severe depressive symptoms and worse EF via higher levels of IL-6, and depressive symptoms also predicted IL-6 increase, and all observed associations were unique to depression but not anxious [34]. Adolescence is widely recognized as a stage of heightened emotional reactions [14], and depressive status measured by DASS-21 in our study likely predispose these adolescents to clinical disorder if not properly addressed, hence warranting much more attention.

Moreover, we observed a quadratic relation between WHtR and EF profile, similar to one recent finding showing a U-shape relationship of BMI with brain volume in children [13]. Besides aforementioned abdominal overweight, we also found adolescent abdominal thinness significantly associated with EF impairment, which other studies also tried to investigate, but all failed, maybe due to the BMI indicator they used [10]. Because of undernutrition in low income region, or self-pressure to restrict dieting fulfilling the thin beauty ideal [35] (oftentimes by unhealthy eating, [36]), individuals with abnormal thinness/underweight probably have lower than desired levels of nutritional and biochemical for optimal EF, e.g., protein, micronutrient, IGF-1 and brain derived neurotrophic factor [37, 38]. Future studies using more sensitive indicators such as body composition and biochemical markers are needed to confirm and extend our finding.

Our study neither found a significant association of overall overweight and EF, nor observed a moderating effect of depression on that association. It is possible that the BMI indicator incorporating both lean and fat mass is an insufficient marker of body fat. Although one adult study found stress significantly moderated the abdominal obesity and EF association [39], our study didn't find stress and anxiety as statistically significant moderators. Our post-hoc analysis indicating that anxiety and stress were not associated with abdominal overweight may be one of the reasons. More mechanism researches of the association between different type of negative emotions and abdominal overweight in adolescents should be conducted in the future.

Limitations

The present study had some limitations. Firstly, our sample was from one of the underdeveloped cities in China with a relatively low obesity prevalence. More samples with diverse socioeconomic status should be performed in future. Secondly, due to low rate of obesity and clinical EF impairment, we utilized the overweight and sub-clinical EF cutoffs. Although the effects estimated were at magnitudes that may be considered sub-clinical in adolescents, these findings are important at the population level. Thirdly, adolescent lifestyle behaviors relied on self-report. Future studies should use more objective measurements, e.g., using actiwatch to assess night sleep duration and physical activity, to reduce reporting bias. And lastly, given the cross-sectional nature, we cannot determine the causal association, and cannot exclude the effects influenced by unmeasured or unknown confounders.

Conclusion

Our study confirmed and extended previous researches by examining the association of overall and abdominal weight spectrum with EF, and determining the moderating effect of negative emotions on the weight-EF link in adolescents. We highlighted WHtR as a more sensitive indicator of EF than BMI that needs more attention, and identified depression as a modifiable target to improve EF in adolescents with abdominal overweight that requires more care.

Abbreviations

BMI, body mass index

BRI, the Behavior Regulation Index

BRIEF, the Behavior Rating Inventory of Executive Function

DASS-21, the 21-item version of the Depression Anxiety and Stress Scales

EF, executive function

GEC, the Global Executive Composite

MI, the Metacognition Index

SCHEDULE-A, study of the Shanghai Children's Health, Education and Lifestyle Evaluation-Adolescents

WHtR, waist-to-height ratio

Declarations

Ethics approval and consent to participate

The study was approved by the Shanghai Children's Medical Center Human Ethics Committee (SCMCIRB-K2018103). All parents and their adolescent children provided written informed consents.

Consent for publication

Not applicable.

Availability of data and materials

Data used for this study were derived from the SCHEDULE-A project. The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was supported by Ministry of Science of Technology of PRC (2016YFC1305203), Shanghai Municipal Health Commission (GWV-10.1-XK07), National Natural Science Foundation (81773443, 81602868, 81602870), Science and Technology Commission Shanghai Municipality (17XD1402800, 2018SHZDZX05, 19YF1442200), Shanghai Municipal Education Commission (D1502), Shanghai Municipal Commission of Health and Family Planning (2017ZZ02026).

Author's contributions

Conceptualization: Lin Q, Jiang F; Data curation: Lin Q, Shan W, Zhu Q; Formal analysis: Lin Q; Funding acquisition: Jiang Y, Zhang Y, Sun W, Jiang F; Investigation: Shan W, Zhao J, Wang X; Methodology: Lin Q, Jiang Y, Zhang Y, Shan W, Zhao J, Wang X, Zhu Q, Sun W, Lu H, Jiang F; Project administration: Zhang Y, Jiang F; Supervision: Zhang Y, Shan W, Zhao J, Wang X, Zhu Q, Jiang F; Validation: Lin Q, Jiang Y, Sun X; Visualization: Lin Q; Writing—original draft: Lin Q; Writing—review & editing: Lin Q, Jiang Y, Sun X, Zhang Y, Shan W, Zhao J, Wang X, Zhu Q, Sun W, Lu H, Jiang F. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Acknowledgments

We would like to thank the research staffs involved in the SCHEDULE-A project who collected and coded the data, and teachers from the selected schools who cooperated with our research work. We also wish to thank the adolescents and their families who participated in the study.

References

1. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A *et al*: Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *N Engl J Med*2017, 377(1):13-27. <https://doi.org/10.1056/NEJMoa1614362>
2. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128·9 million children, adolescents, and adults. *Lancet*2017, 390(10113):2627-2642. [https://doi.org/10.1016/s0140-6736\(17\)32129-3](https://doi.org/10.1016/s0140-6736(17)32129-3)
3. Dong Y, Jan C, Ma Y, Dong B, Zou Z, Yang Y *et al*: Economic development and the nutritional status of Chinese school-aged children and adolescents from 1995 to 2014: an analysis of five successive national surveys. *Lancet Diabetes Endocrinol*2019, 7(4):288-299. [https://doi.org/10.1016/s2213-8587\(19\)30075-0](https://doi.org/10.1016/s2213-8587(19)30075-0)
4. Pan XF, Wang L, Pan A: Epidemiology and determinants of obesity in China. *Lancet Diabetes Endocrinol*2021, 9(6):373-392. [https://doi.org/10.1016/s2213-8587\(21\)00045-0](https://doi.org/10.1016/s2213-8587(21)00045-0)

5. Sung H, Siegel RL, Torre LA, Pearson-Stuttard J, Islami F, Fedewa SA *et al*: Global patterns in excess body weight and the associated cancer burden. *CA: a cancer journal for clinicians* 2019, 69(2):88-112. <https://doi.org/10.3322/caac.21499>
6. Bluher M: Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 2019, 15(5):288-298. <https://doi.org/10.1038/s41574-019-0176-8>
7. Diamond A: Executive functions. *Annu Rev Psychol* 2013, 64:135-168. <https://doi.org/10.1146/annurev-psych-113011-143750>
8. Mamrot P, Hanć T: The association of the executive functions with overweight and obesity indicators in children and adolescents: A literature review. *Neurosci Biobehav Rev* 2019, 107:59-68. <https://doi.org/10.1016/j.neubiorev.2019.08.021>
9. Chen S, Jia Y, Woltering S: Neural differences of inhibitory control between adolescents with obesity and their peers. *International Journal of Obesity* 2018, 42(10):1753-1761. <https://doi.org/10.1038/s41366-018-0142-x>
10. Groppe K, Elsner B: Executive function and weight status in children: A one-year longitudinal perspective. *Child Neuropsychol* 2017, 23(2):129-147. <https://doi.org/10.1080/09297049.2015.1089981>
11. Chrysaidou K, Kotsis V, Chainoglou A, Tzovaras F, Gidaris D, Chatzipapa N *et al*: Impact of ambulatory SBP and overweight on executive function performance in children and adolescents. *J Hypertens* 2020, 38(6):1123-1130. <https://doi.org/10.1097/hjh.0000000000002371>
12. Brambilla P, Bedogni G, Heo M, Pietrobelli A: Waist circumference-to-height ratio predicts adiposity better than body mass index in children and adolescents. *Int J Obes (Lond)* 2013, 37(7):943-946. <https://doi.org/10.1038/ijo.2013.32>
13. Bohon C, Welch H: Quadratic relations of BMI with depression and brain volume in children: Analysis of data from the ABCD study. *J Psychiatr Res* 2021, 136:421-427. <https://doi.org/10.1016/j.jpsychires.2021.02.038>
14. Blakemore SJ: Adolescence and mental health. *Lancet* 2019, 393(10185):2030-2031. [https://doi.org/10.1016/s0140-6736\(19\)31013-x](https://doi.org/10.1016/s0140-6736(19)31013-x)
15. Lindberg L, Hagman E, Danielsson P, Marcus C, Persson M: Anxiety and depression in children and adolescents with obesity: a nationwide study in Sweden. *BMC Med* 2020, 18(1):30. <https://doi.org/10.1186/s12916-020-1498-z>
16. Schwartz BS, Glass TA, Pollak J, Hirsch AG, Bailey-Davis L, Moran TH *et al*: Depression, its comorbidities and treatment, and childhood body mass index trajectories. *Obesity (Silver Spring)* 2016, 24(12):2585-2592. <https://doi.org/10.1002/oby.21627>

17. Ashwell M, Gibson S: A proposal for a primary screening tool: 'Keep your waist circumference to less than half your height'. BMC Med2014, 12:207. <https://doi.org/10.1186/s12916-014-0207-1>
18. Zhang YX, Wang ZX, Chu ZH, Zhao JS: Profiles of body mass index and the nutritional status among children and adolescents categorized by waist-to-height ratio cut-offs. Int J Cardiol2016, 223:529-533. <https://doi.org/10.1016/j.ijcard.2016.07.303>
19. Wang K, Shi HS, Geng FL, Zou LQ, Tan SP, Wang Y *et al*: Cross-cultural validation of the Depression Anxiety Stress Scale-21 in China. Psychol Assess2016, 28(5):e88-e100. <https://doi.org/10.1037/pas0000207>
20. Baron IS: Behavior rating inventory of executive function. Child Neuropsychol2000, 6(3):235-238. <https://doi.org/10.1076/chin.6.3.235.3152>
21. Qian Y, Wang YF: [Reliability and validity of behavior rating scale of executive function parent form for school age children in China]. Beijing Da Xue Xue Bao Yi Xue Ban2007, 39(3):277-283.
22. Torske T, Naerland T, Bettella F, Bjella T, Malt E, Hoyland AL *et al*: Autism spectrum disorder polygenic scores are associated with every day executive function in children admitted for clinical assessment. Autism research : official journal of the International Society for Autism Research2020, 13(2):207-220. <https://doi.org/10.1002/aur.2207>
23. Yang L, Cao C, Kantor ED, Nguyen LH, Zheng X, Park Y *et al*: Trends in Sedentary Behavior Among the US Population, 2001-2016. Jama2019, 321(16):1587-1597. <https://doi.org/10.1001/jama.2019.3636>
24. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L *et al*: National Sleep Foundation's updated sleep duration recommendations: final report. Sleep Health2015, 1(4):233-243. <https://doi.org/10.1016/j.sleh.2015.10.004>
25. Macfarlane DJ, Lee CC, Ho EY, Chan KL, Chan DT: Reliability and validity of the Chinese version of IPAQ (short, last 7 days). J Sci Med Sport2007, 10(1):45-51. <https://doi.org/10.1016/j.jsams.2006.05.003>
26. Frye SS, Fernandez-Mendoza J, Calhoun SL, Gaines J, Sawyer MD, He F *et al*: Neurocognitive and behavioral functioning in adolescents with sleep-disordered breathing: a population-based, dual-energy X-ray absorptiometry study. Int J Obes (Lond)2018, 42(1):95-101. <https://doi.org/10.1038/ijo.2017.229>
27. Hartanto A, Yong JC: Measurement matters: Higher waist-to-hip ratio but not body mass index is associated with deficits in executive functions and episodic memory. PeerJ2018, 2018(9). <https://doi.org/10.7717/peerj.5624>
28. Deng T, Lyon CJ, Bergin S, Caligiuri MA, Hsueh WA: Obesity, Inflammation, and Cancer. Annu Rev Pathol2016, 11:421-449. <https://doi.org/10.1146/annurev-pathol-012615-044359>

29. Blaut M: Gut microbiota and energy balance: role in obesity. *Proc Nutr Soc*2015, 74(3):227-234. <https://doi.org/10.1017/s0029665114001700>
30. Lowe CJ, Reichelt AC, Hall PA: The Prefrontal Cortex and Obesity: A Health Neuroscience Perspective. *Trends Cogn Sci*2019, 23(4):349-361. <https://doi.org/10.1016/j.tics.2019.01.005>
31. Hidese S, Ota M, Matsuo J, Ishida I, Hiraishi M, Yoshida S *et al*: Association of obesity with cognitive function and brain structure in patients with major depressive disorder. *J Affect Disord*2018, 225:188-194. <https://doi.org/10.1016/j.jad.2017.08.028>
32. Sutaria S, Devakumar D, Yasuda SS, Das S, Saxena S: Is obesity associated with depression in children? Systematic review and meta-analysis. *Arch Dis Child*2019, 104(1):64-74. <https://doi.org/10.1136/archdischild-2017-314608>
33. Fourrier C, Sampson E, Hori H, Schubert KO, Clark S, Mills NT *et al*: Exploratory study of association between blood immune markers and cognitive symptom severity in major depressive disorder: Stratification by body mass index status. *Brain Behav Immun*2020, 88:242-251. <https://doi.org/10.1016/j.bbi.2020.06.007>
34. Mac Giollabhui N, Swistun D, Murray S, Moriarity DP, Kautz MM, Ellman LM *et al*: Executive dysfunction in depression in adolescence: the role of inflammation and higher body mass. *Psychol Med*2020, 50(4):683-691. <https://doi.org/10.1017/s0033291719000564>
35. Caqueo-Úrizar A, Ferrer-García M, Toro J, Gutiérrez-Maldonado J, Peñaloza C, Cuadros-Sosa Y *et al*: Associations between sociocultural pressures to be thin, body distress, and eating disorder symptomatology among Chilean adolescent girls. *Body Image*2011, 8(1):78-81. <https://doi.org/10.1016/j.bodyim.2010.11.004>
36. Zarychta K, Chan CKY, Kruk M, Luszczynska A: Body satisfaction and body weight in under- and healthy-weight adolescents: mediating effects of restrictive dieting, healthy and unhealthy food intake. *Eat Weight Disord*2020, 25(1):41-50. <https://doi.org/10.1007/s40519-018-0496-z>
37. Okereke O, Kang JH, Ma J, Hankinson SE, Pollak MN, Grodstein F: Plasma IGF-I levels and cognitive performance in older women. *Neurobiol Aging*2007, 28(1):135-142. <https://doi.org/10.1016/j.neurobiolaging.2005.10.012>
38. Zwipp J, Hass J, Schober I, Geisler D, Ritschel F, Seidel M *et al*: Serum brain-derived neurotrophic factor and cognitive functioning in underweight, weight-recovered and partially weight-recovered females with anorexia nervosa. *Prog Neuropsychopharmacol Biol Psychiatry*2014, 54:163-169. <https://doi.org/10.1016/j.pnpbp.2014.05.006>
39. Lasikiewicz N, Hendrickx H, Talbot D, Dye L: Exploring stress-induced cognitive impairment in middle aged, centrally obese adults. *Stress*2013, 16(1):44-53. <https://doi.org/10.3109/10253890.2012.682109>

Tables

Table 1. Participant characteristics.

Characteristic	Participants, mean $\pm$ SD / No. (%)
Age, y	15.32 $\pm$ 1.79
Sex	
Boys	963 (49.8)
Girls	972 (50.2)
Household factors	
Parental highest education	
Lower than high school	1 374 (73.2)
High school or higher	503 (26.8)
Family income (RMB)	
<50 000	939 (58.7)
$\geq$ 50 000	661 (41.3)
Individual behaviors	
Screen exposure time	
Sitting&Watching, $\geq$ 2 h/day	868 (44.9)
Playing games, $\geq$ 2 h/day	769 (39.8)
Sleep duration, short	978 (51.1)
Physical activity	
Low	557 (28.8)
Moderate	710 (36.7)
High	666 (34.5)
Predictors	
Overall weight spectrum (BMI z-score)	
Underweight, <-1	436 (22.5)
Overweight, $\geq$ 1	215 (11.1)
Obesity, $\geq$ 2	32 (1.7)
Abdominal weight spectrum (WHtR)	
Thinness, <0.40	708 (36.6)
Overweight, $\geq$ 0.46	442 (22.8)
Obesity, $\geq$ 0.50	188 (9.7)
Potential Moderators	
Depression, $\geq$ 14	492 (25.4)
Anxiety, $\geq$ 10	1 022 (52.8)
Stress, $\geq$ 19	379 (19.6)
Executive function problems	
GEC score	
Sub-clinical problem, >60	393 (20.3)
Clinical problem, >65	176 (9.1)
BRI score	
Sub-clinical problem, >60	340 (17.6)
Clinical problem, >65	224 (11.6)
MI score	
Sub-clinical problem, >60	399 (20.6)
Clinical problem, >65	127 (6.6)

Abbreviations: BMI, body mass index; BRI, the Behavioral Regulation Index; GEC, the Global Executive Composite; MI, the Metacognition Index; WHtR, waist-to-height ratio.

Table 2. Association of overall and abdominal weight spectrum, as well as negative emotion with executive function problems in adolescents.

		GEC problem		BRI problem		MI problem	
		OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
Model a-e¹							
Overall weight status							
Normal		Ref.		Ref.		Ref.	
Overweight		1.01 (0.65, 1.58)	0.949	0.92 (0.53, 1.61)	0.782	1.17 (0.63, 2.17)	0.613
Abdominal weight spectrum							
Normal		Ref.		Ref.		Ref.	
Thinness		1.31 (0.88, 1.95)	0.179	1.19 (0.57, 2.48)	0.645	1.33 (1.04, 1.72)	0.025
Overweight		1.59 (1.07, 2.35)	0.021	1.59 (0.93, 2.70)	0.089	1.55 (1.21, 1.98)	< 0.001
Depression							
No		Ref.		Ref.		Ref.	
Yes		3.38 (1.35, 8.44)	0.009	2.95 (1.08, 8.09)	0.036	3.03 (1.45, 6.36)	0.003
Anxiety							
No		Ref.		Ref.		Ref.	
Yes		2.78 (1.65, 4.70)	< 0.001	3.04 (1.85, 4.99)	< 0.001	2.44 (1.36, 4.39)	0.003
Stress							
No		Ref.		Ref.		Ref.	
Yes		1.75 (1.07, 2.87)	0.026	1.44 (1.04, 1.97)	0.026	1.48 (0.96, 2.29)	0.076

Abbreviations: BRI, the Behavioral Regulation Index; GEC, the Global Executive Composite; MI, the Metacognition Index.

¹ All models were adjusted for social-demographic factors (i.e. age, sex, parental highest education, and family gross income) and individual lifestyle behaviors (i.e. screen exposure, nighttime sleep duration, and physical activity).

Table 3. Simple effects of abdominal weight spectrum on executive function problems stratified by depression condition in adolescents¹.

		GEC problem		BRI problem		MI problem	
		OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
No depression							
Abdominal weight spectrum							
Normal		Ref.		Ref.		Ref.	
Thinness		1.51 (1.10, 2.08)	0.010	1.37 (0.67, 2.80)	0.386	1.30 (1.01, 1.66)	0.042
Overweight		1.14 (0.80, 1.63)	0.472	1.09 (0.73, 1.64)	0.666	0.96 (0.72, 1.26)	0.757
Depression							
Abdominal weight spectrum							
Normal		Ref.		Ref.		Ref.	
Thinness		1.01 (0.52, 1.95)	0.974	0.94 (0.41, 2.15)	0.887	1.46 (0.90, 2.36)	0.122
Overweight		1.92 (1.26, 2.93)	0.003	2.17 (1.30, 3.63)	0.003	2.44 (1.82, 3.26)	< 0.001

Abbreviations: BRI, the Behavioral Regulation Index; GEC, the Global Executive Composite; MI, the Metacognition Index.

¹ Adjusted for social-demographic factors (i.e. age, sex, parental highest education, and family gross income) and individual lifestyle behaviors (i.e. screen exposure, nighttime sleep duration, and physical activity).

Figures

Figure 1

Flowchart of the participants.

Abbreviations: BMI, body mass index; BRIEF, the Behavior Rating Inventory of Executive Function; DASS, the Depression Anxiety and Stress Scales with 21 items; SCHEDULE-A, study of the Shanghai Children’s Health, Education and Lifestyle Evaluation-Adolescents; WHtR, waist-to-height ratio.

Figure 2

Moderating effects of negative emotions on the association between abdominal weight spectrum and executive function problems in adolescents.

Note: All models were adjusted for sociodemographic characteristics (i.e., age, sex, parental education level, and gross family income) and individual behaviors (screen time, night sleep duration, and physical activity), and the P values indicated whether each interaction term of negative emotions (i.e., depression, anxiety, and stress) and abdominal weight spectrum (i.e., abdominal thinness, and overweight) on executive dysfunction reached statistical significance.

Abbreviations: BRI, the Behavioral Regulation Index; GEC, the Global Executive Composite; MI, the Metacognition Index.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementary.docx](#)