Expression of long noncoding RNAs in the ovarian granulosa cells of women with diminished ovarian reserve using high-throughput sequencing

Li Dong

Shandong University of Traditional Chinese Medicine

Xin Xin

Shandong University of Traditional Chinese Medicine

Hsun-Ming Chang

University of British Columbia

Peter C.K. Leung

University of British Columbia

Chen Yu

Shandong University of Traditional Chinese Medicine

Fang Lian

Affiliated Hospital of Shandong University of Traditional Chinese Medicine

Haicui Wu (✉ haicui_w@163.com)

Affiliated Hospital of Shandong University of Traditional Chinese Medicine

Research Article

Keywords: Diminished ovary reserve, Ovarian granulosa cells, Long non-coding RNA, High-throughput sequencing

Posted Date: May 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1276715/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Infertility is a global reproductive-health problem, and diminished ovarian reserve (DOR) is one of the common causes of female infertility. Long noncoding RNAs (lncRNAs) are crucial regulators of numerous physiological and pathological processes in humans. However, whether lncRNAs are involved in the development of DOR remains to be elucidated.

Methods

Ovarian granulosa cells (OGCs) extracted from infertile women with DOR and from women with normal ovarian reserve (NOR) were subjected to high-throughput sequencing. Comprehensive bioinformatics analysis was conducted to identify the differential expression of messenger RNAs (mRNAs) and lncRNAs. Sequencing results were validated by the selection of random lncRNAs using real-time reverse transcription-quantitative polymerase chain reaction.

Results

Compared with the NOR group, a total of 244 lncRNAs were upregulated (53 known and 191 novel), and 222 lncRNAs were downregulated (36 known and 186 novel) in the DOR group. Similarly, 457 mRNAs had differential expression between the two groups. Of these, 169 were upregulated and 288 were downregulated. Bioinformatics analysis revealed that the differentially expressed genes of mRNA and lncRNAs were considerably enriched in “cell adhesion and apoptosis”, “steroid biosynthesis”, and “immune system”. A co-expression network comprising lncRNAs and their predicted target genes revealed the possible involvement of the “thyroid hormone signaling pathway” and “protein binding, digestion and absorption” in DOR pathogenesis. The expression of SLC16A10 was positively regulated by multiple lncRNAs. Seven differentially expressed lncRNAs (NEAT1, RAB5A, MEG3, GNG12, MST1L, GCN1, and ZEB2-AS1) were identified, among which the expression of NEAT1, GNG12 and ZEB2-AS1 was in accordance with RNA-sequencing.

Conclusions

We presented the first data showing that the expression profiles of lncRNA and mRNA in OGCs between NOR and DOR patients using RNA sequencing. The lncRNAs that we identified may serve as novel diagnostic biomarkers for patients with DOR.

Background

Infertility is a worldwide problem involving couples of reproductive ages. It has become the third most common disease after cancer and cardiovascular diseases [1]. As a serious medical problem, infertility can have destructive effects on the quality of life and marital satisfaction. As economic and social pressures increase, the age at which women have children continues to increase. The proportion of infertile women due to diminished ovarian reserve (DOR) is increasing gradually worldwide. The term of DOR was first described by Navot and colleagues in 1987 [2]. It is characterized by a decline in the quantity and/or quality of oocytes, with increased serum levels of follicle-stimulating hormone (FSH) and decreased serum levels of anti-Müllerian hormone (AMH) [3]. The main clinical manifestations are irregular menstruation, perimenopause syndrome, and infertility. The United States Assisted Reproductive Technology Population Survey revealed an increase in DOR prevalence from 19% in 2004 up to 26% in 2011 [3]. Studies have demonstrated that DOR typically develops into premature ovarian failure (POF) in 1–6 years [4].

Women with DOR usually result in poor pregnancy outcomes and present major challenges in reproductive medicine, particularly receiving *in vitro* fertilization/intracytoplasmic sperm injection (IVF/ICSI) treatment [5]. However, the etiology and pathogenesis of DOR are not clear. The key etiological factors are age, genetics, iatrogenicity (surgery, radiotherapy, or chemotherapy), and immune and environmental factors. Therefore, to improve pregnancy outcomes in women with DOR, the clinicians should focus on the molecular mechanisms underlying DOR occurrence. In this way, timely prediction of the diagnosis, necessary interventions, and therapeutic measures through identification of clinical biomarkers can be undertaken.

Long noncoding RNAs (lncRNAs) are functional RNA molecules of > 200 nucleotides in length that cannot be translated into proteins. Recently, studies have revealed that lncRNAs regulate the expression of genes in terms of transcriptional, posttranscriptional, and epigenetic modifications [6]. These actions may affect the growth, development, aging, and disease progression of organisms [7]. Additionally, lncRNAs are involved in the regulation of stem cell differentiation, embryonic development, cell proliferation, apoptosis, cell metabolism, and immune reaction [8, 9]. Recent studies have shown differential expression of lncRNAs in several gynecological disorders, such as endometriosis [10], premature ovarian insufficiency (POI) [11], and polycystic ovary syndrome (PCOS) [12]. Although several lncRNAs have been found to be involved in the abovementioned ovarian diseases, the association between lncRNAs and DOR is largely unknown.

Technologies based on transcription, gene chips, and high-throughput sequencing, have found that various lncRNAs are related to the occurrence, development, and treatment of diseases. The use of biological methods to study differences in the expression of lncRNAs between patients and healthy individuals and prescreening lncRNAs by high-throughput sequencing can aid in the discovery of new targets related to many diseases. In-depth study of the biomolecular mechanisms of lncRNAs can help explain the complexity and pathological states of diseases and support the diagnosis and treatment of refractory diseases.

In this study, we aimed to employ transcriptome analysis to investigate the biological role of lncRNAs during DOR progression. Ovarian granulosa cells (OGCs) from women with DOR and those from women with normal ovarian reserve (NOR) were obtained to perform RNA-sequencing (RNA-seq) and analyze the expression profiles between two groups. Bioinformatics analysis was undertaken to identify differentially expressed (DE) lncRNAs and messenger (m)RNAs and their associated functions and signaling pathways. In addition, we applied real-time reverse transcription-quantitative polymerase chain reaction (RT-qPCR) to verify the differential expression of seven lncRNAs. These aberrantly expressed lncRNAs may serve as potential biomarkers for the development of DOR and may introduce new routes for the diagnosis of women with POF.

Results

Baseline characteristics of patients used for sequencing

As presented in Table 1, patients in the DOR group had low AMH levels, low antral follicle count (AFC), low number of oocytes obtained and high basal FSH levels compared to the NOR group, the differences in the above characteristics being indicators of ovarian reserve ($P < 0.001$). No other features were significantly different ($P > 0.05$).

Table 1

Comparison of the baseline characteristics between the NOR group and DOR group used for sequencing

	NOR (n = 6)	DOR (n = 6)	P
Patient age (years)	30.50 ± 1.76	35.17 ± 4.88	0.068
Infertility duration (years)	4.50 ± 2.95	5.17 ± 2.40	0.677
Infertility type (%)			0.567
Primary	4/6 (66.7)	2/6 (33.3)	
Secondary	2/6 (33.3)	4/6 (66.6)	
BMI (kg/m ²)	22.72 ± 0.98	21.75 ± 1.33	0.181
AMH (ng/mL)	4.64 ± 1.11	0.79 ± 0.23	< 0.001
Basal FSH (mIU/mL)	6.99 ± 1.40	13.57 ± 2.04	< 0.001
AFC (n)	25.67 ± 2.73	6.83 ± 1.94	< 0.001
Previous IVF/ICSI attempts (n)	0.00 ± 0.00	1.17 ± 1.60	0.135
Number of oocytes retrieved	16.17 ± 3.49	4.00 ± 2.28	< 0.001
Data are the mean ± SD and proportion (%).			
AMH, anti-Müllerian hormone; AFC, antral follicle count; BMI, body mass index; FSH, follicle-stimulating hormone; ICSI, intracytoplasmic sperm injection; IVF, <i>in vitro</i> fertilization.			

Sequencing of complementary (c)DNA libraries and transcriptome profiles of OGCs

A total of 999,067,624 raw reads were acquired using the NovaSeq™ 6000 platform (Illumina, San Diego, CA, USA). After excluding adapter contamination, undetermined bases, and low-quality bases, 902,773,866 clean reads were obtained, accounting for 135.42 GB. Although the GC content of the clean data was ~ 42%, the quality scores of clean reads were > 99.9 and > 98.83% for Q20 and Q30, respectively. In summary, these results indicated that the reliability and quality of the sequencing data were sufficient for further analyses (Table 2). Over 96% of clean reads were mapped to the reference human genome using HISAT (<http://daehwankimlab.github.io/hisat2/>), including uniquely mapped reads from 84.10–86.76% (**Supplemental Table 1**).

Table 2
Statistical data of the reads for 12 cDNA libraries

Sample	Raw reads	Raw bases	Clean reads	Clean bases	Valid ratio (reads)	Q20%	Q30%	GC content (%)
DOR_1	95746384	14.36G	87242788	13.09G	91.12	99.99	98.96	42
DOR_2	90791244	13.62G	79121838	11.87G	87.15	99.99	98.88	43
DOR_3	72114510	10.82G	66912764	10.04G	92.79	99.99	98.96	41.50
DOR_4	70676974	10.60G	65727486	9.86G	93.00	99.99	98.97	41
DOR_5	91948332	13.79G	85217766	12.78G	92.68	99.99	98.93	42
DOR_6	81704846	12.26G	74061172	11.11G	90.64	99.99	98.83	44
NOR_1	88597962	13.29G	81949398	12.29G	92.50	99.99	98.91	42
NOR_2	67940108	10.19G	62960406	9.44G	92.67	99.99	98.93	41.50
NOR_3	66981682	10.05G	61871236	9.28G	92.37	99.99	98.98	42
NOR_4	88814944	13.32G	73435164	11.02G	82.68	99.99	98.91	43.50
NOR_5	90275194	13.54G	77710494	11.66G	86.08	99.99	98.88	43
NOR_6	93475444	14.02G	86563354	12.98G	92.61	99.99	98.97	42

Identification of lncRNAs and mRNAs

To further recognize the protein-coding ability of unknown transcripts, Coding Potential Calculator (CPC; <http://cpc2.gao-lab.org/>) and Coding-Non-Coding Index (CNCI; <https://github.com/www-bioinfo-org/CNCI/>) were designed to exclude those with coding potential. As a result, a total of 45,333 novel lncRNAs were detected from 12 cDNA libraries (**Supplemental Table 2**). These lncRNAs were dispersed evenly across the 46 human chromosomes (Fig. 1a). With regard to the genomic location of the lncRNAs, the types of lncRNAs were class_code “i” (intraintronic transcript, 72.24%), class_code “u” (intergenic transcript, 18.37%), class_code “o” (sense transcript, 3.55%), class_code “j” (bidirectional transcript, 3.25%), and class_code “x” (antisense transcript, 2.59%). These lncRNAs showed no apparent bias for genomic location (**Supplemental Fig. 1** and Fig. 1b). Additionally, a total of 47,520 known lncRNAs were identified between the two groups.

We compared lncRNAs with mRNAs to illustrate the structural features of lncRNAs. The expression and number of lncRNAs were comparable to those of mRNAs (Fig. 1c). Furthermore, the median length of lncRNA transcripts was 1504 bp, which was shorter than the median length of 2317 bp for mRNA transcripts. These findings suggest that these lncRNAs are shorter than the mRNAs (Fig. 1d). There were fewer exons in lncRNAs (mean, 3) than in mRNAs (mean, 9). In total, 68.86% of mRNAs had ≥ 5 exons,

whereas 77.02% of lncRNAs had ≤ 3 exons (Fig. 1e). Moreover, in OGCs, the predicted open reading frame (ORF) length of lncRNAs was shorter than that of mRNAs (Fig. 1f).

Expression profile of mRNAs

A total of 457 DE mRNA genes were found between the OGCs of the two groups. Of these genes, 169 genes were upregulated, and 288 genes were downregulated (Fig. 2a-b). To further explore the functions of these DE mRNA genes, analyses of functional enrichment were performed using the Gene Ontology (GO) (<http://geneontology.org/>) database. Among the 457 DE mRNA genes, 3404 GO terms with functional information were enriched (Fig. 2c-d). A total of 542 GO terms were enriched significantly ($P < 0.05$). For biological process (BP), 359 terms were found, including “cell adhesion” (GO:0007155), “positive regulation of apoptotic process” (GO:0043065) and “steroid biosynthetic process” (GO:0006694). For molecular function (MF), 137 terms were discovered, including “oxidoreductase activity” (GO:0016491). For “cellular component” (CC), 42 terms were found, including “endoplasmic reticulum lumen” (GO:0005788) (**Supplemental Table 3**).

We used the Kyoto Encyclopedia of Genes and Genomes (KEGG) (www.genome.jp/kegg/) database to identify which genes had significantly enriched signaling pathways. Nineteen significantly enriched signaling pathways related to the DE mRNA genes ($P < 0.05$) were found, including “steroid biosynthesis”, “focal adhesion”, “ECM–receptor interaction”, “PPAR signaling pathway”, and “PI3K-AKT signaling pathway” (Fig. 2e).

Expression profile of lncRNAs

We discovered that 466 lncRNAs had differential expression between the DOR group and NOR group. Of these, 244 were upregulated (53 known and 191 novel), and 222 were downregulated (36 known and 186 novel) (**Supplemental Table 4**, Fig. 3a-b). To further probe the functions of these lncRNAs, we forecasted the cis-regulated target genes of the DE lncRNAs between the DOR group and NOR group. A total of 52 lncRNAs were found to have target genes if 100 kilobase pairs (kbp) was used as the cutoff. Some lncRNAs had 2–3 target genes, and 57 probable lncRNA target genes were identified (**Supplemental Table 5**). Based on these cis-regulated target genes, we used the GO database to identify 76 important GO terms: 57 BP terms (e.g., “stem cell proliferation, GO:0072089), 6 CC terms (e.g., “proteasome activator complex”, GO:0008537), and 13 MF terms (e.g., “CXCR3 chemokine receptor binding”, GO:0048248) (Fig. 3c-d). Analyses of signaling pathway enrichment using the KEGG database revealed that these lncRNA target genes were primarily enriched in “ferroptosis”, “fatty acid biosynthesis”, and the “JAK-STAT signaling pathway” (Fig. 3e).

We discovered the predicted outcomes of the DE lncRNAs with cis-regulated genes. We listed the first-six and last-four lncRNA–gene pairs according to the Pearson correlation coefficient. The first-six lncRNA–gene pairs were regulated in the same direction, whereas the last-four lncRNA–gene pairs were in the opposite direction (Table 3).

Table 3
Differentially expressed lncRNA–gene pairs between the DOR group and NOR group.

Gene	lncRNA transcript	Cis location (bp)	Pearson correlation coefficient
CXCL10	MSTRG.57192.1	1K	1.00
TNFSF15	MSTRG.77544.1	100K	1.00
PGBD5	MSTRG.9312.2	100K	1.00
CNTN3	MSTRG.52583.1	100K	0.99
SLC16A10	MSTRG.66173.1	10K	0.99
SLC16A10	MSTRG.66167.1	100K	0.99
BNIP2	MSTRG.27592.2	100K	-0.31
SYNCRIP	ENST00000656092	100K	-0.32
NPIP15	MSTRG.31257.1	10K	-0.33
C2CD2	MSTRG.48668.1	100K	-0.35

Coenriched GO terms of the DE lncRNAs and mRNAs

We then aimed to identify the key pathways that regulate the ovarian reserve. Hence, we identified five significantly enriched GO terms in the enrichment of the DE mRNAs and the enrichment of the lncRNA target genes. The significantly enriched GO terms were “CXCR chemokine receptor binding”, “CXCR3 chemokine receptor binding”, “positive regulation of transforming growth factor-beta production”, “regulation of bile acid biosynthetic processes”, and “cellular response to organonitrogen compound”. Three pathways were involved in BP. and the other two pathways were involved in MF (Table 4).

Table 4
Coenriched GO terms of DE lncRNAs and mRNAs.

GO Term	GO function	P
GO:0048248 (CXCR3 chemokine receptor binding)	molecular_function	0.002
GO:0071636 (positive regulation of transforming growth factor beta production)	biological_process	0.005
GO:0045236 (CXCR chemokine receptor binding)	molecular_function	0.010
GO:0070857 (regulation of bile acid biosynthetic process)	biological_process	0.027
GO:0071417 (cellular response to organonitrogen compound)	biological_process	0.034

Construction of a co-expression network of DE lncRNAs and mRNAs

We performed comprehensive analyses of the lncRNA–mRNA regulatory network to further explore the potential regulatory mechanism. Co-expression analyses between the DE mRNA genes and lncRNAs were performed first. Twenty-four positively correlated lncRNA–mRNA co-expression pairs containing 24 DE lncRNAs and 19 DE mRNAs were acquired at the verge of a Pearson correlation coefficient ($r > 0.40$) (**Supplemental Table 6**). Then, a lncRNA–mRNA regulatory network comprising dysregulated lncRNAs and their cis-target mRNAs was visualized using Cytoscape v3.7.1 (<https://cytoscape.org/>). Solute carrier family 16 member 10 (SLC16A10) was included in this network and was regulated by five lncRNAs: MSTRG.66173, MSTRG.66167, MSTRG.66177, MSTRG.66174, and MSTRG.66170 (Fig. 4). Among them, SLC16A10 was enriched in several GO terms, including “protein binding” (GO:0005515), “thyroid hormone generation” (GO:0006590), and “cell junction” (GO:0030054). SLC16A10 was also enriched in the “thyroid hormone signaling pathway” and “protein digestion and absorption”.

Verification of the DE lncRNAs by RT–qPCR

To verify the RNA-seq results, RT–qPCR was performed to examine seven of the identified DE lncRNAs between the OGCs from the DOR group and NOR group. Table 5 displayed the baseline characteristics of the case used for RT–qPCR validation, with patients in the DOR group not significantly different from those in the NOR group ($P > 0.05$), except for indicators related to ovarian reserve such as AMH level, AFC, and number of oocytes retrieved ($P < 0.05$). These seven lncRNAs include nuclear rich abundant transcript 1 (NEAT1), Ras-related protein in brain 5A (RAB5A), maternally expressed gene 3 (MEG3), guanine nucleotide-binding protein subunit gamma-12 (GNG12), macrophage stimulating 1 like (MST1L), general control nonderepressible 1 (GCN1), zinc-finger E-box binding homeobox 2-AS1 (ZEB2-AS1). Table 6 presented information on the relevant sequencing results for the above lncRNAs.

Table 5

Comparison of the baseline characteristics between the NOR group and DOR group used for RT-qPCR

	NOR (n = 6)	DOR (n = 6)	P
Patient age (years)	30.50 ± 2.88	35.50 ± 4.72	0.057
Infertility duration (years)	3.00 ± 1.67	3.50 ± 2.43	0.687
Infertility type (%)			1.000
Primary	2/6 (33.3)	1/6 (16.7)	
Secondary	4/6 (66.7)	5/6 (83.3)	
BMI (kg/m ²)	20.20 ± 2.61	21.52 ± 2.86	0.424
AMH (ng/mL)	4.13 ± 0.59	0.78 ± 0.22	< 0.001
Basal FSH (mIU/mL)	6.41 ± 1.48	11.02 ± 4.42	0.051
AFC (n)	26.00 ± 2.68	7.00 ± 2.10	< 0.001
Previous IVF/ICSI attempts (n)	0.67 ± 1.21	0.67 ± 0.82	1.000
Number of oocytes retrieved	16.50 ± 7.79	4.17 ± 1.84	0.011
Data are the mean ± SD and proportion (%).			
AMH, anti-Müllerian hormone; AFC, antral follicle count; BMI, body mass index; FSH, follicle-stimulating hormone; ICSI, intracytoplasmic sperm injection; IVF, <i>in vitro</i> fertilization.			

Table 6

Sequencing information of lncRNAs used for RT-qPCR verification

Transcript name	Gene name	log ₂ (fold_change)	Regulation	P
MSTRG.15237.9	NEAT1	-4.69	Down	0.041834004
MSTRG.50996.10	RAB5A	-3.07	Down	0.003518952
MSTRG.25915	MEG3	-3.83	Down	0.176775112
MSTRG.5280.1	GNG12	-4.85	Down	0.0003983567
MSTRG.3657.2	MST1L	-4.88	Down	0.000306066
MSTRG.20933.1	GCN1	5.98	Up	0.000182295
MSTRG.43755	ZEB2-AS1	-13.50	Down	0.000166045

Of these, the expression of NEAT1, GNG12, and ZEB2-AS1 was consistent with the RNA-seq results. The expression of the remaining four lncRNAs was not significantly different but showed a consistent trend

with the RNA-seq data (Fig. 5). These results indicate that the seq-RNA data were reliable.

Discussion

Follicular reserve and ovarian function are measures of reproductive endocrine functionality and fertility potential [13]. OGCs play important roles in the regulation of follicle activation, growth, development, and atresia [14]. Recently, several studies targeting specific genes and regulating various signaling pathways have shown that lncRNAs have vital roles in the formation and development of ovarian follicles [15–17].

We report, for the first time, the expression profiles of lncRNAs and mRNAs in OGCs of DOR patients and NOR patients, as identified by high-throughput sequencing. We identified 466 DE lncRNAs and 457 mRNAs associated with ovarian reserve. We showed that these dysregulated genes were involved in multiple biological processes, including cell adhesion and apoptosis, the immune system, and signaling pathways such as steroid biosynthesis. Notably, multiple lncRNAs could contribute to DOR by targeting and regulating signaling pathways mediated by SLC16A10, such as thyroid hormone synthesis and protein digestion and absorption. Overall, our results indicate the potential regulatory role of related lncRNAs and mRNAs in DOR.

In recent years, several studies have identified an important role for lncRNAs in the regulation of follicle development [18, 19]. Studies have identified differential expression of lncRNAs associated with reproductive function by using RNA-seq of OGCs obtained from humans [20], pigs [21], and chickens [22]. Most of those studies have focused on PCOS [23], endometriosis [24, 25], and POF [26, 27]. The lncRNAs we identified had fewer exons, shorter ORFs, and shorter transcript lengths compared with those in protein-coding transcripts, and these characteristics were similar to those from porcine OGCs [28] and human placentas at term [29]. These data demonstrate that the lncRNA outcomes we obtained were reliable.

Several research teams have reported aberrant lncRNA expression in POI and identified hundreds of lncRNAs associated with POI. Some of these lncRNAs could be used as biomarkers for the diagnosis or prognosis of DOR [16, 30]. The discrepancies between the DE lncRNAs identified in our study and those in other studies are due (at least in part) to differences in sample selection and sequencing methods. Moreover, several researchers have conducted functional studies on these DE lncRNAs [31–33] and reported different underlying molecular mechanisms. However, most of those studies concentrated on a particular gene or signaling pathway and did not analyze them from a holistic perspective. Additionally, most of the identified lncRNAs and mRNAs were associated with proliferative and apoptotic processes. The DE genes identified in our study were mostly involved in the signaling pathways related to the adhesion, proliferation, and apoptosis of cells, steroid biosynthesis and metabolism, and the immune response. All of these are biological processes involved in follicular atresia induced by OGC apoptosis [34, 35].

Folliculogenesis is regulated by multiple mechanisms involving endocrine and intraovarian signaling pathways [36, 37], and changes caused by the differential expression of genes may further affect ovarian

reserve. We revealed several DE lncRNAs that could be associated with ovarian reserve (NEAT1, MEG3, GNG12, MST1L, RAB5A, and ZEB2-AS1), all of which have vital roles in the proliferation and apoptosis of cells and merit further investigation. Among these lncRNAs, NEAT1 is a relatively well-studied lncRNA involved in the formation and maintenance of a “nuclear paraspeckle” (a nuclear body with multiple gene-expression functions). NEAT1 can promote disease through regulation of mitochondrial function [38]. NEAT1 is expressed abundantly in oocytes as well as in primordial, primary, and small antral follicles. It is involved mainly in the expression of genes related to apoptosis and extracellular matrix-related functions to achieve epigenetic control of follicular development [39, 40]. Interference with NEAT1 alleviates the pathological changes of PCOS by reducing the expression level of insulin-like growth factor 1. Therefore, it suppresses OGC apoptosis and enhances cell proliferation [41]. Some studies have analyzed publicly available RNA-seq data and found that NEAT1 and MEG3 (which are expressed abundantly in oocytes) are essential for oocyte growth [39], which we also discovered in this study. MEG3 can inhibit the proliferation of OGCs and POF through the p53-p66Shc pathway [15], which differs from our findings. Therefore, further studies are needed to confirm the mechanism of action of MEG3 in oocytes. RAB5A-positive vesicles modulate the localization and levels of centromere protein F at centromeres, likely through interaction with nuclear lamina. Therefore, an appropriate spindle length and kineto-microtubule attachment are ensured in meiotic oocytes [42]. GNG12 [43] and ZEB2-AS1 [44, 45] can promote the growth of tumor cells by regulating the proliferation, apoptosis, and migration of cells. MST1L [46] can regulate disease development through an inflammatory response pathway. GNG12, ZEB2-AS1, and MST1L tended to have low expression in the OGCs from DOR patients. Such low expression level may contribute to DOR development by increasing apoptosis and inhibiting cell proliferation in OGCs. In particular, GNG12 and ZEB2-AS1 have been insufficiently investigated with respect to follicle growth. However, we observed significant downregulation of GNG12 and ZEB2-AS1 expression by RNA-seq and RT-qPCR, suggesting that GNG12 and ZEB2-AS1 may be closely related to DOR function.

We assessed the enrichment of functions and signaling pathways using the GO database and KEGG database, respectively, to reveal the biological functions of the DE lncRNA target genes and mRNAs. The DE genes were involved mainly in cell adhesion, steroid biosynthesis, and pathways related to immunity and apoptosis, which are the functional pathways of OGCs [34, 47, 48]. Cell adhesion is involved in the in vitro culture of porcine [49] and human [50] OGCs and can affect the proliferative potential and survival capability. Pashaiasl and collaborators measured the mRNA expression profile of the OGCs of DOR patients. Similar to our findings, they identified DE genes involved in focal adhesion [51], as well as DE genes involved in ossification, ovarian-follicle development, vasculogenesis, and sequence-specific activity of DNA-binding transcription factors. OGCs are the most important somatic cells for the synthesis of steroid hormones. We found that the DE genes could affect steroid biosynthesis and thus regulate the proliferation and apoptosis of OGCs through multiple pathways. These findings have been documented in studies in sheep [52], mice [53] and women with POF [54].

Autoimmunity is responsible for approximately 4–30% of POI cases [55]. Women suffering from POI carry a high risk of having autoimmune disorders [56]. Chemokines play an important role in reproductive immunology [57] and regulate leukocyte migration by attracting cells that express their cognate ligands.

C-X-C motif chemokine ligand 12 (CXCL12), C-X-C motif chemokine receptor 4 (CXCR4), and CXCR7 can influence the development of endometriosis, Asherman's syndrome, endometrial cancer, and ovarian cancer [58]. CXCR3, which was identified in the present study, has been investigated less thoroughly. Therefore, women should be screened for common autoantibodies (e.g., steroid cell, anti-ovarian, and anti-thyroid autoantibodies), which can aid in the early diagnosis and treatment of DOR and the prevention of POF. A prospective cohort study involving 164 women found a higher prevalence of OGC apoptosis, worse ovarian response, lower oocyte count, lower embryo count, and poor pregnancy outcomes in patients with DOR [59].

We found that the phosphoinositide 3-kinase/protein kinase B (PI3K-AKT) and Janus kinase/signal transducer and activator of transcription (JAK-STAT) pathways enriched by the DE genes were associated with apoptotic processes. Several studies have demonstrated that the inhibition of the PI3K-AKT pathway leads to apoptosis and autophagy in OGCs [53, 54]. The importance of the JAK/STAT signaling pathway in ovarian development and folliculogenesis in horses [60] and humans [61] has been demonstrated. Polyfluoroalkyl substances reduce the ovarian reserve and decrease endogenous hormone synthesis by activating the peroxisome proliferator-activated receptor (PPAR) pathway [62], which has been indicated as a new biomarker of follicular capacity [63]. By constructing a lncRNA-mRNA co-expression network, we found that SLC16A10 was positively regulated by multiple lncRNAs. Additionally, the function of the target gene SLC16A10 was enriched mainly in thyroid hormone synthesis as well as the binding, digestion, and absorption processes of proteins. One retrospective study found that increased levels of thyrotropic hormone may be associated with decreased serum levels of AMH [64], whereas the ovarian reserve was not associated with anti-thyroid peroxidase antibodies or anti-thyroglobulin antibodies. Transplantation of human amniotic epithelial cells has been shown to restore ovarian function and improve ovarian reserve and fertility in cyclophosphamide-induced POI rats by affecting protein digestion/absorption, and steroid-biosynthesis pathways [65].

Conclusions

The expression profiles of lncRNAs and mRNAs in OGCs from women with DOR and NOR were mined using transcriptome sequencing. A total of 466 lncRNAs and 457 mRNAs were identified to have differential expression in the transcripts. Bioinformatic analysis revealed that most of these lncRNAs and mRNAs were involved in important pathways and biological processes related to cell adhesion, apoptosis, steroid biosynthesis, and the immune system. Dysregulated lncRNAs could serve as diagnostic biomarkers. Further studies should concentrate on the elucidation of specific molecular mechanisms to provide a new understanding for the diagnosis and treatment of DOR.

Methods

Ethical approval of the study protocol

The research protocol was approved by the Reproductive Medicine Ethics Committee of the Affiliated Hospital of Shandong University of Traditional Chinese Medicine (Ref: SZ2018090103). All patients provided written informed consent before sample collection.

Patients and sample collection

Samples and clinical data were collected at the Center for Integrative Reproduction and Genetics, Affiliated Hospital of Shandong University of Traditional Chinese Medicine, from September 2018 to October 2020. Twenty-four patients were recruited in our study. These patients underwent IVF/ICSI using a gonadotrophin-releasing hormone antagonist for controlled ovarian hyperstimulation. All patients satisfied the following criteria: age of 25–40 years; body mass index (BMI) of 18–25 kg/m²; and no endocrine disease, uterine anomaly, endometriosis, or chromosomal abnormalities.

The study group (n = 12) comprised infertile women with DOR. The diagnostic criteria adopted an amended version of the Bologna standard [66], with participants fulfilling a minimum of two of the following conditions: (i) bilateral AFC < 6; (ii) AMH < 1.10 ng/mL; (iii) menstrual basal FSH level of 10–40 mIU/mL. The control group (n = 12) was women with NOR and pure male factor (azoospermia or severe oligospermia/aspermia) infertility. Six patient samples from each group were selected randomly for lncRNA sequencing. The other six samples underwent RT–qPCR.

OGCs were collected and purified by density-gradient centrifugation from follicular fluid, which was obtained by ultrasound-guided transvaginal ovarian puncture for oocyte retrieval from the first aspiration follicle of each ovary. Oocyte retrieval was performed 36 h after injection of human chorionic gonadotropin once follicles > 17 mm in diameter. Follicular fluid was centrifuged at 2000 rpm for 5 min at room temperature and the supernatant was removed. Phosphate-buffered saline (5 mL) was added to the precipitate and mixed. Then, 5 mL of Ficoll-Paque (GE Healthcare, Chicago, IL, USA) was added to a 15-mL centrifuge tube. The suspension was added slowly to the upper layer of Ficoll-Paque, and centrifugation at 2000 rpm for 20 min at room temperature was carried out. The OGC layer was aspirated and transferred to a 1.5-mL centrifuge tube, and repeated pipetting was undertaken followed by centrifugation at 2000 rpm for 3 min at room temperature. Finally, the supernatant was removed and the OGCs were stored in a – 80°C for further analyses.

Preparation and sequencing of RNA libraries

TRIzol® Reagent (Invitrogen, Carlsbad, CA, USA) was employed to isolate and purify total RNA according to the manufacturer's instructions. The amount and purity of total RNA in each sample was analyzed using a bioanalyzer (2100 series; Agilent Technologies, Santa Clara, CA, USA) and an RNA Nano1000 Assay Kit (Agilent Technologies) so that the RNA integrity number (RIN) > 7.0. Ribosomal RNA was removed according to the instructions of the Ribo-Zero™ rRNA Removal Kit (Illumina). Under a high temperature, the remaining RNA was broken into short fragments using divalent cations. Subsequently, cDNA was created by reverse transcription using fragmented RNA as a template, followed by addition of

deoxynucleoside triphosphate, RNase H, and *Escherichia coli* DNA polymerase I to synthesize the second strand. After purification of AMPure XP beads (Beckman Coulton, Fullerton, CA, USA) and end repair, poly(A) was added, and sequencing connectors were attached. Then, cDNA of a certain length range was extracted. PCR amplification was performed to obtain a cDNA library. The average insert size of the final cDNA library was 300 ± 50 bp. Twelve separate RNA-seq libraries were generated for the study group and control group.

RNA-seq

High-throughput sequencing of samples was executed on a NovaSeq 6000 using 2×150 -bp paired-end sequences (PE150) according to a standard procedure. First, reads containing adapter contamination, low-quality bases, or unidentified bases were excluded by applying cutadapt v1.9 (<https://cutadapt.readthedocs.io/en/stable/#>). Then, the quality of reads was authenticated using FastQC v0.10.1 (www.bioinformatics.babraham.ac.uk/projects/fastqc/). The Q20, Q30, and GC contents of clean data were calculated, and subsequent analyses were based on these data. Sequencing and data acquisition were undertaken by LC-Bio Technology (Hangzhou, China).

Transcript assembly

HISAT (v2.0.4) was utilized to preprocess raw data and map the processed valid data to the human reference genome (GRCh38). Annotation files in Gene Transfer Format were created to identify which genes these reads mapped to. The mapped reads from each sample were assembled using StringTie v1.3.0 (<https://ccb.jhu.edu/software/stringtie/index.shtml/>) with default parameters. Then, all transcripts in the sample were joined using gffcompare (<https://ccb.jhu.edu/software/stringtie/gffcompare.shtml>) to build a whole transcriptome. Then, StringTie was used to calculate fragments per kilobase of exon model per million mapped reads (FPKM) to quantify the expression of the transcripts.

lncRNA identification

First, transcripts that overlapped with known mRNAs and transcripts shorter than 200 bp were removed. If the remaining transcripts duplicated the lncRNA, then these transcripts were deemed to be “known lncRNAs”. The rest of the non-annotated transcripts were assumed to recognize the underlying novel lncRNAs. Then, we employed CPC (0.9-r2) and CNCI (v2.0) to forecast the coding potential of the remaining transcripts of length ≥ 200 bp, mapped read coverage ≥ 3 , and exon number ≥ 1 . Transcripts with a CPC score < -1 and CNCI score < 0 were assumed to be potential novel lncRNAs.

Differential expression of mRNAs and lncRNAs

StringTie was adopted to evaluate the expression of mRNAs and lncRNAs by calculation of FPKM. Multiple values were analyzed by the packages -edgeR or DESeq2 within R (R Institute for Statistical Computing, Vienna, Austria). $P < 0.05$ and $|\log_2(\text{fold change})| \geq 1$ were used as the basis for screening DE mRNAs and lncRNAs between the DOR group and NOR group.

Target-gene prediction and functional analyses of lncRNAs

The interaction between lncRNAs and their adjacent genes is known as “*cis*-regulation”. We used Python scripts to predict the *cis*-regulatory relationships between mRNAs and lncRNAs in the 100-kbp range upstream and downstream of chromosomes. We undertook analyses of enrichment of the functions and signaling pathways of lncRNA-targeted mRNAs using GO and KEGG databases, respectively.

Validation of lncRNA expression by RT-qPCR

Based on analysis of sequencing results and literature review, the expression of seven lncRNAs (GNG12, MEG3, NEAT1, MST1L, RAB5A, GCN1 and ZEB2-AS1) was measured by RT-qPCR in OGC samples from 12 patients to validate the RNA-seq results. Total RNA was extracted from samples using the RNeasy Micro Kit following the manufacturer’s instructions (Qiagen, Stanford, VA, USA). The content and purity of RNA were measured at an absorbance of 260/280 nm using Scandrop™ 100 (Analytik Jena, Jena, Germany). Total RNA was reverse-transcribed into cDNA using the TURScript First Strand cDNA Synthesis Kit (Aidlab, Beijing, China) according to the manufacturer’s instructions. The specific primers designed for RT-qPCR using Primer-Blast (National Center for Biotechnology Information) and primer-primer6 together are listed in Table 7. The analytikjena-qTOWER2.2 PCR System (Analytik Jena) was applied to conduct RT-qPCR. The reaction protocol was as follows: initial denaturation at 95°C for 3 min, denaturing at 95°C for 10 s, and annealing at 60°C for 30 s for 40 cycles. The expression of glyceraldehyde-3-phosphate dehydrogenase (an endogenous reference transcript) was used to quantify relative lncRNA expression using the $2^{-\Delta\Delta C_t}$ method. For each reaction, three independent biological replicates were employed.

Table 7
Primer sequences used for RT–qPCR

Primer	Sequence (5'–3')
GAPDH-F	TGCACCACCAACTGCTTAGC
GAPDH-R	GGCATGGACTGTGGTCATGAG
GNG12-F	TCCCTGTGTCTTTCTTCTC
GNG12-R	TAGCCTATGTGCCTCTTG
MEG3-F	TGGCATAGAGGAGGTGAT
MEG3-R	AGGTGAGGAAGGAAGACA
NEAT1-F	AGTGTGAGTCCTAGCATTG
NEAT1-R	GAACCTCCTCCTCCTAAGC
RAB5A-F	CGGTAACGAGTCCATTGTAG
RAB5A-R	CCTTGATGAGTGCTGTATCC
ZEB2-AS1-F	CTGGATAGCAAAGGACACC
ZEB2-AS1-R	GTTTCTTCGCTTCCACCT
MST1L-F	GAGCCTGGAAATGGTTGA
MST1L-R	AAGTTCTCCTCCAGTTGTG
GCN1-F	AAGGGTAGGAAAGTCTGGTTA
GCN1-R	ATTGAGAATCGAACCTGTTATCA

Statistical analyses

SPSS 23.0 (IBM, Armonk, NY, USA) was used for statistical analyses. Prism 8.0.1 (GraphPad, San Diego, CA, USA) was employed for statistical analyses of RT–qPCR results and graphs. Measurement data are expressed as the mean $\pm$ SD. Student's t test was undertaken to determine a discrepancy in the expression of lncRNAs between the DOR group and NOR group. $P < 0.05$ was deemed significant.

Abbreviations

AFC
Antral follicle count
AMH
Anti- Müllerian hormone
BMI
Body Mass Index

BP
Biological process
CC
Cellular component
cDNA
Complementary DNA
CNCI
Coding-Non-Coding Index
CPC
Coding Potential Calculator
DE
Differentially expressed
DOR
Diminished ovarian reserve
FPKM
Fragments per kilobase of exon model per million mapped reads
FSH
Follicle-stimulating hormone
GO
Gene Ontology
ICSI
Intracytoplasmic sperm injection
IVF
In vitro fertilization
KEGG
Kyoto Encyclopedia of Genes and Genomes
lncRNAs
Long noncoding RNAs
MF
Molecular function
NOR
Normal ovarian reserve
OGCs
Ovarian granulosa cells
ORF
Open reading frame
PCOS
Polycystic ovary syndrome
POF
Premature ovarian failure

POI
Premature ovarian insufficiency
PPAR
Peroxisome proliferator-activated receptor
RT–qPCR
Reverse transcription-quantitative polymerase chain reaction
RNA-seq
RNA-sequencing.

Declarations

Ethics approval and consent to participate

The study was conducted in accordance with the Declaration of Helsinki, and approved by the Ethics Committee of the Reproductive Medicine Ethics Committee of the Affiliated Hospital of the Shandong University of Traditional Chinese Medicine (Ref. SZ2018090103). Written informed consent was obtained from all the participants.

Consent for publication

Not applicable.

Availability of Data and materials

All the raw data has been stored in GEO (Gene Expression Omnibus) with the login number GSE193136 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE193136>).

Competing interests

The authors declare that they have no commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

This research was funded by Natural Science Foundation of Shandong Province (NO. ZR2021MH255), National Natural Science Foundation of China (NO. 81703958, 81774355 and 81974577) and Key Technology Research and Development Program of Shandong (NO. 2019GSF108067).

Authors' Contributions

FL and HW conceived and designed this study. LD contributed to the acquisition and analyses of data. LD and XX contributed to drafting the manuscript. CY was responsible for sample collection. HMC and PCKL contributed to manuscript revision. All authors read and approved the final submitted version of this manuscript.

Acknowledgments

The authors thank the nurses and laboratory staff of the Department of Assisted Reproduction of the Affiliated Hospital of Shandong University of Traditional Chinese Medicine for their technical assistance.

References

1. Qi Z, Liu Y, Yang H, Yang X, Wang H, Liu B, Yuan Y, Wang G, Xu B, Liu W, Xu Z and Deng Y (2021) Protective role of mA binding protein YTHDC2 on CCNB2 in manganese-induced spermatogenesis dysfunction. *Chemico-biological interactions*:109754. doi: 10.1016/j.cbi.2021.109754
2. Navot D, Rosenwaks Z and Margalioth E (1987) Prognostic assessment of female fecundity. *Lancet* (London, England) 2:645–7. doi: 10.1016/s0140-6736(87)92439-1
3. Devine K, Mumford SL, Wu M, DeCherney AH, Hill MJ and Propst A (2015) Diminished ovarian reserve in the United States assisted reproductive technology population: diagnostic trends among 181,536 cycles from the Society for Assisted Reproductive Technology Clinic Outcomes Reporting System. *Fertil Steril* 104:612 – 19.e3. doi: 10.1016/j.fertnstert.2015.05.017
4. Balmaceda J and Schwarze J (2007) Assessment of ovarian reserve - should we perform tests of ovarian reserve routinely? *Human reproduction* (Oxford, England) 22:1492; author reply 1492-3. doi: 10.1093/humrep/del509
5. Cohen J, Chabbert-Buffet N and Darai E (2015) Diminished ovarian reserve, premature ovarian failure, poor ovarian responder—a plea for universal definitions. *Journal of assisted reproduction and genetics* 32:1709–12. doi: 10.1007/s10815-015-0595-y
6. Ponting C and Belgard T (2010) Transcribed dark matter: meaning or myth? *Human molecular genetics* 19:R162-8. doi: 10.1093/hmg/ddq362
7. Kumar M and Goyal R (2017) LncRNA as a Therapeutic Target for Angiogenesis. *Current topics in medicinal chemistry* 17:1750–1757. doi: 10.2174/1568026617666161116144744
8. Kong Y, Lu Z, Liu P, Liu Y, Wang F, Liang E, Hou F and Liang M (2019) Long Noncoding RNA: Genomics and Relevance to Physiology. *Comprehensive Physiology* 9:933–946. doi: 10.1002/cphy.c180032
9. McDonel P and Guttman M (2019) Approaches for Understanding the Mechanisms of Long Noncoding RNA Regulation of Gene Expression. *Cold Spring Harbor perspectives in biology* 11. doi: 10.1101/cshperspect.a032151
10. Cui D, Ma J, Liu Y, Lin K, Jiang X, Qu Y, Lin J and Xu K (2018) Analysis of long non-coding RNA expression profiles using RNA sequencing in ovarian endometriosis. *Gene* 673:140–148. doi: 10.1016/j.gene.2018.06.046
11. Pankiewicz K, Laudański P and Issat T (2021) The Role of Noncoding RNA in the Pathophysiology and Treatment of Premature Ovarian Insufficiency. *International journal of molecular sciences* 22. doi: 10.3390/ijms22179336

12. Huang X, Hao C, Bao H, Wang M and Dai H (2016) Aberrant expression of long noncoding RNAs in cumulus cells isolated from PCOS patients. *Journal of assisted reproduction and genetics* 33:111–21. doi: 10.1007/s10815-015-0630-z
13. Jiang Z, Jin L, Shi W, Xi J, Hu Y, Liu X, Sheng J, Ding G and Huang H (2019) A combination of follicle stimulating hormone, estradiol and age is associated with the pregnancy outcome for women undergoing assisted reproduction: a retrospective cohort analysis. *Science China. Life sciences* 62:112–118. doi: 10.1007/s11427-018-9322-6
14. Chen S, Wang Y, Liao L, Meng L, Li J, Shi C, Han H, Zheng X and Shen H (2020) Similar Repair Effects of Human Placenta, Bone Marrow Mesenchymal Stem Cells, and Their Exosomes for Damaged SVOG Ovarian Granulosa Cells. *Stem cells international* 2020:8861557. doi: 10.1155/2020/8861557
15. Xiong Y, Liu T, Wang S, Chi H, Chen C and Zheng J (2017) Cyclophosphamide promotes the proliferation inhibition of mouse ovarian granulosa cells and premature ovarian failure by activating the lncRNA-Meg3-p53-p66Shc pathway. *Gene* 596:1–8. doi: 10.1016/j.gene.2016.10.011
16. Yao G, He J, Kong Y, Zhai J, Xu Y, Yang G, Kong D, Dong F, Shi S, Yang Q and Sun Y (2019) Transcriptional profiling of long noncoding RNAs and their target transcripts in ovarian cortical tissues from women with normal menstrual cycles and primary ovarian insufficiency. *Molecular reproduction and development* 86:847–861. doi: 10.1002/mrd.23158
17. Zhao W and Dong L (2018) Long non-coding RNA HOTAIR overexpression improves premature ovarian failure by upregulating Notch-1 expression. *Experimental and therapeutic medicine* 16:4791–4795. doi: 10.3892/etm.2018.6750
18. Bouckenheimer J, Assou S, Riquier S, Hou C, Philippe N, Sansac C, Lavabre-Bertrand T, Combes T, Lemaître JM, Boureux A and De Vos J (2016) Long non-coding RNAs in human early embryonic development and their potential in ART. *Hum Reprod Update* 23:19–40. doi: 10.1093/humupd/dmw035
19. Tu J, Chen Y, Li Z, Yang H, Chen H and Yu Z (2020) Long non-coding RNAs in ovarian granulosa cells. *J Ovarian Res* 13:63. doi: 10.1186/s13048-020-00663-2
20. Zheng Y, Bian Y, Wu R, Chen W, Fu L, Li P, Wang Y, Yang X, Zhao S and Shi Y (2021) High-Throughput Sequencing Profiles About lncRNAs and mRNAs of Ovarian Granulosa Cells in Polycystic Ovary Syndrome. *Frontiers in medicine* 8:741803. doi: 10.3389/fmed.2021.741803
21. Meng L, Zhao K, Wang CC, Tao J, Wu Z, Teerds K and Zhang S (2021) Characterization of Long Non-Coding RNA Profiles in Porcine Granulosa Cells of Healthy and Atretic Antral Follicles: Implications for a Potential Role in Apoptosis. *Int J Mol Sci* 22. doi: 10.3390/ijms22052677
22. Wang Y, Shi H, Zhang G, Wu P, Chen L, Shen M, Li T, Lv X, Gu Y and Wang J (2021) Transcriptome Analysis of Long Noncoding RNAs and mRNAs in Granulosa Cells of Jinghai Yellow Chickens Illuminated With Red Light. *Front Genet* 12:563623. doi: 10.3389/fgene.2021.563623
23. Wang L, Fan H, Zou Y, Yuan Q, Hu X, Chen X, Zhu C, Zhang X and Cui H (2020) Aberrant Expression of Long Non-coding RNAs in Exosomes in Follicle Fluid From PCOS Patients. *Front Genet* 11:608178. doi: 10.3389/fgene.2020.608178

24. Khalaj K, Miller JE, Lingegowda H, Fazleabas AT, Young SL, Lessey BA, Koti M and Tayade C (2019) Extracellular vesicles from endometriosis patients are characterized by a unique miRNA-lncRNA signature. *JCI Insight* 4. doi: 10.1172/jci.insight.128846
25. Wu J, Huang H, Huang W, Wang L, Xia X and Fang X (2020) Analysis of exosomal lncRNA, miRNA and mRNA expression profiles and ceRNA network construction in endometriosis. *Epigenomics* 12:1193–1213. doi: 10.2217/epi-2020-0084
26. Guo S, Quan S and Zou S (2021) Roles of the Notch Signaling Pathway in Ovarian Functioning. *Reprod Sci* 28:2770–2778. doi: 10.1007/s43032-021-00610-6
27. Zheng C, Liu S, Qin Z, Zhang X and Song Y (2021) lncRNA DLEU1 is overexpressed in premature ovarian failure and sponges miR-146b-5p to increase granulosa cell apoptosis. *J Ovarian Res* 14:151. doi: 10.1186/s13048-021-00905-x
28. Ruszkowska M, Nynca A, Pauksztó L, Sadowska A, Swigonska S, Orlowska K, Molcan T, Jastrzebski JP and Ciereszko RE (2018) Identification and characterization of long non-coding RNAs in porcine granulosa cells exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Anim Sci Biotechnol* 9:72. doi: 10.1186/s40104-018-0288-3
29. Majewska M, Lipka A, Pauksztó L, Jastrzebski JP, Gowkielewicz M, Jozwik M and Majewski MK (2018) Preliminary RNA-Seq Analysis of Long Non-Coding RNAs Expressed in Human Term Placenta. *Int J Mol Sci* 19. doi: 10.3390/ijms19071894
30. Dang Y (2016) The role and mechanism of non-coding RNA in primary ovarian insufficiency. Ph.D. thesis. Shandong University.
31. Li D, Wang X, Dang Y, Zhang X, Zhao S, Lu G, Chan WY, Leung PCK and Qin Y (2021) lncRNA GCAT1 is involved in premature ovarian insufficiency by regulating p27 translation in GCs via competitive binding to PTBP1. *Mol Ther Nucleic Acids* 23:132–141. doi: 10.1016/j.omtn.2020.10.041
32. Li D, Xu W, Wang X, Dang Y, Xu L, Lu G, Chan WY, Leung PCK, Zhao S and Qin Y (2021) lncRNA DDGC participates in premature ovarian insufficiency through regulating RAD51 and WT1. *Mol Ther Nucleic Acids* 26:1092–1106. doi: 10.1016/j.omtn.2021.10.015
33. Wang F, Chen X, Sun B, Ma Y, Niu W, Zhai J and Sun Y (2021) Hypermethylation-mediated downregulation of lncRNA PVT1 promotes granulosa cell apoptosis in premature ovarian insufficiency via interacting with Foxo3a. *J Cell Physiol* 236:5162–5175. doi: 10.1002/jcp.30222
34. Harvey CN, Esmail M, Wang Q, Brooks AI, Zachow R and Uzumcu M (2009) Effect of the methoxychlor metabolite HPTE on the rat ovarian granulosa cell transcriptome in vitro. *Toxicol Sci* 110:95–106. doi: 10.1093/toxsci/kfp089
35. Wang X, Meng K, Wang H, Wang Y, Zhao Y, Kang J, Zhang Y and Quan F (2021) Identification of small extracellular vesicle subtypes in follicular fluid: Insights into the function and miRNA profiles. *J Cell Physiol* 236:5633–5645. doi: 10.1002/jcp.30251
36. Chen Y, Chang H, Cheng J, Tsai H, Wu C and Leung P (2015) Transforming growth factor- β 1 up-regulates connexin43 expression in human granulosa cells. *Human reproduction (Oxford, England)* 30:2190–201. doi: 10.1093/humrep/dev175

37. Craig J, Orisaka M, Wang H, Orisaka S, Thompson W, Zhu C, Kotsuji F and Tsang B (2007) Gonadotropin and intra-ovarian signals regulating follicle development and atresia: the delicate balance between life and death. *Frontiers in bioscience: a journal and virtual library* 12:3628–39. doi: 10.2741/2339
38. Wang Y, Hu SB, Wang MR, Yao RW, Wu D, Yang L and Chen LL (2018) Genome-wide screening of NEAT1 regulators reveals cross-regulation between paraspeckles and mitochondria. *Nat Cell Biol* 20:1145–1158. doi: 10.1038/s41556-018-0204-2
39. Bouckenheimer J, Fauque P, Lecellier CH, Bruno C, Commes T, Lemaître JM, De Vos J and Assou S (2018) Differential long non-coding RNA expression profiles in human oocytes and cumulus cells. *Sci Rep* 8:2202. doi: 10.1038/s41598-018-20727-0
40. Ernst EH, Nielsen J, Ipsen MB, Villesen P and Lykke-Hartmann K (2018) Transcriptome Analysis of Long Non-coding RNAs and Genes Encoding Paraspeckle Proteins During Human Ovarian Follicle Development. *Front Cell Dev Biol* 6:78. doi: 10.3389/fcell.2018.00078
41. Zhen J, Li J, Li X, Wang X, Xiao Y, Sun Z and Yu Q (2021) Downregulating lncRNA NEAT1 induces proliferation and represses apoptosis of ovarian granulosa cells in polycystic ovary syndrome via microRNA-381/IGF1 axis. *J Biomed Sci* 28:53. doi: 10.1186/s12929-021-00749-z
42. Ma R, Hou X, Zhang L, Sun SC, Schedl T, Moley K and Wang Q (2014) Rab5a is required for spindle length control and kinetochore-microtubule attachment during meiosis in oocytes. *Faseb j* 28:4026–35. doi: 10.1096/fj.14-250886
43. Xiang Z, Lv Q, Chen X, Zhu X, Liu S, Li D and Peng X (2020) Lnc GNG12-AS1 knockdown suppresses glioma progression through the AKT/GSK-3 β / β -catenin pathway. *Biosci Rep* 40. doi: 10.1042/bsr20201578
44. Mahboobeh Z, Pegah M, Fatemeh S, Elham K, Hanieh A, Milad R and Mohammad S (2020) lncRNA ZEB2-AS1: A promising biomarker in human cancers. *IUBMB Life* 72:1891–1899. doi: 10.1002/iub.2338
45. Zhang B, Fan DB, Liu L, Qin Y and Feng DQ (2020) Knockdown of ZEB2-AS1 inhibits cell proliferation, invasion and induces apoptosis in osteosarcoma by combining with EZH2. *Eur Rev Med Pharmacol Sci* 24:6533–6539. doi: 10.26355/eurev_202006_21637
46. Sun F, Tan W, Dan Y, Wang X, Guo Y and Deng G (2020) Copy number gain of pro-inflammatory genes in patients with HBV-related acute-on-chronic liver failure. *BMC Med Genomics* 13:180. doi: 10.1186/s12920-020-00835-5
47. Guerrero-Netro HM, Chorfi Y and Price CA (2015) Effects of the mycotoxin deoxynivalenol on steroidogenesis and apoptosis in granulosa cells. *Reproduction* 149:555–61. doi: 10.1530/rep-15-0018
48. Wang H, He Y, Cheng D, Pu D, Tan R, Gao L, Cui Y and Wu J (2019) Cypermethrin exposure reduces the ovarian reserve by causing mitochondrial dysfunction in granulosa cells. *Toxicol Appl Pharmacol* 379:114693. doi: 10.1016/j.taap.2019.114693

49. Ożegowska K, Brązert M, Ciesiołka S, Nawrocki MJ, Kranc W, Celichowski P, Jankowski M, Bryja A, Jeseta M, Antosik P, Bukowska D, Skowroński MT, Bruska M, Pawelczyk L, Zabel M, Nowicki M and Kempisty B (2019) Genes Involved in the Processes of Cell Proliferation, Migration, Adhesion, and Tissue Development as New Potential Markers of Porcine Granulosa Cellular Processes In Vitro: A Microarray Approach. *DNA Cell Biol* 38:549–560. doi: 10.1089/dna.2018.4467
50. Kranc W, Brązert M, Budna J, Celichowski P, Bryja A, Nawrocki MJ, Ożegowska K, Jankowski M, Chermuła B, Dyszkiewicz-Konwińska M, Jeseta M, Pawelczyk L, Bręborowicz A, Rachoń D, Bruska M, Nowicki M, Zabel M and Kempisty B (2019) Genes responsible for proliferation, differentiation, and junction adhesion are significantly up-regulated in human ovarian granulosa cells during a long-term primary in vitro culture. *Histochem Cell Biol* 151:125–143. doi: 10.1007/s00418-018-1750-1
51. Pashaiasl M, Ebrahimi M and Ebrahimie E (2016) Identification of the key regulating genes of diminished ovarian reserve (DOR) by network and gene ontology analysis. *Mol Biol Rep* 43:923–37. doi: 10.1007/s11033-016-4025-8
52. Esmaeili-Fard SM, Gholizadeh M, Hafezian SH and Abdollahi-Arpanahi R (2021) Genome-wide association study and pathway analysis identify NTRK2 as a novel candidate gene for litter size in sheep. *PLoS One* 16:e0244408. doi: 10.1371/journal.pone.0244408
53. Qazi IH, Cao Y, Yang H, Angel C, Pan B, Zhou G and Han H (2021) Impact of Dietary Selenium on Modulation of Expression of Several Non-Selenoprotein Genes Related to Key Ovarian Functions, Female Fertility, and Proteostasis: a Transcriptome-Based Analysis of the Aging Mice Ovaries. *Biol Trace Elem Res* 199:633–648. doi: 10.1007/s12011-020-02192-x
54. Yang X, Zhou Y, Peng S, Wu L, Lin HY, Wang S and Wang H (2012) Differentially expressed plasma microRNAs in premature ovarian failure patients and the potential regulatory function of mir-23a in granulosa cell apoptosis. *Reproduction* 144:235–44. doi: 10.1530/rep-11-0371
55. Kirshenbaum M and Orvieto R (2019) Premature ovarian insufficiency (POI) and autoimmunity—an update appraisal. *J Assist Reprod Genet* 36:2207–2215. doi: 10.1007/s10815-019-01572-0
56. Sharif K, Watad A, Bridgwood C, Kanduc D, Amital H and Shoenfeld Y (2019) Insights into the autoimmune aspect of premature ovarian insufficiency. *Best Pract Res Clin Endocrinol Metab* 33:101323. doi: 10.1016/j.beem.2019.101323
57. Lim W, Bae H, Bazer FW and Song G (2018) Cell-specific expression and signal transduction of C-C motif chemokine ligand 2 and atypical chemokine receptors in the porcine endometrium during early pregnancy. *Dev Comp Immunol* 81:312–323. doi: 10.1016/j.dci.2017.12.020
58. Matsuda F, Inoue N, Manabe N and Ohkura S (2012) Follicular growth and atresia in mammalian ovaries: regulation by survival and death of granulosa cells. *J Reprod Dev* 58:44–50. doi: 10.1262/jrd.2011-012
59. Fan Y, Chang Y, Wei L, Chen J, Li J, Goldsmith S, Silber S and Liang X (2019) Apoptosis of mural granulosa cells is increased in women with diminished ovarian reserve. *J Assist Reprod Genet* 36:1225–1235. doi: 10.1007/s10815-019-01446-5

60. Hall SE, Upton RMO, McLaughlin EA and Sutherland JM (2018) Phosphoinositide 3-kinase/protein kinase B (PI3K/AKT) and Janus kinase/signal transducer and activator of transcription (JAK/STAT) follicular signalling is conserved in the mare ovary. *Reprod Fertil Dev* 30:624–633. doi: 10.1071/rd17024
61. Frost ER, Ford EA, Peters AE, Reed NL, McLaughlin EA, Baker MA, Lovell-Badge R and Sutherland JM (2020) Signal transducer and activator of transcription (STAT) 1 and STAT3 are expressed in the human ovary and have Janus kinase 1-independent functions in the COV434 human granulosa cell line. *Reprod Fertil Dev* 32:1027–1039. doi: 10.1071/rd20098
62. Ding N, Harlow SD, Randolph JF, Jr., Loch-Caruso R and Park SK (2020) Perfluoroalkyl and polyfluoroalkyl substances (PFAS) and their effects on the ovary. *Hum Reprod Update* 26:724–752. doi: 10.1093/humupd/dmaa018
63. Tatone C, Benedetti E, Vitti M, Di Emidio G, Ciriminna R, Vento ME, Cela V, Borzì P, Carta G, Lispi M, Cimini AM and Artini PG (2016) Modulating Intrafollicular Hormonal Milieu in Controlled Ovarian Stimulation: Insights From PPAR Expression in Human Granulosa Cells. *J Cell Physiol* 231:908–14. doi: 10.1002/jcp.25182
64. Osuka S, Iwase A, Goto M, Takikawa S, Nakamura T, Murase T, Kato N, Kotani T and Kikkawa F (2018) Thyroid Autoantibodies do not Impair the Ovarian Reserve in Euthyroid Infertile Women: A Cross-Sectional Study. *Horm Metab Res* 50:537–542. doi: 10.1055/a-0637-9430
65. Zhang Y, Ouyang X, You S, Zou H, Shao X, Zhang G, Zhang C and Hu L (2020) Effect of human amniotic epithelial cells on ovarian function, fertility and ovarian reserve in primary ovarian insufficiency rats and analysis of underlying mechanisms by mRNA sequencing. *Am J Transl Res* 12:3234–3254.
66. Ferraretti AP, La Marca A, Fauser BCJM, Tarlatzis B, Nargund G and Gianaroli L (2011) ESHRE consensus on the definition of 'poor response' to ovarian stimulation for in vitro fertilization: the Bologna criteria. *Human Reproduction* 26:1616–1624. doi: 10.1093/humrep/der092

Figures

Figure 1

Comparison of the properties of mRNA and lncRNA from women with NOR or DOR. **a** Genomic mapping of lncRNAs in different samples. **b** Location of lncRNA types on 46 chromosomes. Each circle represents a type of lncRNA and corresponds to “i”, “j”, “o”, “u”, or “x” from the outer circle to the inner circle. **c** The expression and amount of lncRNAs and mRNAs. **d** Distribution of the transcript length of lncRNAs and mRNAs. **e** Distribution of the exon number of lncRNAs and mRNAs. **f** Distribution of the ORF length of lncRNAs and mRNAs

Figure 2

Identification and characterization of the DE mRNA genes between women in the DOR group and NOR group. **a** Number of DE mRNA genes showing upregulated and downregulated expression. **b** Volcano plot of the DE mRNA genes. **c** Histogram of the GO-enriched terms of DE mRNA genes. **d** Scatter plot of the GO-enriched DE mRNA genes. **e** Scatter plot of the signaling pathway enrichment of DE mRNA genes using the KEGG database

Figure 3

Identification and characterization of the DE lncRNAs between women in the DOR group and NOR group. **a** Number of DE lncRNAs with upregulated and downregulated expression. **b** Volcano plot of the DE lncRNAs. **c** Histogram of the GO-enriched terms for the target genes of the DE lncRNAs. **d** Scatter plot of the GO enrichment for the target genes of the DE lncRNAs. **e** Scatter plot of the signaling pathway enrichment for the target genes of the DE lncRNAs using the KEGG database

Figure 4

The lncRNA-mRNA regulatory network. Each lncRNA is a red round rectangle. Each mRNA is a green ellipse

Figure 5

Expression of key lncRNAs as determined by RT-qPCR. *P < 0.05; **P < 0.01

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementaryfile.zip](#)