Everolimus-induced Hyperpermeability of Endothelial Cells Causes Lung Injury

Xiaolin Chen (✉ cxl888hhh@163.com)

Pingxiang People's Hospital

Jianhui Chen

Pingxiang People's Hospital

Shuihong Liu

Pingxiang People's Hospital

Xianfan Li

Pingxiang People's Hospital

Research Article

Keywords: Endothelial cell, Everolimus, Permeability, Lung injury, Myosin light chain, Ca²⁺.

Posted Date: January 25th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1280276/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The mammalian target of rapamycin (mTOR) inhibitors, everolimus (but not dactolisib), is frequently associated with lung injury in clinical therapies. However, the underlying mechanisms remain unclear. Endothelial cell barrier dysfunction plays a major role in the pathogenesis of the lung injury. This study hypothesizes that everolimus increases pulmonary endothelial permeability, which leads to lung injury.

Methods: We tested the effects of everolimus and dactolisib on human pulmonary microvascular endothelial cell (HPMEC) permeability by measuring transendothelial electrical resistance (TEER) and the passage of horseradish peroxidase-albumin (Pa). F-actin remodeling visualized using immunofluorescent imaging and its related key enzymes were measured by western blotting. Then, we established a mouse model using intraperitoneal injections of everolimus and dactolisib, and the total protein in bronchoalveolar lavage fluid (BALF) and the accumulation of Evans blue dye in lungs were used to examine the lung vascular permeability. The protective effects of myosin light chain (MLC) related pathway inhibitors on everolimus-induced hyperpermeability was observed.

Results: Everolimus increased HPMEC permeability, which was confirmed by decreased TEER and an increased transendothelial BSA flux, which was associated with MLC phosphorylation and F-actin stress fiber formation. In contrast, dactolisib did not have effect on the permeability of HPMECs. In addition, inhibition, or knockdown of PKC α or MLC kinase attenuated everolimus-induced hyperpermeability and MLC phosphorylation in HPMECs. Furthermore, everolimus induced an increasing concentration of intracellular calcium Ca²⁺ leakage in HPMECs and this was normalized with ryanodine pretreatment. Additionally, ryanodine decreased everolimus-induced phosphorylation of PKC α and MLC, and barrier disruption in HPMECs. Consistent with in vitro data, everolimus treatment caused a visible lung-vascular barrier dysfunction, including an increase in protein in BALF and lung capillary-endothelial permeability, which was significantly attenuated by pretreatment with an inhibitor of PKC α , MLCK, and ryanodine.

Conclusions: This study shows that everolimus induced pulmonary endothelial hyper-

permeability, at least partly, in an MLC phosphorylation-mediated EC contraction which is influenced in a Ca²⁺-dependent manner and can lead to lung injury.

Introduction

Based on the function of mammalian target of rapamycin (mTOR) signaling in physiological and pathological processes in the cell, pharmacological inhibitors are often used as disease interventions[1]. Rapamycin and its analogues (such as everolimus and temsirolimus) selectively inhibit mTOR by binding to FK506-binding protein 12 (FKBP12) and have been widely used in clinical studies[2, 3]. In addition, a newer generation of selective ATP-competitive inhibitor of mTOR, dactolisib (NVP-BEZ235), has been approved for clinical cancer treatment[2, 4]. Over the past few years, adverse effects such as pneumonitis[5-7], have been reported in most large clinical trials of everolimus or temsirolimus. However,

the incidence and risk of pneumonitis has not been reported in dactolisib clinical trials and case reports[4, 6, 8]. The exact mechanisms underlying these two generations of mTOR inhibitor-induced lung toxicities remain unclear.

Although a few studies have investigated the mechanism by which mTOR inhibitors lead to lung injury, the results are poorly defined and controversial[9-11]. Our group has recently published a paper showing that rapamycin aggravated lipopolysaccharide (LPS)-induced pulmonary endothelial cell (EC) hyperpermeability and lung injury in mice[12]. The endothelial cells play a critical role in maintaining the integrity of the lung capillary-alveolar barrier[13], which is critically dependent on the stabilization of the F-actin cytoskeleton and the assembly of cell junctions[14, 15]. Activation of F-actin fiber formation and cytoskeletal contractility represents an important mechanism underlying the induction of intercellular gap formation and endothelial hyperpermeability[15, 16]. One of the determinant events in endothelial cell contraction is the phosphorylation of myosin light chain (MLC) at Ser19 and/or Thr18[17], which is mainly regulated by MLC kinase (MLCK) and PKC α [16, 18, 19]. Intercellular Ca²⁺ plays an important regulatory role in the disruption of endothelial barrier function by activation of PKC α or MLCK[20]. Previous data has shown that displacement of FKBP12/12.6 from ryanodine receptors 2 (RyR2) induces an intracellular Ca²⁺ leak, leading to the activation of PKC α in sirolimus- treated human aortic endothelial cells (HAECs)[21, 22].

In this study, we hypothesized that everolimus induces an intracellular calcium release and the activation of PKC α and MLCK, and that this has consequences for barrier dysfunction in human pulmonary microvascular endothelial cells (HPMECs). To test this, we compared the effects of everolimus and dactolisib on lung endothelial permeability using a mouse model to identify an MLC-dependent mechanism that was associated with an intercellular Ca²⁺ release.

Methods

Chemicals and reagents

Antibodies against p-MLC (3674), MLC (8505), and PKC α (2056) were purchased from Cell Signaling Technology (Beverly, MA, USA). MLCK Antibody (ab232949) was obtained from Abcam (Waltham, MA, USA). FITC-Phalloidin (P5282) and Evans blue (EB) were obtained from Sigma Aldrich (Saint Louis, MO, USA). Everolimus (HY-10218), ryanodine (HY-103306), dactolisib (HY-50673), GF109203X (HY-13867), and ML-7 (HY-15417) were provided by Med Chem Express (Newark, NJ, USA). The Lipofectamine 2000 reagent was purchased from Invitrogen (Carlsbad, CA, USA).

Cell culture and treatment

HPMECs (ScienCell Research Laboratories, USA) were cultured in extracellular matrix (ScienCell, America) complemented with 10 % fetal bovine serum (FBS; Hyclone, Logan, USA), at 37 °C and 5 % CO₂ incubator. For all experiments, cells were grown to 80 % confluence and stimulated with PBS, everolimus, or dactolisib, with or without inhibitors (GF109203X, ML-7, Ryanodine).

Western blotting

Western blotting was performed as described previously[23]. The treated HPMEC lysates were collected and subjected to SDS-PAGE and immunoblotting. After being blocked with 5 % bovine serum albumin (BSA), The blots were incubated overnight at 4 °C with primary antibodies. The immunoreactive bands were then detected using ECL reagent (Pierce, 34078) and analyzed with Image J software (BIO RAD, Hercules, CA, USA).

F-actin staining assay

Immunofluorescence was detected using an F-actin assay. Confluent monolayers of HPMECs were grown on glass coverslips and fixed with 4 % paraformaldehyde, and blocked with 5 % BSA and probed with FITC-Phalloidin (1:200). Then, cell monolayers were stained with DAPI for 5 min. A confocal microscope with a laser-scanner (Tokyo, Japan) was used to perform and analyze the stained actin structures.

Transendothelial electrical resistance (TEER) assay

Based on previous studies[12, 23], HPMECs were cultured in transwell inserts (Millipore, Burlington, MA, USA) and grown to complete confluence. The resistance values of the endothelial cell monolayer were detected by a Millicell-ERS Voltohmmeter (Millipore, Burlington, MA, USA) and were recorded in $\Omega \cdot \text{cm}^2$. The TEER data for each group was normalized to the medium (control) group.

Transendothelial albumin leakage assay

Confluent monolayers of HPMECs were seeded on a 0.4 μm pore 24-well transwell insert and grown to complete confluence. The medium containing horseradish peroxidase (HRP)-labeled albumin (50 $\mu\text{g}/\text{mL}$; Solarbio, Beijing, China) was added to the upper chamber, followed by incubation for treatment with everolimus or dactolisib, with or without inhibitors for 24 h. The content of albumin in the down chambers was measured by the TMB Soluble Substrate kit (Solarbio,Beijingm,China), and recorded with a microplate reader (Elx 800, BioTek). The leakage of HRP-BSA of each of the treated groups was normalized to the control group.

Cell transfection

HPMECs were transfected with siRNA-MLCK and si-PKC α (Santa Cruz Biotech

Nology) using Lipofectamine 2000. HPMECs were plated at 80 % confluence for the transfection experiments using siRNA-MLCK AND si-PKC α , following manufacturer's instructions. The transfected complex was subsequently added to each well of a 6-well plate and supplemented with serum-free OptiMEM, and the plate was incubated for 6 h at 37 °C. The culture medium was replaced after 48 h, and TEER measurements, and immunoblotting were performed at 24 h after transfection. Target proteins for siRNA were confirmed with western blots.

Intracellular calcium leak measurement

Intracellular calcium [Ca^{2+}] was quantified using the fluorescent calcium indicator Fluo-3 AM (Beyotime, Shanghai, China). After being treated, groups of HPMCEs were added with the Fluo-3 AM and were maintained at 37 °C in Hanks balanced buffered solution for 1 h prior to fluorescence being detected. The fluorescence intensities were acquired for each group using a BD FACSCalibur flow cytometer, and a qualitative assessment was conducted for intracellular calcium flux analysis using FlowJo software.

Experimental animals

C57BL/6 J (B6) mice were purchased from Nanchang University and approval was granted by the Animal Studies Ethics Committee of Pingxiang people's Hospital. The mice (aged 6–8 weeks, weighing 20–25 g) were randomly assigned into six groups: (1) control group (PBS), (2) everolimus, (3) dactolisib, (4) everolimus + Ryanodine, (5) everolimus + GF109203X (GF), (6) everolimus + ML-7. Groups 1–3 were treated daily with a single intraperitoneal (i.p.) injection of PBS, everolimus at 1 mg/kg/day, or dactolisib at 1 mg/kg/day, respectively, for 8 weeks[9]. Based on the findings of previous studies, the inhibitor groups were pre-administered with GF (1 mg/kg/day)[24], ML-7 (1 mg/kg/day)[18], or ryanodine (2 mg/kg/day)[25] for 2 days per week, 1 day prior to intraperitoneal injection with everolimus.

Detection of protein in bronchoalveolar lavage fluid (BALF)

All mice were anaesthetized with isoflurane[12]. After anesthesia, BALF was obtained, as per our previous description, and 2.0 mL of BALF was centrifuged at 3000 RPM for 10 min at 4 °C. The concentration of total protein in the BALF was measured using BCA assay kits (Beyotime, Shanghai, China).

Evaluation of lung permeability

EB (5mg/mL, dissolved in PBS) was injected intravenously into the tail vein, 30 min before exsanguination. The left lung was collected and weighed. Then, 0.5 mL of formamide (Sigma-Aldrich) was added and incubated at 60 °C for 24 h. The intensity of EB in the lung tissue was determined by detecting absorbance at wavelength 610 nm. The extravasated EB concentration in the lung tissue was calculated against a standard curve (micrograms of EB dye per lung).

Statistical analysis

All data were analyzed by GraphPad Prism v6.0 software and SPSS statistics 17.0 (IBM Inc., Chicago, IL). The mean values of the groups were expressed as mean $\pm$ SD and were compared using one-way ANOVA followed by post hoc analysis. Results were considered statistically significant at $p < 0.05$.

Results

Everolimus induced lung endothelial hyperpermeability

Two generations of mTOR inhibitors, everolimus and dactolisib, were used in this study. TEER and the permeability of albumin (Pa) of the HPMEC monolayer were measured to evaluate the endothelial barrier function. As shown in Figures 1a and 1b, 24 h treatment with 10 nM and 100 nM everolimus induced a decrease in TEER values (Fig. 1a). Consistent with the TEER values, 10 nM, and 100 nM everolimus lead to an increase in Pa by 60 % and 80 %, respectively (Fig. 1b). In contrast, dactolisib had no effect on the TEER and Pa values in the endothelial cells (Fig. 1a-b). These data demonstrated that everolimus exhibits a dose- dependent hyperpermeability in HPMEC monolayers, while dactolisib does not.

Everolimus induced F-actin fiber formation and MLC phosphorylation in endothelial cells

Formation of actin stress fibers, which was mainly generated by MLC phosphorylation, was closely linked to an increased endothelial permeability[26, 27]. An immunofluorescent assay, with FITC-phalloidin to label F-actin, was used to detect the formation of stress fibers and paracellular gaps in everolimus or dactolisib-treated HPMECs. Consistent with the results in permeability detection, everolimus induced alterations in the distribution of F-actin stress fibers and lead to gap formation in the HPMECs (Fig. 2a-b). In addition, we also observed a slight increase in MLC phosphorylation in everolimus-treated cells (Fig. 2c-d). In contrast, dactolisib had no effect on the MLC phosphorylation, morphological changes, and gap formation on the monolayer of HPMECs (Fig. 2a-d). These results suggest that everolimus- induced phosphorylation of MLC and F-actin formation are accompanied with increased cell contraction in HPMECs.

PKC α and MLCK are required for everolimus-induced MLC phosphorylation

PKC α and MLCK are the main regulators of MLC phosphorylation and have been reported to mediate cytoskeletal contraction regulation of endothelial permeability[17, 26]. Therefore, specific pharmacological inhibitors (ML-7 and GF109203X, GF) were used to examine the involvement of MLCK and PKC α in everolimus-induced endothelial hyperpermeability. As shown in Figure 3, everolimus-induced MLC phosphorylation was attenuated by GF (5 mM) and ML-7 (10 mM) in HPMECs (Fig.3c and 3g). Pretreatment with GF and ML-7 consistently abolished EC hyperpermeability responses to everolimus, which was evidenced by an increase in TEER (Fig.3a) and a decrease in the permeability coefficient for Pa (Fig.3b).

Moreover, we utilized siRNAs to reduce PKC α and MLCK expression in the everolimus- treated HPMECs. In accordance with the effects of pharmacological inhibition, knockdown of PKC α and MLCK significantly abolished everolimus-induced MLC phosphorylation (Fig.3f) and hyperpermeability (Fig.3d-e). These results indicate that both MLCK and PKC α contribute to MLC phosphorylation and EC contraction, which resulted in hyperpermeability in everolimus-induced EC.

Everolimus impairs endothelial cell barrier function via an increase in Ca²⁺ release

Intracellular calcium [Ca²⁺] binding to the CaM-binding domain of MLCK and PKC α resulted in activation of PKC α and MLCK[20, 28, 29]. Previous studies have shown that everolimus can lead to intracellular

calcium Ca^{2+} release via displacement of FKBP12.6 from the ryanodine receptor 2 (RyR2). We examined the intracellular $[\text{Ca}^{2+}]$ concentration in the HPMEC monolayers after treatment with everolimus or dactolisib. Compared with the control group, we observed a significant increase in intracellular calcium Ca^{2+} release after everolimus treatment for up to 24 h (Fig.4a). Contrastingly, dactolisib did not effect on Ca^{2+} release (Fig.4a). Moreover, pretreatment with ryanodine (a stabilizer of RyR2 channels) ameliorated the everolimus-induced increase in intracellular Ca^{2+} levels (Fig.4c) and cell barrier dysfunction, which was evidenced by a reduced the passage of BSA-leakage (Fig.4c), and the alleviated

drop of TEER (Fig.4d) in HPMECs. Moreover, the everolimus-induced phosphorylation of MLC was attenuated in the ryanodine pretreatment group (Fig.4 e-f).

Collectively, these data suggest that the disruption of calcium release through the RYR2 channel by everolimus induced an intracellular Ca^{2+} leakage and enhanced phosphorylation of PKC α and MLC, resulting in an impaired lung endothelial barrier function.

Everolimus increases lung capillary-alveolar permeability in vivo

To determine whether the toxicity effects of 2 generations of mTOR inhibitors (everolimus and dactolisib) on the lung injury, we treated the mice with PBS, everolimus, or dactolisib as per the previously described methods. Capillary-alveolar barrier injures were attributable to lung injury. We examined the concentration of total protein in BALF and the amount of EB in the lungs to assess the lung vascular permeability. As shown Figure 5, we found that everolimus (EVE) significantly increased the amount of EB in the lungs (Fig.5a), and the total protein in BALF (Fig.5b) was increased compared to both the control and dactolisib groups. Compared to everolimus treatment alone, pretreatment with GF, ML-7, and ryanodine dramatically decreased the leakage of protein in BALF, as well as the accumulation of EB in the lungs, indicating an improved barrier function (Fig.5a-b). These data indicate that everolimus induces pulmonary vascular hyperpermeability and lung injury through an endothelial barrier dysfunction-dependent mechanism.

Discussion

While mTOR inhibitors are an effective method of systemic treatment for tumor and organ transplantations, their side effects, such as causing lung injury, have drawn extensive concerns[30-32]. However, the underlying molecular mechanisms of everolimus on pulmonary toxicity remains largely unclear. Herein, we reported a novel role for everolimus in inducing endothelial cell barrier dysfunction and lung injury by promoting MLC phosphorylation and EC hyperpermeability, which was strongly dependent on the activation of PKC α and MLCK. Furthermore, we showed that the harmful effects of everolimus appeared to be mediated by destabilization of the intracellular Ca^{2+} release channel RyR2, which altered intracellular $[\text{Ca}^{2+}]$ stores. Moreover, we established the first animal model of everolimus-related lung injury, by treating mice with a daily regimen of everolimus. However, there was no damage of barrier function in the dactosilib-treated pulmonary endothelial cells and mice. Based on these data, we

demonstrated that everolimus has the potential to induce endothelial barrier dysfunction, which can induce pulmonary vascular hyperpermeability and lung injury.

Previous studies have shown that lung injury is a frequent side-effect of mTOR inhibitors, such as everolimus and temsirolimus, that are used in clinical treatments[6, 7, 31]. In vivo studies have shown that rapamycin augments lung injury induced by cigarette smoke in mice [33], and similar result was observed in temsirolimus-treated mice[9]. Other studies have also demonstrated that rapamycin aggravates LPS-induced lung apoptosis by suppression of mTOR activation in mice[11]. Clinically, serum Krebs von den Lungen 6, a biomarker of lung capillary-alveolar permeability injury, is elevated in patients with everolimus-induced lung disease[30, 34]. Consistent with these studies, our findings demonstrate that mice with a daily regimen of everolimus treatment have an altered endothelial barrier integrity, which results in increasing pulmonary vascular endothelial permeability and lung injury.

Pulmonary endothelial barrier dysfunction and plasma protein infiltration into the pulmonary alveoli occurs in the initial stages of the pathogenesis of pneumonitis[35, 36]. The hyperpermeability of pulmonary vasculature may lead to the leakage of plasma proteins within the alveolar lumen, which is a sign of pulmonary vascular endothelial injury[37-39]. Previous studies have showed that rapamycin enhances LPS-induced pulmonary edema, an indication of disruption to the capillary-alveolar barrier[11]. Furthermore, it has been reported that temsirolimus-induced lung injury can be induced following capillary-alveolar barrier injury[9]. Consistent with previous studies, we highlighted a possible connection between everolimus-induced endothelial hyperpermeability and the development of lung injury through finding an increase in the concentrations of protein in BALF and EB in the lungs in mice treated with everolimus. This showed that there was detectable damage in the capillary-alveolar barrier. Furthermore, we performed in vitro studies using HPMECs and found that everolimus increased endothelial permeability by highlighting the alteration of TEER and Pa values. Contrastingly, dactolisib did not demonstrate this effect. Based on these data, our study demonstrated that everolimus has the potential to cause pulmonary endothelial injury, which could result in lung injury.

The critical role for MLC phosphorylation and F-actin stress fiber formation in EC contraction and barrier dysfunction has been well documented[15, 16]. Our data demonstrated that everolimus was associated with an increase in F-actin formation and endothelial hyperpermeability, whereas dactolisib did not demonstrate this effect. This was supported by a recent study which showed that dactolisib had no effect on HPMEC viability and LPS induced-lung injury in mice [3]. Previous studies demonstrated that an upregulation in the activity of MLCK and PKC α resulted in increased MLC phosphorylation, leading to disruption of the vascular barrier integrity in a murine model of LPS-induced lung injury[40-42]. The in vitro data showed that the increased activity in the MLCK/MLC pathway lead to paracellular gap formation and endothelial cell barrier disruption[26, 43]. PKC α has been shown to modulate endothelial barrier function by activating MLC through inhibition of MLC phosphatase (MLCP) by direct phosphorylation of PKC-potentiator protein phosphatase inhibitory protein-17 (CPI-17)[26]. Blocking PKC α /CPI-17 pathways minimized the LPS-induced endothelial barrier dysfunction[15]. We have previously shown that rapamycin induced PKC α phosphorylation in HUVECs[44]. In this study, we noted

that PKC α plays an essential role in everolimus-induced barrier disruption, through enhanced MLC phosphorylation and actin stress fiber formation. Furthermore, we also found that everolimus-induced MLC phosphorylation was dependent on the activation of both MLCK and PKC α , which was confirmed by the pretreatment inhibition or knockdown of MLCK or PKC α . Pretreatment with the inhibitor anti-MLCK and -PKC α in vivo prevents the everolimus-induced endothelial hyperpermeability and the development of lung injury. These results indicate that both PKC α and MLCK are responsible for the phosphorylation of MLC and the endothelial barrier dysfunction induced by everolimus.

Everolimus inhibits mTOR through binding to FKBP12, a key stabilizing component of RyR2 calcium release channels in various cell types[5, 45]. Dactolisib, which also named NVP-BEZ235, does not show this effect[45, 46]. Previous reports have shown that rapamycin induced Ca²⁺ release via RyR2 in endothelial cells[21]. Studies have also demonstrated that low concentrations of [Ca²⁺] induced PKC α and MLCK activation, which lead to phosphorylation of MLC[28, 29, 47]. Herein, we found a different toxicity effect on HPMECs exposed to everolimus and dactolisib, which may have been associated with the Ca²⁺ leakage from the ryanodine receptors. Our data showed that everolimus induced an intracellular Ca²⁺ release in HPMECs, which was blocked by ryanodine. However, dactolisib, a selective ATP-competitive inhibitor of the mTOR that does not bind FKBP12/12.6, did not affect endothelial barrier function and intracellular Ca²⁺ leakage. Furthermore, we found that inhibition of RyR2 by ryanodine not only blocked intracellular Ca²⁺ leakage, but also attenuated MLC phosphorylation and hyperpermeability in the everolimus-treated HPMECs. In addition, similar results were seen in vivo. Collectively, these results suggest that everolimus-induced leakage of calcium from RyR2 may enhance MLC phosphorylation, leading to impaired endothelial barrier function.

Conclusion

In conclusion, our results demonstrate that everolimus promotes endothelial hyperpermeability, at least partly through MLC phosphorylation-mediated EC contraction by PKC α and MLCK activation, which may be dependent on an endothelial intracellular [Ca²⁺] leakage. The effect was also observed in vivo, as mice treated with everolimus displayed pulmonary vascular leakage and lung injury. This study elucidated a molecular mechanism that may contribute to everolimus-induced endothelial barrier dysfunction and help to ameliorate the management of mTOR inhibitor-associated side effects.

Abbreviations

HAECs: human aortic endothelial cells; EC: Endothelial cell; TEER: transendothelial electrical resistance; HPMECs: human pulmonary microvascular endothelial cells; MLC: myosin light chain; PKC α : Protein Kinase C- α ; MLCK: myosin light chain kinase; LPS: Lipopolysaccharide; BALF: Bronchoalveolar lavage fluid; RyR2: Ryanodine Receptor 2; mTOR: mammalian target of rapamycin; FKBP12:FK506-binding protein12; GF:GF109203X

Declarations

Acknowledgements

Not applicable.

Ethics approval and consent to participate

The ethics approval was obtained from the Animal Studies Ethics Committee of Pingxiang people's Hospital.

Consent for publication

All authors provided consent for publication

Availability of data and materials

All data generated or analyzed during this study are included in this published Article.

Competing interests

The authors declare that they have no competing interests.

Funding

This project was supported by the Natural Science Foundation of Jiangxi Province (Grant No. 20202BABL216005).

Authors' contributions

CJH and LXF designed and performed the experiments; Funding acquisition; YZP contributed reagents/materials/analysis tools; CXL designed the experiments and wrote the paper. All authors read and approved the final manuscript.

References

1. Saxton RA, Sabatini DM: mTOR Signaling in Growth, Metabolism, and Disease. *Cell* 2017, 168(6):960-976.
2. Wise-Draper TM, Moorthy G, Salkeni MA, Karim NA, Thomas HE, Mercer CA, Beg MS, O'Gara S, Olowokure O, Fathallah H *et al*: A Phase Ib Study of the Dual PI3K/mTOR Inhibitor Dactolisib (BEZ235) Combined with Everolimus in Patients with Advanced Solid Malignancies. *Target Oncol* 2017, 12(3):323-332.
3. Ustun S, Lassnig C, Preitschopf A, Mikula M, Muller M, Hengstschlager M, Weichhart T: Effects of the mTOR inhibitor everolimus and the PI3K/mTOR inhibitor NVP-BEZ235 in murine acute lung injury

- models. *Transpl Immunol* 2015, 33(1):45-50.
4. Lang F, Wunderle L, Badura S, Schleyer E, Bruggemann M, Serve H, Schnittger S, Gokbuget N, Pfeifer H, Wagner S *et al*: A phase I study of a dual PI3-kinase/mTOR inhibitor BEZ235 in adult patients with relapsed or refractory acute leukemia. *BMC Pharmacol Toxicol* 2020, 21(1):70.
 5. Siddiqui AS, Zimmerman JL: Everolimus associated interstitial pneumonitis in a liver transplant patient. *Respir Med Case Rep* 2016, 19:15-17.
 6. Willemsen AE, Grutters JC, Gerritsen WR, van Erp NP, van Herpen CM, Tol J: mTOR inhibitor-induced interstitial lung disease in cancer patients: Comprehensive review and a practical management algorithm. *Int J Cancer* 2016, 138(10):2312-2321.
 7. Garrean S, Massad MG, Tshibaka M, Hanhan Z, Caines AE, Benedetti E: Sirolimus-associated interstitial pneumonitis in solid organ transplant recipients. *Clin Transplant* 2005, 19(5):698-703.
 8. Salazar R, Garcia-Carbonero R, Libutti SK, Hendifar AE, Custodio A, Guimbaud R, Lombard-Bohas C, Ricci S, Klumpen HJ, Capdevila J *et al*: Phase II Study of BEZ235 versus Everolimus in Patients with Mammalian Target of Rapamycin Inhibitor-Naive Advanced Pancreatic Neuroendocrine Tumors. *Oncologist* 2018, 23(7):766-e790.
 9. Washino S, Ando H, Ushijima K, Hosohata K, Kumazaki M, Mato N, Sugiyama Y, Kobayashi Y, Fujimura A, Morita T: Temsirolimus induces surfactant lipid accumulation and lung inflammation in mice. *Am J Physiol Lung Cell Mol Physiol* 2014, 306(12):L1117-1128.
 10. Jia X, Cao B, An Y, Zhang X, Wang C: Rapamycin ameliorates lipopolysaccharide-induced acute lung injury by inhibiting IL-1beta and IL-18 production. *Int Immunopharmacol* 2019, 67:211-219.
 11. Fielhaber JA, Carroll SF, Dydensborg AB, Shourian M, Triantafillopoulos A, Harel S, Hussain SN, Bouchard M, Qureshi ST, Kristof AS: Inhibition of mammalian target of rapamycin augments lipopolysaccharide-induced lung injury and apoptosis. *J Immunol* 2012, 188(9):4535-4542.
 12. Chen X, Hu C, Fan X, Wang Y, Li Q, Su YQ, Zhang DM, Yang Q, Passerini AG, Sun C: mTOR Inhibition Promotes Pneumonitis through Inducing Endothelial Contraction and Hyperpermeability. *Am J Respir Cell Mol Biol* 2021, 65(6):646-657.
 13. Dolmatova EV, Forrester SJ, Wang K, Ou Z, Williams HC, Joseph G, Kumar S, Valdivia A, Kowalczyk AP, Qu H *et al*: Endothelial Poldip2 regulates sepsis-induced lung injury via Rho pathway activation. *Cardiovasc Res* 2021.
 14. Wang L, Letsiou E, Wang H, Belvitch P, Meliton LN, Brown ME, Bandela M, Chen J, Garcia JGN, Dudek SM: MRSA-induced endothelial permeability and acute lung injury are attenuated by FTY720 S-phosphonate. *Am J Physiol Lung Cell Mol Physiol* 2022, 322(1):L149-L161.
 15. Kasa A, Csontos C, Verin AD: Cytoskeletal mechanisms regulating vascular endothelial barrier function in response to acute lung injury. *Tissue Barriers* 2015, 3(1-2):e974448.
 16. Yin J, Lv L, Zhai P, Long T, Zhou Q, Pan H, Botwe G, Wang L, Wang Q, Tan L *et al*: Connexin 40 regulates lung endothelial permeability in acute lung injury via the ROCK1-MYPT1-MLC20 pathway. *Am J Physiol Lung Cell Mol Physiol* 2019, 316(1):L35-L44.

17. Tinsley JH, Teasdale NR, Yuan SY: Myosin light chain phosphorylation and pulmonary endothelial cell hyperpermeability in burns. *Am J Physiol Lung Cell Mol Physiol* 2004, 286(4):L841-847.
18. Luh C, Kuhlmann CR, Ackermann B, Timaru-Kast R, Luhmann HJ, Behl C, Werner C, Engelhard K, Thal SC: Inhibition of myosin light chain kinase reduces brain edema formation after traumatic brain injury. *J Neurochem* 2010, 112(4):1015-1025.
19. Yang G, Li T, Xu J, Liu L: PKC plays an important mediated effect in arginine vasopressin induced restoration of vascular responsiveness and calcium sensitization following hemorrhagic shock in rats. *Eur J Pharmacol* 2010, 628(1-3):148-154.
20. Mizuno Y, Isotani E, Huang J, Ding H, Stull JT, Kamm KE: Myosin light chain kinase activation and calcium sensitization in smooth muscle in vivo. *Am J Physiol Cell Physiol* 2008, 295(2):C358-364.
21. Long C, Cook LG, Hamilton SL, Wu GY, Mitchell BM: FK506 binding protein 12/12.6 depletion increases endothelial nitric oxide synthase threonine 495 phosphorylation and blood pressure. *Hypertension* 2007, 49(3):569-576.
22. Habib A, Karmali V, Polavarapu R, Akahori H, Cheng Q, Pachura K, Kolodgie FD, Finn AV: Sirolimus-FKBP12.6 impairs endothelial barrier function through protein kinase C- α activation and disruption of the p120-vascular endothelial cadherin interaction. *Arterioscler Thromb Vasc Biol* 2013, 33(10):2425-2431.
23. Chen X, Chen J, Li X, Yu Z: Activation of mTOR mediates hyperglycemia-induced renal glomerular endothelial hyperpermeability via the RhoA/ROCK/pMLC signaling pathway. *Diabetol Metab Syndr* 2021, 13(1):105.
24. Vetri F, Qi M, Xu H, Oberholzer J, Paisansathan C: Impairment of neurovascular coupling in Type 1 Diabetes Mellitus in rats is prevented by pancreatic islet transplantation and reversed by a semi-selective PKC inhibitor. *Brain Res* 2017, 1655:48-54.
25. Meissner G: Ryanodine activation and inhibition of the Ca^{2+} release channel of sarcoplasmic reticulum. *J Biol Chem* 1986, 261(14):6300-6306.
26. Kolosova IA, Ma SF, Adyshev DM, Wang P, Ohba M, Natarajan V, Garcia JG, Verin AD: Role of CPI-17 in the regulation of endothelial cytoskeleton. *Am J Physiol Lung Cell Mol Physiol* 2004, 287(5):L970-980.
27. Zhao MJ, Jiang HR, Sun JW, Wang ZA, Hu B, Zhu CR, Yin XH, Chen MM, Ma XC, Zhao WD *et al*: Roles of RAGE/ROCK1 Pathway in HMGB1-Induced Early Changes in Barrier Permeability of Human Pulmonary Microvascular Endothelial Cell. *Front Immunol* 2021, 12:697071.
28. Sandoval R, Malik AB, Minshall RD, Kouklis P, Ellis CA, Tiruppathi C: Ca^{2+} signalling and PKC α activate increased endothelial permeability by disassembly of VE-cadherin junctions. *J Physiol* 2001, 533(Pt 2):433-445.
29. Tiruppathi C, Minshall RD, Paria BC, Vogel SM, Malik AB: Role of Ca^{2+} signaling in the regulation of endothelial permeability. *Vascul Pharmacol* 2002, 39(4-5):173-185.
30. Mizuno R, Asano K, Mikami S, Nagata H, Kaneko G, Oya M: Patterns of interstitial lung disease during everolimus treatment in patients with metastatic renal cell carcinoma. *Jpn J Clin Oncol* 2012,

31. Gartrell BA, Ying J, Sivendran S, Boucher KM, Choueiri TK, Sonpavde G, Oh WK, Agarwal N, Galsky MD: Pulmonary complications with the use of mTOR inhibitors in targeted cancer therapy: a systematic review and meta-analysis. *Target Oncol* 2014, 9(3):195-204.
32. David S, Kumpers P, Shin H, Haller H, Fliser D: Everolimus-associated interstitial pneumonitis in a patient with a heart transplant. *Nephrol Dial Transplant* 2007, 22(11):3363-3364.
33. Wang Y, Liu J, Zhou JS, Huang HQ, Li ZY, Xu XC, Lai TW, Hu Y, Zhou HB, Chen HP *et al*: MTOR Suppresses Cigarette Smoke-Induced Epithelial Cell Death and Airway Inflammation in Chronic Obstructive Pulmonary Disease. *J Immunol* 2018, 200(8):2571-2580.
34. Inoue Y, Barker E, Daniloff E, Kohno N, Hiwada K, Newman LS: Pulmonary epithelial cell injury and alveolar-capillary permeability in berylliosis. *Am J Respir Crit Care Med* 1997, 156(1):109-115.
35. Strieter RM: Pathogenesis and natural history of usual interstitial pneumonia: the whole story or the last chapter of a long novel. *Chest* 2005, 128(5 Suppl 1):526S-532S.
36. Gutbier B, Jiang X, Dietert K, Ehrler C, Lienau J, Van Slyke P, Kim H, Hoang VC, Maynes JT, Dumont DJ *et al*: Vasculotide reduces pulmonary hyperpermeability in experimental pneumococcal pneumonia. *Crit Care* 2017, 21(1):274.
37. Jobe AH: Pathogenesis of respiratory failure in the preterm infant. *Ann Med* 1991, 23(6):687-691.
38. Matthay MA, Zemans RL: The acute respiratory distress syndrome: pathogenesis and treatment. *Annu Rev Pathol* 2011, 6:147-163.
39. Ouwendijk WJ, van den Ham HJ, Delany MW, van Kampen JJ, van Nierop GP, Mehraban T, Zaaraoui-Boutahar F, van IWF, van den Brand JM, de Vries RD *et al*: Alveolar barrier disruption in varicella pneumonia is associated with neutrophil extracellular trap formation. *JCI Insight* 2020, 5(21).
40. Kovacs-Kasa A, Gorshkov BA, Kim KM, Kumar S, Black SM, Fulton DJ, Dimitropoulou C, Catravas JD, Verin AD: The protective role of MLCP-mediated ERM dephosphorylation in endotoxin-induced lung injury in vitro and in vivo. *Sci Rep* 2016, 6:39018.
41. Yu Y, Lv N, Lu Z, Zheng YY, Zhang WC, Chen C, Peng YJ, He WQ, Meng FQ, Zhu MS *et al*: Deletion of myosin light chain kinase in endothelial cells has a minor effect on the lipopolysaccharide-induced increase in microvascular endothelium permeability in mice. *FEBS J* 2012, 279(8):1485-1494.
42. Adyshev DM, Moldobaeva N, Mapes B, Elangovan V, Garcia JG: MicroRNA regulation of nonmuscle myosin light chain kinase expression in human lung endothelium. *Am J Respir Cell Mol Biol* 2013, 49(1):58-66.
43. Ding J, Li Z, Li L, Ding Y, Wang D, Meng S, Zhou Q, Gui S, Wei W, Zhu H *et al*: Myosin light chain kinase inhibitor ML7 improves vascular endothelial dysfunction and permeability via the mitogen-activated protein kinase pathway in a rabbit model of atherosclerosis. *Biomed Pharmacother* 2020, 128:110258.
44. Fan X, Chen X, Feng Q, Peng K, Wu Q, Passerini AG, Simon SI, Sun C: Downregulation of GATA6 in mTOR-inhibited human aortic endothelial cells: effects on TNF-alpha-induced VCAM-1 expression and monocytic cell adhesion. *Am J Physiol Heart Circ Physiol* 2019, 316(2):H408-H420.

45. van Rossum HH, Romijn FP, Smit NP, de Fijter JW, van Pelt J: Everolimus and sirolimus antagonize tacrolimus based calcineurin inhibition via competition for FK-binding protein 12. *Biochem Pharmacol* 2009, 77(7):1206-1212.
46. Tao Z, Barker J, Shi SD, Gehring M, Sun S: Steady-state kinetic and inhibition studies of the mammalian target of rapamycin (mTOR) kinase domain and mTOR complexes. *Biochemistry* 2010, 49(39):8488-8498.
47. De Bock M, Wang N, Decrock E, Bol M, Gadicherla AK, Culot M, Cecchelli R, Bultynck G, Leybaert L: Endothelial calcium dynamics, connexin channels and blood-brain barrier function. *Prog Neurobiol* 2013, 108:1-20.

Figures

Figure 1

Everolimus-induced lung endothelial hyperpermeability

Human pulmonary microvascular endothelial cell (HPMEC) monolayers were treated with 10 nM and 100 nM of everolimus or dactolisib for 24 h. The transendothelial electric resistance (TEER) (a) and horseradish peroxidase (HRP)-albumin permeability (Pa) of transendothelial flux (b) were measured. TEER was normalized to the baseline before treatment (TEER0) (n =4). ** p <0.01; *** p <0.001.

Figure 2

Everolimus induced endothelial cell F-actin fiber formation and MLC phosphorylation

(a) F-actin stress fiber formation was assessed via FITC-phalloidin staining, followed by the incubation of human pulmonary microvascular endothelial cells (HPMECs) with 100 nM everolimus or dactolisib for 24 h. DAPI was used to stain the nuclei. Representative images are shown. Scale bar: 30 μ m. (b) The quantification of the intercellular gap area in each group; n = 4. (c-d) The levels of myosin light chain (MLC) phosphorylation were analyzed using western blotting. * p < 0.05; ** p < 0.01; *** p < 0.001.

Figure 3

PKCα and MLCK are required for everolimus-induced MLC phosphorylation

Human pulmonary microvascular endothelial cells (HPMECs) were pretreated with ML-7 (10 mM) and GF109203X (GF, 5 mM) for 2 h (a-c) or were transfected with siPKCα and siMLCK (d-f) and incubated with 100 nM everolimus for 22 h. This was followed by western blotting of myosin light chain (MLC) phosphorylation (c, f, g and h), and the permeability of EC monolayers by transendothelial electrical resistance (TEER) measurements (a and d) and the horseradish peroxidase (HRP)-albumin leakage assay (b and e); n = 4. * p < 0.05; ** p < 0.01; *** p < 0.001.

Figure 4

Everolimus impairs endothelial cell barrier function via an increase in Ca^{2+} release, whereas dactolisib does not

(a) Human pulmonary microvascular endothelial cells (HRGECs) were treated with 10 and 100 nM everolimus or dactolisib for 24 h and intracellular Ca^{2+} leak levels were measured via the fluorescent dye Fluo-3 AM (n=4). (b) Two-hour pretreatment with the ryanodine receptor 2 channel (RYR2) stabilizer ryanodine (10mM) blocked the EC intracellular calcium leak Ca^{2+} , myosin light chain (MLC) phosphorylation (e), and hyperpermeability response (c-d) elicited by everolimus for 22 h (n=3). * p < 0.05; ** p < 0.01; *** p < 0.001.

Figure 5

Chronic treatment with everolimus increases lung capillary-alveolar permeability in vivo

Mice were treated with daily i.p. injections of vehicle (PBS), everolimus (EVE) at 1 mg/kg/day, or dactolisib at 1 mg/kg/day, for 8 weeks. In the inhibitor intervention in vivo experiments, mice were pretreated with GF, ML-7, and ryanodine, as described in Methods. (a) Quantification of vascular permeability, assessed by Evans blue accumulation in the lung tissue, and (b) total protein in BALF, measured by BCA protein assay kit. Data presented as mean $\pm$ SEM (n = 5–8). * p < 0.05; ** p < 0.01; *** p < 0.001.