Prevalence, Parents' Hesitancy, and Determinants of COVID-19 Vaccination Among Children: A Multi-Country Study From the Eastern Mediterranean Region

Moawiah Khatatbeh (✉ Moawia.m@yu.edu.jo)

Department of Basic Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan.

Samir Albalas

Department of Health Services Administration, Yarmouk University, Irbid, Jordan.

Haitham Khatatbeh

Doctoral School of Health Sciences, Faculty of Health Sciences, University of Pécs, Pécs, Hungary.

Waleed Momani

Department of Basic Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan.

Omar Melhem

Department of Nursing, Fatima College of Health Sciences, Abu Dhabi, United Arab Emirates.

Omar Al Omari

College of Nursing, Sultan Qaboos University, Muscat, Oman.

Zeinab Tarhini

Institute of Epidemiology and Tropical Neurology, GEIST, Limoges, France.

Ashraf A'aqoulah

Department of Health Systems Management, College of Public Health and Health Informatics, King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia.

Mohammed Al-Jubouri

College of Nursing, University of Baghdad, Iraq.

Abdulqadir J. Nashwan

Hamad Medical Corporation, Doha, Qatar.

Ghaleb Adwan

Department of Biology and Biotechnology, An-Najah National University, Nablus, Palestine.

Zaid Altaany

Department of Basic Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan.

Ayat Nashwan

Department of Sociology and Social Work, Yarmouk University, Irbid, Jordan.

Khaled Al-Waqfi

Department of Basic Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan.

Lujain Abuirsheid

Department of Basic Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan.

Raghad Ayasreh

Department of Basic Medical Sciences, Faculty of Medicine, Yarmouk University, Irbid, Jordan.

Mohammed Al Mutairi

Ministry of Health, Kuwait.

Ala'a B. Al-Tammemi

Department of Family and Occupational Medicine, Faculty of Medicine, University of Debrecen, Debrecen, Hungary.

Research Article

Keywords: Arab countries, attitude, COVID-19, hesitancy, prevalence, vaccination, EMR

Posted Date: February 1st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1280929/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Huge efforts are being made to control the spread and impacts of the coronavirus pandemic using vaccines. However, willingness to be vaccinated depends on factors beyond the availability of vaccines. The aim of this study was three-folded: to assess the prevalence of children's vaccination against COVID-19, to explore parents' attitudes towards children's vaccination, and to discover the factors associated with parents' hesitancy towards children's vaccination in several countries in the Eastern Mediterranean Region (EMR).

Methods: This study utilized a cross-sectional descriptive design. A sample of 3744 parents from eight countries, namely, Iraq, Jordan, Kuwait, Lebanon, Palestine, Qatar, Saudi Arabia (KSA), and the United Arab Emirates (UAE), was conveniently approached and surveyed using Google forms. The participants have responded to a 42-item questionnaire pertaining to socio-demographics, children vaccination status, knowledge about COVID-19 vaccines, and attitudes towards vaccinating children and the vaccine itself.

Results: The prevalence of vaccinated children against COVID-19 was 32% as indicated by the parents. Concerning parents' attitudes towards vaccines safety, about one third of participants (32.5%) believe that all vaccines are not safe. In the regression analysis, children's vaccination was significantly correlated with parents' age, education, occupation, parents' previous COVID-19 infection, and their vaccination status. Participants aged ≥ 50 years and those aged 40-50 years had an odds ratio of 17.9 (OR=17.9, CI: 11.16-28.97) and 13.2 (OR=13.2, CI: 8.42-20.88); respectively, for vaccinating their children compared to those aged 18-29 years. Parents who had COVID-19 vaccine were about five folds more likely to vaccinate their children compared with parents who did not receive the vaccine (OR=4.9, CI: 3.12-7.70). The prevalence of children's vaccination in the participating Arab countries is still not promising.

Conclusion: To encourage parents, vaccinate their children against COVID-19, Arab governments should strategize accordingly. Reassurance of the efficacy and effectiveness of the vaccine should target the general population using educational campaigns, social media, and official TV and radio channels.

1. Background

The Coronavirus Disease-2019 (COVID-19) was announced as a Public Health Emergency of International Concern (PHEIC) in January 2020, with 5.5 million deaths reported until January 2022 [1]. Most countries, around the World, have witnessed large numbers of cases despite the precautions taken [2], leading to severe impact including psychological burden [3, 4]. Efficacious vaccine is an essential tool in the fight against the current COVID-19 pandemic to achieve collective immunity, which can help reduce transmission, hospitalizations, and intensive care utilization, as well as prevent additional mortality [5].

Several COVID-19 vaccines are now available for public. However, major issues have been identified to affect vaccine coverage, including limited manufacturing capacity, global inequality in the distribution of COVID-19 vaccines, and most importantly vaccine hesitancy which could represent a major challenge in

the global efforts to control the pandemic [6, 7]. Vaccine hesitancy refers to a delay in accepting or refusing vaccination despite its availability [8].

Many studies around the world have reported vaccine hesitancy among the public and this was reflected in a systematic review involved 31 published studies [9]. One-third of participants in a study from four African and Middle Eastern countries were hesitant about getting vaccinated against the COVID-19 [10]. Similar results were reported from Bangladesh, India, Pakistan, Nepal [11], and Japan [12]. Arab countries also reported similar trends [13–15]. Certainly, individuals who are hesitant in getting vaccinated query the usefulness and safety of vaccines for their children [16]. In Italy, a cross-sectional study was conducted to evaluate the knowledge, attitude, and intention to vaccinate children < 18 years. They found that 41.2% of families of children ≥ 12 years and 36.1% among families of children < 12 years would not vaccinate their children [16]. Perceived vaccine safety and efficacy and perceived risk of transmission of infection to adults were the two determinants of intention to vaccinate for both age groups [16].

Many factors influence vaccination decisions, including trust in the government and health care professionals, social influences, high levels of knowledge about the vaccine, and general positive attitudes toward vaccines [17]. Another study found that parents with different levels of education and employment status had significant differences in vaccine knowledge and awareness; similarly, these two factors also significantly influenced parental vaccine hesitancy. The knowledge score of parents was inversely correlated with vaccine hesitancy and moderately associated with their awareness score [18].

In a study conducted in Poland regarding children vaccination, the main concerns expressed by parents were the possibility of adverse effects, the lack of adequate testing of the preparations in children, and the fear of future complications [19]. Therefore, the COVID-19 vaccine must have proven safety and efficacy in preventing complications as well as transmission to ensure vaccination of children [20].

In the Arab countries, like Jordan, commitment to precaution measures against COVID-19 virus such as hand washing, wearing a face mask, and social distancing is not optimal [21]. This was evident from a previous study which found that contaminated hands can act as carriers for different infectious microbes [22]. Though, figures of the vaccinated population against COVID-19 in Arab countries, including children, are still not promising. To our knowledge, no studies have been conducted in these countries to assess the attitudes of parents towards children vaccination against COVID-19 infection. The aims of this study were to (1) assess the prevalence of children's vaccination against COVID-19, (2) assess parents' attitudes toward children's vaccination, and (3) study factors associated with vaccination's status and hesitancy of parents towards children's vaccination in several Arab countries.

2. Methods And Materials

2.1 Study Setting, Design, and Participants

Our study was conducted in eight countries located in the Eastern Mediterranean Region (EMR), namely: Iraq, Jordan, Kuwait, Lebanon, Palestine, Qatar, Saudi Arabia (KSA), and the United Arab Emirates (UAE).

A descriptive questionnaire-based cross-sectional online survey was employed in this study focusing on married individuals having children (parents). A convenience sample was approached through social media platforms. A questionnaire was firstly developed in English, then it was translated into Arabic (the native language of our respondents) by two bilingual specialists. The questionnaire was then uploaded to Google Form® and disseminated to those who could access the online survey. The inclusion criteria included being a citizen of any participating country, aged 18 or older, married and having children, reading, and understanding Arabic, and being willing to fill the online questionnaire. The google form led participants who were single or not having children to submit their responses at early stage and their responses were excluded. An online link of the questionnaire, along with an introductory letter about the study, was sent to participants via media platforms like Facebook®, WhatsApp®, Twitter®, and email addresses, for a period of four weeks; 15-November- 13-December 2021. In addition, respondents were asked to share the questionnaire link with their relatives, friends, and social networks.

2.2 Study Instrument

The authors have reviewed the available literature and developed a 42-item questionnaire composed of three main sections. The first section consisted of 12 questions on participants' socio-demographic characteristics such as age, gender, marital status, having children, education, nature of work or study, and country of residence. The second section comprised four questions that solicited general knowledge about the COVID-19 vaccine. The third section of the questionnaire consisted of 26 items and inquired about participants' attitudes towards COVID-19 vaccination of their children and the vaccine itself.

2.3 Validity and Reliability

Two researchers checked the content of the questionnaire and its face validity before the final approval. To ensure its reliability, the questionnaire was pilot tested with the first 40 responses. Based on these responses and the feedback, refinements were made. The Cronbach's alpha score was found to be 0.81. The responses of the pilot-testing were excluded from the final analysis.

2.4 Ethical Considerations

All participants have given their informed consent through reading the following statement and ticking a box next to it: "Completing the questionnaire would be considered consent to voluntary participation". Participants were informed that the study would not disclose any personal information and that their data would be stored under high-security settings with only the research team having access to these data.

2.5 Statistical Analysis

The Statistical Package for Social Sciences (SPSS- IBM, Chicago, IL, USA) was used to analyze the data. Categorical variables were reported as frequency counts and percentages. A cross-tabulation analysis using the chi-square test was employed to assess significant differences between categorical variables. Finally, all statistically significant factors revealed from the cross-tabulation analysis were subjected to a

backward Wald stepwise binary logistic regression analysis to assess the independent effect of each factor after controlling for potential confounders. A P value < 0.05 was set for statistical significance.

3. Results

A total of 3744 participants with valid responses were involved in the final analysis. More than half of the sample (55.5%) was females (n=2078) and 44.5% (n=1666) was males. Interestingly, about 40% of the study population have been diagnosed with COVID-19 infection and the majority (83%, n=3108) have been vaccinated against the infection. The socio-demographic characteristics of the study population are shown in Table 1.

Table 1
Socio-Demographic Characteristics of the Study
Participants (N = 3744)

Characteristic	n (%)
Gender	
Male	1666 (44.5)
Female	2078 (55.5)
Age (years)	
18–29	331 (8.8)
30–39	1561 (41.7)
40–49	1340 (35.8)
> 50	512 (13.7)
Work or study field	
None health-related	2679 (71.6)
Health-related	1065 (28.4)
Education	
Secondary or less	544 (14.5)
≥ Graduate	3200 (85.5)
Received a COVID-19 vaccine	
No	256 (6.8)
Yes	3108 (83.0)
I intend to get the vaccine	173 (4.6)
I do not intend to get the vaccine	207 (5.5)
Country of Residence	
Jordan	997 (26.6)
KSA	419 (11.2)
UAE	333 (8.9)
Kuwait	475 (12.7)
Lebanon	424 (11.3)
Palestine	422 (11.3)

Characteristic	n (%)
Qatar	306 (8.2)
Iraq	368 (9.8)
Previous Diagnosis of COVID-19	
No	2223 (59.4)
Yes	1521 (40.6)

Table 2 shows the number of children, their age, and beliefs of the study population towards the COVID-19 vaccine for children. About one-third of the sample (32.5%) reported that all COVID-19 vaccines are not safe and close percentage reported that their children received the vaccine (31.9%, n=1194). Figure 1 shows children's vaccination prevalence per country and figure 2 shows the overall prevalence of children's vaccination in eight Arab countries.

Table 2

Number of children, their age, and beliefs of parents towards COVID-19 vaccine for the children (N = 3744)

Characteristic	n (%)
Number of children	
1	578 (15.4)
2	1022 (27.3)
≥3	2144 (57.3)
Age of children/year	
All kids are younger than 12	1817 (48.5)
All kids between 12-17	579 (15.5)
I have kids in both above categories	1348 (36.0)
If any of your kids had COVID-19 vaccine	
No	2550 (68.1)
Yes	1194 (31.9)
Will the freedom to choose the type of vaccine affect your decision to vaccinate your children?	
No	1211 (32.3)
Yes	1361 (36.4)
Maybe	1172 (31.3)
In your opinion, are COVID-19 vaccines safe?	
All vaccines are safe	2527 (67.5)
All vaccines are not safe	1217 (32.5)

To assess the impact of socio-demographic characteristics on the vaccination against COVID-19 among children, a cross-tabulation analysis was performed and revealed that all these factors were significantly

correlated with children’s vaccination status ($p \leq 0.05$). Interestingly, parents whose work or study was health-related showed a higher tendency for not vaccinating their children compared to others. Moreover, willingness to vaccinate increased with an increased parents’ age. Remarkably, less than 4% of parents whose children were younger than 12 years have vaccinated their children. The results of this analysis are shown in Table 3.

Table 3
Cross-tabulation of socio-demographic factors associated with COVID-19
vaccination among children (N = 3744)

Variable	Children vaccination status		P value
	No (%)	Yes (%)	
Gender			0.037
Male	1105 (66.3)	561 (33.7)	
Female	1445 (69.5)	633 (30.5)	
Participant's education			< 0.001
Secondary or less	311 (57.2)	233 (42.8)	
≥ Graduate	2239 (70.0)	961 (30.0)	
Work or study field			< 0.001
None health-related	1770 (66.1)	909 (33.9)	
Health-related	780 (73.2)	285 (26.8)	
Participant's age/year			< 0.001
18-29	309 (93.4)	22 (6.6)	
30-39	1335 (85.5)	226 (14.5)	
40-40	690 (51.5)	650 (48.5)	
≥50	216 (42.2)	296 (57.8)	
Number of children			< 0.001
1	458 (79.2)	120 (20.8)	
2	837 (81.9)	185 (18.1)	
≥3	1255 (58.5)	889 (41.5)	
Children age			< 0.001
All kids are younger than 12	1747 (96.1)	70 (3.9)	
All kids are between 12-17	254 (43.9)	325 (56.1)	
I have kids in both above categories	549 (40.7)	799 (59.3)	
If the parent had COVID-19 before			< 0.001
No	1462 (65.8)	761 (34.2)	
Yes	1088 (71.5)	433 (28.5)	

Variable	Children vaccination status		P value
	No (%)	Yes (%)	
If the parent had COVID-19 vaccine			< 0.001
No	231 (90.2)	25 (9.8)	
Yes	2011 (64.7)	1097 (35.3)	
I intend to get the vaccine	134 (77.5)	39 (22.5)	
I do not intend to get the vaccine	174 (84.1)	33 (15.9)	

As seen in Table 3, parents who have been vaccinated against COVID-19 and those who intend to get the vaccine in the future were more likely to have vaccinated children. With respect to the country of residence and perceived necessity for COVID-19 vaccine with vaccination status among children, parents living in KSA, UAE, Kuwait, and Iraq were more willing to vaccinate their children compared to parents from Jordan, Lebanon, Palestine, and Qatar. Furthermore, parents who believed that the vaccine is necessary for individuals younger than 18 years were more willing to vaccinate their children than those who didn't have the same belief. Table 4 illustrates these associations.

Table 4
Cross-tabulation of vaccine related factors and country of residence with COVID-19 vaccination among children (n = 3744)

Variable	Children vaccination status		P value
	No (%)	Yes (%)	
Country of residence			< 0.001
Jordan	817 (81.9)	180 (18.1)	
KSA	213 (50.8)	206 (49.2)	
UAE	170 (51.1)	163 (48.9)	
Kuwait	242 (50.9)	233 (49.1)	
Lebanon	348 (82.1)	76 (17.9)	
Palestine	330 (78.2)	92 (21.8)	
Qatar	234 (76.5)	72 (23.5)	
Iraq	196 (53.3)	172 (46.7)	
I advise other parents to vaccinate their children against COVID-19			< 0.001
No	1020 (83.6)	200 (16.4)	
Yes	706 (48.6)	747 (51.4)	
Neutral	824 (76.9)	247 (23.1)	
COVID-19 vaccine is necessary for those younger than 18 years			< 0.001
No	1318 (81.1)	308 (18.9)	
Yes	654 (49.8)	659 (50.2)	

Table 5 presents parents' attitudes towards COVID-19 vaccine and children's vaccination status. Amazingly, all attitudes' items had significant statistical associations with vaccination status ($p \leq 0.05$). About half of parents who believed that COVID-19 vaccines are safe for children had vaccinated their offspring compared to only about 20% of parents who did not have the same belief. In addition, parents who were against children's vaccines due to the unknown long-term side effects of the vaccine or perceived a lack of scientific research about the effects of the vaccine on children were less likely to vaccinate their children. Most importantly, willingness to vaccinate children decreased among parents who think that the vaccine may change human genes.

Table 5

Parents' attitudes towards COVID-19 vaccination of children and children vaccination status (n = 3744)

Variable	Children vaccination status		Total (n)	P value
	No (%)	Yes (%)		
COVID-19 vaccine protects children from the infection				< 0.001
No	1092 (81.0)	256 (19.0)	1348	
Yes	646 (51.7)	633 (48.3)	1249	
Neutral	812 (70.8)	335 (29.2)	1147	
COVID-19 vaccines are safe to be used on children				< 0.001
No	1115 (80.4)	271 (19.6)	1386	
Yes	468 (50.2)	465 (49.8)	933	
Neutral	967 (67.9)	458 (32.1)	1425	
I encourage vaccination of children against COVID-19				< 0.001
No	1225 (80.9)	290 (19.1)	1515	
Yes	647 (52.0)	598 (48.0)	1245	
Neutral	678 (68.9)	306 (31.1)	948	
A child can be infected with COVID-19 even after vaccination				< 0.001
No	122 (68.5)	56 (31.5)	178	
Yes	1957 (69.6)	856 (30.4)	2813	
Neutral	471 (62.5)	282 (37.5)	753	
A child who had COVID-19 infection do not need the vaccine				< 0.001
No	646 (58.4)	460 (41.6)	1106	
Yes	1017 (74.9)	341 (25.1)	1358	
Neutral	887 (69.3)	393 (30.7)	1280	
Natural immunity of the infection is better than vaccination				< 0.001
No	450 (56.2)	351 (43.8)	801	
Yes	1539 (76.0)	487 (24.0)	2026	

Variable	Children vaccination status		Total (n)	P value
	No (%)	Yes (%)		
Neutral	561 (61.2)	356 (38.8)	917	
A healthy child does not require COVID-19 vaccine				< 0.001
No	672 (57.5)	496 (42.5)	1168	
Yes	1250 (76.0)	395 (24.0)	1645	
Neutral	628 (67.5)	303 (32.5)	931	
COVID-19 is not serious and does not require a vaccine				< 0.001
No	1115 (60.6)	725 (39.4)	1840	
Yes	606 (77.5)	176 (22.5)	782	
Neutral	829 (73.9)	293 (26.1)	1122	
Face masks, social distancing, and sterilization are enough to prevent spread of COVID-19				< 0.001
No	644 (61.9)	397 (38.1)	1041	
Yes	1384 (72.1)	535 (27.9)	1919	
Neutral	522 (66.6)	262 (33.4)	784	
I'm against all children vaccines, not only the COVID-19 vaccine				< 0.001
No	1582 (65.0)	852 (35.0)	2434	
Yes	466 (77.8)	133 (22.2)	599	
Neutral	502 (70.6)	209 (29.4)	711	
COVID-19 vaccine should only be given to elderly and children with diseases				< 0.001
No	943 (59.9)	631 (40.1)	1574	
Yes	841 (74.2)	293 (25.8)	1134	
Neutral	766 (73.9)	270 (26.1)	1036	
The benefit of COVID-19 vaccine outweighs its side effects				< 0.001
No	757 (81.0)	178 (19.0)	935	
Yes	805 (58.2)	577 (41.8)	1382	

Variable	Children vaccination status		Total (n)	P value
	No (%)	Yes (%)		
Neutral	988 (69.2)	439 (30.8)	1427	
I'm against children vaccine as long-term side effects of the vaccine aren't known				< 0.001
No	411 (52.6)	370 (47.4)	781	
Yes	1407 (78.0)	397 (22.0)	1804	
Neutral	732 (63.2)	427 (36.8)	1159	
Children should be vaccinated against COVID-19 only when advised by a physician				< 0.001
No	851 (81.4)	195 (18.6)	1046	
Yes	969 (58.2)	696 (41.8)	1665	
Neutral	730 (70.7)	303 (29.3)	1033	
The government have the right to impose COVID-19 vaccine on all citizens				< 0.001
No	1471 (78.3)	408 (21.7)	1879	
Yes	570 (55.4)	458 (44.6)	1028	
Neutral	509 (60.8)	328 (39.2)	837	
COVID-19 vaccine will end the pandemic				< 0.001
No	1193 (78.7)	322 (21.3)	1515	
Yes	481 (54.4)	404 (45.6)	885	
Neutral	876 (65.2)	468 (34.8)	1344	
Children should not be vaccinated against COVID-19 because vaccination is painful				< 0.001
No	1276 (62.2)	777 (37.8)	2053	
Yes	536 (77.8)	153 (22.2)	689	
Neutral	738 (73.7)	264 (26.3)	1002	
Children should not be vaccinated against COVID-19 due to lack of scientific research about effects of the vaccine on children				< 0.001
No	363 (52.2)	332 (47.8)	695	
Yes	1568 (77.8)	448 (22.2)	2016	

Variable	Children vaccination status		Total (n)	P value
	No (%)	Yes (%)		
Neutral	619 (59.9)	414 (40.1)	1033	
COVID-19 vaccine is not effective because of the frequent mutations of the virus				< 0.001
No	525 (57.5)	388 (42.5)	913	
Yes	1131 (78.5)	309 (21.5)	1440	
Neutral	894 (64.3)	497 (35.7)	1391	
COVID-19 vaccine changes human genes				< 0.001
No	891 (63.0)	523 (37.0)	1414	
Yes	472 (77.8)	135 (22.2)	607	
Neutral	1187 (68.9)	536 (31.1)	1723	

In the binary regression analysis, the vaccination status was categorized into two main levels; (Yes) and (No). Children's vaccination was significantly correlated with parents' age, education, occupation, parents' previous COVID-19 infection, and their vaccination status. Participants aged ≥ 50 years and those aged 40-50 years had an OR of 18 (OR=17.9, CI: 11.16-28.97) and 13 (OR=13.2, CI: 8.42-20.88) for vaccinating their children compared to those aged 18-29 years. Significantly, parents who had COVID-19 vaccine were five folds more likely to vaccinate their children compared with parents who didn't receive the vaccine (OR=4.9, CI: 3.12-7.70). Table 6 presents the results of the binary logistic regression analysis.

Table 6
Logistic regression analysis of factors associated with increased odds of having vaccinated children

Variable	Odds ratio	95% Confidence interval		P value
		Lower	Upper	
Participant's age				
18-29	Ref *			
30-39	2.2	1.43	3.62	<0.001
40-40	13.2	8.42	20.88	<0.001
≥50	17.9	11.16	28.97	<0.001
Participant's education				
Secondary or less	2.1	1.46	2.79	<0.001
≥ Graduate	Ref *			
Work or study field				
None health-related	1.3	1.06	1.52	0.009
Health-related	Ref *			
If the parent had COVID-19 infection before				
No	1.3	1.07	1.47	0.005
Yes	Ref *			
If the parent had COVID-19 vaccine				
No	Ref *			
Yes	4.9	3.12	7.70	<0.001
Intend to get the vaccine	2.7	1.49	4.94	0.001
Don't intend to get the vaccine	1.8	1.01	3.35	0.044
* Reference for other categories.				

4. Discussion

The first aim of this study was to assess the prevalence of vaccinated children against COVID-19 in eight Arab countries located in the Eastern Mediterranean Region. The results of this study showed that the prevalence of vaccinated children is 31.9%. This preliminary finding is similar to results from a previous study in the Singapore population [23]. Literature about the vaccine coverage amongst children is dearth

and most published studies have assessed the parents' willingness or intention to vaccinate children rather than true vaccination status. The second aim was to explore parents' attitudes towards vaccinating children against COVID-19 in the Arab countries. The results of this study showed that around one third of the participants (32.5%) believe that all vaccines are not safe. Despite the majority of the parents (83.0%) have been already vaccinated against COVID-19, most of them didn't vaccinate their children against COVID-19 and this correlated with many factors as discussed below.

The third aim of this study was to explore factors associated with vaccination and hesitancy of parents towards children's vaccination in several Arab countries. The results of this study showed significant correlations between children's vaccination status and participants' sociodemographic characteristics. For instance, older parents showed extreme higher readiness to vaccinate their children against COVID-19 than younger counterparts. This result is congruent with a previous study which found that parents younger than 40 are more hesitant to vaccinate children [24]. However, this finding mismatches a previous study which showed that parents younger than 35 in Latin American countries are less likely to be hesitant to vaccinate children [25]. In terms of children's number, parents with three or more children were more willing to vaccinate them than their counterparts. Participants of children aged between 12 and 17 years were more ready to vaccinate them than their counterparts. Our findings might be explained by the fact that younger parents usually have younger children (less than 12 years). Despite its approval, vaccination of the children younger than 12 years is still questionable among public.

In terms of gender, fathers showed a higher tendency towards vaccinating their kids than mothers. This result is consistent with a previous study from Lebanon [14] and US [26]. However, the result was different from a previous study in Poland which showed that mothers are more enthusiastic to vaccinate children than fathers [19]. This may be explained by the difference in the social role of fathers or mothers between communities.

Interestingly, parents whose work or study was health-related showed a lower tendency towards vaccinating their children than parents working in non-health-related fields. This result is consistent with a previous study stated that fewer caregivers plan to vaccinate their children against COVID-19 [27]. Also, parents with a lower educational level showed higher readiness towards vaccinating their children than parents with a higher educational level. This finding is consistent with a previous study which showed that undergraduate parents are more enthusiastic to vaccinate children than parents with higher education [10, 25, 28]. This finding is also congruent with previous studies which concluded that parents' educational level is one of the determinants of willingness to vaccinate children [18, 29].

Parents who haven't been infected with COVID-19 were more eager to vaccinate their children than other participants. This finding can be explained by their carefulness from being infected with COVID-19. Another explanation to this result may refer to the fear of the vaccine's side effects perceived by those who have already got the infection before as the side effects can be similar to symptoms of the infection itself. This result is supported by previous studies. For example, a study showed that parents who are anxious from COVID-19 are keener to vaccinate children [28]. Another study showed that the perceived

danger of being infected with COVID-19 is one of the determinants for parents' willingness to vaccinate children [16]. Additionally, another research found that the readiness to get the COVID-19 vaccine increases with a higher perceived susceptibility to COVID-19 [30]. Vaccinated parents and those who intend to get the vaccine in the future were more willing to vaccinate their children than unvaccinated parents and those who do not intend to get the vaccine. The reason behind this difference can be explained by awareness of vaccinated parents about the COVID-19 vaccines which is supported by an earlier study [28].

Results of the current study showed that parents who had received COVID-19 vaccine or those who intend to get the vaccine are more likely to vaccinate their children. These results are consistent with results from another study stated that the likelihood of child vaccination was greater among parents who had already received or were likely to receive a COVID-19 vaccine [31].

Moreover, parents' attitudes towards vaccinating children are significantly associated with their children's vaccination status. For example, almost half of parents who believe that COVID-19 vaccines are safe on children had vaccinated their children. This result is similar to the rates found in a previous scoping review [29]. On the other hand, only one-fifth of parents who don't trust safety of COVID-19 vaccines on children had vaccinated their children.

The results showed that parents who were against children vaccination because of the unknown long-term side effects of the vaccine were less willing to vaccinate their children. This finding matches previous research which found that one of the main parents' concerns is the possible future complications of the vaccine [19]. Similarly, parents who think that the vaccine may change human genes were less eager to vaccinate their children. These findings were consistent with a previous study which showed that vaccination concerns determine parents' willingness to vaccinate children against COVID-19 [23]. These results are consistent with results from Palestine which reported that adequate information about vaccines and their risk–benefit ratios are important to build trust and favorable attitudes towards vaccines [15].

The results of the current study showed also that willingness towards vaccinating children is significantly different depending on the country of residence. For instance, parents from KSA, UAE, Kuwait, and Iraq were more enthusiastic to vaccinate their children than those from Jordan, Lebanon, Palestine, and Qatar. A previous study has shown high rates of vaccine hesitancy in Jordan and Kuwait [32].

5. Conclusion

The prevalence of children vaccinated against COVID-19 in Arab countries is not optimal. Moreover, parents in the studied countries are still hesitant to vaccinate their children against the COVID-19. This hesitancy is led by certain parents' socio-demographic characteristics, having COVID-19 infection before, and their vaccination status. To increase COVID-19 vaccination rates in the countries located in the EMR, it is essential to address variables connected to parents' hesitancy to vaccinate children. A probable cause of vaccination hesitancy may be also linked to misleading information shared on media platforms.

This explanation was addressed in a previous study done on adolescents and parents of adolescents in the USA which showed that social media negatively impacted the opinions of adolescents and parents on COVID-19 vaccination [33].

Face-to-face information or educational interventions should be used to help parents understand why vaccines are important; explain where, how, and when to access service and respond to hesitations and concerns about vaccine safety or effectiveness. Such interventions are interactive and can be tailored to target specific populations or identified barriers. Concise and clear vaccine information should be provided in multiple languages to improve vaccine confidence. Targeted campaigns should be implemented for parents that are socioeconomically disadvantaged and that are less educated.

Abbreviations

COVID-19: Coronavirus Disease-2019; EMR: Eastern Mediterranean Region; KSA: Kingdom of Saudi Arabia; OR: Odd Ratio; PHEIC: Public Health Emergency of International Concern; UAE: United Arab Emirates.

Declarations

Ethics approval and consent to participate:

The protocol of this study was reviewed and approved by Yarmouk University's Institutional Review Board (IRB) Committee (IRB/2021/47). Written Informed consent (electronically) was obtained from all participants. In addition, all the methods were performed conforming to the World Medical Association Declaration of Helsinki. Participation in this study was voluntary. Informed consent was ensured by the presence of an introductory section in the survey used in this study, with the submission of responses implying an agreement to participate.

Consent for publication:

Not Applicable.

Availability of data and materials:

The dataset generated and analyzed in this study is not publicly available due to institutional guidelines but is available from the corresponding author on reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding:

This research project did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions:

Conceptualization: MK, HK, ABA, WM, AA.; Methodology: MK, HK, ABA, AA; Validation, MK; Formal analysis: MK.; Investigation: all authors; Data curation: all authors; Writing—original draft preparation: MK, HK, ZT.; Writing—review and editing: all authors; Visualization: MK.; Supervision: MK, AA, ABA; Project administration: MK, M-AM. All authors have read and approved to the submitted version of the manuscript.

Acknowledgments:

We would like to thank all the participants in this survey.

References

1. Johns Hopkins University Center. COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins. 2022. <https://coronavirus.jhu.edu/map.html>. Accessed on January 12, 2022.
2. Khatatbeh M. The Battle Against COVID-19 in Jordan: From Extreme Victory to Extreme Burden. *Front Public Heal*. 2021;8.
3. Khatatbeh M, Alhalaiqa F, Khasawneh A, Al-tammemi AB, Khatatbeh H, Alhassoun S, et al. The Experiences of Nurses and Physicians Caring for COVID-19 Patients: Findings from an Exploratory Phenomenological Study in a High Case-Load Country. *Int J Environ Res Public Health*. 2021;18:1–17.
4. Khatatbeh M, Khasawneh A, Hussein H, Altahat O, Alhalaiqa F. Psychological Impact of COVID-19 Pandemic Among the General Population in Jordan. *Front Psychiatry*. 2021;12:618993.
5. Hodgson SH, Mansatta K, Mallett G, Harris V, Emary KRW, Pollard AJ. What defines an efficacious COVID-19 vaccine? A review of the challenges assessing the clinical efficacy of vaccines against SARS-CoV-2. *The Lancet Infectious Diseases*. 2021;21:e26–35.
6. Al-Tammemi AB, Tarhini Z. Beyond equity: Advocating theory-based health promotion in parallel with COVID-19 mass vaccination campaigns. *Public Heal Pract*. 2021;2:100142.
7. Neumann-Böhme S, Varghese NE, Sabat I, Barros PP, Brouwer W, van Exel J, et al. Once we have it, will we use it? A European survey on willingness to be vaccinated against COVID-19. *European Journal of Health Economics*. 2020;21:977–82.
8. MacDonald NE, Eskola J, Liang X, Chaudhuri M, Dube E, Gellin B, et al. Vaccine hesitancy: Definition, scope and determinants. *Vaccine*. 2015;33:4161–4.
9. Sallam M. Covid-19 vaccine hesitancy worldwide: A concise systematic review of vaccine acceptance rates. *Vaccines*. 2021;9:160.

10. Faezi NA, Gholizadeh P, Sanogo M, Oumarou A, Mohamed MN, Cissoko Y, et al. Peoples' attitude toward COVID-19 vaccine, acceptance, and social trust among African and Middle East countries. *Heal Promot Perspect*. 2021;11:171–8.
11. Hawlader MDH, Rahman ML, Nazir A, Ara T, Haque MMA, Saha S, et al. COVID-19 vaccine acceptance in South Asia: a multi-country study. *Int J Infect Dis*. 2022;114:1–10.
12. Machida M, Nakamura I, Kojima T, Saito R, Nakaya T, Hanibuchi T, et al. Acceptance of a covid-19 vaccine in japan during the covid-19 pandemic. *Vaccines*. 2021;9:210.
13. Alduwayghiri EM, Khan N. Acceptance and Attitude toward COVID-19 Vaccination among the Public in Saudi Arabia: A Cross-sectional Study. *J Contemp Dent Pract*. 2021;22:730–4.
14. Kasrine Al Halabi C, Obeid S, Sacre H, Akel M, Hallit R, Salameh P, et al. Attitudes of Lebanese adults regarding COVID-19 vaccination. *BMC Public Health*. 2021;21:1–7.
15. Kateeb E, Danadneh M, Pokorná A, Klugarová J, Abdulqader H, Klugar M, et al. Predictors of willingness to receive covid-19 vaccine: Cross-sectional study of palestinian dental students. *Vaccines*. 2021;9:954.
16. Russo L, Croci I, Campagna I, Pandolfi E, Villani A, Reale A, et al. Intention of Parents to Immunize Children against SARS-CoV-2 in Italy. *Vaccines*. 2021;9:1469.
17. Smith LE, Amlôt R, Weinman J, Yiend J, Rubin GJ. A systematic review of factors affecting vaccine uptake in young children. *Vaccine*. 2017;35:6059–69.
18. Voo JYH, Lean QY, Ming LC, Hanafiah NHM, Al-Worafi YM, Ibrahim B. Vaccine knowledge, awareness and hesitancy: A cross sectional survey among parents residing at sandakan district, sabah. *Vaccines*. 2021;9:1348.
19. Babicki M, Pokorna-Kałowak D, Doniec Z, Mastalerz-Migas A. Attitudes of parents with regard to vaccination of children against covid-19 in Poland. A nationwide online survey. *Vaccines*. 2021;9:1192.
20. Eberhardt CS, Siegrist CA. Is there a role for childhood vaccination against COVID-19? *Pediatric Allergy and Immunology*. 2021;32:9–16.
21. Khatatbeh M, Al-Maqableh HO, Albalas S, Al Ajlouni S, A'aqoulah A, Khatatbeh H, et al. Attitudes and Commitment Toward Precautionary Measures Against COVID-19 Amongst the Jordanian Population: A Large-Scale Cross-Sectional Survey. *Front Public Heal*. 2021;9.
22. Al Momani W, Khatatbeh M, Altaany Z. Antibiotic susceptibility of bacterial pathogens recovered from the hand and mobile phones of university students. *GERMS*. 2019;9:9–16.
23. Griva K, Tan KYK, Chan FHF, Periakaruppan R, Ong BWL, Soh ASE, et al. Evaluating Rates and Determinants of COVID-19 Vaccine Hesitancy for Adults and Children in the Singapore Population: Strengthening Our Community's Resilience against Threats from Emerging Infections (SOCRATES) Cohort. *Vaccines*. 2021;9:1415.
24. Zona S, Partesotti S, Bergomi A, Rosafio C, Antodaro F, Esposito S. Anti-COVID vaccination for adolescents: A survey on determinants of vaccine parental hesitancy. *Vaccines*. 2021;9:1309.

25. Urrunaga-Pastor D, Herrera-Añazco P, Uyen-Cateriano A, Toro-Huamanchumo CJ, Rodriguez-Morales AJ, Hernandez A V., et al. Prevalence and factors associated with parents' non-intention to vaccinate their children and adolescents against COVID-19 in Latin America and the Caribbean. *Vaccines*. 2021;9:1303.
26. Litaker JR, Tamez N, Bray CL, Durkalski W, Taylor R. Sociodemographic Factors Associated with Vaccine Hesitancy in Central Texas Immediately Prior to COVID-19 Vaccine Availability. *Int J Environ Res Public Health*. 2022;19:368.
27. Goldman RD, Krupik D, Ali S, Mater A, Hall JE, Bone JN, et al. Caregiver willingness to vaccinate their children against COVID-19 after adult vaccine approval. *Int J Environ Res Public Health*. 2021;18:10224.
28. Bono SA, Siau CS, Chen WS, Low WY, Faria E, Villela DM, et al. Adults ' Acceptance of COVID-19 Vaccine for Children in Selected Lower- and Middle-Income Countries. *Vaccines*. 2022;10:11.
29. Pan F, Zhao H, Nicholas S, Maitland E, Liu R, Hou Q. Parents' Decisions to Vaccinate Children against COVID-19: A Scoping Review. *Vaccines*. 2021;9:1476.
30. Patwary MM, Bardhan M, Disha AS, Hasan M, Haque MZ, Sultana R, et al. Determinants of COVID-19 Vaccine Acceptance among the Adult Population of Bangladesh Using the Health Belief Model and the Theory of Planned Behavior Model. *Vaccines*. 2021;9:1393.
31. Szilagyi PG, Shah MD, Delgado JR, Thomas K, Vizueta N, Cui Y, et al. Parents' intentions and perceptions about COVID-19 vaccination for their children: Results from a national survey. *Pediatrics*. 2021;148.
32. Sallam M, Dababseh D, Eid H, Al-Mahzoum K, Al-Haidar A, Taim D, et al. High rates of covid-19 vaccine hesitancy and its association with conspiracy beliefs: A study in jordan and kuwait among other arab countries. *Vaccines*. 2021;9:42.
33. Middleman AB, Klein J, Quinn J. Vaccine Hesitancy in the Time of COVID-19: Attitudes and Intentions of Teens and Parents Regarding the COVID-19 Vaccine. *Vaccines*. 2022;10:4.

Figures

Figure 1

Prevalence of children vaccination against COVID-19 per country.

Overall prevalence of children's vaccination in eight Arab countries

Figure 2

Overall prevalence of children vaccination against COVID-19 in eight Arab countries.