

Establishment and evaluation of specific antibiotic-induced Inflammatory bowel disease (IBD) model in rats

Jun Guo Tong (✉ tjg97114@sina.com)

Huzhou Central Hospital <https://orcid.org/0000-0001-7560-8641>

Hai Qian

Huzhou Central Hospital

Dongli Li

Huzhou Central Hospital

Jing Li

Huzhou Central Hospital

Jing Chen

Huzhou Central Hospital

Xiongfeng Li

Huzhou Central Hospital

Research article

Keywords: Antibiotic, Enteritis model, Rats, PCR, IBD

Posted Date: January 29th, 2020

DOI: <https://doi.org/10.21203/rs.2.22238/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at PLOS ONE on February 22nd, 2022. See the published version at <https://doi.org/10.1371/journal.pone.0264194>.

Abstract

Background: Physical and chemical methods have been established for rat enteritis model, but antibiotic induction has been relatively rare. This article aims to establish and evaluate rat model of Inflammatory bowel disease (IBD) using antibiotics.

Methods: Fourteen SD rats were divided into A-G group according to the dosage and method of antibiotics, among which group A was the control group. The drug was stopped on the 7th day, the modeling period was 1-7 days, and the recovery period was 8-15 days. Half of the animals were dissected on 11 th day and the other animals were dissected on 15 th day . Record the food and water intake, body weight, and fecal weight for 2 hours on different days. Nine intestinal flora were analyzed by bacterial culture and three strains were analyzed by quantitative PCR. TNF- α , IL1- β , IL-6 and CRP in abdominal aorta blood were detected and analyzed. Colon and rectal tissues were pathologically examined for inflammation and scored.

Results: Rat weight, food intake, water intake, and two-hour feces were significantly different ($P = .04$, $.016$, $< .001$, $.009$). Compared with group A, there were significant differences in 9 kinds of flora in the experimental group (all $P <.001$). *Bacteroides* , *Faecalibacterium prausnitzii* , and *Dialister invisus* concentrations were analyzed by Quantitative real time polymerase chain reaction (q-PCR) and showed significant differences in groups A, C, and F ($p = .033$). There were significant differences about TNF- α , IL1- β , IL-6 and CRP between the groups ($P = .016$, $.042$, $.037$, $.012$). The colonic and rectal pathological inflammation scores of other groups were significantly different from those of the control group (all $P <.001$).

Conclusion : Specific antibiotic-induced IBD model in SD rats is feasible.

Background

Inflammatory bowel disease (IBD) includes Crohn's disease (CD) and ulcerative colitis (UC), which are common intestinal diseases of patients, and they are increasing year by year.^[1] Clinically, this type of patients has a slow treatment effect and a long treatment period, which have attracted the attention of gastrointestinal physicians. The disease is considered a patient's immune system disorder, and changes in the intestinal flora and its metabolites usually affect the patient's immune system, thereby inducing the disease.^[2, 3] In order to have a deeper understanding of the disease, many scholars use animal models to study the disease.^[4] Rats are good mammals for research.^[5] Sprague Dawley (SD) rats are easy to raise and are easily controlled due to their mild temperament. They have been used for the establishment of various disease models. At present, rat enteritis models mostly use physical and chemical factors.^[6, 7] However, these methods easily lead to intestinal perforation and intestinal necrosis in rats, which can lead to failure of modeling. New modeling methods need to be studied and discussed. Because intestinal flora imbalance may cause enteritis, we thought of using flora imbalance for modeling. Antibiotics often cause imbalances in the intestinal flora.

Therefore, in this article, we used different doses of clindamycin (single use) and different doses of clindamycin plus ampicillin and streptomycin (combination) to cause intestinal flora disturbance in rats. The inflammatory factors in the abdominal aorta and the inflammation of the colon and rectum were analyzed in order to assess whether modeling is feasible.

Methods

Rats

Sprague Dawley (SD) female rats ($n=14$), obtained from Liaoning Changsheng Biotechnology Co., Ltd. ranged in size from 172.4 to 179.5 g. All animals were introduced to quarantine and adaptive feeding in the Specific Pathogen Free (SPF) barrier system of this institution for 9 days. All procedures were performed in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals (NIH Publications No. 80-23) revised in 1996. All experimental procedures and animal handling were performed in accordance with the guidelines of the International Association for the Study of Pain and the animal protocols were approved by Xi'an United Nations Quality Detection Technology CO.,Ltd Animal Committee. The authors tried all efforts to minimize the number of animals used.

SD rats were anesthetized intraperitoneally with 2% pentobarbital sodium (0.2 ml / 100 g). After about 15 ml of arterial blood was drawn from the abdominal aorta using a syringe (of which 5 ml of arterial blood was collected for inflammatory factor detection), the rat's abdominal aorta stopped beating, the pupils were dilated and the SD rats were ensured to be euthanized.

Experimental drugs, reagents, and instruments

The used drugs were used in our experiments: clindamycin hydrochloride(Shanghai Maclean Biotechnology Co., Ltd.) and 99% purity, ampicillin (Hebei Bailingwei Superfine Material Co., Ltd.) and streptomycin (Tianjin Guangxia Fine Chemicals Institute) with \geq 90% purity.

We also used rat IL-1 β , rat IL-6, rat TNF- α , and rat CRP reagents from Bioswamp (Wuhan, China). We purchased soil genomic DNA rapid extraction kit, rapid competent cell preparation kit (one-step method), SanPrep column plasmid DNA small amount extraction kit, SanPrep column DNA gel recovery kit from Sangon Biotech (Shanghai), purchased Taq Plus DNA Polymerase, Agarose B, 4S Red Plus Nucleic Acid Stain (10,000X Aqueous Solution) from BBI, and GeneRuler DNA Ladder Mix (from Thermo Scientific), pMD® 18-T Vector (from TAKARA Bio Ink)

Bacteroides-Bile-Enterprise Agar (BBE), *Lactobacillus* selective agar, Anaerobic bacteria agar, TPY agar medium, mannitol sodium chloride agar medium, Eosin Methylene Blue agar (EMB), CATC agar, Reinforced Clostridium Culture Medium were purchased and utilized for bacterial culture.

The electric day constant temperature incubator was purchased from Tianjin Taisite Instrument Co., Ltd. (Tianjin City, China). The electric day constant temperature incubator was purchased from Shanghai Yiheng Scientific Instrument Co., Ltd. (Shanghai, China). The Lab systems Multiskan MS micro plate

reader was purchased from Thermo Fisher (Pittsburgh, PA, USA). The low-speed condensation centrifuge, was purchased from Shanghai Luxiangyi Centrifuge Instrument Co., Ltd. (Shanghai, China). The electronic balance was purchased from Yuyao Jinnuo Tianping Instrument Co., Ltd. (Zhejiang Sheng, China). The electronic balance was purchased from Sedolis Instrument Co., Ltd. (China). The upright microscope was purchased from Japan Nikon Guangxuan Microscope Manufacturing Co., Ltd. (Tokyo, Japan).

In order to do real-time quantitative PCR, we used the following instruments: clean bench (purchased Jiangsu Su Clean Chemical Equipment Factory), high-speed refrigerated centrifuge (Anhui Zhongke Zhongjia Instrument Co., Ltd.), electrophoresis instrument (Beijing Liuyi), electrophoresis tank (Shanghai Jingyi Plexiglass Products Instrument Factory), Gel Imaging System (Shanghai Furi Technology Co., Ltd.), Micro-Spectrophotometer (Merinton Instrument, Inc), PCR Reaction Amplifier (BIO), Pipette (range 100-1000ul, 20-200ul, 0.5-10ul) (BBI, Canada), sequencer (ABI, Foster, CA, USA), StepOne fluorescence quantitative PCR instrument (ABI, Foster, CA, USA).

Animal grouping and modeling

Group A was the control group, group B was the low-dose clindamycin group (250 mg/kg), group C was the middle-dose clindamycin group (500 mg/kg), and group D was the high-dose clindamycin group (750 mg/kg). Group E was the low-dose triple antibiotic group (clindamycin, ampicillin, and streptomycin; 250 mg/kg, 272.1 mg/kg, and 136.1 mg/kg, respectively). Group F was the medium-dose triple antibiotic group (clindamycin, ampicillin, and streptomycin; 500 mg/kg, 563.7 mg/kg, and 281.8 mg/kg, respectively). Group G was the high-dose triple antibiotic group (clindamycin, ampicillin, and streptomycin; 750 mg/kg, 835.8 mg/kg, and 417.9 mg/kg, respectively). The experiment was divided into two stages: the modeling period (days 1-7) and the recovery period (days 8-15). The administration volume was 10 ml/kg once a day through stomach feeding by oral needle during the modeling period between 8:30-10:00 AM. The intragastric administration was stopped at 8th day. The weight, food-intake volume, water-intake volume, and stool samples were taken on days 1, 3, 5, 7, 9, 11, and 14 within 2 hours were collected. For each rat, the fecal microbial flora on the 1st, 4th, 8th, 11th, and 14th days were examined. On day 11 and day 15, half of each animal was dissected with 2% pentobarbital sodium (0.2 ml/100 g) by intraperitoneal anesthesia injection. After SD rats were euthanized, we quickly removed the colon and rectal tissue for next experiments. Rat (D2300) abnormally died on day 13 and we failed to collect fecal volume within 2 hours of antibiotic use, food intake and water intake on days 13 and 14, and blood on day 15. (Fig.1A-C)

Bacterial Culture

The microbial species detected were *Staphylococcus aureus*, *Bifidobacterium*, yeast, *Bacteroides*, *Clostridium*, anaerobic bacteria, *E. coli*, *Enterococcus*, and *Lactobacillus*.

We Mixed 1 g of feces with 9 ml of tryptone soy broth, diluted to the appropriate concentration, and took 20 ul of the sample and spread it evenly on the agar medium using a coating bar. Organisms were plated

onto mannitol sodium chloride agar medium plates, EMB, and CATC agar plates under aerobic conditions at 37 °C for 48 hours. Organisms cultured on TPY agar medium plates, BBE agar plates, reinforced Clostridium medium plates, anaerobic agar plates, and lactobacillus selective agar plates were cultured at 37 °C for 48 hours in anaerobic conditions. Organisms inoculated on DRBC agar plates were cultured for 5 days at 28 °C in aerobic conditions. Colonies were enumerated using the following formula: number of colonies (cfu/g) = number of plate colonies × 50 × dilution factor, with $\times 10^6$ (E6) as a uniform unit.

Real-time quantitative PCR analysis

Soil genomic DNA rapid extraction kit (B518233, Shengong Biological Co., Ltd., Shanghai) was used to extract fetal DNA from SD rats. The process is as follows:1. Weigh 400 mg of SD rat feces, add 400 μ l of 65 °C pre-warmed Buffer SCL, mix by shaking, and place in 65 °C water bath for 5 min.2. Centrifuge at 12,000 rpm for 3 minutes at room temperature. Pipette 350 μ l of the supernatant into a clean 1.5 ml centrifuge tube.3. Add equal volume of Buffer SP, mix by inversion, and ice bath for 10 min.4. Centrifuge at 12,000 rpm for 3 minutes at room temperature. Pipette 350 μ l of the supernatant into a clean 1.5 ml centrifuge tube.5. Add 200 μ l of chloroform, mix well, and centrifuge at 12,000 rpm for 5 minutes. Pipette the upper aqueous phase into a clean 1.5 ml centrifuge tube.6. Add 2 volumes of absolute ethanol, invert 8 times to mix thoroughly, and leave at room temperature for 3 min. Centrifuge at 10,000 rpm for 5 min at room temperature and discard the supernatant.7. Add 1 ml of 75% ethanol, rinse by inversion for 3 minutes, centrifuge at 10,000 rpm for 2 minutes, and discard the supernatant.8. Repeat step 7 once.9. Open the lid and invert for 10 minutes at room temperature until the residual ethanol is completely evaporated.10. Dissolve the resulting DNA in 70 μ l TE Buffer. The extracted DNA can be immediately used for further experiments. qPCR experiment was performed by pMD™18-T Vector Cloning Kit(TAKARA BIO INC, Dalian,China) ..Bacteroides, Faecalibacterium prausnitzii, Dialister invisus in rats fecal were collected and analyzed by PCR. After PCR, electrophoresis analysis was performed using 1.5% agarose syrup electrophoresis map.

Analysis of inflammatory factors

On the 11th and 15th days, half of the rats were dissected in each group with 2% pentobarbital sodium (0.2 ml/100 g) by intraperitoneal anesthesia injection, and the abdomen was cut, and 5mL blood was drawn from the abdominal aorta into the blood collection tube using syringe. TNF- α , IL-1 β , IL-6, and CRP was detected in the blood serum without diluting by an enzyme-linked immune sorbent assay (ELISA).

Colon and rectal pathological inflammation assessment

0 points: no inflammation; 1 point: a small amount of multifocal neutrophil infiltration (<10 per HPF); 2: a moderate multifocal neutrophil infiltration (more submucosal involvement) (10 – 50 / HPF); 3 points: a large number of multifocal and even aggregated neutrophils infiltration (more submucosa involvement and muscle layer) (> 50 / HPF); 4 points: the lesions involved the same 3 points, but abscesses or more extensive muscle layer involvement occurred.

1 + 2 + 3 Score <3 is grade I: mild; 1+ 2 + 3 Score is 4 to 6 is grade II: moderate; 1+ 2 + 3 A score of 7 to 9 is grade III: severe; 1 + 2 + 3 A score of > 10 is grade IV: extremely severe.

Ethics

This study follows the Basel Declaration 2010 and Institutional Animal Care and Use Committee (IACUC) of Xi'an United Nations Quality Detection Technology CO.,Ltd. We use animals to a minimum in terms of animal welfare principles and without affecting the accuracy of the experiment. All applicable international ,national and/or institutional guidelines for care and use of animals were followed.

Statistical analysis

Primer Premier 5.0 software was used for sequencing primer design. SPSS 21 software (IBM, USA)was utilized to analyze all data. Measured data were analyzed by ANOVA and F-test. Crosstabs were also used for measurement data analysis. Categorical variables were used by crosstabs and chi-square test. The independent sample T-test was used for comparative analysis between two sets of measured variables.

Results

Comparison of basic indice

The average starting weight of all rats was 17.26 ± 2.49 g, and there was no significant difference in weight between groups A-G ($P <0.05$). There were significant differences from groups A (control) to G (treated) in weight, food intake, water intake, and stool 2 hours post antibiotic use ($P =0.04, 0.016, <0.01$, and 0.009, respectively). Means and standard deviations are shown in Table 1 and Fig.2(A-D).

Comparison of nine species bacterial between groups

Staphylococcus aureus, *Bifidobacterium*, yeast, *Bacteroides*, *Clostridium*, anaerobic bacteria, *E. coli*, *Enterococcus*, *Lactobacillus* were cultured and counted using special medium.(Fig.3). All collected bacterial loads for these 9 species according to above methods were counted and compared between A group to B-G groups. The unit of bacteria in the stool is (CFU / g).The details were as followed: (A): 1376.7 ± 3683.8 (95% CI: 562.15-2191.26); (B): 687.06 ± 1498.74 (95% CI: 355.65-1018.45); (C): 1474.89 ± 4187.53 (95% CI: 548.96-2400.83); (D): 478.17 ± 1758.11 (95% CI: 65.03-891.31); (E): 664.50 ± 1567.91 (95% CI: 317.80-1011.19); (F): 403.77 ± 1171.99 (95% CI: 144.62-662.92); (G): 3609.76 ± 21206.52 (95% CI: -1079.38 ± 8298.91).The compared results were all $<.001$ (Table 2). Fig. 4 (A-E) showed the comparison results of the nine species of bacteria in each group on days 1, 4, 8, 11, and 14 (all $P<.001$).

Real-time quantitative PCR analysis of three strains

Bacteroides forward-primer (5'-3') is TTAAGTATTCCACCTGGGGAGT and reverse-primer(5'-3') is TTAAGCCCGGGTAAGGTTCCCT with product size of 156bp. *Faecalibacterium prausnitzii* forward-primer (5'-3') is CACGGCTCTGGAAATCTATGT and reverse-primer(5'-3') is GCACAATGAGCATACCGAGTT with

product size of 140bp. *Dialister invisus* forward-primer (5'-3') is AGACGGAAACGACTGCTAATACC and reverse-primer(5'-3') is

CAGCTAACATCAGACGCAAACCC with product size of 116bp. The three strain gene fragments were retrieved from the NCBI GenBank database. (listed in Table 3). Figure 5 (A-C) showed amplification plot of the three strains, and the agarose syrup electrophoresis picture is shown in Figure 5 (D). OD and concentration of the three strains were compared between A,C,F groups showing there were no significant difference about OD ($p=.550$),while significant difference existed about concentration ($p=.033$).Mean and standard deviations were listed in Table 4.

Changes of abdominal aortic inflammatory factors

Mean, standard deviation and 95%CI(Confidence Interval) of TNF- α , IL1- β , IL-6, CRP were counted and compared between A to G. showing all existed significant difference($P=.016, .042, .037, .012$).The values were shown in Table 5.

Comparison of pathological inflammation scores

Colonic and rectal tissues were scored and compared for pathological inflammation according to the methods described above. For colon, inflammation scores of A, B,C,D,E,F,G were 0, 2.5 ± 0.24 , 3.5 ± 0.61 , 8 ± 0.26 , 5.5 ± 0.27 , 9.5 ± 0.22 and 10.5 ± 0.35 respectively. Results of comparison with A group showed significant difference(all $P<.001$).For rectum, inflammation scores of A, B,C,D,E,F,G were 0, 3 ± 0.27 , 5 ± 0.15 , 10 ± 0.13 , 6 ± 0.32 , 10 ± 0.48 , 10 ± 0.59 and significance difference existed compared with A group. (all $P<.001$), Table 6 listed the details. The line chart shows a significant increase in colonic and rectal pathological inflammation scores. However, there was no significant difference in the degree of inflammation between the two tissues ($P=.710$).(see Fig.5(E)). Figures 6 and 7 showed the colon and rectal pathological slice pictures of the rats (* 100) between A to G group.

Discussion

This article has made an important discussion on antibiotic-induced rat IBD model. It is important to choose the appropriate animal model of IBD to evaluate the non-clinical anti-inflammatory effects of the drug. Some scholars have conducted comparative research on rat IBD models caused by different chemical factors, and believe that each has its own advantages and disadvantages.^[6] A highly reversible and reliable IBD rat model has broad application prospects for new drug treatment of IBD. Inflammatory bowel disease (IBD) is a group of chronic inflammatory disorders that affect individuals throughout life. Although the etiology and pathogenesis of IBD are largely unknown, studies with animal models of colitis indicate that dysregulation of host/microbial interactions are requisite for the development of IBD.^[8]

The modeling method of this article is from the 2015 master's thesis of Wendi Zhang of Southern Medical University.^[9] In her paper, he compared and analyzed the imbalance of rat enteritis flora induced by antibiotics. Although we used her modeling method, the research focuses are different. We focused on

the changes of 9 kinds of flora and the changes of inflammatory factors in the blood. At the same time, we compare and analyze the colon and rectal pathological inflammation in the two stages of model recovery period.

In our paper, we analyzed the changes of 9 intestinal flora during the modeling period (days 1 and 4) and the recovery period (days 8, 11 and 14). *Staphylococcus aureus* is a major human pathogen that causes a wide range of clinical infections.^[10, 11] we noticed that as the dose of the antibiotic increased, the increase in bacterial load was not obvious, indicating that the drug inhibited the strain. *Bifidobacterium* are defined as a group of living microorganism supplements that confer health benefits on the host when administered in adequate amounts.^[12] The beneficial bacterium was significantly reduced in B, D,E,F group relative to the control group. *Yeast* cells are often employed in industrial fermentation processes for their ability to efficiently convert relatively high concentrations of sugars into ethanol and carbon dioxide.^[13] *Yeast* is not a common intestinal bacterium , and it is absent in the control group. It was only found in the low- and middle-doses of the triple-agent groups. *Bacteroides* is a gram-negative, non-spore, obligate anaerobic bacillus. Our study showed a significant reduction in the high-dose medication group, indicating that it was sensitive to high-dose and combination drugs and it was relative with type 2 diabetes.^[14] *Clostridium* species are anaerobic, Gram-positive, rod-shaped, endospore-forming bacteria belonging to the phylum *Firmicutes*, and they constitute both a class and a genus in the phylum.^[15] *Clostridium* decreased in the single-agent group and the combined drug group increased. *Anaerobic bacteria* have pivotal roles in the microbiota of humans and they are significant infectious agents involved in many pathological processes.^[16] Pathogenic variants of *E. coli* (pathovars or pathotypes) cause much morbidity and mortality worldwide.^[17] *Enterococcus* strains that adhere strongly to the intestinal epithelium, form biofilms and possess antioxidant defence mechanisms seem to have the greatest influence on the inflammatory process.^[18] The genus *Lactobacillus* consists of 173 species and many genomes are available to study taxonomy and evolutionary events.^[19] In another article, we elaborate on the changes of these nine floras. In general, with the increasing dose of antibiotics and the enhanced degree of combination, the beneficial bacteria decreased significantly, while the pathogenic bacteria increased significantly.

We performed real-time quantitative PCR analysis on three strains of *Bacteroides*, *Faecalibacterium prausnitzii*, and *Dialister invisus* (control group, medium-dose alone group, and medium-dose combined group). The results showed that the lower the number of copies, the higher the dose and the combination. *Bacteroides* species make up a significant fraction of the human gut microbiome, and can be probiotic and pathogenic, depending upon various genetic and environmental factors. These can cause disease conditions such as intra-abdominal sepsis, appendicitis, bacteremia, endocarditis, pericarditis, skin infections, brain abscesses and meningitis.^[20-22] There is an increasing interest in *Faecalibacterium prausnitzii*, one of the most abundant bacterial species found in the gut. *F.prausnitzii* phylogroups can be found within this species and their distribution is different between healthy subjects and patients with gut disorders. It also remains unknown whether or not there are other phylogroups within this species, and

also if other *Faecalibacterium* species exist.^[23, 24] *Dialister invisus* is reported to be low or not expressed in IBD patients.^[25] Our research showed that *Dialister invisus* is little expressed in the model.

Our study found that TNF- α and IL-6 were significantly higher than the control group except for group D. However, there was no significant relationship with the dose administered and cytokine levels. TNF- α has been reported as a potent stimulator of IL-6 production.^[26-28] Inflammation induces IL-1 β production in Kupffer cells and hepatocytes.^[29] In our study, IL-1 β was significantly increased compared with the control group, showing elevated value in dose administered group. CRP is currently a hot spot for studying inflammation and related diseases.^[30-32] Our results showed that elevated CRP was associated with those groups that received antibiotic compared with the control group.

Pathological examination is an important method to judge the degree of tissue inflammation. Colon and rectal tissues were scored based on the degree of neutrophil invasion. Our study found that the degree of inflammation of the intestinal mucosa of rats dissected on day 11 was more severe than that of rats dissected on day 15, which may be related to the self-recovery of the intestinal flora. Deregulation of host-microbiota interactions in the gut is a pivotal characteristic of Crohn's disease. It remains unclear, however, whether commensals and/or the dysbiotic microbiota associated with pathology in humans are causally involved in Crohn's pathogenesis.^[33] We investigated the degree of inflammation of colon and rectal tissues in antibiotic-induced SD rat enteritis models with drug doses and combined doses significantly increasing.. And we found that the greater the degree of flora disturbance, the more severe the inflammation of the colon and rectal tissues. This may be related to the release of inflammatory factors by the intestinal disorder flora.^[34] It may also be related to the damage of intestinal mucosa caused by the release of metabolites by pathogenic bacteria to induce humoral and cellular immunity.^[34, 35].

However, there are some shortcomings in this paper, such as not enough experiment rats, not deep discussion about inflammation factors with intestinal mucosa necrosis.

Conclusion

Specific antibiotic-induced IBD model in SD rats is feasible.

Declarations

Ethics approval and consent to participate

This study follows the Basel Declaration 2010. Most of the authors of this article have been trained in animal experiments and have obtained a certificate of competency. We use animals to a minimum in terms of animal welfare principles and without affecting the accuracy of the experiment. Xi'an United Nations Quality Detection Technology laboratory was commissioned to perform our experiments under

his IACUC permission. All applicable international ,national and/or institutional guidelines for care and use of animals were followed.

Consent for publication

Not applicable.

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author.

Competing interests

The contents of this manuscript have not been copyrighted or published previously. There are no directly related manuscripts or abstracts, published or unpublished, by any authors of this manuscript. The authors indicated no conflicts of interests.

Funding

This study was supported by Project 2018C37090 to Guojun Tong. The fund is mainly used for experiment, data collection, experimental indice detection, statistical analysis, expert communication and so on.

Authors' contribution

All authors discussed the methodology and considered it available. GJT ,HQ analyzed all experimental data. XFL, DLL, JL, JC reviewed the statistical results and participated in all Figures drawing and stitching with GJT. GJT wrote the paper. The research teams of Xi'an United Nations Quality Detection Technology Laboratory performed the experiment. All authors reviewed the manuscript.

Acknowledgments

The authors thank Professor Licheng Dai, Researcher Liqing Li , Yingrong Chen, Pingping Niu, Xuting Xu for critical revising the paper.

Authors' Information

Guojun Tong* MD^{1,2}, Hai Qian MD¹, Dongli Li, BS², Jing Li, MD², Jing Chen, MD², Xiongfeng Li, PhD³,

1 General Surgery, Central Laboratory, Huzhou Central Hospital, Hongqi Road198#, Huzhou, China,
313000

2 Central Laboratory, Huzhou Central Hospital, Hongqi Road 198#, Huzhou, China, 313000

3 Orthopedic Surgery, Huzhou Central Hospital, Hongqi Road 198#, Huzhou, China, 313000

References

- [1] Jacobs J. P., Goudarzi M., Singh N., Tong M., McHardy I. H., Ruegger P., Asadourian M., Moon B. H., Ayson A., Borneman J., McGovern D. P., Fornace A. J., Jr., Braun J. and Dubinsky M. A Disease-Associated Microbial and Metabolomics State in Relatives of Pediatric Inflammatory Bowel Disease Patients[J]. *Cell Mol Gastroenterol Hepatol*, 2016, 2(6): 750-766.
- [2] Srikanth C. V. and Cherayil B. J. Intestinal innate immunity and the pathogenesis of *Salmonella enteritis*[J]. *Immunologic research*, 2007, 37(1): 61-78.
- [3] Peloquin J. M. and Nguyen D. D. The microbiota and inflammatory bowel disease: insights from animal models[J]. *Anaerobe*, 2013, 24: 102-106.
- [4] Uzal F. A., McClane B. A., Cheung J. K., Theoret J., Garcia J. P., Moore R. J. and Rood J. I. Animal models to study the pathogenesis of human and animal *Clostridium perfringens* infections[J]. *Veterinary microbiology*, 2015, 179(1-2): 23-33.
- [5] Jiminez J. A., Uwiera T. C., Douglas Inglis G. and Uwiera R. R. E. Animal models to study acute and chronic intestinal inflammation in mammals[J]. *Gut pathogens*, 2015, 7: 29-29.
- [6] Jianmei1 Z., Mincong1 H., Yangling2 L., Bo2 Z., and Nengming1 L. Comparison on the Model Rats with Colitis Induced by 2,4,6-three Nitrobenzene Sulfonic Acid and Dex-tran Sulfate Sodium[J]. *China pharmacy*, 2017, 28(10): 1353-1356.
- [7] HU Tingting C. H. Establishment of acute radiation proctitis model in rats[J]. *Journal of Clinical Oncology*, 2019, 24(7): 584-587.
- [8] Kanneganti M., Mino-Kenudson M. and Mizoguchi E. Animal models of colitis-associated carcinogenesis[J]. *J Biomed Biotechnol*, 2011, 2011: 342637.
- [9] Zhang W. Construction of Antibiotic-Related Diarrhea Model[J]. Master Thesis, 2015, :
- [10] Tong S. Y. C., Davis J. S., Eichenberger E., Holland T. L. and Fowler V. G., Jr. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management[J]. *Clinical microbiology reviews*, 2015, 28(3): 603-661.
- [11] Balasubramanian D., Harper L., Shopsin B. and Torres V. J. *Staphylococcus aureus* pathogenesis in diverse host environments[J]. *Pathogens and disease*, 2017, 75(1): ftx005.
- [12] Kim B. J., Park T., Moon H. C., Park S. Y., Hong D., Ko E. H., Kim J. Y., Hong J. W., Han S. W., Kim Y. G. and Choi I. S. Cytoprotective alginate/polydopamine core/shell microcapsules in microbial encapsulation[J]. *Angew Chem Int Ed Engl*, 2014, 53(52): 14443-14446.

- [13]Dzialo M. C., Park R., Steensels J., Lievens B.andVerstrepen K. J. Physiology, ecology and industrial applications of aroma formation in yeast[J]. FEMS Microbiol Rev, 2017, 41(Supp_1): S95-s128.
- [14]Horie M., Miura T., Hirakata S., Hosoyama A., Sugino S., Umeno A., Murotomi K., Yoshida Y.andKoike T. Comparative analysis of the intestinal flora in type 2 diabetes and nondiabetic mice[J]. Exp Anim, 2017, 66(4): 405-416.
- [15]Paredes C. J., Alsaker K. V.andPapoutsakis E. T. A comparative genomic view of clostridial sporulation and physiology[J]. Nat Rev Microbiol, 2005, 3(12): 969-978.
- [16]Gajdacs M., Spengler G.andUrban E. Identification and Antimicrobial Susceptibility Testing of Anaerobic Bacteria: Rubik's Cube of Clinical Microbiology?[J]. Antibiotics (Basel), 2017, 6(4):
- [17]Croxen M. A., Law R. J., Scholz R., Keeney K. M., Wlodarska M.andFinlay B. B. Recent advances in understanding enteric pathogenic Escherichia coli[J]. Clin Microbiol Rev, 2013, 26(4): 822-880.
- [18]Golinska E., Tomusiak A., Gosiewski T., Wiecek G., Machul A., Mikolajczyk D., Bulanda M., Heczko P. B.andStrus M. Virulence factors of Enterococcus strains isolated from patients with inflammatory bowel disease[J]. World J Gastroenterol, 2013, 19(23): 3562-3572.
- [19]Inglis R. C., Meile L.andStevens M. J. A. Clustering of Pan- and Core-genome of Lactobacillus provides Novel Evolutionary Insights for Differentiation[J]. BMC Genomics, 2018, 19(1): 284.
- [20]Zafar H.andSaier M. H., Jr. Comparative genomics of transport proteins in seven Bacteroides species[J]. PloS one, 2018, 13(12): e0208151-e0208151.
- [21]Vineis J. H., Ringus D. L., Morrison H. G., Delmont T. O., Dalal S., Raffals L. H., Antonopoulos D. A., Rubin D. T., Eren A. M., Chang E. B.andSogin M. L. Patient-Specific Bacteroides Genome Variants in Pouchitis[J]. mBio, 2016, 7(6): e01713-01716.
- [22]Feng S.andMcLellan S. L. Highly Specific Sewage-Derived Bacteroides Quantitative PCR Assays Target Sewage-Polluted Waters[J]. Applied and environmental microbiology, 2019, 85(6): e02696-02618.
- [23]Lopez-Siles M., Duncan S. H., Garcia-Gil L. J.andMartinez-Medina M. *Faecalibacterium prausnitzii*: from microbiology to diagnostics and prognostics[J]. The ISME journal, 2017, 11(4): 841-852.
- [24]Munukka E., Rintala A., Toivonen R., Nylund M., Yang B., Takanen A., Hänninen A., Vuopio J., Huovinen P., Jalkanen S.andPekkala S. *Faecalibacterium prausnitzii* treatment improves hepatic health and reduces adipose tissue inflammation in high-fat fed mice[J]. The ISME journal, 2017, 11(7): 1667-1679.
- [25]Schirmer M., Franzosa E. A., Lloyd-Price J., McIver L. J., Schwager R., Poon T. W., Ananthakrishnan A. N., Andrews E., Barron G., Lake K., Prasad M., Sauk J., Stevens B., Wilson R. G., Braun J., Denson L. A., Kugathasan S., McGovern D. P. B., Vlamakis H., Xavier R. J.andHuttenhower C. Dynamics of

metatranscription in the inflammatory bowel disease gut microbiome[J]. *Nature microbiology*, 2018, 3(3): 337-346.

[26]Leonard M., Ryan M. P., Watson A. J., Schramek H.andHealy E. Role of MAP kinase pathways in mediating IL-6 production in human primary mesangial and proximal tubular cells[J]. *Kidney Int*, 1999, 56(4): 1366-1377.

[27]Amrani Y., Ammit A. J.andPanettieri R. A., Jr. Tumor necrosis factor receptor (TNFR) 1, but not TNFR2, mediates tumor necrosis factor-alpha-induced interleukin-6 and RANTES in human airway smooth muscle cells: role of p38 and p42/44 mitogen-activated protein kinases[J]. *Mol Pharmacol*, 2001, 60(4): 646-655.

[28]Coelho-Santos V., Gonçalves J., Fontes-Ribeiro C.andSilva A. P. Prevention of methamphetamine-induced microglial cell death by TNF- α and IL-6 through activation of the JAK-STAT pathway[J]. *Journal of neuroinflammation*, 2012, 9: 103-103.

[29]Kanamori Y., Murakami M., Sugiyama M., Hashimoto O., Matsui T.andFunaba M. Interleukin-1 β (IL-1 β) transcriptionally activates hepcidin by inducing CCAAT enhancer-binding protein δ (C/EBP δ) expression in hepatocytes[J]. *The Journal of biological chemistry*, 2017, 292(24): 10275-10287.

[30]Sudhakar M., Silambanan S., Chandran A. S., Prabhakaran A. A.andRamakrishnan R. C-Reactive Protein (CRP) and Leptin Receptor in Obesity: Binding of Monomeric CRP to Leptin Receptor[J]. *Frontiers in immunology*, 2018, 9: 1167-1167.

[31]Lee P. T., Bird S., Zou J.andMartin S. A. M. Phylogeny and expression analysis of C-reactive protein (CRP) and serum amyloid-P (SAP) like genes reveal two distinct groups in fish[J]. *Fish & shellfish immunology*, 2017, 65: 42-51.

[32]Søndberg E., Sinha A. K., Gerdes K.andSemsey S. CRP Interacts Specifically With Sxy to Activate Transcription in *Escherichia coli*[J]. *Frontiers in microbiology*, 2019, 10: 2053-2053.

[33]Roulis M., Bongers G., Armaka M., Salviano T., He Z., Singh A., Seidler U., Becker C., Demengeot J., Furtado G. C., Lira S. A.andKollias G. Host and microbiota interactions are critical for development of murine Crohn's-like ileitis[J]. *Mucosal immunology*, 2016, 9(3): 787-797.

[34]Bastaki S. M. A., Al Ahmed M. M., Al Zaabi A., Amir N.andAdeghate E. Effect of turmeric on colon histology, body weight, ulcer, IL-23, MPO and glutathione in acetic-acid-induced inflammatory bowel disease in rats[J]. *BMC complementary and alternative medicine*, 2016, 16: 72-72.

[35]Almohazey D., Lo Y.-H., Vossler C. V., Simmons A. J., Hsieh J. J., Bucar E. B., Schumacher M. A., Hamilton K. E., Lau K. S., Shroyer N. F.andFrey M. R. The ErbB3 receptor tyrosine kinase negatively regulates Paneth cells by PI3K-dependent suppression of Atoh1[J]. *Cell death and differentiation*, 2017, 24(5): 855-865.

Tables

Table 1 Comparison of basic status for rats at total days (g,`X±S)

	Weight	Food intake	Water intake	Stool in 2h
A	204.02±13.5	19.02±3.63	38.86±4.73	0.96±0.48
B	203.31±17.2	19.58±2.83	49.76±11.31	2.20±1.38
C	208.09±22.15	18.93±2.75	44.40±9.24	2.23±1.09
D	188.13±24.05	15.47±5.66	41.96±18.95	2.47±1.53
E	209.00±18.58	18.80±2.11	60.33±9.02	2.82±2.01
F	210.70±19.18	20.31±3.57	52.68±10.23	3.09±1.79
G	212.12±20.31	20.66±3.13	53.47±7.84	2.33±1.55
F	2.313	2.807	6.213	3.043
P	0.04	0.016	<0.001	0.009

Table 2 Comparison of total nine species of bacteria between A and B-G groups (E6)

	A	B	C	D	E	F	G
X±S	1376.7±3683.8	687.1±1498.7	1474.9±4187.6	478.2±1758.1	664.5±1567.9	403.8±1172.0	3609.8±21206.5
95%CI for mean	lower	562.1	355.7	549.0	65.0	317.8	144.6
	upper	2191.3	1018.5	2400.8	891.3	1011.2	662.9
P		<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

Table 3 Sequences of primers used for real-time quantitative PCR

Target bacteria	Forward(5'-3')	Reverse(5'-3')	Product size(bp)	Accession number
<i>Bacteroides</i>	TTAAGTATTCCACCTGGGGAGT	TTAACCGCCGGTAAGGTTCCCT	156	CR626927
<i>Faecalibacterium prausnitzii</i>	CACGGCTCTGAAATCTATGT	GCACAATGAGCATACCGAGTT	140	NZ_PXUP01000071
<i>Dialister invisus</i>	AGACGGAAACGACTGCTAATACC	CAGCTAACGACGCAAACCC	116	LT223661

Table 4 Comparison of *Bacteroides*, *Faecalibacterium prausnitzii*, *Dialister invisus* examination by Realtime PCR between A,C, F groups (ng/ul), ($\bar{X} \pm SD$)

	A group	C group	F group	F Value	P Value
Realtime PCR	21.72±10.60	34.60±19.85	26.14±19.86	10.711	0.033*
OD	1.88±0.18	1.79±0.25	1.79±0.16	0.612	0.550

Table 5 Comparison of inflammation factors from group A to group G (pg/ml)

	N	Mean	Std. Deviation	95% Confidence Interval for Mean			F	P
				Lower Bound	Upper Bound			
TNF- α	A	2	113.61300	44.932393	-190.08854	617.31454	2.343	0.016*
	B	2	185.77900	6.512453	127.26693	244.29107		
	C	2	194.93750	4.417296	155.24967	234.62533		
	D	1	186.35500	.	.	.		
	E	2	198.69700	1.019648	189.53583	207.85817		
	F	2	172.78350	1.989091	154.91222	190.65478		
	G	2	150.13950	4.182537	112.56090	187.71810		
IL1- β	A	2	89.47750	31.312810	-191.85693	370.81193	1.186	0.042*
	B	2	105.95750	14.735398	-26.43480	238.34980		
	C	2	99.08750	.832265	91.60990	106.56510		
	D	1	66.01800	.	.	.		
	E	2	119.33000	33.504134	-181.69270	420.35270		
	F	2	116.66600	.185262	115.00149	118.33051		
	G	2	95.97050	1.694935	80.74211	111.19889		
IL-6	A	2	133.31300	40.869358	-233.88361	500.50961	1.337	0.037*
	B	2	142.08650	1.957979	124.49476	159.67824		
	C	2	146.28750	11.082485	46.71533	245.85967		
	D	1	108.33600	.	.	.		
	E	2	152.85550	1.164605	142.39194	163.31906		
	F	2	151.65200	.895197	143.60897	159.69503		
	G	2	122.04600	1.388758	109.56851	134.52349		
CRP	A	2	11.04950	2.851762	-14.57256	36.67156	2.807	0.012*
	B	2	12.06650	.003536	12.03473	12.09827		
	C	2	11.06000	1.548564	-2.85329	24.97329		
	D	1	7.75200	.	.	.		
	E	2	15.17450	2.004648	-2.83655	33.18555		
	F	2	11.94850	.215668	10.01080	13.88620		
	G	2	11.80500	.449720	7.76443	15.84557		
Total	13	11.84308	2.166844	10.53367	13.15249			

*p<0.05 indicate significant difference

Table 6 Mean inflammation score by pathology ($X \pm S$)

groups	Colon	P	Rectum	P
A	reference		reference	
B	2.5±0.24	0.007	3±0.27	0.004
C	3.5±0.61	0.001	5±0.15	<0.001
D	8±0.26	<0.001	10±0.13	<0.001
E	5.5±0.27	<0.001	6±0.32	<0.001
F	9.5±0.22	<0.001	10±0.48	<0.001
G	10.5±0.35	<0.001	10±0.59	<0.001

Figures

Figure 1

Study flow and method. A: Fourteen SD rats of similar body weight were randomly divided into A-G 7 groups, 2 in each group, of which group A was the control group, B-D group was the low-dose, middle-dose, and high-dose clindamycin administration group alone; E-G The group was a low-dose, middle-dose, high-dose clindamycin, ampicillin, and streptomycin combination administration group. B: SD rats were anesthetized with 2% pentobarbital sodium (0.2 ml / 100 g) by intraperitoneal injection on day 11 and day 15, and 5 ml of abdominal aortic blood was taken. Colon and rectal tissues were taken at the same time.C: The schematic diagram is the modeling process, 1-7 days is the model period for feeding specific antibiotic, and 8-15 is the recovery period after specific antibiotic withdrawal.




Figure 2

Basic indicators of the SD rat feeding process. A: High-low maps of animal weights comparison in each group on 1st, 3rd, 5th, 7th, 9th, 11th, 13th, and 14th days. There were statistical differences between the groups, $P = .04$; B: Histograms of the food intake of the animals in each group on the 1st, 3rd, 5th, 7th, 9th, 11th, 13th, and 14th days. There was statistical difference between the groups, $P < .016$; C: Heat maps of animals in each group on 1st, 3rd, 5th, 7th, 9th, 11th, 13th, and 14th days. There was a statistical difference between the groups, $P < .001$; D: Bar graphs of animals in each group for 2 hours on day 1, 3, 5, 7, 9, 9, 11, 13, 14. There was a statistical difference between the groups, $P = .009$.

Figure 3

A picture of counted bacteria specimen under a microscope on fourth day. a: Staphylococcus aureus; b: Bifidobacterium; c: yeast; d: Bacteroides e: Clostridium; f: anaerobic bacteria; g: E. Coli; h: enterococcus; i: Lactobacillus




Figure 4

Nine kinds of flora were compared in each group on days 1, 4, 8, 11, and 14, respectively. A: comparison on the first day; B: comparison on the 4th day; C: comparison on the 8th day; D: comparison on the 11th day; E: comparison on the 14th day, all P <.001.

Figure 5

Real-time quantitative PCR doubling curve graph and agarose syrup electrophoresis graph of three strains, and line graph of inflammation score of colon and rectal tissue. A: Bacteroides amplification plot; B: Faecalibacterium prausnitzii amplification plot; C: Dialister invisus amplification plot; D: Bacteroides, Faecalibacterium prausnitzii, Dialister invisus agarose syrup electrophoresis picture, each done twice; E: Colon and rectal tissue inflammation score line chart, showing differences in each group, among which colon tissue inflammation score A / B: $P = .007$; A / C: $P = .001$; A / D, A / E, A / F, A / G: all $P < .001$; and rectum tissue inflammation score: A/B: $P = .004$, all others :all $P < .001$. But the score between colon and rectum are not significant, $P = .710$.

Figure 6

Colonic HE staining pathological picture (* 100), the left is the dissection of the rat colon tissue on the 11th day, and the right is the dissection of the rat colon tissue on the 15th day. In group A, there was no obvious neutrophil infiltration and mucosal edema; in group B-G, neutrophil infiltration became more and more serious, mucosal edema became more and more obvious, and even mucosal necrosis and focal ulcer formed.

Figure 7

Rectal tissue HE staining pathological picture (* 100), the left side is the dissection of rat rectum tissue on the 11th day, and the right side is the dissection of rat rectum tissue on the 15th day. In group A, there was no obvious neutrophil infiltration and mucosal edema; in group B-G, neutrophil infiltration became more and more serious, mucosal edema became more and more obvious, and even mucosal necrosis and focal ulcer formed.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- NC3RsARRIVEGuidelines2013.pdf