Aortic root aneurysm with right atrial clot: a rare combination

Debmalya Saha (✉ debmalya.cmc@gmail.com)

AMRI Hospital Dhakuria <https://orcid.org/0000-0002-5350-4005>

Kaushik Mukherjee

AMRI Hospital Dhakuria

Amrita Guha

AMRI Hospital Dhakuria

Research Article

Keywords: aneurysm, aorta, appendage, atrial, Bentall, clot

Posted Date: February 2nd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1282263/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Though the incidence of aneurysm involving the aortic root and/or ascending aorta is common, the combination of aortic root aneurysm and right atrial clot is extremely rare. No such case is reported in literature till date. This case report presents a 52-year gentleman who came to our emergency department with complains of breathlessness, abdominal distention, pedal swelling, and decreased urine output with extremely poor general condition. After hemodynamic stabilization and preoperative optimization and work up he was managed with Bentall procedure with right atrial clot removal. Immediate post-operative course was normal except deranged liver function tests. The patient was discharged on post-operative day ten.

Introduction

Right atrial clot in the background of aortic root aneurysm is an uncommon occurrence. Here we will present a case of 52-year male with aortic root aneurysm and right atrial clot after informed-written consent and institutional ethical approval obtained.

Case Report

A 52-year male with extremely poor general condition showed symptoms of breathlessness, abdominal distension, pedal swelling, and decreased urine output. On general examination, his heart rate was 140/minute, blood pressure of 95/30mm of Hg, respiratory rate of 35/min, saturation of 90% at room air with presence of anasarca. During examination of cardiovascular system, a diastolic murmur was audible at aortic area. On examining the respiratory system, we got crepitations on both lung fields. A diagnosis of acute decompensated heart failure was made on clinical ground.

On routine blood investigations, total leucocyte count was high with polymorphic leukocytosis, and there was deranged LFTs and deranged kidney function tests.

On Chest X ray, we found cardiomegaly with prominent aortic knob along with increased para-hilar vascularity, bilateral blunted costophrenic angles. ECG showed low voltage waves with features of left ventricular diastolic overload. Bedside 2D trans-thoracic ECHO(TTE) revealed poor LVEF of 25%, global hypokinesia, severe aortic regurgitation and dilatation of aortic root and ascending aorta (ascending aorta 6.2cm). CT angiography of heart and great vessels could not be done because of deranged renal function and respiratory distress of the patient.

The patient was managed conservatively initially with inotropes, judicious use of diuretics, IV antibiotics (renally safe), high flow oxygen and intermittent noninvasive positive pressure ventilation (NIPPV/BiPAP) and strict intake-output charting.

We parallelly planned for Bentall procedure on high-risk basis with ongoing resuscitation and stabilization. Our approach was via median sternotomy. The aortic root and ascending aorta were

aneurysmal (Figure 1). Right atrial appendage was firm on palpation. Standard aorto-bicaval cannulation was done. Cardiopulmonary bypass was established. Aortic valve with ascending aorta was replaced with aortic valved conduit with re-implantation of coronary buttons. A large, organized clot occupying right atrial appendage (Figure 2) was removed.

Immediate post-operative period was uneventful other than deranged liver function tests which were in improving trend. The patient was discharged on post-operative day ten.

Conclusion

This type of rare association between ascending aortic aneurysm and right atrial clot must be reported in literature. This may alter the intra-operative strategy and subsequently can improve the operative outcome for this patient.

Declarations

FUNDING: No funding source found.

CONFLICT OF INTERESTS: The authors declare that there is no conflict of interest.

ETHICS APPROVAL: This study was approved by the Institutional Ethical Committee and

CONSENT TO PARTICIPATE: Informed consent to participate was taken from the patient's relatives in their own language.

CONSENT FOR PUBLICATION: Informed consent for publication was taken from the patient's relatives in their own language.

AVAILABILITY OF DATA & MATERIAL: Available

CODE AVAILABILITY: Not applicable.

AUTHORS' CONTRIBUTIONS: Equal.

References

1. Debmalaya Saha, Kaushik Mukherjee, Amrita Guha. Aortic root aneurysm with right atrial clot: a rare combination. *Authorea*. December 27, 2021.
DOI: [10.22541/au.164063678.80860991/v1](https://doi.org/10.22541/au.164063678.80860991/v1)

Figures

Figure 1

Aneurysmal aortic root and ascending aorta

Figure 2

Right atrial appendage clot

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- CAREchecklist.docx