Impacts of skull stripping on construction of three-dimensional T1-weighted imaging-based brain structural network in full-term neonates

Geliang Wang

The First Affiliated Hospital of Xi'an Jiaotong University

Yajie Hu

The First Affiliated Hospital of Xi'an Jiaotong University

Xianjun Li

The First Affiliated Hospital of Xi'an Jiaotong University

Miaomiao Wang

The First Affiliated Hospital of Xi'an Jiaotong University

Congcong Liu

The First Affiliated Hospital of Xi'an Jiaotong University

Jian Yang

The First Affiliated Hospital of Xi'an Jiaotong University

Chao Jin (✉ chaojin@mail.xjtu.edu.cn)

The First Affiliated Hospital of Xi'an Jiaotong University

Research

Keywords: Skull striping, neonatal brain, brain structural networks, 3D T1 Weighted Imaging

Posted Date: March 26th, 2020

DOI: <https://doi.org/10.21203/rs.2.22268/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BioMedical Engineering OnLine on June 3rd, 2020. See the published version at <https://doi.org/10.1186/s12938-020-00785-0>.

Abstract

Background: Skull stripping remains a challenge for neonatal brain MR image analysis. However, little is known about how accuracy of the skull stripping affects the neonatal brain tissue segmentation and subsequent network construction. This paper therefore aimed to clarify this issue by comparing two automatic (FMRIB Software Library's Brain Extraction Tool, BET; Infant Brain Extraction and Analysis Toolbox, iBEAT) and a semiautomatic (iBEAT with manual correction) processes in constructing 3D T1-Weighted Imaging (T1WI)-based brain structural network.

Methods: Twenty-two full-term neonates (gestational age, 37-42 weeks; boys/girls, 13/9) without abnormalities on MRI who underwent brain 3D T1WI were retrospectively recruited. Two automatic (BET and iBEAT) and a semiautomatic preprocessing (iBEAT with manual correction) workflows were separately used to perform the skull stripping. Brain tissue segmentation and volume calculation were performed by a John Hopkins atlas-based method. Sixty-four gray matter regions were selected as nodes; volume covariance network and its properties (clustering coefficient, C_p ; characteristic path length, L_p ; local efficiency, E_{local} ; global efficiency, E_{global}) were calculated by GREYNA. Analysis of variance (ANOVA) was used to compare the differences in the calculated volume between three workflows.

Results: There were significant differences in volumes of 50 brain regions between three workflows ($P < 0.05$). Three neonatal brain structural networks presented small-world topology. The semiautomatic workflow showed remarkably decreased C_p , increased L_p , decreased E_{local} , and decreased E_{global} , in contrast to two automatic ones.

Conclusions: Imperfect skull stripping indeed affected the accuracy of brain structural network in full-term neonates.

Background

It is known that Magnetic Resonance Imaging (MRI) has become a very important tool for investigating the early brain development and injury in neonates [1,2]. In particular, MRI-based brain structural network analysis provides critical ways to understand the topological structure of brain information integration and transmission during the early development [3-5]. As a basic preprocessing, skull stripping, designed to eliminate skull, scalp, dura, and other non-brain tissues and retain brain parenchyma from brain MRI, is an essential process in brain tissue segmentation and subsequent brain network construction [6]. For neonates, numerous efforts have been made to perform the automated processing [7-13], such as FMRIB Software Library's (FSL) Brain Extraction Tool (BET) [14] and infant brain extraction and analysis toolbox (iBEAT developed by the IDEA group at the University of North Carolina at Chapel Hill) [15]. However, skull stripping remains a challenge for neonatal brain MRI analysis due to its low tissue contrast, large within-tissue intensity variations, and regionally heterogeneous [16]. Besides, inaccurate skull stripping, e.g. unremoved non-brain tissues would result in the overestimation of local brain volume and cortical thickness [17]. And it would further affect the construction of brain structural network. To our knowledge,

little is known about the impact of preprocessing accuracy on the accuracy of brain tissue segmentation and structural network construction.

The present study aimed to investigate the effects of skull stripping on brain tissue segmentation and structural network construction based on three-dimensional (3D) T1-Weighted Imaging (T1WI). Firstly, the study constructed the processing flow based on 3D T1-Weighted MRI on 22 term neonates' gray matter; secondly, repeatability and consistency of brain region volume's segmentation were further performed to verify the accuracy; in final part, the study correlated the 64 brain regions volumes of the other two workflows, which did a comparison on three workflows' with the standard volume. Such a workflow could be applied to characterize structural connectivity and may provide valuable anticipatory information about the potential for encountering abnormalities at a later stage in development.

Results

Repeatability and Reliability

The Bland-Altman graph of two repeated measurements indicated that 95% of the differences between two measurements located in the $\text{mean} \pm 1.96\text{SD}$ (standard deviation) range that suggest the good repeatability (Fig. 1A). Meanwhile, the ICC analysis indicated the average correlation coefficient was 0.894. Most of values were greater than 0.8, while those of a few brain regions, e.g. caudate nucleus, precuneus, superior occipital gyrus, and inferior occipital gyrus were relatively lower, but were greater than 0.7 (Fig. 1B).

Comparisons of brain structural network between three workflows

Brain region volume

The differences of volume of 64 brain regions performed through ANOVA analysis among three workflows which are shown below (Fig. 2). There were significant differences of volume between three workflows in 50 brain regions, while no significant difference was found in the remaining 14 brain regions, such as bilateral superior frontal gyrus, bilateral pre- and post-central gyrus, bilateral superior parietal gyrus, bilateral precuneus, bilateral superior occipital gyrus and so on.

Small-world properties

In the defined threshold range, the neonatal brain network exhibited high-efficiency small-world topology (Fig. 3).

Properties of brain structural network

The corrected workflow, compared with the others, showed a significantly decreased C_p and increased L_p . With regard to network efficiency, the corrected workflow showed a significantly decreased E_{local} and E_{global} (Fig. 4).

Discussion

To investigate the impacts of skull stripping on the accuracy of brain tissue segmentation and structural network construction, three workflows (BET, iBEAT and iBEAT with manual correction) were compared to perform the 3D T1WI-based brain structural network. Our results indicated that the iBEAT with manual correction showed a more accurate consistency and repeatability in brain segmentation. Besides, significant differences in calculations of brain volume and structural network properties between the three workflows further implied the importance of accurate skull stripping in brain segmentation and subsequent brain network construction.

The ICC results indicated that the majority of brain regions showed good repeatability and reliability, while the ICC of some regions, such as caudate nucleus, gyrus rectus, left superior parietal gyrus, left precuneus, left superior occipital gyrus and right inferior occipital gyrus, were relatively lower. One reason may be rooted in the inherently low spatial resolution, insufficient tissue contrast, and ambiguous tissue intensity distributions in neonatal MRI [16]. On the other, the adult head coil used in this study may affect the MR image quality in neonates [18,19]. In this regard, MRI acquisition settings including the specific head coil and scanning parameters should be adjusted for neonates. In details, a neonatal head coil with appropriate size and high signal-to-noise ratio, as well as the use of appropriate scanning parameters (e.g. smaller FOV, longer repetition time and echo time than adult scanning) would facilitate the signal acquisition [18,20].

By comparing the three workflows, we found that slight adjustment for skull stripping would remarkably affect the calculations of regional brain volume. Of 64 regions, significant differences existed in brain volume calculations of 50 regions between the three workflows. This further confirmed the difficulty of brain segmentation in neonatal MRI. Specifically, most brain volumes calculated by BET workflow were smaller than those by iBEAT and iBEAT with manual correction. This may be linked with the difference of skull stripping in neonatal MRI between BET and iBEAT [16]. It is worth noting that more unremoved skull components were found in BET than iBEAT; while unremoved components by iBEAT mainly located in the base of skull (Fig. 5). It may be such facts that led to the larger brain volumes by iBEAT than iBEAT with manual correction. From the above, the accuracy of skull stripping would have considerable effects on the neonatal brain segmentation.

By studying the brain structural network, we found the highly small-world topology in full-term neonates. This may suggest that a highly efficient brain network, serving for brain information integration and transfer, has been constructed in the early development. This is also in agreement with prior findings in neonates [5]. Besides, significant differences were found in local (C_p and E_{loc}) and global (E_{glob}) topological properties of brain structural network between the three workflows. These may indicate that these parameters were sensitive to the accuracy of preprocessing (skull stripping).

The study had several limitations. First, the sample size was relatively small in this study, and more data should be acquired to further improve the accuracy. Second, the template and atlas we used are made

by foreign neonates. Given the differences of brain morphology between Western and Eastern neonates [21-24], it may lead to certain errors in the initial estimation of tissue intensity distributions and thus resulted in large deformations in the registration procedure. In this regard, a dedicated atlas and template appropriate for Chinese neonates is needed to be developed in further study. Besides, based on the above, further studies regarding the differences of brain structural network constructed by iBEAT and iBEAT with manual correction between Western and Eastern neonates are also required to facilitate the understandings of preprocessing's impacts on the brain network construction.

Conclusions

In this study, we quantitatively analyzed the influences of skull stripping on brain volume calculation and topological properties of brain structural network in neonates. Our results indicated that the brain networks had robust small-world configuration, in addition, there were significant differences in both local and global topological parameters between the three workflows. These findings enhanced the importance of accurate skull stripping in calculation of brain tissue volume and brain structural network construction.

Methods

This study was approved by the local institutional review board of the first author's affiliation. The parents of the neonates were informed regarding the goals and risks of the MRI scan, and requested for the written consent.

Subjects

This study recruited 22 full-term neonates (13 males and 9 females; gestational age range: 37-42 weeks) without any MRI abnormalities or evidences of any clinical episodes that might cause cerebral damages.

MRI data acquisition

All data were acquired on a 3.0 Tesla scanner (Signa HDxt, General Electric Medical System, Milwaukee, WI, USA) with an 8-channel phase array radio-frequency head coil. To reduce the head movement and complete the MRI procedure, the subjects were sedated with a relatively low dose of oral chloral hydrate (25-50 mg/kg). The potential risks of the chloral hydrate were fully considered. The selection, monitoring, and management of subjects were strictly performed following the guidelines [25]. Neonates were laid in a supine position and snugly swaddled in blankets. A pediatrician was present during the MRI scan. Micro earplugs were inserted into the external auditory canal for hearing protection. Heads of the subjects were immobilized by molded foam, which was placed around the head. The temperature, heart rate, and oxygen saturation were monitored throughout the procedure.

Three-dimensional fast spoiled gradient-recalled echo (3D-FSPGR) T1-Weighted magnetic resonance images were acquired with the parameters: repetition time/echo time = 10.28 ms/4.62 ms, inversion time = 400 ms, field of view = 240 mm, voxel size = 0.94×0.94×1 mm³.

Construction of brain structural network

Figure 6 detailed the workflow for constructing the neonatal brain structural network. The three steps for the workflow were detailed as follows.

Data preprocessing. To investigate the impacts of data preprocessing on network construction, three preprocessing workflows were performed by BET, iBEAT and iBEAT with manual correction.

BET workflow: non-brain tissues in each T1WI image were automatically removed by BET. iBEAT workflow: three automatic preprocessing steps, including brain contrast enhancement and skull stripping were sequentially performed. iBEAT with manual correction: (1) manually align the AC-PC line; (2) strip the skull by using iBEAT; (3) identify the remaining non-brain tissues by the boundaries of brain gray matter and manually remove these tissues. The manual correction was performed by two pediatric radiologists with 5 years of experience and differences in identifying the remaining non-brain tissues were resolved by consensus.

Calculations of brain region volume. To calculate the volume of each brain region, the preprocessed data were firstly registered to a standard 3D-T1WI-based template (Johns Hopkins University) by linear (rigid transformation) and nonlinear (affine transformation) registrations. And then the volume of each brain region can be estimated by Eq.(1):

[Please see the supplementary files section to view this equation.]

Construction of brain structural network. By using the graph theoretic approaches, the cortical and subcortical regions were used as nodes to construct the brain networks, with connections between nodes defined as correlations between regional brain volumes. Here, 64 brain regions that mainly involved the gray matters and several important subcortical regions (thalamus, hippocampus and cerebellum) were selected as network nodes (Table 1). GRETNA (www.nitrc.org/projects/gretna/) was used to construct the network. Partial correlations between all nodes' volumes were firstly estimated as edges of the network, and then network was constructed by binary connective matrices. The network properties, such as clustering coefficient (C_p), characteristic path length (L_p), global (E_{global}) and local efficiency (E_{local}) were calculated. The C_p of a network is defined by the average of the clustering coefficients across nodes, where the C_p of a node is the ratio of the number of actually connection nearest neighbors of this node to the maximum number of possible connection [26]. C_p quantifies the local interconnectivity of a graph. The L_p of a graph is the average of the shortest path length between all pairs of nodes in the network, and it is an indicator of overall routing efficiency of a graph [27]. The E_{local} of a network is the average of the local efficiency over all nodes. It measures the mean local efficiency of the network. The E_{global} of a network is defined by the mean shortest path length [28]. It measures the extent of information propagation through the whole network. Typically, a small-world network should fulfill the following conditions: $C_p/C_{rand} > 1$ and $L_p/L_{rand} \approx 1$.

Statistical analysis

Regarding the manual corrections for anterior commissure-posterior commissure (AC-PC) alignment and skull stripping, the repeatability and consistency of brain volume calculations between two repeated measurements were evaluated by the Bland-Altman graph and intraclass correlation coefficient (ICC). Analysis of variance (ANOVA) was used to compare the differences in the volumes of 64 brain regions between the three workflows.

All the segmentation, calculation of brain region volume and network parameters, and statistical analysis were performed by using the MATLAB R2016b (Mathworks Inc, Natick, MA, USA).

References

- [1] Dubois J, Dehaene-Lambertz G, Kulikova S, Poupon C, Hüppi PS, Hertz-Pannier L. The early development of brain white matter: a review of imaging studies in fetuses, newborns and infants. *Neuroscience*. 2014;276: 48-71.
- [2] Holland D, Chang L, Ernst TM, Curran M, Buchthal SD, Alicata D, Skranes J, Johansen H, Hernandez A, Yamakawa R, Kuperman JM, Dale AM. Structural growth trajectories and rates of change in the first 3 months of infant brain development. *JAMA Neurol*. 2014;71(10):1266-1274.[U1]
- [3] Cao M, Huang H, He Y. Developmental connectomics from infancy through early childhood. *Trends Neurosci*. 2017;40(8):494-506.
- [4] Gao W, Lin W, Grewen K, Gilmore JH. Functional connectivity of the infant human brain: plastic and modifiable. *Neuroscientist*. 2017;23(2):169-184.
- [5] Brown CJ, Miller SP, Booth BG, Andrews S, Chau V, Poskitt KJ, Hamarneh G. Structural network analysis of brain development in young preterm neonates. *Neuroimage*. 2014;101:667-680.[U2]
- [6] Huang H, Shu N, Mishra V, Jeon T, Chalak L, Wang ZJ, Rollins N, Gong G, Cheng H, Peng Y, Dong Q, He Y. Development of human brain structural networks through infancy and childhood. *Cereb Cortex*. 2015;25(5):1389-1404.[U3]
- [7] Weisenfeld NI, Warfield SK. Automatic segmentation of newborn brain MRI. *Neuroimage*. 2009;47(2):564-572.
- [8] Shi F, Fan Y, Tang S, Gilmore JH, Lin W, Shen D. Neonatal brain image segmentation in longitudinal MRI studies. *Neuroimage*. 2010;49(1):391-400.[T4]
- [9] Gui L, Lisowski R, Faundez T, Hüppi PS, Lazeyras F, Kocher M. Morphology-driven automatic segmentation of MR images of the neonatal brain. *Med Image Anal*. 2012;16(8):1565-1579.[U5]
- [10] Mahapatra D. Skull stripping of neonatal brain MRI: using prior shape information with graph cuts. *J Digit Imaging*. 2012;25(6):802-814.[U6]

- [11] Anbeek P, Išgum I, van Kooij BJ, Mol CP, Kersbergen KJ, Groenendaal F, Viergever MA, de Vries LS, Benders MJ. Automatic segmentation of eight tissue classes in neonatal brain MRI. Plos One. 2013;8(12):e81895.
- [12] Gousias IS, Alexander H, Counsell SJ, Latha S, Rutherford MA, Heckemann RA, Hajnal JV, Daniel R, David EA. Magnetic resonance imaging of the newborn brain: automatic segmentation of brain images into 50 anatomical regions. Plos One. 2013;8(4):e59990.
- [13] Cardoso MJ, Melbourne A, Kendall GS, Modat M, Robertson NJ, Marlow N, Ourselin S. AdaPT: An adaptive preterm segmentation algorithm for neonatal brain MRI. Neuroimage. 2013;65(1):97-108.[\[T7\]](#)
- [14] Smith SM. Fast robust automated brain extraction. Hum Brain Mapp. 2002;17(3):143-155.
- [15] Dai Y, Shi F, Wang L, Wu G, Shen D. iBEAT: a toolbox for infant brain magnetic resonance image processing. Neuroinformatics. 2013;11:211-225.
- [16] Li G, Wang L, Yap PT, Wang F, Wu Z, Meng Y, Dong P, Kim J, Shi F, Rekik I, Lin W, Shen D. Computational neuroanatomy of baby brains: A review. Neuroimage. 2019;185:906-925.
- [17] Shi F, Wang L, Dai Y, Gilmore JH, Lin W, Shen D. LABEL: pediatric brain extraction using learning-based meta-algorithm. Neuroimage. 2012;62 (3):[1975-1986](#).[\[U8\]](#)
- [18] Hillenbrand CM, Reykowski A. MR imaging of the newborn: a technical perspective. Magn Reson Imaging Clin N Am. 2012;20(1):63-79.
- [19] Hughes EJ, Winchman T, Padormo F, Teixeira R, Wurie J, Sharma M, Fox M, Hutter J, Cordero-Grande L, Price AN, Allsop J, Bueno-Conde J, Tusor N, Arichi T, Edwards AD, Rutherford MA, Counsell SJ, Hajnal JV. A dedicated neonatal brain imaging system. Magn Reson Med. 2017;78(2):794-804.
- [20] Lopez Rios N, Foias A, Lodygensky G, Dehaes M, Cohen-Adad J. Size-adaptable 13-channel receive array for brain MRI in human neonates at 3T. NMR in biomedicine. 2018;31(8):e3944.
- [21] Hillenbrand CM, Reykowski A. MR imaging of the newborn: a technical perspective. Magn Reson Imaging Clin N Am. 2012;20(1):63-79.
- [22] Pannek K, Guzzetta A, Colditz PB, Rose SE. Diffusion MRI of the Tang Y, Hojatkashani C, Dinov ID, Sun B, Fan L, Lin X, Qi H, Hua X, Liu S, Toga AW. The construction of a Chinese MRI brain atlas: a morphometric comparison study between Chinese and Caucasian cohorts. NeuroImage. 2010;51(1):33-41.
- [23] Tang Y, Hojatkashani C, Dinov ID, Sun B, Fan L, Lin X, Qi H, Hua X, Liu S, Toga AW. The construction of a Chinese MRI brain atlas: a morphometric comparison study between Chinese and Caucasian cohorts. NeuroImage. 2010;51(1):33-41.

- [24] Lee JS, Lee DS, Kim J, Kim YK, Kang E, Kang H, Kang KW, Lee JM, Kim JJ, Park HJ, Kwon JS, Kim SI, Yoo TW, Chang KH, Lee MC. Development of Korean standard brain templates. *Journal of Korean medical science*. 2005;20(3):483-488.
- [25] Bai J, Abdul-Rahman MF, Rifkin-Graboi A, Chong YS, Kwek K, Saw SM, Godfrey KM, Gluckman PD, Fortier MV, Meaney MJ, Qiu A. Population differences in brain morphology and microstructure among Chinese, Malay, and Indian neonates. *PloS one*. 2012;7(10):e47816.
- [26] Côté CJ, Wilson S. Guidelines for monitoring and management of pediatric patients before, during, and after sedation for diagnostic and therapeutic procedures. *Pediatr Dent*. 2019;41(4):26E-52E.
- [27] Watts DJ, Strogatz SH. Collective dynamics of 'small-world' networks. *Nature*. 1998;393(6684):440-442.[T9]
- [28] Rubinov M, Sporns O. Complex network measures of brain connectivity: Uses and interpretations. *Neuroimage*. 2010;52(3):1059-1069.
- [29] Latora V, Marchiori M. Efficient behavior of small-world networks. *Phys Rev Lett*. 2001;87(19):198701.

Declarations

Ethics approval and consent to participate

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

Consent for publication

All authors confirmed the consent for publication.

Availability of data and material

The codes and datasets of this study are available from the corresponding authors upon reasonable request.

Competing interests

The authors declare that they have no conflict of interest.

Funding

This study was funded by National Natural Science Foundation of China (No. 51706178, 81171317, 81971581), National Key Research and Development Program of China (2016YFC0100300), the 2011 New Century Excellent Talent Support Plan of the Ministry of Education, China (NCET-11-0438).

Authors' contributions

G.L.W. and Y.J.H. performed the data acquisition, analysis and manuscript writing. X.J.L. performed the data processing method, M.M.W. and C.C.L. performed the data acquisition, J.Y. and C.J. designed the study, analyzed the data and wrote the manuscript.

Acknowledgements

The authors appreciated Prof. Li Liu, Prof. Xihui Zhou, Dr. Hongxia Song and Gailian Li from the Neonatology Department for preparing and monitoring the neonates before and during imaging. We also thank all participants and their parents for their loyalty and cooperation.

Table

Table 1 The list of 64 brain regions

No.	Region	Hemisphere	No.	Region	Hemisphere
1	thalamus	left	33	angular gyrus	left
2	thalamus	right	34	angular gyrus	right
3	putamen	left	35	superior temporal gyrus	left
4	putamen	right	36	superior temporal gyrus	right
5	globus pallidus	left	37	middle temporal gyrus	left
6	globus pallidus	right	38	middle temporal gyrus	right
7	caudate nucleus	left	39	inferior temporal gyrus	left
8	caudate nucleus	right	40	inferior temporal gyrus	right
9	superior frontal gyrus	left	41	fusiform gyrus	left
10	superior frontal gyrus	right	42	fusiform gyrus	right
11	middle frontal gyrus	left	43	parahippocampal gyrus	left
12	middle frontal gyrus	right	44	parahippocampal gyrus	right
13	inferior frontal gyrus	left	45	entrhinal cortex	left
14	inferior frontal gyrus	right	46	entrhinal cortex	right
15	medial fronto-orbital gyrus	left	47	superior occipital gyrus	left
16	medial fronto-orbital gyrus	right	48	superior occipital gyrus	right
17	lateral fronto-orbital gyrus	left	49	middle occipital gyrus	left
18	lateral fronto-orbital gyrus	right	50	middle occipital gyrus	right
19	gyrus rectus	left	51	inferior occipital gyrus	left
20	gyrus rectus	right	52	inferior occipital gyrus	right
21	precentral gyrus	left	53	cuneus	left
22	precentral gyrus	right	54	cuneus	right
23	postcentral gyrus	left	55	lyngual gyrus	left
24	postcentral gyrus	right	56	lyngual gyrus	right
25	superior parietal gyrus	left	57	amygdala	left
26	superior parietal gyrus	right	58	amygdala	right
27	precuneus	left	59	hippocampus	left
28	precuneus	right	60	hippocampus	right

29	cingular gyrus	left	61	cerebellar hemisphere	left
30	cingular gyrus	right	62	cerebellar hemisphere	right
31	supramarginal gyrus	left	63	insular cortex	left
32	supramarginal gyrus	right	64	insular cortex	right

Figures

Figure 1

Results of repeatability and reliability of brain region volume calculated by iBEAT with correction. (A) Bland-Altman graph of whole brain volume between two measurements with 95% of the differences located in the mean±1.96SD (standard deviation) range showing the good repeatability; (B) Intraclass correlation coefficient (ICC) of volume of 64 brain regions between two measurements.

Figure 2

Comparisons of the calculated volume of 64 brain regions between BET, iBEAT and iBEAT with correction by analysis of variance (ANOVA).

Figure 3

Comparisons of small-world property of brain structural network between BET, iBEAT and iBEAT with correction. A small-world network should fulfill the following conditions: $C_p/C_{rand} > 1$ and $L_p/L_{rand} \approx 1$; C_p and L_p indicates the clustering coefficient and characteristic path length, respectively.

Figure 4

Comparisons of properties of brain structural network between BET, iBEAT and iBEAT with correction. (A) clustering coefficient, (B) characteristic path length, (C) global efficiency, (D) local efficiency.

Figure 5

Illustrations of skull stripping and registration by BET, iBEAT and iBEAT with manual correction.

Step1: Data preprocessing

Step2: Calculations of brain region volume

Step3: Construction of brain structural network

Figure 6

Workflow for construction of neonatal brain structural network. AC-PC align refers to the alignment of data according to anterior commissure-posterior commissure line.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Equation.docx](#)