Impact of Sofosbuvir and Daclastavir on Health-related Quality of Life in Patients Co-infected With Hepatitis C and Human Immunodeficiency Virus

Evy Yuniastuti (✉ evy.yuniastuti@gmail.com)

Faculty of Medicine Universitas Indonesia / Cipto Mangunkusumo Hospital <https://orcid.org/0000-0001-6650-0559>

Fhadilla Amelia

HIV Integrated Services, Ciptomangunkusumo Hospital, Jakarta, Indonesia

Arini Ika Hapsari

HIV Integrated Services, Ciptomangunkusumo Hospital, Jakarta, Indonesia

Bramantya Wicaksana

HIV Integrated Services, Ciptomangunkusumo Hospital, Jakarta, Indonesia

Veritea Natali

HIV Integrated Services, Ciptomangunkusumo Hospital, Jakarta, Indonesia

Alvina Widhani

Allergy and Clinical Immunology Division, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia/Ciptomangunkusumo Hospital, Jakarta, Indonesia

Andri Sanityoso Sulaiman

Hepatobiliary Division, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia

Teguh Harjono Karjadi

Allergy and Clinical Immunology Division, Department of Internal Medicine, Faculty of Medicine, Universitas Indonesia/Ciptomangunkusumo Hospital, Jakarta, Indonesia

Research

Keywords: health-related quality of life, hepatitis C, HCV, HIV, sofosbuvir, daclastavir

Posted Date: December 16th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-128529/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

We conducted a real-life study of health-related quality of life (HRQoL) transformation before and 12 weeks after sofosbuvir and daclatasvir therapy in HCV/HIV co-infected patients. Factors related to the significant changes of each HRQoL domain were also evaluated.

Methods

A prospective study was performed in the HIV integrated clinic at Cipto Mangunkusumo Hospital, Jakarta. HCV/HIV co-infected patients who started sofosbuvir and daclatasvir from government free DAA program in 2017-2019. WHOQoL-HIV BREF and RAND SF-36 questionnaires were recorded at baseline and post-treatment week 12.

Results

145 patients with mean age of 37.8 years (SD=4.2) were included in the analysis. Most of patients were male (89%), previous IVDU (89%), active smoker (50.4%) and non-cirrhosis (80%). SVR12 was achieved in 95.5% of patients. Sofosbuvir and daclatasvir treatments showed positive impacts on 4 domains of WHOQoL-HIV BREF and 3 domains of SF-36. Predicting factors of significant increase in each domain were: male and normal body mass index (BMI) for level of independence (RR 4.01,95% CI 1.09-14.74 and 4.80,95% CI 1.79-12.81); higher HCV-RNA for overall perception of QoL (RR 0.42,95% CI 0.18-0.94); non-smoking status for overall perception of health (RR 0.32,95% CI 0.15-0.66); male and fibrosis stage 0-1 for general health (6.21,95% CI 1.69-22.88 and 2.86,95% CI 1.16-7.00). Spiritual/personal belief decline was predicted by non-smoking status (RR 0.49,95% CI 0.24-0.98). Treatment success was not associated with any changes of HR-QoL.

Conclusions

HCV/HIV co-infected patients were successfully treated with sofosbuvir and daclatasvir and experienced improvement of HRQoL 12 weeks after treatment completion.

Introduction

HCV and HIV co-infection is a public health problem affecting more than 2 million people worldwide [1]. Evidence shows that HCV/HIV co-infection cause several negative impacts on the patients, including persistent HCV viremia, higher HCV viral load, and faster fibrosis progression [2]. The global use of combination antiretroviral therapy (cART) has made significant improvements in AIDS-related morbidity and mortality, but for HCV/HIV co-infected patients, liver-related mortality remains the common cause of death [3–5]. The reported prevalence of HCV among HIV-infected patients in Indonesia was 17.9% (95% CI 15.0-20.5), even higher in intravenous drug user (IVDU) population (81.6%; 95% CI 71.1–90.3), indicating that Indonesia has one of the highest rates of HCV/HIV co-infection in South East Asia [6].

The World Health Organization (WHO) called for the elimination of viral hepatitis by 2030 via recommendation of hepatitis treatment for all HCV-infected patients, including HCV/HIV co-infected patients [7]. The new HCV drugs, direct-acting antivirals (DAAs), has revolutionized the clinical management of HCV-infected patients. The introduction of these drugs has made HCV the first chronic viral infection that can be cured. This can be achieved in more than 90% of infected individuals, including HCV/HIV-coinfected patients, with limited side effects [8, 9]. Moreover, DAAs also showed promising results in reducing morbidity, mortality, extrahepatic manifestation, and progression to hepatocellular carcinoma [10].

In recent years, health-related quality of life (HRQoL) has gained worldwide recognition as the gold standard of patient-reported outcome (PRO) [11]. The WHO defines HRQoL as an individual's perception of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectations, standards and concerns". It is a broad-ranging concept affected in a complex way by the person's physical health, psychological state, personal beliefs, social relationships and their relationship to salient features of their environment" [12]. Chronic diseases such as both HIV and HCV are strongly related to patients' quality of life. Moreover, HCV infection was related to extrahepatic manifestation that could worsen patient HRQoL and may cause depression in some severe cases, further resulting in disruption of work production and daily activities [13]. The transformation of HRQoL before and after an intervention may help clinicians to understand a patient's perspective. Furthermore, improvement in HRQoL in accordance with patient's well-being may also be related to economical gain [14, 15].

Clinical trials of DAAs therapy have exhibited improvement in HRQoL, mostly in HCV mono-infected patients [13, 16–20]. Few studies evaluated HRQoL transformation in HCV/HIV co-infected patients after DAA treatments. HCV/HIV co-infected patients were known to have a lower quality of life and lower QoL gain after DAA treatment [21, 22]. Furthermore, only limited studies are available on the impact of DAAs on HRQoL in HCV/HIV co-infected patients in real-life settings [22, 23].

In 2017, the Indonesian government started a free DAA program for HCV mono-infected and HCV/HIV co-infected patients, mainly using a combination of sofosbuvir and daclatasvir [24]. Therefore, we conducted a real-life observational study to evaluate HRQoL transformation 12 weeks after sofosbuvir and daclatasvir therapy in HIV and HCV co-infected patients. We also evaluated factors related to the significant changes in each HR-QoL domain.

Material And Methods

Patients and study design

This prospective observational study was performed in HIV integrated clinic Cipto Mangunkusumo Hospital, Jakarta that provide multidisciplinary HIV/HCV care. HCV/HIV-infected patients who started the government's free DAA treatment program between September 2017 and July 2019 were invited to join the study. Eligibility criteria included patients' age more than 18 years old, stable using cART and no active opportunistic infections. Pregnant or breastfeeding women, diagnosis of diabetes mellitus and chronic kidney disease patients were excluded from the study. Prospective data were recorded at baseline and post-treatment week 12. Patients who did not complete the full course of DAA treatment or post-treatment week 12 evaluation were not included in the analysis.

HCV Treatment

Non-cirrhotic patients (fibrosis stage 0–3) were treated with oral daily sofosbuvir (400 mg) and daclatasvir (60 or 90 mg) for 12 weeks while cirrhotic patients (fibrosis stage 4) were treated with daily sofosbuvir and daclatasvir for 24 weeks. Daclatasvir 90 mg was used in patients with efavirenz or nevirapine-based ART and daclatasvir 60 mg in other regimens. Undetectable HCV-RNA 12 weeks after treatment completion was defined as a successful treatment response or sustained virologic response (SVR12).

Sociodemographic and clinical variables

Sociodemographic variables were collected through personal interviews. These included education, marital status, employment status, tobacco use, and mode of HCV acquisition. Recent CD4⁺ T-cell counts, cART combination, prior interferon (IFN) treatment failure, Hepatitis B co-infection, haemoglobin levels, and BMI were taken from the patient's medical records. Before starting DAA treatment, all patients were required to have HCV-RNA quantification and fibrosis staging using transient elastography to define the duration of treatment. The following cut-offs were used to stage the liver fibrosis: F0-F1 < 7.1 kPa, F2 7.1–9.4 kPa, F3 9.5–12.4 kPa, and F4 ≥ 12.5 kPa [25].

Assessment of HRQoL

HRQoL was measured using the WHO Quality of Life for HIV, in its abbreviated version (WHOQoL-HIV BREF) and 36-items RAND Short Form survey (RAND SF-36) before and after treatment completion at 12 weeks. Patients were not aware of their final HCV-RNA results before completing these HRQoL questionnaires.

WHOQoL-HIV BREF is a disease-specific instrument which assesses HRQoL of HIV-infected patients. It consists of 31 questions that are grouped into six domains: physical, psychological, level of independence, social relations, environment, and spiritual/religion/personal beliefs. There are two other items that ask about an individual's overall perception of quality of life and overall perception of health. The point for each answer ranges from 1 to 5. Most items are scored in the positive direction. The mean score of each question within each domain is used as the domain score. The mean scores were then multiplied by 4 resulting in final scores ranging between 4 and 20 [26, 27]. WHOQoL-HIV BREF had been translated into the Indonesian language with internal consistency (Cronbach's alpha) of 0.513–0.798 [28].

The RAND 36-Item Health Survey is a generic quality of life instrument that measures eight domains: physical functioning, role limitations due to physical health problems, bodily pain, general health, energy/fatigue, social functioning, role limitations due to personal or emotional problems, and emotional well-being. Moreover, it also includes one question that indicates perceived change in health. The items score range from 0–100 with greater scores indicating better health. Scores show the percentage of total possible score achieved. Items in the same domain are averaged to get the final score for each domain [29, 30]. The internal consistency of the Indonesian version of RAND SF-36 was 0.789 [31].

Statistical analysis

The data were analysed using the Statistical Package for Social Sciences (SPSS Inc, Chicago, IL) version 20.0 and GraphPad Prism 7 for Windows (GraphPad Software, Inc., La Jolla, CA, USA). Descriptive characteristics were presented with frequency and percentage for categorical variables and mean with SD or median with IQR (Q1–Q3) for continuous variables. Comparative analyses between each QoL component in WHOQoL-HIV-BREF and SF-36 were performed using *t*-tests for analyses of variance. Wilcoxon-rank test were used for non-parametric test. The outcome was dichotomized as improved vs not-improved to evaluate the predictors of each improved QoL component. Bivariate analysis was done using Chi-square test. Independent predictors of domain improvement were assessed using multivariate logistic regression binary model for all variables that had *p*-value < 0.25. A *p*-value of less than 0.05 was considered significant and risk ratio (RR) with 95% CI was calculated to determine the association.

Results

Clinical and demographic characteristics

Overall, 179 HCV/HIV co-infected patients were invited to participate before starting DAA treatment between September 2017 to July 2019, 173 of those patients fulfilled the inclusion criteria and signed the informed consent (97%). 145 (81%) patients completed evaluation and were finally analysed (Fig. 1).

Baseline demographic and clinical characteristics are summarized in Table 1 (Additional File 1). The mean age was 37.8 years (SD = 4.2) and 89% of patients were male. Previous IVDU represented majority of patients (89%). About half of patients (50.4%) were actively smoking while 45.5% were previous smokers. Most of the patients were married (58.6%) and employed (84.2%). The median duration of cART used was 9 years (IQR 4–12) and median CD4⁺ T-cell count was 485 cells/μL (IQR 284–676). Only 13 patients (9%) had been treated with previous HCV treatment standard (pegylated interferon and ribavirin combination). None of the patients ever used DAA treatment before. 4.1% patient were also hepatitis B virus (HBV) co-infected and using tenofovir disoproxil fumarate (TDF). Liver fibrosis degree was F0-F1 in 86 (59.3%) patients, F2-F3 in 30 (20.7%) and F4 in 29 (20%) patients. Sofosbuvir with 90 mg daclatasvir was used in majority of the patients (79.3%). Eighty percents of patients used sofosbuvir and daclatasvir for 12 weeks. Treatment success or SVR12 was achieved in 138 of 145 patients (95.5%).

Table 1
Clinical and demographic characteristics

Characteristics		N = 145
Male gender	n(%)	129 (89.0)
Age, years	Mean (SD)	37.8 (4.2)
Previous IVDU	n(%)	129 (89.0)
Education	University, n(%)	64 (44.1)
	Secondary, n(%)	79 (54.5)
	Primary, n(%)	2 (1.4)
Marital status	Married, n(%)	85 (58.6)
	Widow/widower/divorce, n(%)	22 (15.2)
	Not married, n(%)	38 (26.2)
Employment status	Regular employment, n(%)	67 (46.2)
	Non-regular employment, n(%)	56 (38.6)
	Not working, n(%)	22 (15.2)
Religion	Moslem, n(%)	114 (78.6)
	Christian	29 (20.0)
	Buddhist	2 (1.4)
Alcohol use	active, n(%)	25 (17.2)
	Past, n(%)	111 (76.6)
	Never, n(%)	9 (6.2)
Tobacco use	Active smoker, n(%)	73 (50.4)
	Past smoker, n(%)	66 (45.5)
	Never smoke, n(%)	6 (4.1)
BMI	Mean (SD)	22.3 (3.5)
Hemoglobin, g/dL	Mean (SD)	14.9 (4.3)
Recent CD4 ⁺ T cell count, cells/ μ L	Median (IQR)	485 (392–675)
cART duration, years	Median (IQR)	9 (4–12)
HBV co-infection	n(%)	6 (4.1)
HCV treatment history	Treatment naïve, n(%)	132 (91)
	Interferon failure, n(%)	13 (9)
Fibrosis stage	F0-F1, n(%)	86 (59.3)
	F2-F3, n(%)	30 (20.7)
	F4, n(%)	29 (20)
HCV-RNA, IU/mL	> 800,000, n(%)	110 (75.9)
	< 800,000, n(%)	35 (24.1)
	Mean (SD), log10	6.22 (0.83)
HCV genotype	1, n(%)	23 (15.8)
	2, n(%)	4 (2.8)
	3, n(%)	3 (2.1)
	4, n(%)	3 (2.1)
	Not available, n(%)	112 (77.2)

IVDU: intravenous drug user; BMI: body mass index; cART: combination antiretroviral therapy; HBV: hepatitis B virus; HCV: hepatitis C virus; HCV-RNA: hepatitis C virus-ribonucleic acid; DAA: direct-acting antiviral; SD: standard deviation; IQR: interquartile range; SOF + DAC: sofosbuvir and daclastavir; SVR12: sustained virological response after 12 weeks of treatment

Characteristics		N = 145
DAA combination	SOF + DAC60, n(%)	30 (20.7)
	SOF + DAC90, n(%)	115 (79.3)
Duration of DAA treatment	12 weeks, n(%)	116 (80)
	24 weeks, n(%)	29 (20)
DAA treatment response	SVR12, n(%)	138 (95.5)
	Non-responder, n(%)	7 (4.5)
IVDU: intravenous drug user; BMI: body mass index; cART: combination antiretroviral therapy; HBV: hepatitis B virus; HCV: hepatitis C virus; HCV-RNA: hepatitis C virus-ribonucleic acid; DAA: direct-acting antiviral; SD: standard deviation; IQR: interquartile range; SOF + DAC: sofosbuvir and daclatasvir; SVR12: sustained virological response after 12 weeks of treatment		

Impact of sofosbuvir and daclatasvir treatment to HR-QoL changes

Sofosbuvir and daclatasvir treatment showed positive impacts on 4 of 8 domains in WHOQoL-HIV BREF (level of independence, environment, overall perception of quality of life and overall perception of health) and negative impact to 1 domain (spiritual/religion/personal belief) as seen in Fig. 2. Statistically non-significant improvements were observed in the physical and psychological domains. Significant increments were observed in 3 of 9 domains in SF-36: general health, energy/fatigue, and health change (Fig. 2). However, domain of pain indicated significant deterioration. We also observed improvements in physical functioning and mental health that were aligned with WHOQoL-HIV BREF assessment, although these domains did not reach statistical significance. In both tools, perception of health shown the highest significant improvement in 12 weeks after the end of treatment.

Factors related to the change in HR-QoL domains

Increasing level of independence can be predicted by gender and baseline BMI in WHOQoL-HIV BREF. It increased 4.01 times higher in male patients (95% CI 1.09–14.74) and 4.80 times higher in patients with normal BMI than patients with obesity (95% CI 1.79–12.81) (Table 2) (Additional File 2). Higher baseline HCV-RNA independently predicted lower improvement of overall perception of QoL (RR 0.42, 95% CI 0.18–0.94). Improvement of overall perception of health was associated with non-smoking status of the patients. It increased 0.32 times lower in non-smoker than smoker (RR 0.32, 95% CI 0.15–0.66). In the area of spiritual/religion/personal belief, non-smoking status was associated with lower spirituality decline (RR 0.49, 95% CI 0.24–0.98). We did not find any factor that can predict improvement of environmental component.

Table 2
Predictors of WHOQoL-HIV BREF component 12 weeks after DAA treatment

	Level of Independence				Environment				Spiritual/religion/personal beliefs				Overall Perception of health			
	Bivariate		Multivariate		Bivariate		Multivariate		Bivariate		Multivariate		Bivariate		Multivariate	
	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	P	RR (95% CI)	P	RR (95% CI)	P	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	P
Male gender	2.56 (0.91–7.22)	0.050	4.01 (1.09–14.74)	0.037	0.86 (0.57–1.29)	0.677			1.12 (0.52–2.40)	0.992			1.49 (0.52–4.28)	0.559		
Age ≤ 40 year	1.02 (0.69–1.52)	1.000			1.09 (0.78–1.53)	0.735			1.50 (0.85–2.63)	0.203	1.80 (0.81–4.03)	0.151	0.84 (0.48–1.48)	0.702		
University education	0.84 (0.58–1.23)	0.462			0.96 (0.71–1.29)	0.901			0.84 (0.53–1.34)	0.581			0.71 (0.40–1.25)	0.306		
Married vs not married	1.50 (0.69–3.26)	0.407			0.83 (0.39–1.79)	0.334			1.06 (0.47–2.42)	0.496			0.97 (0.42–2.25)	0.604		
Married vs Widow/divorce	1.71 (0.65–4.50)	0.407			0.48 (0.18–1.29)	0.334			0.59 (0.23–1.53)	0.496			1.77 (0.54–5.77)	0.604		
Regular worker vs non-regular	0.57 (0.28–1.17)	0.127	0.62 (0.29–1.31)	0.215	1.25 (0.62–2.55)	0.735			1.13 (0.54–2.34)	0.933			1.46 (0.66–3.22)	0.383		
Regular worker vs not working	1.45 (0.52–4.06)	0.127	0.77 (0.23–2.56)	0.663	0.90 (0.33–2.41)	0.735			0.96 (0.35–2.65)	0.933			2.14 (0.64–7.17)	0.383		
CD4 > 500 vs 350–500	1.47 (0.57–3.82)	0.636			0.71 (0.28–1.81)	0.772			1.04 (0.39–2.79)	0.944			0.65 (0.24–1.80)	0.787		
CD4 > 500 vs < 350	0.79 (0.37–1.72)	0.636			0.77 (0.35–1.67)	0.772			0.82 (0.37–1.85)	0.944			0.71 (0.30–1.69)	0.787		
HCV-RNA > 800000 IU/mL	1.02 (0.66–1.57)	1.000			1.04 (0.73–1.49)	0.992			0.87 (0.53–1.44)	0.749			0.55 (0.32–0.93)	0.054		
Interferon failure	1.03 (0.56–1.91)	1.000			0.54 (0.24–1.24)	0.132	0.34 (0.10–1.15)	0.083	0.42 (0.12–1.54)	0.220	0.40 (0.08–1.97)	0.261	1.16 (0.49–2.75)	0.748		
Achieving SVR12	1.60 (0.49–5.23)	0.460			1.29 (0.54–3.06)	0.702			0.58 (0.29–1.16)	0.234	0.24 (0.05–1.24)	0.087	1.93 (0.31–12.07)	0.675		
Non-smoker	0.92 (0.64–1.33)	0.796			0.81 (0.60–1.09)	0.214	0.65 (0.33–1.26)	0.197	0.62 (0.39–0.99)	0.234	0.49 (0.24–0.98)	0.043	1.18 (0.69–2.03)	0.671		
BMI normal vs underweight	0.71 (0.27–1.91)	0.001	0.75 (0.28–2.03)	0.572	0.63 (0.23–1.72)	0.554			0.70 (0.26–1.90)	0.685			0.60 (0.22–1.64)	0.108		
BMI normal vs overweight/obese	4.99 (1.88–13.19)	0.001	4.80 (1.79–12.81)	0.002	1.17 (0.53–2.57)	0.554			0.76 (0.33–1.72)	0.685			2.32 (0.81–6.65)	0.108		
Non-IVDU	0.67 (0.32–1.42)	0.373			0.91 (0.54–1.52)	0.908			0.70 (0.29–1.69)	0.571			0.92 (0.38–2.25)	1.000		
Fibrosis stage F0-F1 vs F2-F3	1.09 (0.47–2.51)	0.982			1.01 (0.44–2.34)	0.506			0.69 (0.29–1.62)	0.631			1.06 (0.40–2.83)	0.323		
Fibrosis stage F0-F1 vs F4	1.02 (0.44–2.38)	0.982			1.63 (0.70–3.80)	0.506			0.75 (0.31–1.80)	0.631			0.53 (0.22–1.30)	0.323		
Non-anemia	1.30 (0.74–2.26)	0.451			1.29 (0.81–2.06)	0.349			0.95 (0.53–1.69)	1.000			1.15 (0.54–2.44)	0.912		

HCV-RNA: hepatitis C virus ribonucleic acid; SVR12: sustained virological response at 12 weeks post treatment; BMI: body mass index; IVDU: intravenous drug use

Male patients exhibited 6.21 times higher general health increase than female patients (95% CI 1.69–22.88, $p = 0.006$) (Table 3) (Additional File 3). Patients with fibrosis stage 0–1 indicated 2.85 times higher general health increase than those with fibrosis stage 4 (95% CI 1.16–7.00, $p = 0.023$) We did not find any significant predictor of improvement of energy and health change domains, nor predictor of pain worsening in this study. Only history of interferon failure showed a better trend of health change improvement (RR 3.09, 95% CI 0.96–10.01, $p = 0.059$).

Table 3
Predictors of SF-36 component 12 weeks after DAA treatment

	Pain		General Health				Energy/fatigue				Health Change			
	Bivariate		multivariate		Bivariate		multivariate		Bivariate		multivariate		Bivariate	
	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p
Male gender	1.05 (0.43– 2.58)	1.000			3.14 (1.12– 8.80)	0.005	6.21 (1.69– 22.88)	0.006	1.79 (0.85– 3.75)	0.111	2.78 (0.91– 8.46)	0.072	2.07 (0.73– 5.87)	0.196
Age ≤ 40 year	0.72 (0.42– 1.26)	0.356			0.91 (0.67– 1.25)	0.695			1.12 (0.79– 1.60)	0.627			1.07 (0.66– 1.73)	0.935
University education	1.03 (0.59– 1.77)	1.000			0.96 (0.71– 1.29)	0.901			0.95 (0.70– 1.30)	0.871			0.83 (0.53– 1.29)	0.511
Married vs not married	1.56 (0.63– 3.88)	0.574			1.05 (0.49– 2.27)	0.648			0.57 (0.26– 1.25)	0.351			1.59 (0.71– 3.57)	0.178
Married vs Widow/divorce	1.42 (0.47– 4.26)	0.574			1.56 (0.61– 4.00)	0.648			0.98 (0.38– 2.50)	0.351			2.50 (0.84– 7.40)	0.178
Regular worker vs non-regular	1.76 (0.78– 3.96)	0.356			0.90 (0.44– 1.85)	0.020	1.08 (0.50– 2.35)	0.843	1.26 (0.62– 2.56)	0.380			0.97 (0.47– 2.02)	0.880
Regular worker vs not working	1.42 (0.47– 4.26)	0.356			3.69 (1.27– 10.71)	0.020	2.52 (0.76– 8.39)	0.132	2.00 (0.74– 7.37)	0.380			1.26 (0.45– 3.56)	0.880
CD4 > 500 vs 350–500	1.06 (0.36– 3.10)	0.957			0.60 (0.23– 1.55)	0.676			0.84 (0.33– 2.12)	0.914			0.62 (0.24– 1.60)	0.206
CD4 > 500 vs < 350	0.86 (0.36– 2.06)	0.957			0.84 (0.39– 1.83)	0.676			0.84 (0.39– 1.83)	0.914			0.62 (0.28– 1.37)	0.206
HCV-RNA > 800000 IU/mL	0.99 (0.52– 1.88)	1.000			0.97 (0.69– 1.37)	1.000			0.94 (0.66– 1.32)	0.861			0.78 (0.49– 1.23)	0.399
Interferon failure	0.56 (0.15– 2.08)	0.515			0.99 (0.58– 1.67)	1.000			1.02 (0.60– 1.72)	1.000			1.81 (1.11– 2.95)	0.069
Achieving SVR12	0.91 (0.27– 3.05)	1.000			0.75 (0.46– 1.23)	0.455			1.90 (0.58– 6.20)	0.253			0.85 (0.35– 2.05)	0.706
Non-smoker	1.01 (0.59– 1.75)	1.000			0.89 (0.66– 1.20)	0.564			0.89 (0.66– 1.21)	0.564			1.05 (0.69– 1.62)	0.950
BMI normal vs underweight	1.20 (0.40– 3.64)	0.659			0.66 (0.24– 1.80)	0.437			0.60 (0.22– 1.65)	0.440			0.40 (0.15– 1.08)	0.177
BMI normal vs overweight/obese	1.54 (0.60– 3.98)	0.659			1.37 (0.62– 3.02)	0.437			1.26 (0.57– 2.77)	0.440			0.90 (0.39– 2.06)	0.177
Non-IVDU	1.22 (0.56– 2.68)	0.763			0.32 (0.11– 0.89)	0.005	0.27 (0.07– 1.13)	0.072	0.94 (0.56– 1.57)	1.000			1.23 (0.67– 2.24)	0.720
Fibrosis stage F0-F1 vs F2-F3	1.00 (0.37– 2.66)	0.276			1.28 (0.55– 2.94)	0.243	1.69 (0.70– 4.08)	0.240	1.20 (0.52– 2.75)	0.298			0.76 (0.32– 1.80)	0.690
Fibrosis stage F0-F1 vs F4	0.50 (0.20– 1.22)	0.276			2.06 (0.88– 4.85)	0.243	2.85 (1.16– 7.00)	0.023	0.55 (0.23– 1.32)	0.298			0.72 (0.30– 1.71)	0.690

HCV-RNA: hepatitis C virus ribonucleic acid; SVR12: sustained virological response at 12 weeks post treatment; BMI: body mass index; IVDU: intravenous drug

	Pain				General Health				Energy/fatigue				Health Change			
	Bivariate		multivariate		Bivariate		multivariate		Bivariate		multivariate		Bivariate			
	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p	RR (95% CI)	p		
Non-anemia	0.58 (0.33– 1.04)	0.141	0.46 (0.19– 1.13)	0.090	1.16 (0.75– 1.80)	0.621			1.40 (0.84– 2.32)	0.221	1.68 (0.68– 4.12)	0.261	1.63 (0.79– 3.39)	0.228	1	(
HCV-RNA: hepatitis C virus ribonucleic acid; SVR12: sustained virological response at 12 weeks post treatment; BMI: body mass index; IVDU: intravenous drug																

Sustained virologic response-achievers and non-responders did not reach significant change on any domain of HR-QoL in both questionnaires.

Discussion

Our study is one of few cohorts evaluating impact of DAA treatment in quality of life of HCV/HIV co-infected patients. Early on, improved HRQoL was shown to be associated with a SVR of IFN-based treatment that has high toxicity, more complexity and lower treatment uptake in many low-middle income countries [32]. In HCV/HIV co-infected patients, SVR rates are historically 20–30% lower than in HCV mono-infected patients [33]. The introduction of interferon-free DAA treatment has been a significant breakthrough since this combination is likely to close the gap of SVR between HCV/HIV co-infection and HCV mono-infection [8, 9].

Results from clinical trials of DAA treatments have shown improvements in the physical and psychological components of HRQoL, mostly in HCV mono-infected patients [16, 19, 34]. Many clinicians were concerned that clinical trial results cannot be generalized to real-world situations due to the studies' strict selection criteria [35]. Several researchers have attempted to investigate the impact of DAA treatments on HR-QoL of HCV/HIV co-infected patients, with various treatment combinations and tools, with conflicting results [23, 34, 36, 37]. In studies comparing HCV mono-infected and HCV/HIV co-infected patients, HCV/HIV co-infected patients were associated with significantly lower HR-QoL and lower gain in the HR-QoL scores [21, 22]. Our study design aimed to inform real-world impact of sofosbuvir and daclatasvir combination treatment in a younger and mainly former IVDU population.

We found that the improvements were observed across most of HR-QoL domains in both WHOQOL-HIV BREF and SF-36. This finding provides additional evidence that DAA treatment has a positive influence on HCV/HIV co-infected patients on cART [13, 23, 34, 37, 38]. Improvements were not observed in some domains when comparing 12 weeks after treatment scores with baseline scores. Longer study duration and more follow-up timepoints could show potential benefits in more domains.

Using multivariate analyses, we determined several baseline predictors of several WHOQoL-HIV BREF domain changes 12 weeks after treatment completion. We found that male patients had improved their level of independence 4 times higher than female patients after DAA treatment, which is consistent with previous studies in male HIV patients [39–41]. Most female patients in our study were married and not working despite being highly educated. Tesfay, et al [42] showed that monthly income was an independent predictor of poor HR-QoL among female HIV patients. We also found that having normal BMI was associated with better improvement in level of independence compared to being overweight/obese. This result is aligned with a study in Southern Ethiopia that showed normal BMI significantly improved QoL score of HIV-infected patients. Protection from infectious diseases, improvement of health status, and the ability to live a productive life are promoted by better nutritional status.[43] Moreover, people with obesity have higher risk of having mobility disability, eventuating a higher risk of becoming unemployed [44, 45].

Lower improvement of overall perception of QoL was noticed in patients with higher baseline HCV-RNA. HCV viremia has been associated with depression and fatigue [46]. Younossi, et al [34]. found that in HCV/HIV co-infected patients, continuous viremia was associated with substantial impairment in QoL. However, we did not observe differences of any HR-QoL changes in SVR responders compared to non-responders. All patients were well-informed of the high success rate of these expensive but free drugs. Though the final questionnaires were delivered before HCV-RNA results came out, many patients were confident of their treatment success. Yeung, et al, demonstrated that those achieving an SVR had higher HR-QoL scores over time. Only 38% participants in that study achieved SVR, 30% did not respond, 13% had ongoing treatment, and 17% had unknown treatment response [38]. Our study had a much higher treatment response (95.5%), but the shorter period of HR-QoL evaluation might be insufficient. A longer duration of evaluation might be needed to see further impact of SVR as another study with high treatment response found modest immediate improvement following SVR, then continued thereafter [38].

Overall perception of health significantly improved after treatment in our study, and this effect was more prevalent in smokers. Our finding contradicted with Toghianifar, et al [47] study that showed overall QoL among current and past smokers was decreased relative to non-smokers. Some evidence of disparaging behaviour towards relative risk of lung cancer on smoker population might explain this finding [48]. We also observed a significant reduction on spiritual/religion belief domain compared to baseline, and this effect was higher in smokers. Highly successful treatment belief might influence our participants' perception of health and spiritual, since patients did not know their HCV-RNA results when they filled in their final questionnaires. As they had used cART for a median of 9 years and knew their untreated HCV status for a long time, completing DAA treatment would be considered as morbidity risk reduction for these patients despite their smoking status. Evidence has shown that smoking behaviour was significantly related to religious involvement (religious attendance, importance, religious/spiritual comfort-seeking, and religious/spiritual decision-making). Higher religious involvement was linked with a lower risk of being persistent or ex-smoker [49]. Islam as a dominant religion among participants (78.6%) might also play a role in this finding as smoking is considered as a discouraged act (mukrooh) in Islamic law [50].

We proceeded to determine baseline predictors of several SF-36 domain changes 12 weeks after treatment completion in the studied population. Similar to other reports, we demonstrated a significant rise in the general health domain [34, 51]. Positive relationships between general health increase with male gender and stage F0-F1 fibrosis were also noticed. This gender association was in line with the level of independence domain in WHOQoL-HIV BREF. Younossi et al also confirmed a similar increase of general health among stage F0-F1 fibrosis in HCV mono-infected patients treated with sofosbuvir and ledipasvir [52].

Limitations

There are several limitations to our study. Firstly, the study was done in a tertiary center and the population size was not large enough to acquire conclusive results of certain subanalyses. However, we were able to evaluate 81% of the patients who received treatment during the study period. Since it was done in the early phase of free DAA treatment program in Indonesia, we believe that these encouraging results would endorse the expansion of the program. Secondly, we only used WHOQoL-HIV BREF and RAND SF-36 whereas many studies use multiple instruments [53–57]. Moreover, we did not evaluate depression and anxiety as other factors that could be potentially related to QoL.

Despite those limitations, our study could give a standing point for future research on QoL studies and health outcome improvement among HCV/HIV co-infected patient. These findings provide information about QoL and some influencing factors among HCV/HIV co-infected patients in Indonesia where studies in these cohorts are still limited.

Conclusions

In summary, our study indicates that treatment with sofosbuvir and daclatasvir is associated with improvement of quality of life 12 weeks after treatment completion in HCV/HIV co-infected patients. Our data support the fact that treating HCV, including in HIV co-infected patients, will lead to substantial PRO improvement in addition to the possibility of curing HCV. Expanding free access to this simple and highly active treatment is important for HIV-infected patients on cART.

Abbreviations

HRQoL: Health-Related Quality of Life; DAA: Direct-Acting Antiviral; IVDU: Intravenous Drug User; WHOQoL-HIV BREF: WHO Quality of Life for HIV; cART: Combination Antiretroviral Therapy; PRO: Patient-Reported Outcome; SVR12: Sustained virological response at 12 weeks post treatment; IFN: Interferon; RAND SF-36: 36-items RAND Short Form survey; TDF: Tenofovir Disoproxil Fumarate

Declarations

Ethics approval and consent to participate

Ethics Committee at the Faculty of Medicine Universitas Indonesia approved the study before initiation. Each participant gave their written informed consent after explanation of the study procedure and purpose. Permission was given for the collection of medical data from medical records. Trained researchers were available to assist when needed.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

Not Applicable

Authors' contributions

EY, AW, ASS designed the study. EY, FA, VN, AW, ASS, THK collected and managed the data. EY, FA, AIH, BW performed statistical analysis and draft the manuscript. EY was a major contributor in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors thank Salfia Lastari and Utami Susilawati for their statistical analysis, Jacquita Suemarni Affandi for the valuable input, Madyaningati, all HIV integrated center staffs, and all patients who participated in the study.

References

1. Platt L, Easterbrook P, Gower E, Al E. Prevalence and burden of HCV co-infection in people living with HIV: a global systematic review and meta-analysis. *Lancet Infect Dis* 2016; 16: 797–808.
2. Hernandez, MD. Sherman K. HIV/HCV coinfection natural history and disease progression, a review of the most recent literature. *Curr Opin HIV AIDS* 2011; 6: 478–82.
3. Smith CJ, Ryom L, Weber R, Morlat P, Pradier C, Reiss P, et al. Trends in underlying causes of death in people with HIV from 1999 to 2011 (D:A:D): a multicohort collaboration. *Lancet* 2014; 384: 241–8.
4. Croxford S, Kitching A, Desai S, Kall M, Edelstein M, Skingsley A, et al. Mortality and causes of death in people diagnosed with HIV in the era of highly active antiretroviral therapy compared with the general population: an analysis of a national observational cohort. *Lancet Public Health* 2017; 2: 35–46.
5. Hernando V, Perez-Cachafeiro S, Lewden C et al. All-cause and liver-related mortality in HIV positive subjects compared to the general population: differences by HCV co-infection. *J Hepatol* 2012; 57: 743–51.
6. Martinello M, Amin J, Matthews GV, Dore GJ. Prevalence and disease burden of HCV coinfection in HIV cohorts in the Asia Pacific region: A systematic review and meta-analysis. *AIDS Rev* 2016; 18: 68–80.
7. World Health Organization. Hepatitis C. 2020. <https://www.who.int/en/news-room/fact-sheets/detail/hepatitis-c>.
8. Chung RT, Baumert TF. Curing chronic hepatitis C – the arc of a medical triumph. *N Engl J Med* 2014; 370: 1576–8.
9. Abutaleb A, Sherman KE. A changing paradigm: management and treatment of the HCV/HIV-co-infected patient. *Hepatol Int* 2018; 12: 500–9.
10. World Health Organization. Guidelines for the care and treatment of persons diagnosed with chronic hepatitis C virus infection. Geneva: WHO; 2018.
11. Bullinger M, Quitmann J. Quality of life as patient-reported outcomes: Principles of assessment. *Dialogues Clin Neurosci* 2014; 16: 137–45.
12. World Health Organization. WHOQOL: measuring quality of life. <https://www.who.int/healthinfo/survey/whoqol-qualityoflife/en/>.
13. Younossi ZM, Stepanova M, Sulkowski M, Naggie S, Henry L, Hunt S. Sofosbuvir and ledipasvir improve patient-reported outcomes in patients co-infected with hepatitis C and human immunodeficiency virus. *J Viral Hepat* 2016; 23: 857–65.
14. Groen WG, Kuijpers W, Oldenburg HS, Wouters MW, Aaronson NK, van Harten WH. Supporting lung cancer patients with an interactive patient portal: feasibility study. *JMIR Cancer* 2017; 3: e10.
15. Muragundi P, Tumkur A, Shetty R, Naik A. Health-related quality of life measurement. *J Young Pharm* 2012; 4: 54.
16. Younossi ZM, Henry L. Systematic review: patient-reported outcomes in chronic hepatitis C—the impact of liver disease and new treatment regimens. *Aliment Pharmacol Ther* 2015; 41: 497–520.
17. Younossi ZM, Stepanova M, Nader F et al. Patient-reported outcomes in chronic hepatitis C patients with cirrhosis treated with sofosbuvir-containing regimens. *Hepatology* 2014; 59: 2161–9.
18. Stepanova M, Nader F, Cure S, Bourhis F, Hunt S, Younossi Z. Patients' preferences and health utility assessment with SF-6D and EQ-5D in patients with chronic hepatitis C treated with sofosbuvir regimens. *Aliment Pharmacol Ther* 2014; 40: 676–85.
19. Younossi ZM, Stepanova M, Feld J et al. Sofosbuvir/velpatasvir improves patient-reported outcomes in HCV patients: results from ASTRAL-1 placebo-controlled trial. *J Hepatol* 2016; 65: 33–9.
20. Younossi ZM, Stepanova M, Sulkowski M, Foster GR, Reau N, Mangia A, et al. Ribavirin-free regimen with sofosbuvir and velpatasvir is associated with high efficacy and improvement of patient-reported outcomes in patients with geno- types 2 and 3 chronic hepatitis C: results from Astral-2 and -3 clinical trials. *Clin Infect Dis* 2016; 63: 1042–8.
21. Pereira M, Fialho R. Assessment of factors associated with the quality of life of patients living with HIV/HCV co-infection. *J Behav Med* 2016; 39: 767–81.
22. Goñi-Esarte S, Juanbeltz R, Martínez-Baz I, Castilla J, Miguel RS, Herrero JI, et al. Long-term changes on health-related quality of life in patients with chronic hepatitis C after viral clearance with direct-acting antiviral agents. *Rev Esp Enfermedades Dig* 2019; 111: 445–52.
23. Scheiner B, Schwabl P, Steiner S, Bucsics T, Chromy D, Aichelburg MC, et al. Interferon-free regimens improve health-related quality of life and fatigue in HIV/HCV-coinfected patients with advanced liver disease. *Medicine (Baltimore)* 2016; 95: e4601.
24. World Health Organization. Progress report on access to hepatitis C treatment. Focus on overcoming barrier in low- and middle-income countries. 2018. <https://www.who.int/hepatitis/publications/hep-c-access-report-2018/en/>.
25. Truong TN, Laureillard D, Lacombe K, Thi HD, Hanh PPT, Xuan LTT, et al. High proportion of HIV-HCV coinfecting patients with advanced liver fibrosis requiring hepatitis C treatment in Haiphong, Northern Vietnam (ANRS 12262). *PLoS One* 2016; 11: e0153744.
26. Khademi N, Saeidi S, Zangeneh A, Saeidi F, Choobtashani M. The relationship between life satisfaction and quality of life among people living with HIV/AIDS in Kermanshah-West of Iran. *Health Soc Care Community* 2020; 28: 2218–24
27. World Health Organization. *WHOQOL-HIV instrument users manual*. 2002.
28. Muhammad NN, Shatri H, Djoerban Z, Abdullah M. Uji kesahihan dan keandalan kuesioner World Health Organization Quality of Life-HIV Bref dalam bahasa Indonesia untuk mengukur kualitas hidup pasien HIV/AIDS. *J Penyakit Dalam Indones* 2017; 4: 112.
29. van Westrienen PE, Pisters MF, Toonders SAJ, Gerrits M, de Wit NJ, Veenhof C. Quality of life in primary care patients with moderate medically unexplained physical symptoms. *Qual Life Res* 2020; 29: 693–703.
30. Ware JE, Gandek B. Overview of the SF-36 Health Survey and the International Quality of Life Assessment (IQOLA) project. *J Clin Epidemiol* 1998; 51: 903–12.
31. Salim S, Yamin M, Alwi I, Setiati S. Validity and reliability of the Indonesian version of SF-36 quality of life questionnaire on patients with permanent pacemakers. *Acta Medica Indones* 2017; 49: 10–6.

32. Bonkovsky HL WJ. Reduction of health-related quality of life in chronic hepatitis C and improvement with interferon therapy. The Consensus Interferon Study Group. *Hepatology* 1999; 29: 264–70.
33. Torriani FJ, Rodriguez-Torres M, Rockstroh JK, Lissen E. Peginterferon Alfa-2a plus ribavirin for chronic hepatitis C virus infection in HIV- infected patients. *N Engl J Med* 2004; 351: 438–50.
34. Younossi ZM, Stepanova M, Sulkowski M, Wyles D, Kottlil S, Hunt S. Patient-reported outcomes in patients co-infected with hepatitis C virus and human immunodeficiency virus treated with sofosbuvir and velpatasvir: The ASTRAL-5 study. *Liver Int* 2017; 37: 1796–804.
35. Saeed S, Strumpf EC, Walmsley SL, Rollet-Kurhajec K, Pick N, Martel-Laferrière V, et al. How generalizable are the results from trials of direct antiviral agents to people coinfectd with HIV/HCV in the real world? *Clin Infect Dis* 2016; 62: 919–26.
36. Younossi ZM, Stepanova M, Sulkowski M et al. Sofosbuvir and ledipasvir improve patient-reported outcomes in patients co-infected with hepatitis C and human immunodeficiency virus. *J Viral Hepat* 2016; 23: 857–65.
37. Yeung MW, Young J, Moodie E, Rollet-Kurhajec KC, Schwartzman K, Greenaway C, et al. Changes in quality of life, healthcare use, and substance use in HIV/hepatitis C coinfectd patients after hepatitis C therapy: A prospective cohort study. *HIV Clin Trials* 2015; 16: 100–10.
38. Saeed S, Moodie EEM, Strumpf E, Gill J, Wong A, Cooper C, et al. Real-world impact of direct acting antiviral therapy on health-related quality of life in HIV/Hepatitis C co-infected individuals. *J Viral Hepat* 2018; 25: 1507–14.
39. Bakiono F, Ouédraogo L, Sanou M, Samadoulougou S, Kirakoya-Samadoulougou PWLGF, Robert A. Quality of life in people living with HIV: a cross-sectional study in Ouagadougou, Burkina Faso. *Springerplus* 2014; 3: 372.
40. Nobre N, Pereira M, Roine RP, Sintonen H, Sutinen J. Factors associated with the quality of life of people living with HIV in Finland. *AIDS Care* 2017; 29: 1074–8.
41. Arjun BY, Unnikrishnan B, Ramapuram JT, Thapar R, Mithra P, Kumar N, et al. Factors influencing quality of life among people living with HIV in coastal South India. *J Int Assoc Provid AIDS Care* 2015; 16: 247–53.
42. Tesfay A, Gebremariam A, Gerbaba M, Abrha H. Gender differences in health related quality of life among people living with HIV on highly active antiretroviral therapy in Mekelle Town, Northern Ethiopia. *Biomed Res Int* 2015; 2015: 516369.
43. Tesemma AL, Abate MG, Abebo ZH, Madebo WE. Determinants of poor quality of life among adults living with HIV and enrolled in highly active antiretroviral therapy at public health facilities of Arba Minch Town Administration in Southern Ethiopia. *HIV AIDS* 2019; 11: 387–94.
44. de Munter JS, Tynelius P, Ahlström G, Rasmussen F. The bidirectional association between body weight and mobility disability: A population-based cohort. *Disabil Health J* 2016; 9: 632–7.
45. Mattias Norrbäck, Tynelius P, Ahlström G, Rasmussen F. The association of mobility disability and obesity with risk of unemployment in two cohorts from Sweden. *BMC Public Health* 2019; 19: 347.
46. Thein H-H, Butler T, Krahn M, Rawlinson W, Levy MH, Kaldor JM, et al. The effect of hepatitis C virus infection on health-related quality of life in prisoners. *J Urban Health* 2006; 83: 275–88.
47. Toghianifar N, Najafian J, Pooya A, Rabiei K, Eshtrati B, Anaraki J, et al. Association of smoking status with quality of life in a cross-sectional population-based sample of iranian adults: Isfahan healthy heart program. *Asia-Pacific J Public Health* 2012; 24: 786–94.
48. Kroshnick JA, Malhotra N, Mo CH, Bruera EF, Chang LC, Pasek J, et al. Perceptions of health risks of cigarette smoking: A new measure reveals widespread misunderstanding. *PLoS One* 2017; 12: 1–23.
49. Zinzi D Bailey, Slopen N, Albert M, Williams DR. Multidimensional religious involvement and tobacco smoking patterns over 9–10 years: A prospective study of middle-aged adults in the United States. *Soc Sci Med* 2015; 138: 128–35.
50. Ghouri N, Atcha M, Sheikh A. Influence of Islam on smoking among Muslims. *BMJ Clin Res* 2006; 332: 291.
51. Younossi ZM, Stepanova M, Omata M, Mizokami M, Walters M, Hunt S. Quality of life of Japanese patients with chronic hepatitis C treated with ledipasvir and sofosbuvir. *Medicine (Baltimore)* 2016; 95: e4243.
52. Younossi ZM, Stepanova M, Afdhal N, Kowdley K, Zeuzem S, Henry L, et al. Improvement of health-related quality of life and work productivity in chronic hepatitis C patients with early and advanced fibrosis treated with ledipasvir and sofosbuvir. *J Hepatol* 2015; 63: 337–45.
53. Ichikawa T, Miyaaki H, Miuma S, Taura N, Motoyoshi Y, Akahoshi H, et al. Hepatitis C virus-related symptoms, but not quality of life, were improved by treatment with direct-acting antivirals. *Hepatol Res* 2018; 48: 232–9.
54. Pereira L, Medeiros J De, Barreto M, Lima C, Maria M, Pires A, et al. Treatment with sofosbuvir and daclatasvir (with or without ribavirin) improves patient reported outcomes in hepatitis C. *Osong Public Heal Res Perspect* 2018; 9: 50–8.
55. Younossi ZM, Stepanova M, Esteban R, Jacobson I, Zeuzem S, Sulkowski M, et al. Superiority of interferon-free regimens for chronic hepatitis C. *Medicine (Baltimore)* 2017; 96: e5914.
56. Younossi ZM, Stepanova M, Feld J, Zeuzem S, Sulkowski M, Foster GR, et al. Sofosbuvir and velpatasvir combination improves patient-reported outcomes for patients with HCV infection, without or with compensated or decompensated cirrhosis. *Clin Gastroenterol Hepatol* 2017; 15: 421-430.e6.
57. Doyle JS, Grebely J, Spelman T, Alavi M, Matthews G, Thompson AJ, et al. Quality of life and social functioning during treatment of recent hepatitis C infection: A multi-centre prospective cohort. *PLoS One* 2016; 11: 1–16.

Figures

Figure 1

Flowchart of the study

Figure 2