The foot drawing method: Reliability of measuring foot length and outward rotation in children with clubfoot.

Evgenia Manousaki (✉ evgenia.manousaki@med.lu.se)

Lund University

Hanneke Andriesse

Lund University

Gunnar Hägglund

Lund University

Axel Ström

Skåne University Hospital

Anna-Clara Esbjörnsson

Lund University

Research Article

Keywords: Clubfoot, Foot length, Foot rotation, Reliability

Posted Date: February 1st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1288602/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The Ponseti method is the gold standard for clubfoot treatment. However, relapse and residual gait deviations are common, and follow-up until 7 years of age is recommended. We evaluated the reliability of the foot drawing method, a new instrument for the follow-up of clubfoot. The method uses drawings of the foot in the neutral position and external rotation to measure foot length and outward rotation.

Methods: Nineteen children aged 2.5–7 years who were treated with the Ponseti method for congenital clubfoot were included. Two raters made the drawings twice (D1 and D2). Each rater measured foot length, foot rotation, and foot–tibial rotation independently (D1). Later, the raters repeated the measurements (D2). Interrater reliability was assessed using the D1 from each rater. Intrarater reliability was assessed using the measurements from each rater's D1 and D2. Bland–Altman plots were used to visualize the limits of agreement (LoA). The mean, 95% confidence interval, and one standard deviation of the differences in all measurements were calculated.

Results: The mean differences between and within raters were: foot length <1 mm, foot rotation <1°, and foot–tibia rotation <2°, which indicated no systematic differences. The LoA for foot length were: 4.5 mm and 5.9 mm between raters for D1, –4.8 mm and 5.9 mm for rater 1 (D1–D2), and –5.1 mm and 5 mm for rater 2 (D1–D2). The LoA for foot rotation: were –12° and 10.6° between raters (D1), –8.4° and 6.6° for rater 1 (D1–D2), and –14° and 14.1° for rater 2 (D1–D2). The LoA for foot–tibia rotation were: –17.8° and 14.3° between raters (D1), –12° and 12.2° for rater 1 (D1–D2), and –12.7° and 13.6° for rater 2 (D1–D2).

Conclusions: The absence of systematic differences between and within raters, and LoA observed indicate that the foot drawing method is applicable in clinical practice and research. However, the results of the foot and foot–tibia rotation analyses imply that caution is needed when interpreting changes in foot rotation in feet with higher degrees of rotation.

Background

The Ponseti method is considered to be the gold standard for clubfoot treatment [1–3]. However, relapse and residual gait deviations are common, even after successful initial treatment [3–5], and children with clubfoot are recommended to undergo clinical follow-up at least until the age of 7 years [6]. Different measures are used to evaluate clubfoot status [7–13], but there is a lack of agreement about which measurements are the most appropriate [14].

Since 1995, we have performed a continuous follow-up of all children with clubfoot according to a standardized protocol [15]. This follow-up has included drawings on paper of the foot in a neutral position and externally rotated. The drawings are used to measure foot length (FL) and foot outward rotations, and provide a simple visual method for evaluating the development of FL and mobility. The procedure is named the foot drawing method (FDM).

A recent study of FL showed that feet with clubfoot normally grow regularly over time and that feet with a shorter length than expected are more susceptible to relapse [16]. To our knowledge, no studies have reported on the development of foot rotations in children treated for clubfoot. Foot abduction (called foot rotation from this point) is defined by Ponseti as the foot's ability to rotate outward and is a key correction parameter [2]. Casting-and-stretching techniques are intended to mobilize the clubfoot, and orthotic treatment is used to maintain mobility [2]. Foot rotation depends on the mobility of the tibiotalar and subtalar joints in combination with mid-and forefoot abduction. In addition, rotation in the knee and hip joints, as well as tibia and femur torsion may influence foot rotation. Regular assessment of the range of foot rotation may provide a simple method for detecting changes in mobility.

The aim of this study was to evaluate the intra- and interrater reliability of the FDM.

Patients And Methods

Patients

The study was approved by the National Ethical Review Authority (Dnr 2020-03008). Twenty children who were being treated for idiopathic congenital clubfoot were invited and all agreed to participate in the study. The children were 2.5–7 years of age and were scheduled to have their usual follow-up visits during October and November 2020. One child canceled the visit because of illness. All parents were verbally informed in advance about the procedures and provided written informed consent. All children were treated according to the Ponseti method [2]. Foot abduction orthosis (FAO) was used until at least the age of 4 years and longer if indicated [3].

Methods

The foot drawing method

Foot drawing

The child sits on a chair with the hips and knees in 90° of flexion, and the full foot in contact with the floor. The examiner sits in front of the child and places a paper on the floor parallel to the front of the chair. The examiner holds the foot lightly on the outside with one hand and moves the lower leg into flexion and extension in the knee. The other hand adjusts the thigh so that the tibial tuberosity is pointing straight forward. The foot is then placed down on the paper in its natural position [12, 16]. A line is drawn around the foot holding the pen vertically, and the medial and lateral malleoli are marked on the paper. Then, while holding the lower leg with one hand, the examiner rotates the foot into maximal outward rotation of the subtalar joint. The medial foot margin is then drawn again. Thereafter, the examiner fixes the distal femur. The foot is then rotated further outward with the other hand, which induces tibial–fibular outward rotation at the knee level. The medial foot margin is marked again (Figure 1). Both feet are drawn using this procedure, even the contralateral foot in unilateral cases.

Foot measurements

The FL and outward rotations are measured as follows.

Foot length: FL is measured as described by Manousaki et al. in 2021 [16] as the distance between two parallel lines, drawn distally and proximally on the footprint: The proximal line is perpendicular to an imaginary line passing through the middle of the hindfoot. The distal line is parallel to the proximal line, including the footprint (Figure 2).

Foot outward rotation (FR)

FR is defined as the angle between the medial foot margin (not including forefoot adduction or abduction) with the foot in maximal outward rotation as described above and a line drawn vertically to the long side of the paper (Figure 3). The angle is measured with a goniometer (Figure 3).

Foot and tibial–fibular outward rotation (FTR)

FTR is defined as the angle between the medial foot margin with the foot and lower leg in maximal outward rotation as described above and the line drawn vertically to the long side of the paper. The angle is measured with a goniometer (Figure 3).

Procedures

Two experienced examiners, rater 1 and rater 2, made drawings (Ds) of the children's feet twice (D1 and D2), independently of each other. A short break between each drawing was taken to allow the child to walk around. First, one rater made both drawings, and then, after a short break, the other rater repeated the procedure. The starting order of raters alternated for every new patient.

Each of the two raters scanned their drawings (D1 and D2) and measured FL, FR, and FTR directly on the copy of their first drawing (D1). Thereafter both the originals and copies were given to an independent person. After a period of 3–4 weeks, the raters received their D2 drawing, which was anonymized, and they repeated the measurements of FL, FR, and FTR. These were collected by an independent person and forwarded to another independent person for further analysis.

Statistical analysis

Power analysis was performed for both FL and foot rotations to estimate the number of feet needed to identify systematic differences. For the power analysis, we estimated a priori that, to detect a clinically significant change between assessments, systematic differences of 4 mm and assuming a standard deviation (SD) of 2 mm in FL and 5° (assuming SD 3°) in foot rotations would be acceptable. The measurements of foot rotation required more feet, at least seven, to investigate systematic differences with a power of 80% and $\alpha = 5$. To study the possible effects of age, we chose to have two age groups (4.5 years or younger and older than 4.5 years of age). Therefore, we decided to include at least 14 children but increased this further to account for dropouts.

The D1 from each rater was used to assess interrater reliability. The measurements from each rater's D1 and D2 were used to assess intrarater reliability. Bland–Altman plots were used to visualize the limits of agreement (LoA). The mean, 95% confidence interval (CI), and SD of the differences were calculated for all measurements.

Both feet, with and without clubfoot for children with unilateral involvement, were included in the analysis. Before including all feet, we performed a sensitivity analysis, in which we included unilateral clubfoot and randomly one clubfoot from each of the children with clubfoot in both feet. We also performed the analysis including all feet but excluded the contralateral unaffected foot in participants with unilateral clubfoot. The results from the two different approaches (one foot from each child vs one foot from those with unilateral clubfoot and two feet from those with bilateral clubfoot) were similar to the results from the analysis that included all feet. Hence, the results presented in this article include both feet from all included children.

All analyses were performed using R version 3.6.3 [17].

Results

Nineteen children (15 boys, 12 with unilateral clubfoot) aged 2.5–7 years were included. Eleven children were older than 4.5 years and eight children were 4.5 years or younger.

The average FR was greater for children aged ≤ 4.5 years (21° (SD 6°)) than for those >4.5 years (15° (SD 5°)) ($p = 0.001$). The average FTR was also greater for the younger children (52° (SD 13°)) than the older children (40° (SD 10°)) ($p = 0.005$). The average FR for the contralateral unaffected feet (all ages) was 19° (SD 3°), and the average FTR was 48° (SD 7°).

Increased intra- and interrater differences were observed with increasing degrees of rotation for FR and FTR. Neither age nor clubfoot laterality had an effect on these differences. Thus, we present the following results for the entire group without dividing the children into age groups.

The mean difference between and within raters was <1 mm for FL, $<1^\circ$ for FR, and $<2^\circ$ for FTR (Tables 1 and 2). These small differences did not differ statistically or clinically.

Foot length

The LoA for FL between raters for the first measurement (D1) were -4.5 mm to 5.9 mm (Table 1, Figure 4). The intrarater LoA between D1 and D2 were -4.8 mm to 5.9 mm for rater 1 and -5.1 mm to 5 mm for rater 2 (Table 2).

Table 1
Mean differences between each rater's first measurement D1 (interrater reliability)

Measurement	Mean (CI)	SD	LoA
Foot length (mm)	0.7 (−0.2; 1.6)	2.6	−4.5; 5.9
Foot rotation (°)	−0.7 (−2.6; 1.2)	5.8	−12; 10.6
Foot–tibia rotation (°)	−1.7 (−4.4; 0.7)	8.2	−17.8; 14.3
<i>CI: confidence interval, SD: standard deviation, LoA: limits of agreement</i>			

Table 2
Mean differences between first and second measurements for each rater

Measurement	Intrarater R1			Intrarater R2		
	Mean (CI)	SD	LoA	Mean (CI)	SD	LoA
Foot length (mm)	−0.6 (−1.5; 0.3)	2.7	−5.9; 4.8	0 (−0.8; 0.9)	2.6	−5; 5.1
Foot rotation (°)	−0.9 (−2.2; 0.4)	3.8	−8.4; 6.6	0.1 (−2.3; 2.4)	7.2	−14; 14.1
Foot–tibia rotation (°)	0.1 (−1.9; 2.1)	6.2	−12; 12.2	0.4 (−1.8; 2.7)	6.7	−12.7; 13.6
<i>R1: rater 1, R2: rater 2, CI: confidence interval, SD: standard deviation, LoA: limits of agreement</i>						

Foot outward rotation

The LoA for FR between raters' D1 were −12° to 10.6° (Table 1, Figure 5). The intrarater LoA were −6.6° to 8.4° for rater 1 and −14.1° to 14° for rater 2 (Table 2).

Foot and tibia–fibular outward rotation

The LoA for FTR between raters' D1 were −17.8° to 14.3° (Table 1, Figure 6). The intrarater LoA were −12° to 12.2° for rater 1 and −12.8° to 13.6° for rater 2 (Table 2).

Discussion

We evaluated the reliability of the FDM, a new instrument for follow-up of children with clubfoot. The systematic differences between raters were small (<1 mm for FL and <2° for FR and TR), which indicated that no significant bias was observed. The LoA was <6 mm for FL and <18° for FR and FTR. The intra- and interrater differences were more pronounced in feet with higher degrees of rotation.

The reliability of the FDM depends on the child's cooperation, ability to understand instructions, and attention span. Other factors that are important to consider when testing reliability are sample size, statistical method, assessment procedure, and experience of the assessors. The FDM comprises three procedures that all affect reliability: positioning of the foot on the paper, drawing around the foot, and measuring the parameters. When the FDM is used clinically, all procedures are completed during the visit. Thus, we chose not to divide the procedures in this reliability analysis.

Small clubfeet are associated with difficulties at the initial correction and a higher tendency to relapse [10, 11, 18]. In a previous study, we found that FL at the age of 2 years could be used as a prognostic tool because children with unilateral clubfoot with smaller FL compared with the contralateral foot at 2 years of age had a higher tendency to relapse and poorer motion quality at 7 years of age [16]. Therefore, it is important to use a reliable method to measure FL.

Different methods to measure FL have been reported as reliable [7, 9]. In our study, the small mean differences (<1 mm (SD <3 mm)) between measurements and between and within raters indicate that FL could be used in research to compare different treatment groups. The Bland–Altman plots for FL showed good intra- and interrater LoA (Table 1, Figure 4). Until the age of 7 years, feet with clubfeet grow about 1.2 cm per year [16]. The LoA observed in the Bland–Altman analysis indicate that changes >6 mm are clinically detectable. This implies that the FL could be useful for follow-up as the child grows.

To our knowledge, the longitudinal development of FR and FTR in children with clubfoot has not been investigated. Gupta et al. [19] showed that, in healthy children aged 0–8 years, foot abduction (external rotation of the foot, with flexed knee, simulating the Ponseti maneuver) ranged from 20° to 101° (mean 61° (SD 14°)). Outward rotation of the clubfoot is important during the initial correction with Ponseti casting. After the initial correction, the acquired outward rotation (i.e., abduction) is maintained with the FAO by positioning the feet in 60° of outward rotation. This outward rotation is a combination of mobility from different joint levels such as foot, ankle-tibia, knee, and even hip joint. Older children with clubfeet had significantly smaller FR and FTR angles. The use of FAO generally ends by the age of 4 years, and the lower outward rotations found in older children in this study reflect this pattern.

The estimated error in goniometer measurements used for clinical assessments range from 5° to 25° [20–23]. The variations depend on the body segment measured, procedure, and experience of the examiners [20–24]. Intrarater reliability is usually better than interrater reliability. The small mean differences for the FR and FTR measurements ($<2^\circ$ (SD 4°)) and the absence of systematic bias found in our study indicate that the method may be useful in research involving group comparisons. The LoA according to Bland–Altman analysis were relatively high: up to 14° for intrarater and 18° for interrater reliability. Nevertheless, the Bland–Altman plots showed that more pronounced differences were observed when the FR and FTR angles were high ($>20^\circ$ for FR and 45° for FTR, Figures 5 and 6). This implies the presence of proportional bias. Because most functional problems appear when the feet are becoming stiff (i.e., developing less rotation), this bias has little impact on the usefulness of the FDM to measure clinically important ranges of motion.

Several sources of error may have contributed to the differences observed, for example, the positioning of the foot and the force each rater used to rotate the foot. In a clinical follow-up, we suggest comparing drawings between visits because the starting position of the foot can affect the measurements.

Given that the measurements were continuous variables, we found it appropriate to use Bland–Altman plots and systematic differences for the evaluation. Bland–Altman plots give the opportunity to assess visually the differences between measurements, which we believe is valuable when evaluating a method for the first time.

We included both feet from each child, even the unaffected foot for children with unilateral clubfoot. Because the feet from the same child are related and the aim of our study was to analyze the reliability of the method and not the outcome of a treatment, we felt it was important to include both feet. In the clinical setting, the appearance and mobility of both feet, including the unaffected in children with unilateral clubfoot, are drawn and analyzed. Both raters had long experience with the FDM and with clubfoot treatment, both of which can affect reliability [24]. By standardizing the method as described previously, we expect that less experienced colleagues will be able to use the method. The reliability of the method with inexperienced examiners is yet to be determined.

Conclusions

We conclude that the FDM is applicable in clinical practice and research in the context for which it is aimed to be used. By comparing measurements during follow-up, developments in foot growth and foot mobility can be monitored. However, the results from the foot and foot–tibia rotation analyses imply that caution is needed when interpreting changes in foot rotation in feet with higher degrees of rotation.

Abbreviations

D1: drawing 1

D2: drawing 2

FL: foot length

FDM: foot drawing method

FR: Foot outward rotation

FTR: Foot and tibial–fibular outward rotation

SD: standard deviation

LoA: limits of agreement

CI: confidence interval

Declarations

Ethics approval and consent to participate

All methods were carried out in accordance with relevant guidelines and regulations. All experimental protocols were approved by the Swedish Ethical Review Authority (Dnr 2020-03008). Written informed consent was obtained from all subjects and their legal guardians

Consent for publication

Not Applicable.

Availability of data and materials

All data generated or analyzed during this study are included in this published article and its supplementary information files.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was supported by grants from: the Faculty of Medicine, Lund University, Region Skåne, Region Kronoberg, Södra sjukvårdsregionen, Promobilia Stiftelsen, Greta and Johan Kocks stiftelser and” Stiftelsen för bistånd åt rörelsehindrade i Skåne”. The foundations had no role in the design or conduct of this study.

Authors' contributions

EM: concept and design of the study, collection of data, interpretation of data, drafting of the manuscript.

HA: concept and design of the study, collection of data, interpretation of data, critical review of the manuscript.

GH: concept and design of the study, collection of data, interpretation of data, critical review of the manuscript.

AS: design of the study, statistical analysis, interpretation of data, critical review of the manuscript.

ACE: concept and design of the study, collection of data, interpretation of data, critical review of the manuscript.

All authors have read and approved the manuscript.

Acknowledgements

Not Applicable.

References

1. Zionts LE, Sangiorgio SN, Ebrahimzadeh E, Morcuende JA. The current management of idiopathic clubfoot revisited: results of a survey of the POSNA membership. *J Pediatr Orthop*. 2012;32(5):515–20.
2. Ponseti IV. *Congenital clubfoot: fundamentals of treatment*. Oxford University Press; 1996.
3. Ponseti IV, Smoley EN. The classic: congenital club foot: the results of treatment. *Clin Orthop Relat Res*. 2009;467(5):1133–45.
4. Thomas HM, Sangiorgio SN, Ebrahimzadeh E, Zionts LE. Relapse rates in patients with clubfoot treated using the Ponseti method increase with time: a systematic review. *JBJS Rev*. 2019;7(5):e6.
5. Tuinsma ABM, Vanwanseele B, van Oorschot L, Kars HJJ, Grin L, Reijman M, et al. Gait kinetics in children with clubfeet treated surgically or with the Ponseti method: a meta-analysis. *Gait Posture*. 2018;66:94–100.
6. van Praag VM, Lysenko M, Harvey B, Yankanah R, Wright JG. Casting is effective for recurrence following Ponseti treatment of clubfoot. *J Bone Joint Surg Am*. 2018;100(12):1001–8.
7. Gelfer Y, Durham S, Daly K, Ewins D. Intraobserver reliability of static measures in the normally developing infant foot and clubfoot. *J Pediatr Orthop B*. 2009;18(5):214–9.
8. Muller S, Carlsohn A, Muller J, Baur H, Mayer F. Static and dynamic foot characteristics in children aged 1–13 years: a cross-sectional study. *Gait Posture*. 2012;35(3):389–94.
9. Kesemenli CC, Kapukaya A, Subasi M, Necmioglu S, Arslan H, Ozbag D, et al. Anthropometric study of patients treated for clubfoot. *J Pediatr Orthop*. 2003;23(4):498–502.
10. Hemo Y, Yavor A, Gigi R, Wientroub S. The significance of foot length at the initiation of the Ponseti method: a prospective study. *J Child Orthop*. 2019;13(3):252–7.
11. Kumar K. The role of footprints in the management of clubfeet. *Clin Orthop Relat Res*. 1979(140):32–6.
12. Hazlewood ME, Simmons AN, Johnson WT, Richardson AM, van der Linden ML, Hillman SJ, et al. The Footprint method to assess transmalleolar axis. *Gait Posture*. 2007;25(4):597–603.
13. Wallace JJ, White HD, Augsburg SF, Talwalkar VR, Muchow RD, Iwinski HJ, et al. Foot size asymmetry following Ponseti treatment versus comprehensive surgical releases for unilateral clubfeet. *J Pediatr Orthop B*. 2019;28(2):153–8.
14. Gelfer Y, Hughes KP, Fontalis A, Wientroub S, Eastwood DM. A systematic review of reported outcomes following Ponseti correction of idiopathic club foot. *Bone Jt Open*. 2020;1(8):457–64.
15. Andriesse, H. Follow-up of children with congenital clubfoot. Development of a new evaluation instrument. Lund, Sweden: Institution for Health Sciences, Division of Physiotherapy, Lund University Department of Orthopaedics, Lund University Hospital; 2007. Available at: <https://portal.research.lu.se/ws/files/4917201/633402.pdf>. Accessed January 23, 2022.

16. Manousaki E, Esbjornsson AC, Hagglund G, Andriesse H. Development of foot length in children with congenital clubfoot up to 7 years of age: a prospective follow-up study. *BMC Musculoskelet Disord*. 2021;22(1):487.
17. R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
18. Ponseti IV. Relapsing clubfoot: causes, prevention, and treatment. *Iowa Orthop J*. 2002;22:55–6.
19. Gupta P, Mittal N, Jindal N, Verma P, Sharma M. A study of normal foot abduction across various age groups in children. *Int Orthop*. 2017;41(11):2365–9.
20. Rome K. Ankle joint dorsiflexion measurement studies. A review of the literature. *J Am Podiatr Med Assoc*. 1996;86(5):205–11.
21. Menadue C, Raymond J, Kilbreath SL, Refshauge KM, Adams R. Reliability of two goniometric methods of measuring active inversion and eversion range of motion at the ankle. *BMC Musculoskelet Disord*. 2006;7:60.
22. Hayes K, Walton JR, Szomor ZR, Murrell GA. Reliability of five methods for assessing shoulder range of motion. *Aust J Physiother*. 2001;47(4):289–94.
23. McDowell BC, Hewitt V, Nurse A, Weston T, Baker R. The variability of goniometric measurements in ambulatory children with spastic cerebral palsy. *Gait Posture*. 2000;12(2):114–21.
24. Flynn JM, Donohoe M, Mackenzie WG. An independent assessment of two clubfoot-classification systems. *J Pediatr Orthop*. 1998;18(3):323–7.

Figures

Figure 1

A line is drawn around each foot with the pen kept vertical. Two more lines are added. The first line marks the medial foot edge after the foot has been rotated in maximal external rotation in relation to the tibia (red line). The second marks the medial foot edge after the foot has been rotated further outward on the knee level by adding maximal tibial rotation (blue line).

Figure 2

Foot length (blue line) is defined as the distance between two lines. The proximal line (green) is perpendicular to the imaginary line that passes from the middle of the hindfoot (grey line). The distal line (red) is parallel to the proximal line and includes the whole footprint.

Figure 3

Foot rotation (yellow) and foot and tibia rotation (FTR, purple) are measured with a goniometer between each foot margin (red: FR, blue: FTR) and a line (green) drawn vertically to the long side of the paper.

Figure 4

Interrater Bland–Altman plot for foot length. The y-axis shows the difference between the first (D1) measurements of the two assessors. The x-axis shows the mean foot length.

Figure 5

Interrater Bland–Altman plot for foot rotation. The y-axis shows the difference between the first measurement (D1) of the two assessors. The x-axis shows the mean foot rotation.

Figure 6

Interrater Bland–Altman plot for foot and foot-tibia rotation. The y-axis shows the difference between the first (D1) measurement of the two assessors. The x-axis shows the mean foot-tibia rotation.