

# Targeted Interventions to Prevent Chronic Low Back Pain in High Risk Patients: Development and Delivery of a Pragmatic Training Course of Psychologically Informed Physical Therapy for the TARGET Trial

Jason Beneciuk (✉ [beneciuk@phhp.ufl.edu](mailto:beneciuk@phhp.ufl.edu))

University of Florida <https://orcid.org/0000-0002-5888-3630>

Steven Z. George

Duke University, Duke Clinical Research Institute

Carol Greco

University of Pittsburgh, School of Medicine

Michael Schneider

University of Pittsburgh, School of Health and Rehabilitation Sciences

Stephen T. Wegener

Johns Hopkins University School of Medicine

Robert Saper

Boston Medical Center

Anthony Delitto

University of Pittsburgh

---

## Methodology

**Keywords:** Psychologically informed physical therapy, post-professional education, quality improvement

**Posted Date:** March 13th, 2019

**DOI:** <https://doi.org/10.21203/rs.2.129/v3>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on May 6th, 2019. See the published version at <https://doi.org/10.1186/s13063-019-3350-3>.

# Abstract

**Background:** Low back pain (LBP) is a public health concern because it is highly prevalent and the leading cause of disability worldwide. Psychologically Informed Physical Therapy (PIPT) is a secondary prevention approach that first aims to identify individuals at high risk for transitioning to chronicity and then provides tailored treatment to reduce that risk. Training models that are feasible to implement with acceptable training quality are needed to improve scalability for widespread implementation of PIPT. This manuscript describes the PIPT training program that was developed for training physical therapists providing PIPT in the TARGET Trial. **Methods:** The PIPT training program was developed, tested and modified using an iterative process. Content development consisted of stakeholder engagement, beta testing, modification of training and confirmation of final course objectives. Methods of delivery consisted of a website that included brief online educational modules followed by a live 8-hour workshop that included video-based mock case scenarios and case-based role playing. Attitudes, beliefs and confidence in implementing PIPT principles were assessed before and immediately after training to measure training quality and impact. **Results:** Early stakeholder engagement and beta testing indicated the need for increased emphasis on experiential learning opportunities and patient-centered communication training. Booster training varied extensively across TARGET sites with involvement of 'clinician champions' providing brief follow-up sessions identified as best practice. Favorable post training changes in physical therapist attitudes and beliefs toward biopsychosocial treatment orientation and increased confidence in implementing PIPT principles were observed. **Conclusions:** PIPT training for provider participation in the TARGET Trial was feasible to deliver. Course content was acceptable to physical therapists and resulted in improved beliefs and confidence in applying PIPT skills during clinical practice. Ongoing consultation and site-based continuing education were methods by which specific TARGET sites maintained or augmented PIPT skill training, however implementing ongoing training was challenging in general. Due to the pragmatic nature of the TARGET trial, it was not possible to directly measure the effect of PIPT training on treatment fidelity which was a limitation of our approach. Trial registration: ClinicalTrials.gov Identifier: NCT02647658

## Background

Low back pain (LBP) is an imperative public health concern because it is highly prevalent and the leading cause of disability worldwide (1). Global prevalence of LBP has increased by 17.3% from 2005 to 2015 and continues to be a leading cause of global years lived with disability since 1990 (1, 2). Although most individuals will rapidly recover (3), many continue to experience ongoing or chronic pain, accounting for a large proportion of the burden associated with LBP (4, 5). The risk factors for chronic pain are complex and multifactorial including psychological and behavioral features such as pain catastrophizing (6), fear-avoidance beliefs (7), and maladaptive pain coping (8). These factors can be addressed via cognitive-behavioral therapy, typically offered by psychologists and other behavioral healthcare providers, however seldom addressed through initial treatment delivered by primary care providers or physical therapists.

Prevention of chronic pain has emerged as a high priority in the United States, with specific emphasis placed on identifying at-risk patients and offering nonpharmacologic treatments as ‘front-line’ options (9-11). One promising strategy is Psychologically Informed Physical Therapy (PIPT), a secondary prevention approach for LBP that first aims to identify individuals at high risk for transitioning to chronicity and then provides tailored treatment by merging impairment-focused physical therapy with cognitive behavioral therapy methods as needed to reduce that risk (12, 13). The primary goal of PIPT is prevention of future LBP-associated disability via: 1) identification of individuals with elevated pain-associated distress and at high risk for transitioning from acute to chronic LBP, and 2) providing targeted treatment aimed at ameliorating psychological factors linked to prolonging disability in conjunction with traditional impairment-based physical therapy. PIPT optimizes variables predictive of persistent pain and is therefore consistent with a top priority of the Federal Pain Research Strategy (i.e., formalizing individualized treatment recommendations based on risk factors) (14).

Recent systematic review findings indicate psychologically-based treatments can enhance physical therapy interventions for patients at high risk for prolonged LBP related disability while emphasizing the role of risk stratification for acute LBP and specifically recommending the need for determining reproducibility in the United States and optimizing implementation in clinical practice (15, 16). One successful risk stratification approach for LBP uses the 9-item STarT Back Tool (17) to screen for modifiable prognostic factors, determine patient risk for developing persistent LBP related disability and uses that information to match patients with different care pathways. Physical therapists have played an integral role as treatment providers of PIPT in previous studies that found significant improvements in patient LBP disability and quality of life outcomes, while also resulting in less time off work and greater healthcare cost savings when compared to standard care (18, 19). However, the training to deliver these and other psychologically-based interventions has ranged from 2 to 9 days (20-25), which may not be scalable for widespread implementation in many United States health care systems.

There is a need for pragmatic PIPT training models that are feasible to implement with acceptable training quality and impact. These training models should also recognize clinician preferences for continuing education and be scalable to be responsive to national priorities for pain research and practice (14). A study by Beneciuk and George (26) provides ‘proof of concept’ to support the idea that one-day pragmatic PIPT training models are feasible and can result in improved physical therapist attitudes and beliefs about managing back pain using psychological principles. In contrast, minimal changes were observed for clinicians that did not participate in the stratified care and PIPT training module. Furthermore, therapists who received PIPT training had better 4-week patient clinical outcomes for pain intensity and self-reported disability compared to therapists who were not trained (26). There is a timely opportunity for striking a balance between lengthy comprehensive training programs and pragmatic single day courses. The latter are capable of providing a general overview, making such an approach feasible for widespread participation, without drastically compromising training quality and impact. Such pragmatic approaches that encourage efficient treatment delivery strategies may provide a viable option for enhancing clinical practice paradigms aimed at improving patient outcomes through widespread implementation (14).

This manuscript describes the PIPT training that was developed and delivered to prepare physical therapists for providing treatment in the TARGET Trial [<http://www.targettrial.pitt.edu>]. TARGET is a large, pragmatic cluster-randomized clinical trial of patients seen in the primary care setting with acute LBP determined by the STarT Back Tool to be at high risk for persistent LBP related disability. In this manuscript we will first provide a brief overview of the TARGET Trial design and goals to place the purpose of the training in context. Second, we will describe development and delivery of the PIPT training to prepare physical therapists at TARGET Trial clinical treatment sites and describe strategies used to enhance learning. Third, we will provide data on training quality and impact. Finally, we will discuss preliminary lessons learned and provide suggestions for future pragmatically delivered PIPT training initiatives.

## Methods

### TARGET Trial Overview

The Targeted Interventions to Prevent Chronic Low Back Pain in High Risk Patients (TARGET Trial) is a multisite, pragmatic cluster-randomized clinical trial studying patients with acute LBP who seek care from a primary care physician and are at high risk for persistent disability. The study is designed to assess if prompt referral of patients to physical therapists with PIPT training reduces the rate of progression to chronic LBP 6-months later (primary trial outcome) and improves back-related function as compared to guideline-based primary care management. Secondary outcomes include additional healthcare resource utilization. The TARGET Trial enrolls patients from primary care clinics across multiple health systems in five geographic regions across the United States (Pittsburgh, PA; Boston, MA; Baltimore, MD; Salt Lake City, UT; and Charleston, SC) with a total planned sample size (n = 1860) that exceeds or is similar to previously completed studies (18, 19, 27). The TARGET Trial is funded by the Patient Centered Outcomes Research Institute and was prospectively registered with ClinicalTrials.gov (NCT02647658) on January 4, 2016. **Additional trial details can be found at ClinicalTrials.gov registry site.**

### PIPT Training Program

Considering the pragmatic study design, several factors were considered when developing the overall structure of the PIPT training program. First, there was the need for developing a multidisciplinary training team consisting of individuals representing physical therapy and clinical psychology. Second, there was the challenge of addressing the feasibility barrier of providing training to physical therapists from different healthcare systems located across five diverse geographical regions in the United States. Third, there was the importance of identifying facilitators for physical therapists to attend the PIPT training (e.g., cost, continuing education credit, and time commitment). Finally, the potential impact of discussions between physical therapists was not as concerning based on the cluster-randomized clinical trial study design, which decreased likelihood for contamination across clinical sites. Prior to trial initiation, the PIPT training program was developed, tested and modified using an iterative process to

enhance optimal effects during study training that were intended to be implemented during routine clinical practice. (Figure 1)

## **Training Background**

Development of the PIPT training program was guided by previous protocols that have tested risk stratification approaches for LBP using the STarT Back Tool (21, 24). In addition, members of the TARGET intervention team (JMB and SZG) provided PIPT training for a small-scale feasibility study, training providers within a single healthcare system (26). Key developers of the TARGET Trial PIPT training program consisted of physical therapy, clinical psychology and chiropractic providers.

## **Content Development**

PIPT training program content development consisted of stakeholder engagement, beta testing, modification of training and confirmation of final course objectives. Each of these stages is described in greater detail below.

### *1. Stakeholder engagement*

Prior to providing formal PIPT training, feedback and support from key stakeholders was obtained. We initially targeted a single healthcare system (University of Pittsburgh Medical Center, Centers for Rehab Services) to provide an introduction and overview of the TARGET Trial and PIPT training program. Our initial strategy included a formal task force meeting that consisted of representation from TARGET Trial investigators, healthcare system executives, outpatient clinical education, senior management and clinicians. The key objective of this meeting was to prepare a task force of physical therapists within UPMC Centers for Rehab Services to become '*clinical champions*' for implementing a standard biopsychosocial model for patients with musculoskeletal pain conditions.

### *2. Beta testing*

Similar training programs have undergone beta testing to obtain critical feedback for guiding modification and prior to providing actual training in preparation for trial participation (21, 24). Beta testing of the PIPT training program was provided for two separate cohorts of licensed physical therapists over a two-month period in Pittsburgh, PA (n = 40) and at a professional conference in National Harbor, MD (n = 15). Participant feedback was collected through discussion and brief surveys with key suggestions considered during subsequent modifications to the PIPT training program. One important outcome of the beta testing provided by course participants was the need for blended learning opportunities (i.e., strategic use of combined web-based and in-person content with interactive activities to enhance clinician learning), which led to strategies for developing additional and previously recommended (28) delivery platforms (e.g., PIPT website).

### *3. Modification of training*

An iterative process of development for the PIPT training program was used, that incorporated feedback from beta course participants, standardized self-assessments and intervention team discussions. This led to restructuring of the live workshop to include: 1) several interactive breakout sessions, designed for the participants to gain initial experience implementing PIPT skills; 2) video-based mock clinical scenarios suitable for live course learning opportunities; 3) development of online video modules for training maintenance; and 4) increased time allotment and enhanced patient-centered communication training methods dedicated to address barriers to clinical practice integration. These modifications are consistent with enhancements provided following pilot testing of other previous training packages (24).

#### 4. *Final course objectives*

Following beta testing, feedback from participants and modification of training content and methods, final course learning objectives were finalized by the investigator team (**Table 1**). Collectively, the overall objectives of the PIPT training program were to provide participants with a summary of evidence and clinical skills necessary to support implementing PIPT principles into routine clinical practice for patients identified as being at high risk for transitioning from acute to chronic LBP. Methods of delivery (described below) were intended to promote a blended learning (i.e., flipped classroom) environment with instructional strategies guided by specific learning objectives (29, 30). Flipped classroom pedagogy principles included as part of the PIPT training program included: flexible learning, improved preparation for the live workshop, self-reflection, peer-learning and enhanced rapport with instructors.

#### 5. *Final course content*

A description of the final course content is provided in **Table 2**. Course content was broadly described as either '*Overview*' – providing rationale and supporting data for the PIPT approach, or '*PIPT Management*' – providing specific principles or skills with demonstration and practice. '*PIPT Management*' content consisted of: 1) patient-centered communication; 2) pain coping skills; 3) patient education; 4) activity based intervention; 5) impairment based intervention; and 6) treatment monitoring components (**Figure 2**). The course content was provided in sequential order for all training sessions.

### **Methods of Delivery**

Consistent with the pragmatic study design of the TARGET Trial, the training was designed to be easily replicated in clinical settings to augment delivery of PIPT implementation. Flipped classroom instructional methods were integrated to enhance preparation for the live workshop, with the intention that each site would determine specific methods favorable for that specific geographical region and health delivery system.

#### 1. *PIPT website*

The TARGET Trial website [<http://www.targettrial.pitt.edu/>] provided an alternative delivery platform for content resources that included an overview of the TARGET Trial and provider resources (including key recommended readings). Course participants registered for formal training courses were directed to a

separate PIPT website [<http://www.rstce.pitt.edu/pipt/>] that provided a course overview, learning objectives, education modules and additional educational resources for patients. Twelve brief pre-course online educational modules were developed to provide necessary foundational information required to optimize the experiential nature of the 1-day live workshop and promote a flipped classroom learning model. These online educational modules were designed to be viewed in sequence, with each module ranging from 8 to 22 minutes in duration (total viewing time: 150 minutes). Links to voiceover PowerPoint presentations, electronic handouts and audio files were provided for each module to offer course participants different learning platform options. Physical therapists had the opportunity to obtain 2.5 hours of continuing education credit after viewing all the online modules. Viewing online video modules was highly recommended prior to attending the live workshop, however we did not monitor everyone who accessed the website. Physical therapists seeking continuing education credit were required to complete a quiz after each online module, thereby providing a method to monitor online attendance certificate of completion eligibility. In situations where physical therapists were not seeking continuing education credit (as not required in all states), quizzes were not administered and there was no method to objectively monitor online attendance.

## *2. Live workshop*

All sites participating in the TARGET Trial had practicing physical therapists who would deliver PIPT at a local clinic. The sites were required to host live workshops as part of their site participation requirement. However, our ability to monitor which specific physical therapists attended and where they were practicing in a given site was limited as registration processes varied across TARGET site locations. Live one-day workshops were provided by at least one physical therapist and clinical psychologist up to three times at each of the five TARGET sites throughout the United States (Pittsburgh, PA; Boston, MA; Baltimore, MD; Salt Lake City, UT; and Charleston, SC). Combinations of teaching methods (**Table 2**) were used during each 8-hour workshop including: PowerPoint presentations; video based mock case scenarios depicting appropriate and inappropriate communication styles; and case-based role playing.

To enhance clinical skills training, we utilized several structured teaching and learning strategies, including: 1) instructor led teaching on specific clinical skills; 2) instructor led case-based role playing with mock patient; 3) course participant led case-based role playing (i.e., breakout sessions) where smaller groups of 2 to 4 participants each assumed different stakeholder roles (e.g., patient, clinician, and observer) for a variety of clinical scenarios; and 4) class discussion to provide individual learning experience perspectives. To further enhance learning, participants were encouraged to demonstrate newly acquired clinical skills that were evaluated by instructors and other participants for real-time feedback. This case-based role playing was used to develop clinical skills involving self-reflection, motivational interviewing, pain coping skills and activity-based interventions. Barriers and facilitators to implementing PIPT components (**Figure 2**) during routine clinical practice were addressed throughout the live workshops.

## *3. PIPT course materials*

Each physical therapist was provided with course materials at the live one-day workshop that could be referenced afterwards. These materials consisted of workshop content, including specific descriptions and scenarios pertaining to PIPT interventions such as patient-centered communication, pain coping skills, patient education, activity-based intervention, impairment-based intervention and treatment monitoring components.

### **Quality Improvement Strategies to Enhance and Assess Quality and Impact of Provider Training**

Establishing treatment fidelity to insure the reliability and validity of behavioral interventions has been identified as a major challenge (31, 32). The National Institutes of Health Behavior Change Consortium (NIHBCC) has developed and recently updated a treatment fidelity framework consisting of five domains (i.e., study design, training of providers, treatment delivery, treatment receipt, and treatment enactment) (31, 32). Consistent with the pragmatic nature of the TARGET Trial, balance between feasibility and obtaining comprehensive fidelity assessments was considered (33). The PIPT program was designed to promote treatment fidelity by providing quality training that impacted key provider factors and could be replicated. Thus, we incorporated **quality improvement** strategies (PIPT treatment checklist and booster training) and measures (physical therapist attitudes, beliefs and confidence, described in greater detail below) to enhance treatment quality and impact of training.

#### ***Strategies to enhance treatment quality***

##### **PIPT treatment checklist**

To promote treatment fidelity, physical therapists were trained to indicate specific PIPT treatment content delivered during patient care by completing self-report checklists (31-34). Strategies for administering checklists varied across geographical regions ranging from traditional hardcopy methods to direct entry into the electronic medical record.

##### **Booster Training**

Due to the pragmatic nature of the trial, the amount and frequency of follow-up communication and training maintenance was different in each geographical region. As a result, booster or refresher training varied extensively, with course instructors and site coordinators offering several options following the live workshop. All course participants were provided options for continued remote communication with instructors, and were encouraged to submit follow-up questions and testimonials to promote a flipped classroom learning environment. One implementation process that may be described as possible 'best practice' within this trial consisted of one-hour follow-up sessions provided at several clinical sites in the Salt Lake City, UT region that were focused on improving specific PIPT skills that physical therapists found difficult to implement. For example, prior to the course, participants indicated difficulty with initiating PIPT interventions, specifically related to patient-provider communication. Cognitive reassurance (engaging the patient in education) was thoroughly discussed in group settings and motivational interviewing strategies were revisited through case-based role playing using specific patient

scenarios that were led by site mentors. Another strategy in Pittsburgh, PA, Baltimore, MD, and Boston, MA regions consisted of offering brief 45-60 minute webinars where course participants were asked to submit topical questions, with instructors and clinical champions providing strategies to overcome barriers to successful implementation.

### ***Measures to assess training impact***

Physical therapists that attended the live workshop were administered questionnaires to assess attitudes, beliefs and confidence (described below). Course instructors did not provide any instruction or advice for how to respond to individual questionnaire items.

#### **Attitudes and beliefs**

Physical therapist attitudes and beliefs about biomedical and biopsychosocial treatment orientations were assessed before training, immediately after completion of training, and 4-months later using the Pain Attitudes and Beliefs Scale for Physical Therapists (PABS-PT) (35, 36). The PABS-PT consists of 19 items about treatment orientation that are rated using a 6-point Likert scale ranging from “totally disagree” to “totally agree”. The PABS-PT biomedical scale (10 items) has a potential score range from 10 to 60 and the PABS-PT biopsychosocial scale (9-items) has a potential score range from 9 to 54 with higher scores indicating increased biomedical or biopsychosocial treatment orientation depending upon the respective scale. TARGET site leaders initiated a request for follow-up assessment four months after training through email that directed course attendees to a remote website containing an electronic version of the PABS-PT with reminders being sent 2 weeks later.

#### **Confidence in PIPT skill application**

Physical therapist confidence in implementing PIPT principles was assessed before training and upon completion of training (same day). Specifically, participants were asked to “rate your level of confidence with implementing psychological informed principles during clinical practice” using an 11-point scale (range 0 to 10) with “0” indicating “no confidence” and “10” indicating “extreme confidence”.

### **Quality Improvement Evaluation of PIPT Training Program**

A total of four hundred and seventy one outpatient physical therapists attended at least one live workshop and completed pre-training questionnaires. Means and standard deviations were calculated for available continuous variables (i.e., age, years in practice, PABS-PT and confidence scores) for the entire study sample by each TARGET site (Table 3). Paired samples t-tests were used to assess for pre to post-course changes in the scores derived from the PABS-PT (biomedical and behavioral scale) and confidence in applying PIPT questionnaires. Then, three separate multiple regression models were used to explore the contribution of TARGET site location and viewing pre-course online video modules as predictors of post-course scores. For each model, Block 1 accounted for pre-training PABS-PT or confidence scores depending upon the outcome of interest; Block 2 added TARGET site location to Block 1; and Block 3 added self-report response to question about viewing pre-course video modules (Yes or No)

to Block 2. Finally, one-way analysis of variance with Bonferroni post-hoc procedures was used to compare physical therapist PABS-PT (biomedical and behavioral scale) and confidence residualized change scores between TARGET site locations to evaluate for training replicability.

## Result

### Attitudes and beliefs

Follow-up assessments upon completion of training were performed for 91.5% (431/471) of course participants. PABS-PT biomedical scale scores decreased from 31.1 [SD = 6.8] to 25.0 [SD = 7.1] ( $P < .001$ ) and behavioral scale scores increased from 36.8 [SD = 4.8] to 41.4 [SD = 5.2] ( $P < .001$ ). Regression models explained 38% and 17% of the variance in post-course PABS-PT biomedical and behavioral scale scores, respectively. Pre-course PABS-PT biomedical ( $\beta = .62$ ,  $P < .001$ ) and behavioral ( $\beta = .41$ ,  $P < .001$ ) scale scores were the strongest predictors of post-course PABS-PT biomedical and behavioral scale scores, respectively. TARGET site location only added 1% additional variability to prediction of post-course PABS-PT biomedical ( $\beta = .09$ ,  $P = .022$ ) and behavioral ( $\beta = .12$ ,  $P = .018$ ) scale scores. Viewing pre-course online modules did not significantly explain any additional variability in post-course PABS-PT biomedical or behavioral scale scores ( $P > 0.05$ ). After adjustment for pre-course scores by each site location, there were no differences in PABS-PT biomedical ( $P = .140$ ) or behavioral ( $P = .095$ ) scale change scores between TARGET site locations. A total of 134 (28.4%) course participants completed a web-based version of the PABS-PT 4-months after training with biomedical ( $25.4 \pm 7.9$ ) and behavioral ( $40.9 \pm 4.4$ ) scale scores observed, however this data was de-identified which does not allow us to determine if sustained scores were maintained over longer duration of time.

### Confidence in PIPT skill application

Follow-up assessments of confidence upon completion of training (same day) were performed for 96.2% (453/471) of course participants. We were not able to capture any additional assessments of confidence at 4-months. Confidence in the ability to implement PIPT principles increased from 4.8 [SD = 2.2] to 7.3 [SD = 1.9] ( $P < .001$ ). The regression model explained 27% of the variance in post-course confidence scores. Pre-course confidence scores ( $\beta = .46$ ,  $P < .001$ ) was the strongest predictor explaining 20% of the variance in post-course confidence scores. TARGET site location ( $\beta = .21$ ,  $P < .001$ ) added 5% additional variability to the prediction of post-course confidence scores. Viewing pre-course online modules ( $\beta = .12$ ,  $P = .003$ ) explained 2% additional variability in post-course confidence scores. Greater post-course confidence scores were observed for those who viewed modules when compared to those who did not (mean = 7.7 [SD = 1.3] versus 7.2 [SD = 1.7],  $P = .004$ ). After adjustment for pre-course scores, differences in confidence change scores between TARGET site locations were observed ( $P < .001$ ). Specifically, Boston, MA reported less improvement in confidence compared to Pittsburgh, PA (mean difference = 1.2, 95% CI = 0.1-2.3,  $P = .029$ ), Salt Lake City, UT (mean difference = 1.6, 95% CI = 0.5-2.7,  $P = .001$ ), Baltimore MD (mean difference = 1.7, 95% CI = 0.6-2.7,  $P < .001$ ) and Charleston, SC (mean difference = 2.4, 95% CI = 1.3-3.4,  $P < .001$ ). Additional findings indicated Boston, MA participants were younger in age

(32.4 years [SD = 8.1]) and had less experience in clinical practice [6.1 years [SD = 7.0]] compared to all other TARGET site locations ( $P < .001$ ) which may have influenced observed changes in confidence (Table 3).

## Discussion

The overall objectives of the training course were to provide physical therapists with a summary of evidence and clinical skills necessary to support the implementation of PIPT principles into clinical practice for patients identified as being high risk for transitioning from acute to chronic LBP. Our experiences have provided several important “lessons learned” that can be used to guide future study of PIPT implementation for long-term sustainability (37).

### *Emphasis on Experiential Learning*

During beta testing, course participants provided consistent feedback about the need to reduce didactics and increase the amount of time devoted to experiential learning experiences. Therefore, PIPT treatment concepts were introduced using video and live mock case scenarios that transitioned into small group practice sessions during each 8-hour workshop. These teaching principles were also utilized in certain regions following initial training as a component of booster training and could perhaps be described as ‘best practice’ within the TARGET Trial.

### *PIPT Clinician Champions*

Several strategies to enhance routine application of PIPT principles following the live course and during active patient enrollment periods (i.e., booster training) were planned during program development stages; however due to the pragmatic nature of the trial and geographical distribution of health systems, implementing these efforts was associated with considerable heterogeneity. For example, clinician-generated case reports that were intended to be the focus of dynamic learning communities were conceptualized as being a virtuous strategy. However, engaging clinicians to be accountable for active learning initiatives was a difficult process and only resulted in a small number of case examples; potentially due to busy, high volume clinical practices. Therefore, future implementation efforts should focus on strategies to identify and incentivize clinical champions within a health system or small region for leading subsequent active learning initiatives (e.g., webinars, formal mentoring opportunities) following initial training. Optimally, these individuals should demonstrate special interest and skill in PIPT and could be valuable resources for circumstances where continued on-site interaction with primary trainers is not feasible. Personal communication with physical therapists that received PIPT training for a previous smaller scale study and provided similar treatment indicated the need for additional follow-up opportunities to address barriers in clinical practice following the training course (26).

### *Need for Specialized Training*

Despite recent recommendations for increased delivery of psychological-based treatment (9, 10) and enthusiasm for risk stratification approaches to LBP management (14, 15) there are challenges to successful wide spread implementation. For example, a potent barrier to successfully delivering psychological-based treatment is the vital need for additional specific post-professional training (12, 38-41). This dilemma is particularly relevant to health care providers where biomedical or impairment-based perspectives have been predominantly emphasized during entry-level education and clinical practice, thereby resulting in clinicians that are not confident or are unprepared in delivering psychological-based interventions (38, 39, 42). For example, our findings indicated fewer years in clinical practice was associated with less improvement in confidence after attending the live PIPT workshop, potentially suggesting that less experienced physical therapists (i.e., new graduates) may not be adequately prepared to successfully implement PIPT strategies with patients or require additional training. Consequently, gaining additional specific knowledge, problem-solving skills and practical experience through formal mentoring opportunities that incorporate booster training is a vital necessity for beneficial shifts in clinical practice paradigms to occur. However, many PIPT or cognitive behavioral treatment approaches require specialized time-intensive training, which may not be feasible for all clinicians and may perhaps present a significant barrier to successful widespread future clinical implementation. Providing single-day overview courses that are followed by structured mentorship experiences over an extended period of time may perhaps provide a viable model for future PIPT training programs.

### ***Patient-Centered Communication***

In our experience, physical therapists' initial perception was that implementing PIPT strategies in practice would be challenging. Training in patient-centered communication appears to be an important component for integrating PIPT into routine clinical practice. As previously indicated, increased time and enhanced training methods were dedicated to patient-centered communication during the preliminary phases of our training program based on early feedback from participants involving barriers to clinical practice integration. Similar challenges and enhancements to communication content and delivery methods have been acknowledged during development of other PIPT based training programs (24, 43). Clinician challenges to providing patient-centered and biopsychosocial-oriented communication for patients with LBP is common (38-40, 42, 44), which is not surprising considering the lack of content dedicated to this topic during entry-level physical therapy training programs. Specifically, our patient-centered communication interventions were enhanced early during trial training stages by integrating motivational interviewing skill development with significant time permitted for: 1) instructor led case-based role playing with mock patients; 2) course participant led case-based role playing (i.e., breakout sessions) where smaller groups of 2 to 4 participants each assumed different stakeholder roles (e.g., patient, clinician, and observer) for a variety of clinical scenarios; and 3) class discussion to provide individual learning experience perspectives. Future implementation strategies should strongly consider providing direct examples that combine patient-centered communication skills and other PIPT interventions such as graded activity or graded exposure to optimize treatment efficiency. This approach may be particularly beneficial as most patient expectations for physical therapy treatment may not be aligned with PIPT-based principles.

## ***Provider Training***

Training quality and impact assessment could have been further enhanced by including formal assessment of skill acquisition, standardized methods to prevent skills drift and providing accommodations to diverse learning styles (32). Formal assessment to determine if learning objectives were attained would have also potentially strengthened the likelihood of subsequent PIPT implementation during trial participation, however this additional assessment may have also reduced the pragmatic nature of the training program and limited physical therapist engagement. PIPT training program evaluation was not the primary aim of the TARGET Trial, and was completed under the auspices of quality improvement, not educational research. Therefore, we were limited on the amount of data collected from physical therapists who attended the live workshop and the ability to conduct formal outcome assessments.

In our opinion, the PABS-PT and confidence measures provided one viable option for assessment of the impact of PIPT training on clinician attitudes and beliefs. Updated strategies for assessment provided by the National Institutes of Health's Behavioral Change Consortium have indicated the importance of ensuring "buy-in" for treatment (32), with previous studies having considered clinician attitudes and beliefs to guide development of targeted treatment and training packages (21). We observed favorable treatment orientation shifts from predominantly biomedical to biopsychosocial, following attendance at the live workshop which is consistent with previous studies where similar duration of PIPT training was delivered (26, 45). Our viewpoint is that this favorable change in treatment orientation and confidence is a practical indicator that the PIPT training has potential for altering therapists' attitudes, beliefs and confidence. However, we acknowledge that additional assessment through direct observation or formal assessment of competency (which may not be feasible in clinical settings) would be needed to determine if the PIPT training resulted in behavioral change for the provider and if these changes are associated with improvements in patient outcomes.

These findings highlight the need to consider clinicians' attitudes and beliefs regarding treatment orientation when introducing new treatment approaches such as PIPT if they are to be adopted in clinical practice (46). Systematic review findings indicate health care providers with predominantly biomedical treatment orientations are more likely to suggest limited work and physical activity, and are less likely to adhere to clinical practice guidelines that emphasize activation for patients with LBP (46, 47). Changing provider beliefs is critical if we are to optimize care for patients who widely believe their persistent LBP results from anatomical or biomechanical causes (48). Therefore, future studies should consider long-term assessment of PIPT training quality and impact to determine if favorable changes in clinicians' attitudes and beliefs are sustained over time and how - or if - these changes influence patient outcomes.

## ***Long-Term Follow-Up to Sustain Changes***

Considering our strategies for continual engagement with physical therapists following in-person training may have been less than optimal, we suggest several strategies to enhance this process in an effort to sustain beneficial changes in attitudes and beliefs over long-term periods. First, sustained

communication between instructors, site leaders and course participants may enhance PIPT training maintenance opportunities. Second, continuing education credit or organizational quality improvement initiatives may provide clinicians with incentive for participating in maintenance activities. Third, system-level recognition for cohorts achieving specific maintenance participation rates may provide clinicians with a sense of personal satisfaction. Finally, we used the PABS-PT to assess clinician attitudes and beliefs about treatment orientations and a single question to assess confidence in implementing PIPT principles, therefore ongoing assessment and feedback may assist with skill maintenance.

### ***Suggestions to Increase Scalability***

Based on our experiences, future pragmatically delivered PIPT training initiatives should consider providing single-day overview courses that are followed by structured mentorship experiences (i.e., either remotely on-line or in-person) over an extended period of time. This may provide a viable continuing education model for future PIPT training programs that can be led by local clinical champions, moderated by course instructors, and provided using remote learning platforms (e.g., webinars, discussion boards). Based on our findings, we also suggest that future PIPT training programs be tailored to participant characteristics and clinical experience. Specifically, content knowledge, attitudes, beliefs and confidence need to be considered as course content and delivery methods may need to be modified. Providing clinicians with incentives to assume these leadership roles will be important as participation and time spent developing formal case reports for 'real world' learning experiences (as an example) will require personal commitment and most likely dedication of time outside of clinical practice.

## **Conclusions**

The PIPT training in the TARGET Trial, which consisted of online educational modules followed by a 1-day live discussion and skills-based training, was feasible to deliver as part of a large, pragmatic trial. The final format for the course was acceptable to physical therapists and resulted in improvement in biopsychosocial attitudes and beliefs and confidence in applying PIPT skills during clinical practice. Ongoing consultation and site-based continuing education were methods by which specific TARGET sites maintained or augmented the PIPT skill training, however ongoing training was challenging for most TARGET sites in general. Treatment fidelity was not measured directly, which was a limitation to our training approach and will continue to be a struggle for future pragmatic trials that are charged with delivering interventions as part of routine clinical practice.

## **Declarations**

### **Ethics approval and consent to participate**

This specific study describing development and delivery of the TARGET Trial PIPT training program met the University of Pittsburgh Institutional Review Board definition for quality improvement and not research, therefore was not reviewed.

## **Consent for publication**

Not applicable.

## **Availability of data and material**

Not applicable.

## **Competing interests**

The authors declare that they have no competing interests.

## **Funding and disclaimer**

Work reported in this publication was funded through a Patient-Centered Outcomes Research Institute (PCORI) Award (PCS-1402-10867). The statements in this publication are solely the responsibility of the authors and do not necessarily represent the views of PCORI, its Board of Governors, or Methodology Committee.

## **Authors' contributions**

All authors read and approved the final manuscript.

(JMB, SZG, CG, MS, and STW) made substantial contributions to development, enhancement, and delivery of PIPT program. (JMB, SZG, MS, CG, and STW) have been involved in drafting the manuscript and revising it critically for important intellectual content. (RS and AD) reviewed and provided final approval of the version to be published. Each author participated sufficiently in order to take public responsibility for appropriate portions of the content; and agreed to be accountable for all aspects of the work.

## **Acknowledgements**

Mallorey Smith (Brooks Rehabilitation, Jacksonville, FL), Gerard Brennan, Steven Hunter and Kate Minick (Intermountain Healthcare, Salt Lake City, UT), Karthik Hariharan, Joel Stevans, Tara Hankin, Brian Caricato, Linda Hough (University of Pittsburgh, Pittsburgh, PA), Michael Freidman, Patti Ephraim, Michael Albert, LaPricia Lewis Boyer (Johns Hopkins, Baltimore, MD), Chris Joyce (MGH Institute of Health Professions, Boston, MA), David Morrisette, Kit Simpson (Medical University of South Carolina, Charleston, SC).

## **References**

1. Global Burden of Disease 2015 Disability-Adjusted Life-Years (DALYs) and Healthy Life Expectancy (HALE) Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990-2015: a systematic analysis for the

- Global Burden of Disease Study 2015. *Lancet* (London, England). 2016;388(10053):1603-58. Epub 2016/10/14.
2. Global Burden of Disease 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* (London, England). 2016;388(10053):1545-602. Epub 2016/10/14.
  3. da C Menezes Costa L, Maher CG, Hancock MJ, McAuley JH, Herbert RD, Costa LO. The prognosis of acute and persistent low-back pain: a meta-analysis. *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2012;184(11):E613-24. Epub 2012/05/16.
  4. Gore M, Sadosky A, Stacey BR, Tai KS, Leslie D. The burden of chronic low back pain: clinical comorbidities, treatment patterns, and health care costs in usual care settings. *Spine*. 2012;37(11):E668-77. Epub 2011/12/08.
  5. Global Burden of Disease 2013 Disability-Adjusted Life-Years (DALYs) and Healthy Life Expectancy (HALE) Collaborators. Global, regional, and national disability-adjusted life years (DALYs) for 306 diseases and injuries and healthy life expectancy (HALE) for 188 countries, 1990-2013: quantifying the epidemiological transition. *Lancet* (London, England). 2015;386(10009):2145-91. Epub 2015/09/01.
  6. Wertli MM, Eugster R, Held U, Steurer J, Kofmehl R, Weiser S. Catastrophizing-a prognostic factor for outcome in patients with low back pain: a systematic review. *The spine journal : official journal of the North American Spine Society*. 2014;14(11):2639-57. Epub 2014/03/13.
  7. Wertli MM, Rasmussen-Barr E, Weiser S, Bachmann LM, Brunner F. The role of fear avoidance beliefs as a prognostic factor for outcome in patients with nonspecific low back pain: a systematic review. *The spine journal : official journal of the North American Spine Society*. 2014;14(5):816-36 e4. Epub 2014/01/15.
  8. Chou R, Shekelle P. Will this patient develop persistent disabling low back pain? *Jama*. 2010;303(13):1295-302. Epub 2010/04/08.
  9. Department of Health and Human Services. National Pain Strategy: A Comprehensive Population Health Strategy for Pain. 2016; Available from: [https://iprcc.nih.gov/National\\_Pain\\_Strategy/NPS\\_Main.htm](https://iprcc.nih.gov/National_Pain_Strategy/NPS_Main.htm). Accessed June 21, 2017.
  10. Qaseem A, Wilt TJ, McLean RM, Forciea MA, Clinical Guidelines Committee of the American College of P. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians. *Annals of internal medicine*. 2017;166(7):514-30. Epub 2017/02/14.

11. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain—United States, 2016. *Jama*. 2016;315(15):1624-45. Epub 2016/03/16.
12. Main CJ, George SZ. Psychologically informed practice for management of low back pain: future directions in practice and research. *Physical therapy*. 2011;91(5):820-4. Epub 2011/04/01.
13. Keefe FJ, Main CJ, George SZ. Advancing Psychologically Informed Practice for Patients With Persistent Musculoskeletal Pain: Promise, Pitfalls, and Solutions. *Physical therapy*. 2018;98(5):398-407. Epub 2018/04/19.
14. Gatchel RJ, Reuben DB, Dagenais S, Turk DC, Chou R, Hershey AD, et al. Research Agenda for the Prevention of Pain and Its Impact: Report of the Work Group on the Prevention of Acute and Chronic Pain of the Federal Pain Research Strategy. *The journal of pain : official journal of the American Pain Society*. 2018;19(8):837-51. Epub 2018/03/27.
15. Chou R, Deyo R, Friedly J, Skelly A, Hashimoto R, Weimer M, et al. *Noninvasive Treatments for Low Back Pain*. Rockville MD: Agency for Healthcare Research and Quality; 2016.
16. Foster NE, Hill JC, O'Sullivan P, Hancock M. Stratified models of care. *Best practice & research Clinical rheumatology*. 2013;27(5):649-61. Epub 2013/12/10.
17. Hill JC, Dunn KM, Lewis M, Mullis R, Main CJ, Foster NE, et al. A primary care back pain screening tool: identifying patient subgroups for initial treatment. *Arthritis and rheumatism*. 2008;59(5):632-41. Epub 2008/04/29.
18. Hill JC, Whitehurst DG, Lewis M, Bryan S, Dunn KM, Foster NE, et al. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomised controlled trial. *Lancet (London, England)*. 2011;378(9802):1560-71. Epub 2011/10/04.
19. Foster NE, Mullis R, Hill JC, Lewis M, Whitehurst DG, Doyle C, et al. Effect of stratified care for low back pain in family practice (IMPACT Back): a prospective population-based sequential comparison. *Annals of family medicine*. 2014;12(2):102-11. Epub 2014/03/13.
20. Hansen Z, Daykin A, Lamb SE. A cognitive-behavioural programme for the management of low back pain in primary care: a description and justification of the intervention used in the Back Skills Training Trial (BeST; ISRCTN 54717854). *Physiotherapy*. 2010;96(2):87-94. Epub 2010/04/28.
21. Sowden G, Hill JC, Konstantinou K, Khanna M, Main CJ, Salmon P, et al. Targeted treatment in primary care for low back pain: the treatment system and clinical training programmes used in the IMPACT Back study (ISRCTN 55174281). *Family practice*. 2012;29(1):50-62. Epub 2011/06/29.
22. Bryant C, Lewis P, Bennell KL, Ahamed Y, Crough D, Jull GA, et al. Can physical therapists deliver a pain coping skills program? An examination of training processes and outcomes. *Physical therapy*. 2014;94(10):1443-54. Epub 2014/06/07.

23. Overmeer T, Boersma K, Denison E, Linton SJ. Does teaching physical therapists to deliver a biopsychosocial treatment program result in better patient outcomes? A randomized controlled trial. *Physical therapy*. 2011;91(5):804-19. Epub 2011/04/01.
24. Main CJ, Sowden G, Hill JC, Watson PJ, Hay EM. Integrating physical and psychological approaches to treatment in low back pain: the development and content of the STarT Back trial's 'high-risk' intervention (StarT Back; ISRCTN 37113406). *Physiotherapy*. 2012;98(2):110-6. Epub 2012/04/18.
25. Cherkin D, Balderson B, Brewer G, Cook A, Estlin KT, Evers SC, et al. Evaluation of a risk-stratification strategy to improve primary care for low back pain: the MATCH cluster randomized trial protocol. *BMC musculoskeletal disorders*. 2016;17(1):361. Epub 2016/08/25.
26. Beneciuk JM, George SZ. Pragmatic Implementation of a Stratified Primary Care Model for Low Back Pain Management in Outpatient Physical Therapy Settings: Two-Phase, Sequential Preliminary Study. *Physical therapy*. 2015;95(8):1120-34. Epub 2015/04/11.
27. Cherkin D, Balderson B, Wellman R, Hsu C, Sherman KJ, Evers SC, et al. Effect of Low Back Pain Risk-Stratification Strategy on Patient Outcomes and Care Processes: the MATCH Randomized Trial in Primary Care. *Journal of general internal medicine*. 2018;33(8):1324-36. Epub 2018/05/24.
28. Forsetlund L, Bjorndal A, Rashidian A, Jamtvedt G, O'Brien MA, Wolf F, et al. Continuing education meetings and workshops: effects on professional practice and health care outcomes. *The Cochrane database of systematic reviews*. 2009(2):CD003030. Epub 2009/04/17.
29. Tang B, Coret A, Qureshi A, Barron H, Ayala AP, Law M. Online Lectures in Undergraduate Medical Education: Scoping Review. *JMIR medical education*. 2018;4(1):e11. Epub 2018/04/11.
30. Hew KF, Lo CK. Flipped classroom improves student learning in health professions education: a meta-analysis. *BMC medical education*. 2018;18(1):38. Epub 2018/03/17.
31. Bellg AJ, Borrelli B, Resnick B, Hecht J, Minicucci DS, Ory M, et al. Enhancing treatment fidelity in health behavior change studies: best practices and recommendations from the NIH Behavior Change Consortium. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association*. 2004;23(5):443-51. Epub 2004/09/16.
32. Borrelli B. The assessment, monitoring, and enhancement of treatment fidelity in public health clinical trials. *Journal of public health dentistry*. 2011;71 Suppl 1:S52-63. Epub 2011/06/10.
33. Toomey E, Hardeman W. Addressing Intervention Fidelity Within Physical Therapy Research and Clinical Practice. *The Journal of orthopaedic and sports physical therapy*. 2017;47(12):895-8. Epub 2017/12/02.
34. Toomey E, Matthews J, Guerin S, Hurley DA. Development of a Feasible Implementation Fidelity Protocol Within a Complex Physical Therapy-Led Self-Management Intervention. *Physical therapy*.

2016;96(8):1287-98. Epub 2016/03/05.

35. Houben RM, Ostelo RW, Vlaeyen JW, Wolters PM, Peters M, Stomp-van den Berg SG. Health care providers' orientations towards common low back pain predict perceived harmfulness of physical activities and recommendations regarding return to normal activity. *European journal of pain* (London, England). 2005;9(2):173-83. Epub 2005/03/02.
36. Mutsaers JH, Peters R, Pool-Goudzwaard AL, Koes BW, Verhagen AP. Psychometric properties of the Pain Attitudes and Beliefs Scale for Physiotherapists: a systematic review. *Manual therapy*. 2012;17(3):213-8. Epub 2012/01/27.
37. Joyce C, Schneider M, Stevans JM, Beneciuk JM. Improving Physical Therapy Pain Care, Quality, and Cost Through Effectiveness-Implementation Research. *Physical therapy*. 2018;98(5):447-56. Epub 2018/04/19.
38. Alexanders J, Anderson A, Henderson S. Musculoskeletal physiotherapists' use of psychological interventions: a systematic review of therapists' perceptions and practice. *Physiotherapy*. 2015;101(2):95-102. Epub 2014/12/02.
39. Synnott A, O'Keefe M, Bunzli S, Dankaerts W, O'Sullivan P, O'Sullivan K. Physiotherapists may stigmatise or feel unprepared to treat people with low back pain and psychosocial factors that influence recovery: a systematic review. *Journal of physiotherapy*. 2015;61(2):68-76. Epub 2015/03/31.
40. Driver C, Kean B, Oprescu F, Lovell GP. Knowledge, behaviors, attitudes and beliefs of physiotherapists towards the use of psychological interventions in physiotherapy practice: a systematic review. *Disability and rehabilitation*. 2017;39(22):2237-49. Epub 2016/09/17.
41. Hall A, Richmond H, Copsey B, Hansen Z, Williamson E, Jones G, et al. Physiotherapist-delivered cognitive-behavioural interventions are effective for low back pain, but can they be replicated in clinical practice? A systematic review. *Disability and rehabilitation*. 2018;40(1):1-9. Epub 2016/11/23.
42. Foster NE, Delitto A. Embedding psychosocial perspectives within clinical management of low back pain: integration of psychosocially informed management principles into physical therapist practice—challenges and opportunities. *Physical therapy*. 2011;91(5):790-803. Epub 2011/04/01.
43. Johnson RE, Jones GT, Wiles NJ, Chaddock C, Potter RG, Roberts C, et al. Active exercise, education, and cognitive behavioral therapy for persistent disabling low back pain: a randomized controlled trial. *Spine*. 2007;32(15):1578-85. Epub 2007/07/11.
44. Jeffrey JE, Foster NE. A qualitative investigation of physical therapists' experiences and feelings of managing patients with nonspecific low back pain. *Physical therapy*. 2012;92(2):266-78. Epub 2011/12/17.

45. Jacobs CM, Guildford BJ, Travers W, Davies M, McCracken LM. Brief psychologically informed physiotherapy training is associated with changes in physiotherapists' attitudes and beliefs towards working with people with chronic pain. *British journal of pain*. 2016;10(1):38-45. Epub 2016/08/24.
46. Gardner T, Refshauge K, Smith L, McAuley J, Hubscher M, Goodall S. Physiotherapists' beliefs and attitudes influence clinical practice in chronic low back pain: a systematic review of quantitative and qualitative studies. *Journal of physiotherapy*. 2017;63(3):132-43. Epub 2017/06/29.
47. Darlow B, Fullen BM, Dean S, Hurley DA, Baxter GD, Dowell A. The association between health care professional attitudes and beliefs and the attitudes and beliefs, clinical management, and outcomes of patients with low back pain: a systematic review. *European journal of pain (London, England)*. 2012;16(1):3-17. Epub 2011/07/02.
48. Setchell J, Costa N, Ferreira M, Makovey J, Nielsen M, Hodges PW. Individuals' explanations for their persistent or recurrent low back pain: a cross-sectional survey. *BMC musculoskeletal disorders*. 2017;18(1):466. Epub 2017/11/19.

## Tables

**Table 1. Psychologically Informed Physical Therapy Training Course Learning Objectives.**

PIPT = Psychologically Informed Physical Therapy; CBT = Cognitive-Behavioral Therapy; ICF = International Classification of Functioning, Disability, and Health.

1. Summarize relationships between pain neuroscience, pain models, and the development and maintenance of chronic low back pain.

---

2. Identify patients at high risk for transitioning from acute to chronic low back pain.

---

3. Apply targeted treatment for patients at high risk for transitioning from acute to chronic low back pain.

---

4. Differentiate key principles and application between graded activity and graded exposure.

---

5. Understand primary assumptions of CBT and specific skills associated with CBT based interventions.

---

6. Recognize effective communication skills and be able to implement as a key component to PIPT.

---

7. Review the Low Back Pain Clinical Practice Guidelines from the Orthopaedic Section of the American Physical Therapy Association to become familiar with: 1) ICF-based classifications; 2) symptoms; 3) impairments; and 4) suggested intervention strategies.

---

8. Be able to implement PIPT practice principles for patients with low back pain.

---

**Table 2. PIPT Training Course Content and Methods of Delivery.**

## Methods of Delivery

	Approximate time allotment dedicated during live workshop	PowerPoint presentations and instructor lead group discussion	Video-based mock case scenarios	Case-based role playing (breakout sessions)
<b>Overview</b>				
Pain Science Update	30 minutes	X		
PIPT Overview	30 minutes	X		
Risk Stratification		X		
Targeted Treatment		X		
Cognitive Behavioral Therapy	30 minutes	X		
Self-reflection	45 minutes	X	X	X
<b>PIPT Management</b>				
Patient-Centered Communication	1 hour, 45 minutes			
Active Listening		X	X	
Motivational Interviewing		X	X	X
Goal-setting		X	X	
Pain Coping Skills	1 hour, 15 minutes			
Physiologic Relaxation		X		X
Imagery		X		X
Replacing cognitive distortions		X	X	X
Patient Education	15 minutes	X		
Activity Based	60 minutes			
Graded Exercise		X	X	X

Graded Exposure		X		X		X
Impairment Based	30 minutes					
Clinical Practice Guidelines		X				
Treatment Monitoring	30 minutes	X				
Challenges and Opportunities	30 minutes	X		X		

PIPT = Psychologically Informed Physical Therapy

**Table 3. PIPT Course Participant Characteristics (n = 471).**

	Total sample (n = 471)	Pittsburgh, PA (n = 77)	Boston, MA (n = 61)	Salt Lake City, UT (n = 80)	Baltimore, MD (n = 111)	Charleston, SC (n = 142)	P-Value*
Age (years)	38.1 (11.0)	40.3 (11.2)	32.4 (8.1)	39.3 (10.3)	36.9 (11.2)	39.5 (11.4)	<.001
Years in practice	11.4 (10.6)	14.6 (11.5)	6.1 (7.0)	11.0 (10.4)	10.9 (11.4)	12.6 (10.2)	<.001
PABS-PT biomedical (pre-training)	31.0 (6.8)	30.3 (6.5)	30.4 (7.6)	28.2 (6.8)	31.0 (6.8)	33.2 (6.0)	<.001
PABS-PT biomedical (post-training)	25.2 (7.2)	26.0 (7.0)	24.3 (7.6)	23.1 (6.8)	25.2 (7.1)	26.2 (7.4)	.032
PABS-PT behavioral (pre-training)	36.9 (3.7)	36.6 (3.2)	36.9 (4.2)	38.3 (3.8)	37.1 (3.3)	36.1 (3.7)	.001
PABS-PT behavioral (post-training)	41.3 (4.2)	40.1 (3.8)	41.7 (5.2)	41.7 (4.3)	41.7 (4.4)	41.2 (3.5)	.067
Confidence (pre-training)	4.8 (2.2)	4.8 (2.1)	4.6 (2.3)	4.9 (2.2)	5.1 (2.3)	4.5 (2.2)	.292
Confidence (post-training)	7.3 (1.9)	7.0 (1.7)	5.6 (3.1)	7.4 (1.8)	7.8 (1.2)	7.7 (1.4)	<.001

(\* ) One-way analysis of variance to compare between TARGET site locations.

PIPT = Psychologically Informed Physical Therapy; PABS-PT = Pain Attitudes and Beliefs Scale for Physical Therapists.

## Figures



**Figure 1**

PIPT training program iterative process for development, testing, and modification.




Figure 2

Overview of key PIPT treatment components for high risk patients in TARGET Trial.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.docx](#)