The KIDSCREEN-27 scale: translation and validation study of the Slovenian version

Leona Cilar Budler (✉ leona.cilar1@um.si)

University of Maribor

Majda Pajnikhar

University of Maribor

Ulrike Ravens-Sieberer

University Medical Center Hamburg-Eppendorf

Owen Barr

University of Ulster

Gregor Stiglic

University of Maribor

Research Article

Keywords: Social Support, Adolescent, Psychometrics, Factor Analysis, Quality of Life

Posted Date: January 31st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1290683/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: There are many methods available for measuring social support and quality of life (QoL) of adolescents, of these, the KIDSCREEN tools are most widely used. Thus, we aimed to translate and validate the KIDSCREEN-27 scale for the usage among adolescents aged between 10 and 19 years old in Slovenia.

Methods: A cross-sectional study was conducted among 2 852 adolescents in primary and secondary school from November 2019 to January 2020 in Slovenia. 6-steps method of validation was used to test psychometric properties of the KIDSCREEN-27 scale. We checked descriptive statistics, performed a Mokken scale analysis, parametric item response theory, factor analysis, classical test theory and total (sub)scale scores.

Results: All five subscales of the KIDSCREEN-27 formed a unidimensional scale with good homogeneity and reliability. The confirmatory factor analysis showed poor fit in user model versus baseline model metrics ($CFI = 0.847$; $TLI = 0.862$) and good fit in root mean square error ($RMSEA = 0.072$; $p(\chi^2) < 0.001$). A scale reliability was calculated using Cronbach's α (0.93), beta (0.86), G6 (0.95) and omega (0.93).

Conclusions: The questionnaire showed good psychometric properties and can be used among adolescents in Slovenia to find out about their quality of life.

Background

Adolescence is the time between the ages of 10 and 19 when individuals have specific health and developmental needs [1]. The transition from childhood to adulthood often presents a challenging period with decline in emotional and mental well-being [2]. Thus, factors that influence emotional development of adolescents, such as personality changes, physical growth, and social and cognitive developments, must be taken into account [3]. Mental disorders often appear in this time, influencing later life. Investments in adolescents' mental well-being will bring great benefits in the following years and decades [4]. Adolescence provides an opportunity to develop positive mental health and wellbeing, and potentially reduce the costs of in health care caused by mental health disorders in adulthood [5].

Mental well-being compromises positive effects, evaluations of life, and absence of negative effects [6]. Many young people are facing everyday stress due to various stressors. Guo et al. [7] found that factors correlated to mental health are gender, physical activity, family economy, siblings, stress, satisfaction of self-appearance, quality of sleep, social trust, willingness to learn, as well as teachers and parent support. Adolescents spend many hours a day in school; thus, the school climate is of utmost importance for establishing mental well-being. Lester and Cross [8] found that factors like feeling safe, having a good connection to school and receiving peer support are protective factors of mental well-being, while connectedness to teachers is protective factor of mental well-being over the transition from primary to secondary school. Yet, Arabiat et al. [9] in their research in Jordan found that the parents' support is more important than school, and peer support is important for maintaining well-being in adolescence.

Social support is derived from family members, friends, and community, and is closely related to student's quality of life [10]. Social support is a protective factor to stress and contribute to one's well-being [11]. There is evidence that social support perception differs among gender, where female students receive more support from friends than from peers, family, and teachers. Male students receive less support from all sources [12]. Williams et al. [13] stated that social support requires the existence of social relationships with their structure, strength and type. Greater social support is linked to better health related quality of life among adolescents [14]. Moreover, lower social support for adolescents may contribute to emotional distress and internet addiction [15].

There are many methods available for measuring social support and quality of life (QoL) of adolescents, of these, the KIDSCREEN tools are most widely used [16; 17]. The KIDSCREEN-27 is a particularly useful instrument because it is easy to understand and use and because of the available volume of data for further comparisons. As the scale is widely used, it enables international comparison [18]. The validation is a rigorous process and crucial for research in the field of health, social, and behavioural sciences. Scale evaluation can be onerous and is often unfamiliar to researchers because of the lack of graduate training on scale development and validation [19]. The COSMIN taxonomy suggested that for assessing measurement properties the following items must be tested: reliability, construct and content validity, reliability, measurement error and responsiveness [20]. Thus, factor analysis (structural validity) and Cronbach α (internal consistency) are not enough to perform psychometric analysis which will provide meaningful results [21]. Recently, advanced, and more user-friendly software -R has been developed that offers numerous packages for psychometric testing [22]. Researchers now have a program that ensures transparent and reproducible data. Yet, for beginners, writing script in R can be complicated and take much time, thus available scripts such as script by Dima [21] are highly welcome.

Aim

The aim of this study was the translation and validation of the KIDSCREEN-27 questionnaire which was used for measuring social support and quality of life of the school-aged adolescents in Slovenia following the 6-step psychometrics protocol in R.

Methods

Design

We conducted a cross-sectional study among adolescents from November 2019 to January 2020 in 16 primary and 9 secondary schools in Slovenia.

The instrument

The KIDSCREEN questionnaire is available in three versions; the original long version consists of 52 items covering ten dimensions of QoL, a 27-item version covering 5 dimensions of QoL, and a 10-item index version [23]. The KIDSCREEN-27 measures five Rasch scaled dimensions: 1) physical well-being, 2) psychological well-being, 3) autonomy and parent relation, 4) peers and social support, and 5) school environments. The questionnaire was funded by the European Commission and developed within a European project "Screening and Promotion for Health-related Quality of Life in Children and Adolescents – A European Public Health Perspective" [23; 24; 25]. Each item was scored on a 5-point Likert scale ranging from 1 meaning "not at all" to 5 meaning "very much". Certain items (1, 9, 10 and 11) were reversed when scoring the questionnaire. Reverse scoring means that the numerical scoring scale runs in the opposite direction. The total KIDSCREEN score was calculated by summing up all the responses. Higher scores of KIDSCREEN indicated better QoL and social support. Before using the questionnaire, the prior permission to use was sought from the authors.

Sample

According to the Statistical Office of the Republic of Slovenia [26] and the Ministry of Education, Science and Sport of the Republic of Slovenia [27] there were 454 primary and 182 secondary schools in Slovenia at the time of planning this project. We used random sampling based on which the expected sample included all available students from 22 primary schools (5% of all elementary schools) and 12 secondary schools (5% of all secondary schools) in Slovenia. Random sampling was performed using online Research Randomizer site. The required sample was calculated on the basis of data from the Statistical Office of the Republic of Slovenia [28], where it is evident that there were 176 898 enrolled students in elementary school and 74 021 in secondary school in school year 2016/2017. The envisaged final number of participants involved was determined by the size of the total population of students, degree of confidence and margin of error [29], which amounted to 384 students. The sample size was increased to avoid the risk of attrition and drop out during the study. Students who are under 10 years old and older than 19 years and those who are not included in the education system were excluded. Survey questionnaires were distributed to 3 860 students in primary and 3 107 students from secondary schools who consent to participation in the classroom where the classes took place.

The study was conducted in 16 primary and 9 secondary schools in different regions of Slovenia. Out of 6 967 distributed questionnaires a total of 2 972 were completed and returned which represents a response rate of 42.66%. A total of 1489 (50.10%) adolescents from the primary and 1483 (49.90%) from the secondary schools participated in the survey. Research involved primary school (from the 6th to the 9th grade) and secondary school students (from the 1st to the 4th grade). They were between 11 and 19 years old. Other characteristics of the sample are shown in the Table 1.

Table 1: Sample Characteristics

	PS (<i>n</i> = 1489)	SS (<i>n</i> = 1483)
	<i>M</i> (<i>SD</i>)	<i>M</i> (<i>SD</i>)
Age	12.2 (1.5)	16.4 (1.3)
Gender	<i>n</i> (%)	<i>n</i> (%)
Female	768 (51.6%)	911 (66.8%)
Male	721 (48.4%)	452 (33.2%)
School year	<i>n</i> (%)	<i>n</i> (%)
1 st	0 (0.0%)	400 (29.3%)
2 nd	0 (0.0%)	381 (28%)
3 rd	0 (0.0%)	252 (18.5%)
4 th	0 (0.0%)	311 (22.8%)
5 th	264 (17.7%)	19 (1.4%)
6 th	305 (20.5%)	0 (0.0%)
7 th	266 (17.9%)	0 (0.0%)
8 th	277 (18.6%)	0 (0.0%)
9 th	377 (25.3%)	0 (0.0%)
Living environment		
With mother	1343 (90.7%)	1369 (91.9%)
With father	1204 (81.4%)	1205 (80.9%)
With sister/brother	1093 (73.9%)	1045 (70.2%)
With grandmother	316 (21.4%)	329 (22.1%)
With grandfather	214 (14.5%)	198 (13.3%)
Other	64 (4.3%)	59 (4.0%)

PS = primary school students; SS = secondary school students; *n* = number of students; *M* = Mean; *SD* = standard deviation

The mean age among students in primary school was 12.2 (*SD* = 1.5) and 16.4 (*SD* = 1.3) among students in secondary school. Primary and secondary school students most frequently answered that they live with mother (PS: 1343, 90.7%; SS: 1369, 91.9%) or father (PS: 1204, 81.4%; SS: 1205, 80.9%). Students who answered "other" (PS: 48, 3.2%; SS: 55, 3.7%) described that they live either with godfather/godmother, great-grandfather/ great-grandmother, uncle/aunt, nephew, cousin, guardian, foster father/foster mother, stepfather/stepmother, or alone.

Procedure

Data collection took place in the second half of 2019. We used a validated questionnaire KIDSCREEN-27 [30] to collect information on the perception of adolescents on support from family, friends, and teachers. Certain items (1, 9, 10 and 11) were reversed when scoring the questionnaire. The total KIDSCREEN score was calculated by summing up all the responses. Higher scores of KIDSCREEN indicated better QoL. The questionnaire was translated into Slovene following recommended guidelines [31]. The translation process included: (a) translation from English to Slovene by two independent native speakers of whom one knew about the instruments and concept under investigation and the other did not; (b) comparison of translated versions and (c) reaching a consensus; (d) back translation into the original language by two native speakers blind to the original version of the questionnaires; (e) comparison of a translated version and original version and (f) reaching a consensus among the involved translators.

The chosen institutions were contacted and approval for conducting the research was sought. On the agreed days, the questionnaires were distributed and collected. The data collection was performed in their classrooms by the teachers. The adolescents completed the questionnaires in a written (printed) version. It took approximately 20 minutes to complete the questionnaire.

The results of the research were analyzed using the programming language for statistical analysis R v 3.6.1. [22] thus ensuring the reproducibility of the research that is often overlooked in nursing research.

Analysis

The collected questionnaires were analyzed using the statistical program R [22]. A 6-step protocol for conducting psychometric analysis for detecting scale properties was used [21]. The basic descriptive statistics and plotting are often overshadowed by advanced methods. To prevent this, item-level descriptives were analyzed to detect outliers, out-of-range values, missing values, reverse-coded items, and item variations [21].

Examination of score distributions for further planned analyses was performed. Item properties were investigated via non-parametric and parametric item response theory (IRT). IRT is a psychometric theory of measurement that demonstrated reliability and validity of measurement [32]. A Mokken scale analysis (MSA) was performed to examine the homogeneity of items in the scale. MSA is a scaling technique used for ordinal data in scaling tests and questionnaires. It consists of two steps: (1) a selection algorithm that partitions a set of ordinal variables into scales (Mokken scales) and (2) methods to investigate assumptions of nonparametric IRT models [33]. Coefficients of homogeneity (H) for items in each subscale were calculated using mokken package [33]. Moreover, Automatic Item Selection Procedure (AISP) function from the Mokken package was used to explore possible dimensionality solutions at increasing thresholds [34].

The parametric IRT model can be used for further diagnosis of unidimensional scales fit for binary (Rasch model) or ordinal items (Rating scale model). In this study the Rating scale model as implemented in the eRm package [35] was used to calculate item-based characteristics. More specifically, the fit of actual responses and their logistic function was measured. According to Bond and Fox [36] the fit can be measured by putting more weight to responses from participants whose ability levels are closer to the item's difficulty level (infit) or putting more weight to extreme scorers (outfit). Item fit statistics (infit and outfit) indicate the model deviation. The mean squares of item fit should stay in the 0.5 – 1.5 range [37].

Then the scale structure was explored using explanatory factor analysis (EFA) and confirmatory factor analysis (CFA), scale reliability using a classical test theory (CTT), and calculation and description of global scores. Factor analysis is a multivariate statistical procedure that reduces a large number of variables into a smaller set of variables (factors). It establishes dimensions between measured variables and latent constructs, and it provides construct validity evidence [38]. Factor analysis was conducted using package Psych [39] and Lavaan [40], which include various factor analysis options.

In the process of conducting CFA, different estimation methods were used: the root mean square error of approximation - RMSEA [41], comparative fit index - CFI [42], and Tucker–Lewis index - TLI [43; 44]. RMSEA is widely used as an absolute fit index that assesses properties of hypothesized model and compares it to a perfect model. CFI and TLI are incremental fit indices that compare the fit of a hypothesized model with that of a baseline model [45]. Parallel analysis method (PA) [46] and the Minimum Average Partial (MAP) [47] were considered when interpreting the data. PA is often used to determine the number of components to keep in a principal component analysis or factors to keep in an exploratory factor analysis. Moreover, MAP is used to determine the number of components, which focuses on the common variance in a correlation matrix. Very simple structure (VSS) was also considered as an alternative procedure for estimation of the optimal number of factors [48]. The sample size Bayesian information criterion (SABIC) was chosen as a model selection method [49]. CTT was conducted to test reliability for the item (sub)sets using Cronbach's α [50; 51] and other reliability indices such as ω , β , and Guttman's lambda 6 - G6 [21; 52]. Dunn et al. [53] stated that calculating coefficient alpha is not enough to say that the measurement is reliable, thus they proposed calculation confidence intervals (CIs) by a method called bootstrapping, which involves repeated resampling with sample replacements to obtain an empirical distribution of an estimator [54]. Dima [21] also proposed calculation of omega, which makes more realistic assumptions than Cronbach's α alone.

Ethics

Prior to the research, ethical approval was obtained from the Slovenian National Medical Ethics Committee (no. 0120-313/2019/13). Headmasters of schools were contacted and sought for the permission to carry out the research. Participants invited to the study were informed orally and in written form about the purpose of the research, instructions for compliance, confidentiality, anonymity, and volunteering, and the possibility of withdrawing from participation in any research phase. We informed them about the fact that we will use the obtained results exclusively for the purpose of the research. We adhered to the principles of the Helsinki Declaration on Biomedical Research in Human Beings [55], the Oviedo Convention [56], the Convention on the Rights of the Child [57], the Personal Data Protection Act [58], and of the Council and the principles of the Code of Medical Deontology of Slovenia [59]. Ethical principles of the Code of ethics in nursing and healthcare of Slovenia [60] were also considered and applied. Adolescents needed additional parental consent to participate in the study [61]. Additional information about the study were provided for students and parents. The contact information was also provided; thus, parent did have an opportunity to ask additional questions about the study. Both students younger than 16 years and parents signed the consent form. We store the consents to participate in the survey and survey questionnaires in locked rooms [62].

Results

All 6 steps of psychometric testing are displayed in the following text.

Step 1: Descriptive statistics

Firstly, response frequencies were checked for all items to show if items have sufficient variation to respondents. A minimum score for KIDSCREEN-27 was 30 and maximum 135. The mean value was 101.46 (SD = 18.43).

Secondly, subscales correlations were calculated to examine the power of mutual connections between subscales (Figure 1).

There are no negative correlations nor outliers between items in the KIDSCREEN-27 subscales. Based on the statistical parameters and graphical data presentation, the KIDSCREEN-27 is constructed of six subscales.

Step 2: Mokken scale analysis

The scale is homogeneous ($H \geq 0.40$) for all five subscales in KIDSCREEN-27 (Table 2).

Table 2: Homogeneity coefficients for subscales of KIDSCREEN-27

Scale	<i>H (se)</i>
Physical well-being	0.450 (0.010)
Psychological well-being	0.479 (0.010)
Autonomy and parent relation	0.431 (0.010)
Peers and social support	0.664 (0.012)
School environment	0.574 (0.011)

H=homogeneity coefficient; se=standard errors.

All items in KIDSCREEN-27 form a single scale at different homogeneity levels. The subscale physical well-being is unidimensional at the homogeneity level of 0.35, psychological well-being at 0.45, autonomy and parent relation at 0.55, peers and social support at 0.55, and school environment at 0.50 (Supplement 1).

Step 3: Parametric item response theory (IRT)

A fit diagnosis was performed with the Rating scale model. Items infit and outfit present the extent to which responses fit logistic function for each item (Table 3).

Table 3: The mean square of item fit values

Scale	Infit		Outfit	
	min	max	min	max
Physical well-being	0.733	1.061	1.065	0.737
Psychological well-being	0.737	0.988	0.718	1.101
Autonomy and parent relation	0.749	1.045	0.780	1.066
Peers and social support	0.540	0.993	0.545	1.020
School environment	0.675	0.852	0.678	0.861

The KIDSCREEN-27 subscales have range of *infit* between 0.540 and 1.061, and *outfit* between 0.545 and 1.101.

Step 4: Factor analysis

Conducted parallel analysis based on tetrachoric correlations suggested usage of 8 factors and 6 components for the KIDSCREEN-27 scale (Figure 2).

Our results from the VSS analysis point to the optimal number of two factors, but both Velicer MAP and SABIC suggest a 5-factor structure (Figure 3).

The CFA showed poor fit according to user model versus baseline model metrics ($CFI = 0.847$; $TLI = 0.862$) and good fit in root mean square error-based metric ($RMSEA = 0.072$; $p(\chi^2) < 0.001$).

Step 5: Classical test theory (CTT)

In the step 5, a reliability for the KIDSCREEN-27 and sub-sets was calculated using Cronbach's α , beta, G6 lambda and omega (Table 4).

Table 4: Classical test theory values for KIDSCREEN-27 and subscales

	α	β	λ^*	ω
KIDSCREEN-27	0.93	0.86	0.95	0.93
Physical well-being	0.78	0.67	0.75	0.79
Psychological well-being	0.85	0.72	0.86	0.84
Autonomy and parent relation	0.83	0.68	0.84	0.82
Peers and social support	0.87	0.84	0.85	0.87
School environment	0.82	0.79	0.78	0.82

* Guttman's Lambda 6

Step 6: Total (sub)scale scores

In the final step, data distribution was checked. Data in all subscales shows acceptable distribution (Figure 4).

Summary statistics were calculated and are shown in Table 5. There were no outliers detected and all parameters show expected scores.

Table 5: Summary statistics

	M	SD
KIDSCREEN-27	101.23	18.46
Physical well-being	18.38	3.97
Psychological well-being	26.49	5.90
Autonomy and parent relation	26.26	5.76
Peers and social support	16.07	3.74
School environment	14.03	3.57

M = mean; SD = standard deviation.

The Slovenian versions of the KIDSCREEN-27 scale achieved good validity, reliability, and psychometric properties.

Discussion

The KIDSCREEN-27 questionnaire is instrument for monitoring QoL of children and adolescents to provide clarity on which factors from a wider social environment may contribute to children's or adolescents' mental well-being [25; 63]. The KIDSCREEN-27 was the first instrument for children that was tested in several countries [17; 18; 25; 63; 64; 65; 66; 67; 68; 69; 70; 71; 72; 73; 74; 75] and in a large representative and multinational sample of children and adolescents in 13 countries (Austria, Czech Republic, France, Germany, Greece, Hungary, Ireland, Poland, Spain, Sweden, Switzerland, The Netherlands, and the United Kingdom) [76]. To our knowledge, this is the first published validation of the KIDSCREEN-27 among adolescents in Slovenia. The conducted analyses show that the Slovene version of the questionnaire has overall good psychometric properties for the use among adolescents. The questionnaire was translated following all the steps of the translation process. The questionnaire was then analyzed following a 6-step analysis in R: descriptive, non-parametric IRT, parametric IRT, factor analysis, classical test theory and total scores) [21].

According to the analysis, MSA identified one item that did not meet the criteria for ordinal scaling. The physical well-being subscale does not form a unidimensional scale at $H \geq 0.40$. Fit diagnosis was performed using the Rating scale model. The KIDSCREEN-27 subscales have range of *infit* between 0.540 and 1.061, and *outfit* between 0.545 and 1.101. Similar results are reported by Véléz et al. [17]. Parallel analysis suggested 8 factors and 6 components, but both Velicer MAP and SABIC suggest a five-factor structure. The CFA showed average fit in the user model versus baseline model metrics ($CFI = 0.847$; $TLI = 0.862$) and good fit in root mean square error ($RMSEA = 0.072$; $p(\chi^2) < 0.001$). This might be since KIDSCREEN-27 questionnaire already measures QoL. CFA results of KIDSCREEN-27 among adolescents in Turkey [70] showed similar results ($RMSEA=0.07$; $CFI=0.96$). Poor results ($RMSEA=0.10$; $CFI=0.91$) were also reported in a study conducted among Chinese adolescents [69]. Another study suggested a modified 7-factor model, which is better at explaining the data than the original 5-factor model [63]. Ng et al. [69] agreed with a 7-factor model.

However, studies conducted among 13 European countries [64] and one conducted among Chilean adolescents [76] have found a 5-factor model acceptable.

The reliability and internal consistency of the KIDSCREEN-27 were calculated using Cronbach's α , beta, G6 and omega. Cronbach's α was 0.93 and is interpreted as acceptable [77; 78; 79]. Ng et al. [69] also reported acceptable internal consistency (0.84 – 0.89). Similar results were reported in a study among Turkish adolescents where Cronbach's α ranged between 0.78 and 0.84 [70]. They also reported that the dimension self-perception has lower Cronbach's α value (0.69), which confirms the results of Serbian (0.58) [67] and Iranian (0.60) [68] studies. In another study conducted among Irish children [63] Cronbach's α was below the recommended limit. Values of each subscale ranged from 0.56 to 0.74. Values in our study ranged from 0.78 to 0.93. Internal consistency was also evaluated using beta (0.86), lambda G6 (0.95) and omega (0.93), which other studies did not perform. Thus, additional comparisons cannot be performed.

Although our study was conducted following rigorous psychometric analysis and among large student sample, there are some limitations that are worth noting. The test-retest reliability was not conducted, because data was collected in a certain time frame. Moreover, the KIDSCREEN-27 is a self-reported questionnaire measuring social support, meaning that participants may give socially desirable answers. Thus, findings should be interpreted with caution. Although all invited schools had the same study protocol, most of them did not involve all students who are enrolled at their school. There may be a gap in how were students selected. Thus, due to a large sample of schools included in the study the authors of the study did not have the information on the strategies used to approach the students in the participating schools.

Conclusion

Overall, KIDSCREEN-27 is a valid and reliable questionnaire for measuring QoL of adolescents and the wider social factors that influence adolescents' mental well-being. The confirmatory factor analysis showed poor fit in user model versus baseline model metrics and good fit in root mean square error. This might be the consequence of having several latent variables in KIDSCREEN-27 questionnaire that are measuring different aspects of the QoL. The findings from this study add to the international understanding of the KIDSCREEN-27 scale and support the use of the Slovene translation of the scale. Although, scale underwent a rigorous process of psychometric testing following a 6-step analysis in R there are possibilities for improvements. A longitudinal design of a similar study would allow the observation of the temporal complexity in the KIDSCREEN-27 questionnaire results. Also, other study limitations should be considered when performing similar research.

Declarations

Ethics approval and consent to participate

Prior to the research, ethical approval was obtained from the Slovenian National Medical Ethics Committee (no. 0120-313/2019/13).

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and analysed during the current study are not publicly available but are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

LCB: Conceptualization, Writing- Original draft preparation, Methodology, Investigation, Analyses MP: Conceptualization, Writing-Reviewing and Editing, Supervision URS: Writing- Reviewing and Editing OB: Writing- Reviewing and Editing, Supervision GS: Data Visualization, Software, Investigation, Supervision.

Acknowledgements

We would like to thank all students who participated in this study.

References

1. World Health Organization. (2019). *Maternal, newborn, child and adolescent health*. Retrieved July 1, 2019, from https://www.who.int/maternal_child_adolescent/topics/adolescence/development/en/
2. Bluth, K., Campo, R. A., Futch, W. S., & Gaylord, S. A. (2017). Age and gender differences in the associations of self-compassion and emotional well-being in a large adolescent sample. *Journal of youth and adolescence*, 46(4), 840-853. <https://doi.org/10.1007/s10964-016-0567-2>
3. Ronen, T., Hamama, L., Rosenbaum, M., & Mishely-Yarlap, A. (2016). Subjective well-being in adolescence: The role of self-control, social support, age, gender, and familial crisis. *Journal of Happiness Studies*, 17(1), 81-104. <https://doi.org/10.1007/s10902-014-9585-5>
4. Patton, G. C., Sawyer, S. M., Santelli, J. S., Ross, D. A., Afifi, R., Allen, N. B., ... & Kakuma, R. (2016). Our future: a Lancet commission on adolescent health and wellbeing. *The Lancet*, 387(10036), 2423-2478. [https://doi.org/10.1016/S0140-6736\(16\)00579-1](https://doi.org/10.1016/S0140-6736(16)00579-1)
5. World Health Organization. (2018) *Adolescents mental health*. Retrieved July 1, 2019, from <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health>
6. Black, L., Panayiotou, M., & Humphrey, N. (2019). The dimensionality and latent structure of mental health difficulties and wellbeing in early adolescence. *PloS one*, 14(2), e0213018. <https://doi.org/10.1371/journal.pone.0213018>
7. Guo, C., Tomson, G., Keller, C., & Söderqvist, F. (2018). Prevalence and correlates of positive mental health in Chinese adolescents. *BMC Public Health*, 18(1), 263. <https://doi.org/10.1186/s12889-018-5133-2>
8. Lester, L., & Cross, D. (2015). The relationship between school climate and mental and emotional wellbeing over the transition from primary to secondary school. *Psychology of Well-being*, 5(1), 9. <https://doi.org/10.1186/s13612-015-0037-8>
9. Arabiat, D. H., Shaheen, A., Nassar, O., Saleh, M., & Mansour, A. (2018). Social and Health Determinants of Adolescents' Wellbeing in Jordan: Implications for Policy and Practice. *Journal of Pediatric Nursing*, 39, 55-60. <https://doi.org/10.1016/j.pedn.2017.03.015>
10. Alsubaie, M.M., Stain, H. J., Webster L. A. D. & Wadman, R. (2019) The role of sources of social support on depression and quality of life for university students. *International Journal of Adolescence and Youth*, 24(4), 484-496. <https://doi.org/10.1080/02673843.2019.1568887>
11. Lopez-Zafra, E., Ramos-Álvarez, M. M., El Ghoudani, K., Luque-Reca, O., Augusto-Landa, J. M., Zarhbouch, B., ... & Pulido-Martos, M. (2019). Social support and emotional intelligence as protective resources for well-being in moroccan adolescents. *Frontiers in psychology*, <https://doi.org/10.103389/fpsyg.2019.01529>
12. Malecki, C. K., & Demaray, M. K. (2006). Social support as a buffer in the relationship between socioeconomic status and academic performance. *School Psychology Quarterly*, 21(4), 375–395. <https://doi.org/10.1037/h0084129>
13. Williams, P., Barclay, L. & Schmied, V. (2004). Defining Social Support in Context: A Necessary Step in Improving Research, Intervention, and Practice. *Qualitative Health Research*, 14(7), 942-960. <https://doi.org/10.1177/1049732304266997>
14. Vettore, M. V., Ahmad, S. F., Machuca, C., & Fontanini, H. (2019). Socio-economic status, social support, social network, dental status, and oral health reported outcomes in adolescents. *European journal of oral sciences*, 127(2), 139-146. <https://doi.org/10.1111/eos.12605>
15. Karaer, Y., & Akdemir, D. (2019). Parenting styles, perceived social support and emotion regulation in adolescents with internet addiction. *Comprehensive psychiatry*, 92, 22-27. <https://doi.org/10.1016/j.comppsyg.2019.03.003>
16. The KIDSCREEN Group. (2010). *KIDSCREEN-27*. Retrieved July 18, 2019, from <https://www.kidscreen.org/cms/es/node/104>
17. Vélez, C. M., Lugo-Agudelo, L. H., Hernández-Herrera, G. N., & García-García, H. I. (2016). Colombian Rasch validation of KIDSCREEN-27 quality of life questionnaire. *Health and Quality of Life Outcomes*, 14(1), 67. <https://doi.org/10.1186/s12955-016-0472-0>
18. Andersen, J. R., Natvig, G. K., Haraldstad, K., Skrede, T., Aadland, E., & Resaland, G. K. (2015). Is the Kidscreen-27 a valid measure of health-related quality of life in 10-year-old Norwegian children? *PeerJ PrePrints*, 3, e1134v1. <https://doi.org/10.7287/peerj.preprints.1134v1>
19. Boateng, G. O., Neilands, T. B., Frongillo, E. A., Melgar-Quinonez, H. R., & Young, S. L. (2018). Best practices for developing and validating scales for health, social, and behavioral research: a primer. *Frontiers in public health*, 6, 149. <https://doi.org/10.3389/fpubh.2018.00149>
20. Mokkink, L. B., Terwee, C. B., Patrick, D. L., Alonso, J., Stratford, P. W., Knol, D. L., ... & De Vet, H. C. (2010). The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Quality of life research*, 19(4), 539-549. <https://doi.org/10.1007/s11136-010-9606-8>
21. Dima, A. L. (2018). Scale validation in applied health research: tutorial for a 6-step R-based psychometrics protocol. *Health Psychology and Behavioral Medicine*, 6(1), 136-161. <https://doi.org/10.1080/21642850.2018.1472602>
22. R Development Core Team (2007). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
23. Ravens-Sieberer, U., Herdman, M., Devine, J., Otto, C., Bullinger, M., Rose, M., & Klasen, F. (2014). The European KIDSCREEN approach to measure quality of life and well-being in children: development, current application, and future advances. *Quality of life research*, 23(3), 791-803. <https://doi.org/10.1007/s11136-013-0428-3>
24. The KIDSCREEN Group. (2004). *KIDSCREEN instruments: Health-Related Quality of Life Questionnaire for Children and Young People*. Retrieved July 18, 2019, from <https://www.kidscreen.org/english/questionnaires/>
25. Ravens-Sieberer, U., Auquier, P., Erhart, M., Gosch, A., Rajmil, L., Bruil, J., ... & Mazur, J. (2007). The KIDSCREEN-27 quality of life measure for children and adolescents: psychometric results from a cross-cultural survey in 13 European countries. *Quality of Life Research*, 16(8), 1347-

1356. <https://doi.org/10.1007/s11136-007-9240-2>
26. Statistical Office of the Republic of Slovenia. (2018). *SURS*. Retrieved July 25, 2018, from http://pxweb.stat.si/pxweb/Database/Dem_soc/09_izobrazevanje/04_osnovnosol_izobraz/01_09527_zac_sol_leta/01_09527_zac_sol_leta.asp
  27. Ministry of Education, Science and Sport of the Republic of Slovenia. (2018). *Direktorat za srednje in višje šolstvo ter izobraževanje odraslih*. Retrieved July 25, 2018, from http://www.mizs.gov.si/si/delovna_podrocja/direktorat_za_srednje_in_visje_solstvo_ter_izobrazevanje_odraslih/
  28. Statistical Office of the Republic of Slovenia. (2017). *STATOPIS: statistični pregled Slovenije 2017*. Retrieved July 25, 2018, from <https://www.stat.si/StatWeb/File/DocSysFile/9743>
  29. Qualtrics. (2018). *Calculating Sample Size*. Retrieved September 13, 2018, from <https://www.qualtrics.com/blog/calculating-sample-size/>
  30. Ravens-Sieberer, U. et al., 2005. KIDSCREEN-52 quality-of-life measure for children and adolescents. *Expert Review of Pharmacoeconomics & Outcomes Research*, 5(3), pp. 353-364. <https://doi.org/10.1586/14737167.5.3.353>
  31. Sperber, A. D. (2004). Translation and validation of study instruments for cross-cultural research. *Gastroenterology*, 126, S124-S128. <https://doi.org/10.1053/j.gastro.2003.10.016>
  32. Rizopoulos, D. (2010). *Item Response Theory in R using Package ltm*. Department of Statistics and Mathematics, WU Wirtschaftsuniversitat Wien.
  33. Van der Ark, L. A. (2007). Mokken scale analysis in R. *Journal of statistical software*, 20(11), 1-19.
  34. Hemker, B. T., Sijtsma, K., & Molenaar, I. W. (1995). Selection of unidimensional scales from a multidimensional item bank in the polytomous Mokken IRT model. *Applied Psychological Measurement*, 19(4), 337-352. <https://doi.org/10.1177/014662169501900404>
  35. Mair, P., & Hatzinger, R. (2007). Extended Rasch modeling: The eRm package for the application of IRT models in R. *Journal of Statistical Software*, 20(9). <https://doi.org/10.18637/jss.v020.i09>
  36. Bond, T., Yan, Z., & Heene, M. (2020). *Applying the Rasch model: Fundamental measurement in the human sciences*. Routledge.
  37. Boone, W., & Rogan, J. (2005). Rigour in quantitative analysis: The promise of Rasch analysis techniques. *African Journal of Research in Mathematics, Science and Technology Education*, 9(1), 25-38. <https://doi.org/10.1080/10288457.2005.10740574>
  38. Williams, B., Onsman, A., & Brown, T. (2010). Exploratory factor analysis: A five-step guide for novices. *Australasian journal of paramedicine*, 8(3). <https://doi.org/10.33151/ajp.8.3.93>
  39. Revelle, W. (2017). Psych: procedures for psychological, psychometric, and personality research. Retrieved August 4, 2021, from <https://CRAN.R-project.org/package=psych>
  40. Rosseel, Y. (2012). Lavaan: An R package for structural equation modeling and more. Version 0.5–12 (BETA). *Journal of statistical software*, 48(2), 1-36. <https://doi.org/10.18637/jss.v048.i02>
  41. Steiger, J. H. (1990). Structural model evaluation and modification: An interval estimation approach. *Multivariate behavioral research*, 25(2), 173-180. https://doi.org/10.1207/s15327906mbr2502_4
  42. Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological bulletin*, 107(2), 238. <https://doi.org/10.1037/0033-2909.107.2.238>
  43. Bentler, P. M., & Bonett, D. G. (1980). Significance tests and goodness of fit in the analysis of covariance structures. *Psychological bulletin*, 88(3), 588. <https://doi.org/10.1037/0033-2909.88.3.588>
  44. Tucker, L. R., & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, 38(1), 1-10. <https://doi.org/10.1007/BF02291170>
  45. Xia, Y., & Yang, Y. (2019). RMSEA, CFI, and TLI in structural equation modeling with ordered categorical data: The story they tell depends on the estimation methods. *Behavior research methods*, 51(1), 409-428. <https://doi.org/10.3758/s13428-018-1055-2>
  46. Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis. *Psychometrika*, 30(2), 179-185. <https://doi.org/10.1007/BF02289447>
  47. Velicer, W. F. (1976). Determining the number of components from the matrix of partial correlations. *Psychometrika*, 41(3), 321-327. <https://doi.org/10.1007/BF02293557>
  48. Revelle, W., & Rocklin, T. (1979). Very simple structure: An alternative procedure for estimating the optimal number of interpretable factors. *Multivariate Behavioral Research*, 14(4), 403-414. https://doi.org/10.1207/s15327906mbr1404_2
  49. Sclove, S. L. (1987). Application of model-selection criteria to some problems in multivariate analysis. *Psychometrika*, 52(3), 333-343. <https://doi.org/10.1007/BF02294360>
  50. Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, 16(3), 297-334. <https://doi.org/10.1007/BF02310555>
  51. Cronbach, L. J., & Shavelson, R. J. (2004). My current thoughts on coefficient alpha and successor procedures. *Educational and psychological measurement*, 64(3), 391-418. <https://doi.org/10.1177/0013164404266386>
  52. Revelle, W., & Zinbarg, R. E. (2009). Coefficients alpha, beta, omega, and the glb: Comments on Sijtsma. *Psychometrika*, 74(1), 145. <https://doi.org/10.1007/s11336-008-9102-z>

53. Dunn, T. J., Baguley, T., & Brunsden, V. (2014). From alpha to omega: A practical solution to the pervasive problem of internal consistency estimation. *British journal of psychology*, 105(3), 399-412. <https://doi.org/10.1111/bjop.12046>
54. DiCiccio, T. J., & Efron, B. (1996). Bootstrap confidence intervals. *Statistical science*, 189-212.
55. World Medical Association. (2013). World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*, 310(20), pp. 2191–2194.
56. Council of Europe. (1997). *Convention for the protection of human rights and dignity on the human being with regard to the application of biology and medicine: Convention on human rights and biomedicine*. Oviedo: Council of Europe.
57. United Nations General Assembly. (1989). *Convention on the rights of the child, 20 November 1989, United Nations, Treaty Series, vol. 1577, p. 3*. New York: United Nations.
58. *Personal Data Protection Act*. Official Gazette of the RS, no. 86/04.
59. *Code of Medical Deontology of Slovenia*. (1992). Ljubljana: Medical Chamber of Slovenia.
60. *Code of ethics in nursing and healthcare of Slovenia*. Official Gazette of the RS, no. 71/14.
61. Harris, J. & Porcellato, L. (2018). Opt-out parental consent in online surveys: ethical considerations. *Journal of Empirical Research on Human Research Ethics*, 13(3), pp. 223–229. <https://doi.org/10.1177/1556264618766953>
62. Royal College of Nursing. (2009). *Research ethics: RCN guidance for nurses*. London: The Royal College of Nursing.
63. Shannon, S., Breslin, G., Fitzpatrick, B., Hanna, D., & Brennan, D. (2017). Testing the psychometric properties of KIDSCREEN-27 with Irish children of low socio-economic status. *Quality of Life Research*, 26(4), 1081-1089. <https://doi.org/10.1007/s11136-016-1432-1>
64. Robitail, S., Ravens-Sieberer, U., Simeoni, M. C., Rajmil, L., Bruil, J., Power, M., ... & Czimbalmos, A. (2007). Testing the structural and cross-cultural validity of the KIDSCREEN-27 quality of life questionnaire. *Quality of Life Research*, 16(8), 1335-1345. <https://doi.org/10.1007/s11136-007-9241-1>
65. Quintero, C. A., Lugo, L. H., García, H. I., & Sánchez, A. (2011). Validation of questionnaire KIDSCREEN-27 to measure health related quality of life in children and adolescents of medellin, Colombia. *Revista Colombiana de Psiquiatría*, 40(3), [https://doi.org/10.1016/S0034-7450\(14\)60141-4](https://doi.org/10.1016/S0034-7450(14)60141-4)
66. Pardo-Guijarro, M. J., Woll, B., Moya-Martínez, P., Martínez-Andrés, M., Cortés-Ramírez, E. E., & Martínez-Vizcaíno, V. (2013). Validity and reliability of the Spanish sign language version of the KIDSCREEN-27 health-related quality of life questionnaire for use in deaf children and adolescents. *Gaceta Sanitaria*, 27(4), 318-324. <https://doi.org/10.1016/j.gaceta.2012.11.003>
67. Stevanovic, D., Tadic, I., Novakovic, T., Kisc-Tepavcevic, D., & Ravens-Sieberer, U. (2013). Evaluating the Serbian version of the KIDSCREEN quality-of-life questionnaires: reliability, validity, and agreement between children's and parents' ratings. *Quality of life research*, 22(7), 1729-1737. <https://doi.org/10.1007/s11136-012-0286-4>
68. Parizi, A. S., Garmaroudi, G., Fazel, M., Omidvari, S., Azin, S. A., Montazeri, A., & Jafarpour, S. (2014). Psychometric properties of KIDSCREEN health-related quality of life questionnaire in Iranian adolescents. *Quality of Life Research*, 23(7), 2133-2138. <https://doi.org/10.1007/s11136-014-0655-2>
69. Ng, J. Y., Burnett, A., Ha, A. S., & Sum, K. W. (2015). Psychometric properties of the Chinese (Cantonese) versions of the KIDSCREEN health-related quality of life questionnaire. *Quality of Life Research*, 24(10), 2415-2421. <https://doi.org/10.1007/s11136-015-0973-z>
70. Baydur, H., Ergin, D., Gerçeklioğlu, G., & Eser, E. (2016). Reliability and validity study of the KIDSCREEN Health-Related Quality of Life Questionnaire in a Turkish child/adolescent population. *Anatolian Journal of Psychiatry/Anadolu Psikiyatri Dergisi*, 17(6). <https://doi.org/10.5455/apd.214559>
71. Nezu, S., Iwasaka, H., Saeki, K., Obayashi, K., Ishizuka, R., Goma, H., ... & Kurumatani, N. (2016). Reliability and validity of Japanese versions of KIDSCREEN-27 and KIDSCREEN-10 questionnaires. *Environmental health and preventive medicine*, 21(3), 154-163. <https://doi.org/10.1007/s12199-016-0510-x>
72. Farias Júnior, J. C. D., Loch, M. R., Lima Neto, A. J. D., Sales, J. M., & Ferreira, F. E. L. D. L. (2017). Reproducibility, internal consistency, and construct validity of KIDSCREEN-27 in Brazilian adolescents. *Cadernos de Saúde Pública*, 33(9). <https://doi.org/10.1590/0102-311x00131116>
73. Vrkić, A. (2018). *Tjelesna aktivnost i kvaliteta života studentica završnih godina učiteljskog fakulteta (Master thesis)*. Zagreb: University of Zagreb. Faculty of Teacher Education. Chair of Kinesiology Education.
74. Power, R., Akhter, R., Muhit, M., Wadud, S., Heanoy, E., Karim, T., ... & Khandaker, G. (2019). Cross-cultural validation of the Bengali version KIDSCREEN-27 quality of life questionnaire. *BMC pediatrics*, 19(1), 19. <https://doi.org/10.1186/s12887-018-1373-7>
75. Ravens-Sieberer, U., Erhart, M., Rajmil, L., Herdman, M., Auquier, P., Bruil, J., ... & Mazur, J. (2010). Reliability, construct and criterion validity of the KIDSCREEN-10 score: a short measure for children and adolescents' well-being and health-related quality of life. *Quality of life research*, 19(10), 1487-1500. <https://doi.org/10.1007/s11136-010-9706-5>
76. Molina, T. G., Montaña, R. E., González, E. A., Sepúlveda, R. P., Hidalgo-Rasmussen, C., Martínez, V. N., ... & George, M. L. (2014). Psychometric properties of the quality of life questionnaire health related KIDSCREEN-27 in Chilean adolescents. *Revista medica de Chile*, 142(11), 1415-1421. <https://doi.org/10.4067/s0034-98872014001100008>

77. Nunnally, J. C. (1994). *Psychometric theory 3E*. Tata McGraw-hill education.

78. Bland, J. M., & Altman, D. G. (1997). Statistics notes: Cronbach's alpha. *Bmj*, 314(7080), 572. <https://doi.org/10.1136/bmj.314.7080.572>

79. DeVellis, R. F. (2016). *Scale development: Theory and applications*. Sage publications.

Supplementary

Supplemental 1 is not available with this version

Figures

Figure 1

Correlation between KIDSCREEN-27 subscales

Figure 2

Parallel analysis scree plot

Figure 3

Very simple structure plot

Figure 4

Data distribution of KIDSCREEN-27 scale