Lung Response to Prone Positioning in Mechanically-ventilated Patients With COVID-19

Alessandro Protti (✉ alessandro.protti@hunimed.eu)

Humanitas University

Alessandro Santini

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Francesca Pennati

Politecnico di Milano

Chiara Chiurazzi

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Michele Ferrari

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Giacomo E. Iapichino

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Luca Carenzo

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Francesca Dalla Corte

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Ezio Lanza

Humanitas Clinical and Research Center – IRCCS, Rozzano (Milan)

Nicolò Martinetti

Humanitas University

Andrea Aliverti

Politecnico di Milano

Maurizio Cecconi

Humanitas University

Research Article

Keywords: acute respiratory distress syndrome, coronavirus disease 2019, hypoxia, mechanical ventilation, pneumonia, prone positioning

Posted Date: January 28th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1292987/v1>

Abstract

Background

Prone positioning improves survival in moderate-to-severe acute respiratory distress syndrome (ARDS) unrelated to the novel coronavirus disease (COVID-2019). This benefit is probably mediated by a decrease in alveolar collapse and hyperinflation and a more homogeneous distribution of lung aeration, with fewer harms of mechanical ventilation. Herein we aimed to verify whether prone positioning causes analogue changes in lung aeration in COVID-2019. A positive result would support prone positioning even in this other population.

Methods

Fifteen mechanically-ventilated patients with COVID-19 underwent a lung computed tomography in the supine and prone position within three days of endotracheal intubation and with a constant positive end-expiratory pressure (PEEP). Using quantitative analysis, we measured the volume of the non-aerated, poorly-aerated, well-aerated, and over-aerated compartments and the gas-to-tissue ratio of the ten vertical levels of the lung. In addition, we expressed the heterogeneity of lung aeration with the standardized median absolute deviation of the ten vertical gas-to-tissue ratios, with lower values indicating less heterogeneity.

Results

By the time of the study, PEEP was 12 (10-14) cmH₂O and the PaO₂:FiO₂ 107 (84-173) mmHg in the supine position. With prone positioning, the volume of the non-aerated compartment decreased by 82 (26-147) ml, of the poorly-aerated compartment increased by 82 (53-174) ml, of the normally-aerated compartment did not significantly change, and of the over-aerated compartment decreased by 28 (11-186) ml. In eight (53%) patients, the volume of the over-aerated compartment decreased more than the volume of the non-aerated compartment. The gas-to-tissue ratio of the ten vertical levels of the lung decreased by 0.34 (0.25-0.49) ml/g per level in the supine position and by 0.03 (-0.11-0.14) ml/g in the prone position ($p<0.001$). The standardized median absolute deviation of the gas-to-tissue ratios of those ten levels decreased in all patients, from 0.55 (0.50-0.71) to 0.20 (0.14-0.27) ($p<0.001$).

Conclusions

In fifteen patients with COVID-19, prone positioning decreased alveolar collapse and hyperinflation, and homogenized lung aeration. A similar response has been observed in other ARDS, where prone positioning improves outcome. Therefore, our data provide a pathophysiological rationale to support prone positioning even in COVID-19.

Background

Prone positioning was recommended for moderate-to-severe acute respiratory distress syndrome (ARDS) well before the appearance of the novel coronavirus disease (COVID-19) [1]. It improves survival [1,2], possibly by reopening or “recruiting” the dorsal non-aerated but perfused lung tissue and by diminishing ventral hyperinflation [3-6]. As a result, arterial oxygenation almost always increases, and, more important, ventilation becomes more evenly distributed, with fewer harms of mechanical ventilation [2-6]. Prone positioning carries some risks, including endotracheal tube obstruction or displacement (up to accidental extubation), reduced venous return, need for more sedation, vomiting, loss of venous access, and pressure sores [2-6]. The advantages of prone positioning are more likely to outweigh these dangers in moderate-to-severe ARDS, i.e., when the risk of secondary lung damage is higher [1,6,7].

Soon after its appearance, prone positioning was recommended by international guidelines and experts in the field [8,9] and widely used [10,11] even for moderate-to-severe ARDS related to COVID-19. The underlying assumption was that prone positioning is beneficial in ARDS due to COVID-19, as it is in ARDS of other origins. However, whether ARDS due to COVID-19 should be treated as ARDS of other origins remains controversial [9,12,13].

This study aimed to verify whether prone positioning decreases alveolar collapse and hyperinflation and homogenizes lung aeration in patients with early ARDS due to COVID-19. We reasoned that a positive response would support prone positioning even in this novel syndrome.

Methods

Our institutional review board approved this study (Comitato Etico dell'IRCCS Istituto Clinico Humanitas Rozzano; protocol n. 465/20). Informed consent was obtained according to local regulations.

We enrolled fifteen patients with laboratory-confirmed COVID-19 from 1/3/2020 to 09/12/2020. Inclusion criteria were: (i) a diagnosis of ARDS [14]; (ii) ongoing invasive mechanical ventilation with deep sedation and neuromuscular blockade; (iii) prone positioning prescribed by the attending physician within three days of endotracheal intubation. Forty-six patients were excluded because (i) they had already undergone a lung computed tomography (CT) after endotracheal intubation (n=16); (ii) they were too unstable for transfer to the radiology unit (n=9); (iii) their body weight exceeded 100 kg (n=6); or (iii) none of the authors was available for collecting data, due to the exceptional clinical workload at that time (n=15) (Table E1).

Lung morphological response

Following a recruitment manoeuvre at 45 cmH₂O of end-inspiratory airway pressure [15], a static end-expiratory lung CT was taken with the patient in the supine position as described in reference 16. Positive end-expiratory pressure (PEEP) was the same as in the intensive care unit (ICU) prior to the study. Patients were then turned prone. After a new recruitment manoeuvre (as above), a second static CT was taken at the same PEEP level. Thereafter, patients were returned to the ICU in the supine position.

- Global inflation

The total volume, tissue weight, and gas volume of the whole lung and its non-aerated (density above -100 HU), poorly-aerated (from -100 to -500 HU), normally-aerated (from -500 to -900 HU), and over-aerated (below -900 HU) compartments were measured as in reference 15. The premorbid lung weight was estimated from the subjects' height [17]. The average lung aeration was expressed as the ratio of total gas volume to total tissue weight [3]. The global lung morphological response to prone positioning was assessed as the change in the total volume of the non-aerated (alveolar collapse) and over-aerated (hyperinflation) compartments from supine to prone [16].

Regional inflation

These same methods were applied to ten equal vertical and horizontal levels forming each CT slice, from the sternum (ventral) to the vertebra (dorsal) and from the apex (cranial) to the base (caudal) of the lung. The regional aeration was computed as the ratio of gas volume to tissue weight of each vertical and horizontal level [3]. The regional lung morphological response to prone positioning was assessed as the degree of heterogeneity of lung aeration along the sterno-vertebral or cranio-caudal axis. This was quantified with the standardized median absolute deviation of the corresponding ten gas-to-tissue ratios within each subject [18]; higher values indicated more heterogeneity.

The hydrostatic pressure (super)imposed on each vertical level was computed as in reference 19.

Lung functional response

Gas exchange and respiratory system mechanics were measured 20 minutes after a recruitment manoeuvre (as above), immediately before and 60 minutes after prone positioning. All ventilatory settings had to remain reasonably constant in the two positions. PEEP was the same as in the radiology unit.

Association between morphological and functional responses

We studied the association between the change in the volume of the non-aerated or over-aerated compartment, or the change in heterogeneity of lung aeration, and: (i) those volumes and heterogeneity in supine position; (ii) the change in oxygenation, compliance, and PaCO₂ in response to prone positioning.

Statistical analysis

Data are presented as median (Q1-Q3) or proportion. They were analysed with the Mann-Whitney rank-sum test, Wilcoxon signed rank-sum test, Fisher's exact test, and Spearman's rank-order correlation using Sigma Plot 11.0 (Jandel Scientific, San Jose, CA). A two-tailed p-value <0.05 was considered statistically significant.

Results

We enrolled fifteen patients with COVID-19 on invasive mechanical ventilation. Their main characteristics at ICU admission are reported in Table 1 and Table E2. Two (13%) were active smokers, and none (0%) had a history of chronic lung disease.

The study was performed 4 (3-5) days after hospital admission and 2 (1-2) days after endotracheal intubation. By that time, in the supine position and with a PEEP of 12 (10-14) cmH₂O, the ratio of arterial tension (PaO₂) to inspiratory fraction (FiO₂) of oxygen was 107 (84-173) mmHg. One patient had a PaO₂:FiO₂ of 273 mmHg. We decided to prone this patient in the radiology unit after noting significant ventral hyperinflation on the lung CT obtained in the supine position.

Lung morphological response to prone positioning

- Global inflation

In the supine position, the total lung volume was 3277 (2390-3533) ml. Four-hundred-and-seven (238-641) ml or 13 (9-22)% of that total lung volume were in the non-aerated compartment; 729 (563-1181) ml or 25 (19-32)% in the poorly-aerated compartment; 1449 (1189-2142) ml or 50 (42-66)% in the normally-aerated compartment; and 31 (21-376) ml or 1.6 (0.8-10.7)% in the over-aerated compartment. The lung weight was 1434 (1079-1872) g, 619 (221-834) g higher than the estimated premorbid one (Figure E1). The lung gas volume was 1541 (1242-2081) ml.

With prone positioning, lung inflation changed as described in Figure 1 and Table E3. The volume of the non-aerated compartment decreased in twelve (80%) patients, and the volume of the over-aerated compartment in fourteen (93%). The total lung volume and weight did not significantly change while the gas volume decreased by 197 (8-290) ml. The volume of the non-aerated compartment decreased by 82 (26-147) ml or 15 (5-53)%; the volume of the poorly-aerated compartment increased by 82 (53-174) ml (or 9 (6-25)%; the volume of the normally-aerated compartment did not significantly change; the volume of the over-aerated compartment decreased by 28 (11-186) ml or 68 (46-81)% (Figure 2). In eight (53%) patients, hyperinflation decreased more than alveolar collapse. The gas-to-tissue ratio of the whole lung decreased in twelve (80%) patients (Figure 1), and from 1.6 (0.7-2.7) to 1.4 (0.8-1.9) ml/g (p=0.008) in the overall study population.

- Regional inflation

In the supine position, the over-aerated compartment tended to be larger ventrally and cranially while the non-aerated compartment dorsally and caudally (Figure E2 and E3). The gas-to-tissue ratio decreased from the sternum to the vertebra and from the apex to the base of the lung (Figure 3). The distribution of inflation was more heterogeneous along the vertical than horizontal axis, with a standardized median absolute deviation of 0.55 (0.50-0.71) and 0.31 (0.16-0.47), respectively (p=0.012). The superimposed pressure partly explained the vertical gradient of aeration: it progressively increased from 0.5 (0.4-0.6) cmH₂O close to the sternum to 9.1 (7.2-11.3) cmH₂O close to the vertebra (Figure E4).

With prone positioning, the volume of the over-aerated compartment decreased in the ventral regions and from the apex to the base of the lung (Figure E2 and E3). The volume of the non-aerated compartment decreased in the dorsal and caudal regions and increased in the ventral ones, although to a minor degree. The gas-to-tissue ratio remained variable throughout the lung (Figure 3), but the degree of heterogeneity decreased along the vertical axis in all (100%) patients and along the horizontal axis in eight (53%). The standardized median absolute deviation decreased from the sternum to the vertebra (from 0.55 [0.50-0.71] to 0.20 [0.14-0.27], $p < 0.001$); it did not change from the apex to the base of the lung (from 0.31 [0.16-0.47] to 0.29 [0.22-0.46]; $p = 0.934$) (Figure 3). Changes in the vertical gradient of aeration were associated with those of the superimposed pressure (Figure E5).

Lung functional response to prone positioning

On returning to the ICU, fourteen patients were studied supine and prone positions. One patient was not because an acute and severe cardiac arrhythmia contraindicated prone positioning. The ventilatory setting remained exactly the same from supine to prone in ten patients. In four others, it was slightly modified: the FiO_2 was increased ($n=1$) or decreased ($n=2$), or the respiratory rate was increased ($n=1$) during prone positioning (Table E4).

With prone positioning, the $\text{PaO}_2\text{:FiO}_2$ improved in all fourteen (100%) patients and by ≥ 20 mmHg in eleven (79%). Compliance increased in six (43%), remained constant in six (43%), and decreased in two (14%). The arterial carbon dioxide tension (PaCO_2) increased in six (46%), remained constant in three (23%), and decreased in four (31%) of the thirteen patients with unchanged respiratory rate and minute ventilation (Figure 4). On average, the $\text{PaO}_2\text{:FiO}_2$ increased by 41 (21-97) mmHg while compliance and PaCO_2 did not significantly change (Table E4).

Association between the morphological and functional response to prone positioning

The change in volume of the non-aerated compartment was associated with neither the severity of the alveolar collapse in the supine position ($\rho = 0.375$, $p = 0.162$) nor the concomitant change in $\text{PaO}_2\text{:FiO}_2$ ($\rho = -0.415$, $p = 0.134$) (Figure E6), compliance ($\rho = 0.062$, $p = 0.820$) or PaCO_2 ($\rho = 0.094$, $p = 0.751$).

The change in volume of the over-aerated compartment was strongly associated with the degree of hyperinflation in the supine position ($\rho = -0.961$, $p < 0.001$) (Figure 2 and Figure E7). It also tended to be associated with the concomitant change in compliance ($\rho = -0.541$, $p = 0.045$) and PaCO_2 ($\rho = 0.491$, $p = 0.085$), but not $\text{PaO}_2\text{:FiO}_2$ ($\rho = -0.130$, $p = 0.648$) (Figure E8).

The change in the heterogeneity of lung aeration along the vertical axis tended to be associated with its value in the supine position ($\rho = -0.468$, $p = 0.076$) but not with the concomitant change in $\text{PaO}_2\text{:FiO}_2$ ($\rho = 0.108$, $p = 0.704$), compliance ($\rho = 0.021$, $p = 0.940$) or PaCO_2 ($\rho = 0.191$, $p = 0.516$).

Discussion

The lung response to prone positioning was variable in patients with early ARDS due to COVID-19. In general, the volume of the non-aerated and over-aerated tissue decreased, and the distribution of aeration became more homogeneous; arterial oxygenation improved, but compliance and PaCO₂ did not.

In ARDS unrelated to COVID-19, prone positioning decreases the alveolar collapse and hyperinflation and homogenizes the distribution of end-expiratory aeration and tidal inflation [3-5]. As a result, mechanical ventilation generates less alveolar deformation and tension and less pulmonary damage [20-25]. This is the strongest rationale for prone positioning in ARDS: making mechanical ventilation safer [6-25]. Increasing arterial oxygenation is probably less important except for the unusual case of life-threatening hypoxemia. In several studies, the survival benefit of prone positioning was not associated with a higher arterial oxygen tension [26-28]. In some of them [26,27], it was instead associated with a lower PaCO₂, a sign of more homogeneous ventilation.

In early ARDS due to COVID-19, the lung morphological response to prone positioning resembled that in other ARDS. Alveolar collapse and hyperinflation decreased, and the distribution of aeration became more homogeneous. As long as the regional gas content is associated with the local distending pressure, the distribution of pleural and transpulmonary pressure became more homogeneous [29-31]. In the supine position, and from sternum to vertebra, the regional gas-to-tissue ratio ranged from 3.1 (2.5-4.0) to 0.1 (0.1-0.2) ml/g; in the prone position, and from vertebra to sternum, from 1.5 (0.8-1.9) to 0.6 (0.2-1.3) ml/g (Figure 3). Therefore, the peak value and dispersion of inflation along the vertical axis were smaller in prone than supine position. Changes in the horizontal distribution of aeration were usually minor. Based on these findings, prone positioning may protect patients with COVID-19 from secondary lung damage [32], as it does in other ARDS.

Several factors probably contributed to redistributing lung aeration with prone positioning. As shown in Figure E5, one of these factors was the superimposed pressure [19]: the gas-to-tissue ratio increased, did not change, or decreased where the superimposed pressure decreased, remained constant, or increased respectively [3]. Other possible factors include (i) the shape of the lung and the chest wall [29,33,34]; (ii) the compression of the lung by the heart and the abdomen [35,36]; (iii) the compliance of the non-dependent and dependent rib cage [37]; and (iv) the distribution of the lung mass along the vertical axis [6].

With prone positioning, arterial oxygenation almost always increased while the volume of the non-aerated lung decreased. Nonetheless, these two responses were unrelated in magnitude. With ARDS, hypoxemia is primarily due to decreased ventilation-to-perfusion ratio, down to zero in the non-aerated lung [38,39]. The improved oxygenation with prone positioning is classically attributed to net recruitment and better ventilation-to-perfusion matching [40,41]. With COVID-19, the distribution of the pulmonary blood flow can be very heterogeneous [42,43], affecting the response to prone positioning. For a given lung recruitment, oxygenation will increase more or less if the newly aerated alveoli are hyper or hypo-perfused. This can be why, in our study population, the reversal of alveolar collapse was not always associated with a proportional increase in arterial oxygenation.

Changes in compliance and PaCO_2 were partly associated with those in hyperinflation. With a larger decrease in the volume of the over-aerated lung compartment, the respiratory system compliance increased. As the chest wall compliance reasonably decreased [37], lung compliance probably increased even more. At the same time, the PaCO_2 tended to decrease in the face of constant minute ventilation. With ARDS, hypercapnia is primarily due to increased ventilation-to-perfusion ratio up to infinity in the over-distended lung [38,39]. A decrease in PaCO_2 may then signal a decrease in the dead space and predict survival [26,27,44]. These data suggest that hyperinflation at lung CT was associated with overdistention and that prone positioning decreased both. However, several poorly predictable factors can confound the interpretation of the functional response to prone positioning in an individual subject. For example, for a given morphological response, the change in the respiratory system compliance will depend on the behaviour of the chest wall and the change in the dead space and PaCO_2 on the distribution of the pulmonary blood flow [6]. If prone positioning ameliorates survival by diminishing the harms of mechanical ventilation, it may be indicated even in patients with large and potentially reversible alveolar collapse and/or hyperinflation at the lung CT, and no better gas exchange or respiratory system mechanics while prone [28].

Hyperinflation is common in patients with COVID-19, even in those ventilated with low tidal volume and airway pressure [16,45,46]. In the seven patients with a larger (than the median) volume of the over-aerated compartment, tidal volume was 6.1 (5.7-6.5) ml/kg of predicted body weight, and plateau airway pressure 23 (21-23) cmH₂O (Table E5). Hyperinflation is a well-known risk factor for secondary damage [47,48]. In our previous study [16], increasing PEEP from 5 to 15 cmH₂O in the supine position decreased the volume of the non-aerated lung by 168 (110-202) ml but increased the volume of the over-aerated lung by 121 (63-270) ml. Hyperinflation increased with a higher PEEP in all (100%) patients. Herein, prone positioning decreased the volume of the non-aerated lung by 82 (26-147) ml and the volume of the over-aerated compartment by 28 (11-186) ml. Hyperinflation *decreased* in all patients but one (93%), especially in those with a larger over-aerated compartment when supine. Therefore, prone positioning may recruit the lung with less hyperinflation than a higher PEEP.

So far, the morphological and functional response to prone positioning in COVID-19 has been investigated only partially [49,50]. In a study conducted with electrical impedance tomography, ventilation increased dorsally and decreased ventrally in the lung section below the belt, while the local ventilation-to-perfusion matching and arterial oxygenation improved [49]. In another study conducted with the CT, the volume of the non-aerated tissue variably decreased dorsally and increased ventrally, in line with the superimposed pressure. A positive response, defined as a rise in $\text{PaO}_2:\text{FiO}_2 \geq 20$ mmHg, was the rule in the early, but not late, phase of the disease, possibly due to the development of pulmonary fibrosis [50]. Our findings complement these previous ones by showing that with prone positioning: (i) aeration is globally more evenly distributed so that harms of mechanical ventilation should be reduced; (ii) a “beneficial” morphological response cannot be predicted from changes in gas exchange and respiratory system mechanics; (iii) the decrease in hyperinflation is frequently larger than recruitment. This can be

particularly important in patients with COVID-19, who are at an increased risk of ventilator-induced lung damage [46].

Some of the limitations of this study deserve a comment. First, the sample size was based on feasibility limitations. During the first pandemic wave, we could not enrol all consecutive eligible patients, which may have been a source of bias (see Table E1). Second, the lung CTs were obtained at end-expiration, and we did not study the distribution of tidal volume in the supine and prone positions. Third, our findings may not be valid for patients with late COVID-19 [50]. Finally, we did not study the impact of prone positioning on patient-centered outcomes, such as survival or duration of mechanical ventilation. Further studies are needed to address these critical issues.

Conclusions

In fifteen mechanically-ventilated patients with early COVID-19, prone positioning variably decreased the amount of alveolar collapse and hyperinflation and improved the distribution of aeration and arterial oxygenation. A similar response has been observed in other ARDS, where prone positioning improves outcome. Therefore, our data provide a pathophysiological rationale to support prone positioning even in COVID-19.

List Of Abbreviations

- ARDS: acute respiratory distress syndrome
- COPD: chronic obstructive pulmonary disease
- COVID-19: coronavirus disease 2019
- CT: computed tomography
- FiO₂: fraction of inspired oxygen
- ICU: intensive care unit
- PaCO₂: arterial carbon dioxide tension
- PaO₂: arterial oxygen tension
- PEEP: positive end-expiratory pressure
- Q1: first quartile
- Q3: third quartile

Declarations

- Ethics approval and consent to participate: this study was approved by our institutional Ethics Committee (Comitato Etico dell'IRCCS Istituto Clinico Humanitas Rozzano; protocol n. 465/20). Informed consent was obtained according to local regulations.
- Consent for publication: not applicable

- Availability of data and materials: the datasets used and analysed during the current study are available from the corresponding author on reasonable request
- Competing interests: the authors declare that they have no competing interests.
- Funding: support was provided solely from institutional and/or departmental sources.
- Authors' contributions: AP takes responsibility for the content of the manuscript, including the data and analysis. AP contributed to the conception and design of the study, analysis, and interpretation of the data and wrote the manuscript. AS contributed to the conception and design of the study, collection, analysis, and interpretation of the data, and revised the manuscript. FP and CC contributed to the collection, analysis, and interpretation of the data and revised the manuscript. MF, GEI, LC, FDC, EL, and NM contributed to the collection of the data and revised the manuscript. AA, and MC contributed to the conception of the study, interpretation of the data, and revised the manuscript. All authors have read and approved this final version of the manuscript.
- Acknowledgements: we are grateful to all nurses, physicians, and other healthcare professionals who helped us perform this study. The following colleagues contributed to the collection, analysis, or interpretation of the data: Valeria Alicino, M.D., Francesca Collino, M.D., Elena Costantini, M.D., Maxim Neganov, M.D., Giulia Paglialunga, M.D., Alberto Parazzoli, N.P., Ilaria Rivetti, M.D., and Valerio Rendiniello, N.P. (Department of Anesthesia and Intensive Care Units, Humanitas Clinical and Research Center – IRCCS, Rozzano, Milan, Italy); Giorgio Picardo, M.D., and Luca Pugliese, M.D. (Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, Italy).

References

1. Fan E, Del Sorbo L, Goligher EC, Hodgson CL, Munshi L, Walkey AJ, et al. An official American Thoracic Society/European Society of Intensive Care Medicine/Society of Critical Care Medicine clinical practice guideline: mechanical ventilation in adult patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2017;195(9):1253-63.
2. Guérin C, Reignier J, Richard JC, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med*. 2013;368(23):2159-68.
3. Gattinoni L, Pelosi P, Vitale G, Pesenti A, D'Andrea L, Mascheroni D. Body position changes redistribute lung computed-tomographic density in patients with acute respiratory failure. *Anesthesiology*. 1991;74(1):15-23.
4. Galiatsou E, Kostanti E, Svarna E, Kitsakos A, Koulouras V, Efremidis SC, et al. Prone position augments recruitment and prevents alveolar overinflation in acute lung injury. *Am J Respir Crit Care Med*. 2006;174(2):187-97.
5. Cornejo RA, Díaz JC, Tobar EA, Bruhn AR, Ramos CA, González RA, et al. Effects of prone positioning on lung protection in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2013;188(4):440-8.
6. Guérin C, Albert RK, Beitler J, Gattinoni L, Jaber S, Marini JJ, et al. Prone position in ARDS patients: why, when, how and for whom. *Intensive Care Med*. 2020;46(12):2385-96.

7. Gattinoni L, Carlesso E, Taccone P, Polli F, Guérin C, Mancebo J. Prone positioning improves survival in severe ARDS: a pathophysiologic review and individual patient meta-analysis. *Minerva Anestesiol.* 2010;76(6):448-54.
8. Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med.* 2020;46(5):854-87.
9. Fan E, Beitler JR, Brochard L, Calfee CS, Ferguson ND, Slutsky AS, et al. COVID-19-associated acute respiratory distress syndrome: is a different approach to management warranted? *Lancet Respir Med.* 2020;8(8):816-21.
10. Gattinoni L, Chiumello D, Rossi S. COVID-19 pneumonia: ARDS or not? *Crit Care.* 2020;24(1):154.
11. Gattinoni L, Chiumello D, Caironi P, Busana M, Romitti F, Brazzi L, et al. COVID-19 pneumonia: different respiratory treatments for different phenotypes? *Intensive Care Med.* 2020;46(6):1099-102.
12. Langer T, Brioni M, Guzzardella A, Carlesso E, Cabrini L, Castelli G, et al. Prone position in intubated, mechanically ventilated patients with COVID-19: a multi-centric study of more than 1000 patients. *Crit Care.* 2021;25(1):128.
13. Camporota L, Sanderson B, Chiumello D, Terzi N, Argaud L, Rimmelé T, et al. Prone position in coronavirus disease 2019 and non-coronavirus disease 2019 acute respiratory distress syndrome: an international multicenter observational comparative study. *Crit Care Med.* 2021(September).
14. ARDS Definition Task Force, Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA.* 2012;307(23):2526-33.
15. Gattinoni L, Caironi P, Cressoni M, Chiumello D, Ranieri VM, Quintel M, et al. Lung recruitment in patients with the acute respiratory distress syndrome. *N Engl J Med.* 2006;354(17):1775-86.
16. Protti A, Santini A, Pennati F, Chiurazzi C, Cressoni M, Ferrari M, et al. Lung response to a higher positive end-expiratory pressure in mechanically-ventilated patients with COVID-19. *Chest.* 2021(October).
17. Cressoni M, Gallazzi E, Chiurazzi C, Marino A, Brioni M, Menga F, et al. Limits of normality of quantitative thoracic CT analysis. *Crit Care* 2013;17(3):R93.
18. Arachchige CNPG, Prendergast LA, Staudte RG. Robust analogs to the coefficient of variation. *Journal of Applied Statistics.* 2020(August).
19. Pelosi P, D'Andrea L, Vitale G, Pesenti A, Gattinoni L. Vertical gradient of regional lung inflation in adult respiratory distress syndrome. *Am J Respir Crit Care Med.* 1994;149(1):8-13.
20. Broccard A, Shapiro RS, Schmitz LL, Adams AB, Nahum A, Marini JJ. Prone positioning attenuates and redistributes ventilator-induced lung injury in dogs. *Crit Care Med.* 2000;28(2):295-303.
21. Valenza F, Guglielmi M, Maffioletti M, Tedesco C, Maccagni P, Fossali T, et al. Prone position delays the progression of ventilator-induced lung injury in rats: does lung strain distribution play a role? *Crit Care Med.* 2005;33(2):361-7.

22. Mentzelopoulos SD, Roussos C, Zakynthinos SG. Prone position reduces lung stress and strain in severe acute respiratory distress syndrome. *Eur Respir J*. 2005;25(3):534-44.
23. Xin Y, Cereda M, Hamedani H, Pourfathi M, Siddiqui S, Meeder N, et al. Unstable inflation causing injury. Insight from prone position and paired computed tomography scans. *Am J Respir Crit Care Med*. 2018;198(2):197-207.
24. Motta-Ribeiro GC, Hashimoto S, Winkler T, Baron RM, Grogg K, Paula LFSC, et al. Deterioration of regional lung strain and inflammation during early lung injury. *Am J Respir Crit Care Med*. 2018;198(7):891-902.
25. Gattinoni L, Pesenti A, Carlesso E. Body position changes redistribute lung computed-tomographic density in patients with acute respiratory failure: impact and clinical fallout through the following 20 years. *Intensive Care Med*. 2013;39(11):1909-15.
26. Gattinoni L, Vagginelli F, Carlesso E, Taccone P, Conte V, Chiumello D, et al. Decrease in PaCO₂ with prone position is predictive of improved outcome in acute respiratory distress syndrome. *Crit Care Med*. 2003;31(12):2727-33.
27. Charron C, Repesse X, Bouferrache K, Bodson L, Castro S, Page B, et al. PaCO₂ and alveolar dead space are more relevant than PaO₂/FiO₂ ratio in monitoring the respiratory response to prone position in ARDS patients: a physiological study. *Crit Care*. 2011;15(4):R175.
28. Albert RK, Keniston A, Baboi L, Ayzac L, Guérin C; Proseva Investigators. Prone position-induced improvement in gas exchange does not predict improved survival in the acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2014;189(4):494-6.
29. Tawhai MH, Nash MP, Lin CL, Hoffman EA. Supine and prone differences in regional lung density and pleural pressure gradients in the human lung with constant shape. *J Appl Physiol* (1985). 2009;107(3):912-20.
30. Terzi N, Bayat S, Noury N, Turbil E, Habre W, Argaud L, et al. Comparison of pleural and esophageal pressure in supine and prone positions in a porcine model of acute respiratory distress syndrome. *J Appl Physiol* (1985). 2020;128(6):1617-25.
31. Katira BH, Osada K, Engelberts D, Bastia L, Damiani LF, Li X, et al. Positive end-expiratory pressure, pleural pressure, and regional compliance during pronation: an experimental study. *Am J Respir Crit Care Med*. 2021;203(10):1266-74.
32. Mathews KS, Soh H, Shaefi S, Wang W, Bose S, Coca S, et al. Prone positioning and survival in mechanically-ventilated patients with coronavirus disease 2019-related respiratory failure. *Crit Care Med*. 2021;49(7):1026-37.
33. Hubmayr RD. Perspective on lung injury and recruitment: a skeptical look at the opening and collapse story. *Am J Respir Crit Care Med*. 2002;165(12):1647-53.
34. Gattinoni L, Taccone P, Carlesso E, Marini JJ. Prone position in acute respiratory distress syndrome. Rationale, indications, and limits. *Am J Respir Crit Care Med*. 2013;188(11):1286-93.

35. Rouby JJ, Puybasset L, Nieszkowska A, Lu Q. Acute respiratory distress syndrome: lessons from computed tomography of the whole lung. *Crit Care Med*. 2003;31(4 Suppl):S285-95.
36. Albert RK, Hubmayr RD. The prone position eliminates compression of the lungs by the heart. *Am J Respir Crit Care Med*. 2000;161(5):1660-65.
37. Pelosi P, Tubiolo D, Mascheroni D, Vicardi P, Crotti S, Valenza F, et al. Effects of the prone position on respiratory mechanics and gas exchange during acute lung injury. *Am J Respir Crit Care Med*. 1998;157(2):387-93.
38. Radermacher P, Maggiore SM, Mercat A. Fifty years of research in ARDS. Gas exchange in acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2017;196(8):964-84.
39. Petersson J, Glenny RW. Gas exchange and ventilation-perfusion relationships in the lung. *Eur Respir J*. 2014;44(4):1023-41.
40. Lamm WJ, Graham MM, Albert RK. Mechanism by which the prone position improves oxygenation in acute lung injury. *Am J Respir Crit Care Med*. 1994;150(1):184-93.
41. Richter T, Bellani G, Scott Harris R, Vidal Melo MF, Winkler T, Venegas JG, et al. Effect of prone position on regional shunt, aeration, and perfusion in experimental acute lung injury. *Am J Respir Crit Care Med*. 2005;172(4):480-7.
42. Ackermann M, Verleden SE, Kuehnel M, Haverich A, Welte T, Laenger F, et al. Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in Covid-19. *N Engl J Med*. 2020;383(2):120-28.
43. Patel BV, Arachchillage DJ, Ridge CA, Bianchi P, Doyle JF, Garfield B, et al. Pulmonary angiopathy in severe COVID-19: physiologic, imaging, and hematologic observations. *Am J Respir Crit Care Med*. 2020;202(5):690-99.
44. Nuckton TJ, Alonso JA, Kallet RH, Daniel BM, Pittet JF, Eisner MD, et al. Pulmonary dead-space fraction as a risk factor for death in the acute respiratory distress syndrome. *N Engl J Med*. 2002;346(17):1281-86.
45. Rezoagli E, Bastia L, Grassi A, Chiericato A, Langer T, Grasselli G, et al. Paradoxical effect of chest wall compression on respiratory system compliance: a multicenter case series of patients with ARDS, with multimodal assessment. *Chest*. 2021;160(4):1335-39.
46. Protti A, Greco M, Filippini M, Vilaro AM, Langer T, Villa M, et al. Barotrauma in mechanically-ventilated patients with Coronavirus disease 2019: a survey of 38 hospitals in Lombardy, Italy. *Minerva Anestesiol*. 2021(2);87:193-98.
47. Dreyfuss D, Saumon G. Ventilator-induced lung injury: lessons from experimental studies. *Am J Respir Crit Care Med*. 1998;157(1):294-323.
48. Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med*. 2013;369(22):2126-36.
49. Perier F, Tuffet S, Maraffi T, Alcalá G, Victor M, Haudebourg AF, et al. Effect of positive end-expiratory pressure and proning on ventilation and perfusion in COVID-19 acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2020;202(12):1713-17.

50. Rossi S, Palumbo MM, Sverzellati N, Busana M, Malchiodi L, Bresciani P, et al. Mechanisms of oxygenation responses to proning and recruitment in COVID-19 pneumonia. *Intensive Care Med.* 2022;48(1):56-66.

Table

Table 1. Characteristics of the study population at ICU admission

Variable	Study population
N	15
Males (n [%])	11 (73)
Age (years)	69 (65-74)
Body mass index (BMI) (kg/m ²)	29 (25-31)
Tidal volume (ml)	400 (400-435)
Tidal volume (ml/kg of PBW)	6.4 (6.0-7.1)
Respiratory rate (bpm)	18 (16-22)
PEEP (cmH ₂ O)	12 (10-15)
FiO ₂ (%)	70 (60-88)
Minute ventilation (L/min)	7.6 (6.4-9.0)
Plateau airway pressure (cmH ₂ O)	23 (18-25)
Driving airway pressure (cmH ₂ O)	9 (7-12)
Compliance (ml/cmH ₂ O)	49 (35-58)
Arterial pH	7.37 (7.31-7.40)
PaCO ₂ (mmHg)	55 (43-61)
PaO ₂ (mmHg)	83 (71-108)
PaO ₂ /FiO ₂ (mmHg)	123 (91-139)
ICU length of stay (days)	20 (11-42)
Mortality in ICU (n [%])	6 (40)

All data refer to the time of admission to our Intensive Care Unit (ICU), except for ICU length of stay and mortality in ICU. BMI: body mass index. PBW: predicted body weight. PEEP: positive end-expiratory

pressure. FiO_2 : inspiratory fraction of oxygen. PaCO_2 : arterial tension of carbon dioxide. PaO_2 : arterial tension of oxygen. The driving airway pressure was the difference between the plateau airway pressure and total PEEP measured with a 5-second end-inspiratory and end-expiratory pause. The compliance was the ratio of the tidal volume to the driving airway pressure. Data are reported as median (Q1-Q3) or proportion.

Figures

Figure 1

Individual morphological response to prone positioning.

Fifteen mechanically-ventilated patients with COVID-19 underwent a lung computed tomography in the supine and prone position. Panels A-D: we describe the individual changes in the total (tissue and gas) volume of the non-aerated (density above -100 HU), poorly-aerated (from -100 to -500 HU), normally-

aerated (from -500 to -900 HU), and over-aerated (below -900 HU) compartments with prone positioning, in descending order. Panels E-F: each CT slice was divided into ten equal vertical levels, from the sternum (vertical level 1) to the vertebra (vertical level 10), and in ten equal horizontal levels, from the apex (horizontal level 1) to the base (horizontal level 10) of the lung. Herein we describe the individual change in the degree of heterogeneous aeration along the vertical and horizontal axis, expressed with the standardized median absolute deviation of regional gas-to-tissue ratios, and presented in descending order. Negative values indicate that the volume of a given compartment or the degree of heterogeneity decreased with prone positioning. Each bar refers to one patient. The same letter in the six panels refers to the same patient.

Figure 2

Colour-coded analysis of lung computed tomography (CT) data.

Representative CT images taken at the level of carina from three patients with COVID-19 in the supine and prone position, with a very large decrease in the volume of the over-aerated compartment in response

to prone positioning. Upper panels: original lung CT images, with aeration shown on a continuous grayscale. Lower panels: using an automated encoding system, we attributed a specific colour to the non-aerated, poorly-aerated, normally-aerated, and over-aerated compartments. The three patients are identified with the same letters as in other figures. With prone positioning, the volume of the over-aerated lung decreased from 318 to 121 ml in patient J; from 738 to 148 ml in patient C; from 503 to 230 ml in patient F.

Figure 3

Regional lung gas-to-tissue ratio in the supine and prone position.

Fifteen mechanically-ventilated patients with COVID-19 underwent a lung computed tomography (CT) in supine and prone positions. Each CT slice was divided into ten equal vertical levels, from the sternum (vertical level 1) to the vertebra (vertical level 10), and in ten equal horizontal levels, from the apex (horizontal level 1) to the base (horizontal level 10) of the lung. Herein we describe the ratio of the gas volume (ml) to the tissue weight (g) in each of those levels, in the supine and prone positions. Data are reported as median (Q1-Q3). Panel A: vertical gradient of lung inflation. On average, the gas-to-tissue ratio decreased by 0.34 (0.25-0.49) ml/g per level in the supine position and by 0.03 (-0.11-0.14) ml/g in the prone position ($p<0.001$). Panel B: horizontal gradient of lung inflation. On average, the gas-to-tissue ratio decreased by 0.14 (0.04-0.27) ml/g per level in the supine position and by 0.11 (-0.05-0.21) ml/g in the prone position ($p=0.003$).

Figure 4

Individual functional response to prone positioning.

Fourteen mechanically-ventilated patients with COVID-19 were evaluated in the supine and prone positions. Herein we describe the response to prone positioning in terms of change in arterial oxygenation (expressed as the ratio of the arterial tension to the inspiratory fraction of oxygen [$\text{PaO}_2:\text{FiO}_2$]) (n=14) (panel A), respiratory system compliance (n=14) (panel B), and carbon dioxide tension (PaCO_2) for the same minute ventilation (n=13) (panel C), in descending order. Each bar refers to one patient. The same letter in the 3 panels refers to the same patient. Please note that patient D, shown in other figures, did not undergo prone positioning and is not shown in this figure. FiO_2 was decreased in the prone position in patient K and patient F, and increased in patient G. The impact of prone positioning on PaCO_2 could not be assessed in patient L because his minute ventilation was increased during prone positioning.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Onlinedatasupplement.docx](#)