Prevalence of HIV, syphilis, and assessment of the social and structural determinants of sexual risk behaviour and health service utilisation among MSM and TG in Terai highway districts of Nepal: Findings based on an Integrated Biological and Behavioural Surveillance survey using respondent driven sampling

Margrethe Storm

Karolinska Institutet

Keshab Deuba (✉ keshab.deuba@savethechildren.org)

National Centre for AIDS and STD Control <https://orcid.org/0000-0002-9933-2210>

Jose Damas

Karolinska Institutet

Upendra Shrestha

Save the Children

Bir Rawal

National Centre for AIDS and STD Control

Rajan Bhattarai

Save the Children

Gaetano Marrone

Karolinska Institutet

Research article

Keywords: MSM, TG, HIV, STI, Nepal, sexual risk behaviour, health care service uptake, structural factors

Posted Date: June 13th, 2019

DOI: <https://doi.org/10.21203/rs.2.10259/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on June 8th, 2020. See the published version at <https://doi.org/10.1186/s12879-020-05122-3>.

Abstract

Background: MSM and TG in Nepal experience considerable discrimination and marginalisation: they are subject to abuse from legal authorities and suffer from mental health issues and substance abuse. This study aims to assess the prevalence of HIV and syphilis, in addition to how individual and socio-structural factors influence sexual risk behaviour and health care service uptake, among MSM and TG in the Terai highway districts of Nepal. **Methods:** A cross-sectional survey was conducted in June 2016 among MSM and TG in eight Terai highway districts of Nepal, recruiting 340 MSM and TG through respondent driven sampling. The primary outcome variables were HIV and syphilis prevalence. The secondary outcome variables were sexual risk behaviour and health care service uptake. Logistic regression models were used to assess the individual and socio-structural determinants of sexual risk behaviour and health care service uptake. **Results:** The prevalence of HIV among MSM was 5%, whereas it was 13% in TG. The prevalence of active syphilis was 4% in the MSM group and 11% among TG. Among TG, 76% were involved in sex work, and 51% reported having experienced discrimination in one or more settings. In multivariable analysis, having visited an outreach centre was positively associated with condom use in the most recent sexual encounter among both MSM (AOR: 5.37, 95% CI: 2.42 - 11.94, $p < 0.001$) and TG (AOR: 2.81, 95% CI: 1.30 – 6.09, $p = 0.009$). Moreover, in the multivariable regression model assessing factors associated with visited outreach center among MSM, respondents that reported they had been forced to marry a female were 3 times more likely to have visited an outreach centre. Among TG, being open towards family about sexual identity/behaviour were positively associated with having visited an outreach centre. **Conclusions:** The high prevalence levels of HIV and syphilis, as well as indicators of marginalisation and discrimination among TG, highlights the increased burden TG in Nepal are facing and the need for tailored interventions for this group. Moreover, since health care service uptake seems to be an important factor in determining sexual risk behaviour among MSM and TG in Nepal, outreach services should be scaled up.

Background

HIV/AIDS remains an important issue in the world with 1.8 million estimated new infections in 2017 [1]. Key populations such as men who have sex with men (MSM) and transgender (TG) are disproportionately affected by the HIV epidemic and have a higher risk of infection either due to a higher level of risk behaviour, or their vulnerable position in society [2]. MSM is an epidemiological term to describe men who engage in sex with other men regardless of their sexual orientation or gender identity, whereas the term TG is used to describe individuals whose gender identity and/or expression does not correspond with the sex assigned to them at birth [2]. Globally, MSM are estimated to be 19 times more likely to be infected with HIV than the general population and HIV prevalence rates in TG are as high as 40% in some settings [3, 4]. In the South East Asia region, five countries (India, Indonesia, Myanmar, Nepal, and Thailand) account for 99% of the regional HIV burden [5].

Nepal is a low-income country with a population of 29 million people and consists of a mountainous region in the north and open terrain (Terai in local language) in the south [6]. Nepal has in the last 15 years experienced a decline in HIV prevalence, however, the HIV epidemic has evolved from being a “low prevalence” to a “concentrated epidemic”, i.e. HIV transmission largely occurs in one or more defined subpopulation but is not well established in the general population [7, 8]. In 2017, Nepal had an estimated adult HIV prevalence of

0.15% [7]. The key affected populations in Nepal include MSM, TG, people who inject drugs (PWID), male sex workers (MSW), female sex workers (FSW), and male labour migrants and their spouses [7]. Similarly to HIV, sexually transmitted infections (STIs) are also disproportionately represented in the key populations in Nepal [7, 9]. Nepal has in the last decade observed a decline in HIV prevalence among young MSM/TG in the Kathmandu valley which coincided with an increase in condom use in this group [10]. In 2017, the prevalence of HIV was 4.8% and 8.6% among MSM and TG, respectively, in the Kathmandu Valley [7].

The studies to date in Nepal have only focused on MSM and TG in the Kathmandu Valley, however key populations are scattered around the country, and there is limited data on MSM and TG in hard to reach areas of Nepal. The Terai highway districts of Nepal have a high population density and share an open border with India in the south, leading to an increased risk of transmission of HIV and STIs due to sex work and drug trafficking across the border [11]. Furthermore, there is limited data on TG in Nepal both with regards to health indicators, as well as estimates of population numbers, and TG is often subsumed under the MSM term. Although MSM and TG share a number of challenges, several studies indicate that TG faces additional individual and socio-structural obstacles and experience a higher burden of adverse health conditions than MSM [4, 12].

Sexual risk behaviour is now recognized as multifaceted and composed of not only individual, but also social and structural factors such as social networks, stigma, discrimination, policies and laws [13, 14]. MSM and TG in Nepal face several social and structural challenges that significantly influence their health and standing in society, such as discrimination, lack of adequate legal protection, reduced social capital, and mental health issues [15-17]. These conditions can, either alone or together, lead to barriers to access to health care, increased sexual risk behaviour, and ultimately poor health outcomes [16, 17].

Nepal has made some progress in terms of the rights of lesbian, gay, bisexual and transgender (LGBT) with the Supreme Court ruling of 2007, which promoted the abolishment of discriminatory laws against sexual and gender minorities, same-sex marriage, and the official recognition of the third gender [9]. Furthermore, the new constitution of 2015 ensures the rights of sexual and gender minorities [15]. A number of non-governmental organisations also advocate the rights and health needs of the Nepalese LGBT community with the most prominent being the Blue Diamond Society (BDS) [9]. However, the implementation of the third gender, same-sex marriages, and prevention of discrimination in private and public settings are still subject to controversy [16, 18]. A study which assessed risk of HIV infection among MSM in Kathmandu found that a large proportion of the survey participants had experienced verbal, physical and sexual abuse which was associated with increased sexual risk behaviour in the bivariate analysis [19]. Furthermore, a recent study investigating the psychosocial well-being of MSM/TG found a high prevalence of mental health issues which was associated with increased sexual risk behaviour [18]. Sexual risk behaviour was connected to low social support, experienced violence and not participating in HIV prevention [18]. Another study that explored the underpinnings of psychosocial issues among MSM/TG in Nepal found that suicidal ideation was associated with perceived discrimination and increased with the extent of perceived discrimination [20].

Despite extensive research into the individual risk determinants of HIV/STIs among MSM and TG in Nepal, there is a lack of quantitative studies describing how socio-structural factors influence sexual risk behaviour and health service uptake among MSM and TG in Nepal. The objective of this study was to assess the HIV

and STIs prevalence, sexual risk behaviour, health service uptake, and how it relates to individual and socio-structural factors among MSM and TG in the Terai highway districts in order to obtain information for planning of policies and programs for treatment and prevention of HIV and STIs in Nepal.

Methods

Study design and setting

This study was based on secondary analysis of data from a cross-sectional Integrated Biological and Behavioural Surveillance (IBBS) survey conducted in the Terai highway districts of Nepal in 2016 using respondent driven sampling (RDS). This study adheres to the STROBE-RDS guidelines for reporting of respondent-driven sampling studies [21]. The IBBS survey covered eight districts of the Terai region; Jhapa, Morang, Sunsari, Nawalparasi, Rupandehi, Kapilbastu, Kailali, and Kanchanpur. The districts were chosen based on previous mapping and size estimations reports of MSM and TG, as well as stakeholders' suggestions. Survey facilities and clinics were set up in Itahari of Sunsari district, Bhairahawa of Rupandehi district, and Dhangadhi of Kailali district. The National Centre for AIDS and STD Control (NCASC) conducted the survey with technical and financial support from Save the Children International/ Global Fund Programs. A formative assessment was carried out prior to the survey to ensure that the sampling strategy, survey protocol, as well as sample size estimates were appropriate. Personal network size was assessed by the question: "How many MSM/TG do you know who also know you, who have been involved in sexual activities with men in the past 3 months? How many of these MSM/TG have you seen in the past month?". Additional details of the survey design have been published in the IBBS report [22].

Study population

Individuals were eligible to participate if they were 1) biological males (including participants who either identified themselves in a different gender than that assigned to them at birth or identified themselves belonging to transgender community), 2) aged 16 years or above, 3) reported they had sexual relations (either oral or anal) with another biological male at least once in the past year prior to the survey, 4) resided in one of the eight selected districts of the Terai highway, 5) had a valid referral coupon and 6) had not previously participated in the survey. The definition of a sex worker in the survey comprised biological males who had anal or oral sex with another male in exchange for money or other items in the past year prior to the survey.

Sampling strategy

Participants were recruited using respondent driven sampling as both MSM and TG are hard to reach populations [23]. Starting with the purposive sampling of eight seeds from each of the selected survey districts, an additional seed was later added at the Bhairahawa facility to increase coverage. The seeds were selected to represent a diverse sample of sexual identities, age, caste, education, and location sites of MSM and TG in Nepal. Each seed received three coupons for referral and one reward coupon. The referral coupons had unique ID numbers to be able to track recruiters and recruits. Participants recruited by seeds who enrolled in the study were given 3 coupons for further recruitment. Out of the nine seeds, three seeds generated five

recruitment waves, whereas six seeds generated four waves. Respondents were given a financial incentive of 300 Nepalese rupees for participation and 200 Nepalese rupees as a reward for each additional participant they recruited. The IBBS survey was initiated in April 2016 and completed in August 2016. Both biological and behavioural data were collected in June 2016.

Sample size

The following formula was used to calculate sample size in each round of IBBS surveys to detect a 15% change in key indicators [24]:

$$n = D Z_{1-\alpha}^2 P_{hat} (1 - P_{hat}) + Z_{1-\beta}^2 P_1 (1 - P_1) + P_2 (1 - P_2) / P_1^2$$

n = Sample size required per survey round; D = Design effect of 2; $Z_{1-\alpha}$ = The z score for the desired confidence level, 1.960 for 95% confidence level for a two-sided test; $Z_{1-\beta}$ = The z score for the desired power, 0.84 for 80%; P_1 = the estimated proportion at the time of the first survey (it was .78 for non-use of condoms at their last anal sex with another man and .51 for knowledge of HIV/AIDS etc.) ; P_2 = The magnitude of change we want to be able to detect, in our case it is 15%; $P_{hat} = (P_1 + P_2) / 2$.

Measurements

Behavioural data were collected using a structured questionnaire that included questions related to socio-demographic characteristics, sexual risk behaviour, knowledge on HIV and STIs, mobility, experiences of stigma and discrimination, and health service uptake. The questionnaire was pre-tested prior to the survey and a pilot to assess the survey tools was conducted. The questionnaire used can be found in the published IBBS report [22]. The primary outcome variables were HIV and syphilis prevalence. The secondary outcome variables were sexual risk behaviour as assessed by condom use in last sexual encounter and health service uptake which was measured by the question "Have you visited or been to any outreach centre (Drop-In Centre (DIC), Information Centre (IC), Counseling Centre (CC)) in the last 12 months?". HIV knowledge was assessed by computing a knowledge score which was dichotomized into low and high knowledge. The variables selected for the generation of the score were based on previously published studies from Nepal [25, 26] and consisted of: 1) Can people reduce their risk of HIV by using a condom correctly every time they have sex?; 2) Can a person get HIV from mosquito bites?; 3) Can people protect themselves from HIV by having one uninfected faithful sex partner?; 4) Can people protect themselves from HIV by abstaining from sexual intercourse?; 5) Can a person get HIV by sharing meal with someone who is infected?; 6) Can a person get HIV by using a needle that is used by someone else?; 7) Do you think that a healthy-looking person can be infected with HIV, the virus that causes AIDS?; 8) Can a pregnant woman infected with HIV transmit the virus to her unborn child?; 9) Can women with HIV transmit the virus to her newborn child through breastfeeding?.

Discrimination was assessed based on the Experience of Discrimination (EOD) tool [27], which entails a self-reported experience of discrimination in eight different situations, in addition to the reaction to being treated unfairly. The eight situations included school, getting hired, work, getting housing, getting medical care, getting service in store/ restaurant, public settings, and from police or other security personnel and has previously been used in the Nepalese setting [20]. "No response"/"Don't remember"/"Don't know" were set to missing.

Laboratory methods

After completion of the questionnaire, participants were tested for HIV and syphilis according to Nepal's HIV testing and counselling protocol. The survey participants underwent counselling prior to and post blood sample collection, along with a clinical examination for STIs symptoms and a basic health examination. On-site rapid screening for HIV1/2 was performed using Determine HIV 1/2 (Abbott, Japan). Reactive samples were confirmed using Uni-Gold HIV ½ (Trinity Biotech, Ireland). Samples that were negative on the first test were considered HIV negative, while samples that tested positive on the two first initial tests were considered HIV positive. Any discrepant results from the two first tests were retested with Stat-Pak HIV 1/2 (Chembio Diagnostic system, Inc. USA). Syphilis was tested using Rapid Plasma Reagin (RPR) test (Becton, Dickson, and company, USA). If the first test was positive the sample was retested using Serodia Treponema Pallidum Particle Agglutination (TPPA) test (Fujirebio Inc.). Based on the RPR test a titer value $\geq 1:8$ was categorized as current active syphilis infection, while a titer of $<1:8$ was considered as having a history of syphilis. Symptomatic treatment for STIs was given to respondents following the National STI Case Management Guidelines [22].

Statistical analysis

Data analysis was performed using IBM SPSS Statistics for Windows, Version 24.0 (Armonk, NY: IBM Corp.) and STATA version 14 (StataCorp, College Station, Texas, USA). Data were summarized with descriptive statistics (mean, median, and standard deviation for numerical variables, frequencies and percentages for categorical variables) using both RDS-adjusted and unadjusted values. Values were adjusted for the RDS sampling method using the RDS II estimator. A correct answer to one of the HIV knowledge questions was given the value 1, while a wrong answer was given 0. Scores between 0-5 were categorised as low knowledge, while scores of 6-9 were categorised as high knowledge. A discrimination score was generated indicating discrimination in any of the eight situations and dichotomized into never experienced discrimination and experienced discrimination in one or more situations. Bivariate associations between independent categorical variables and sexual risk behaviour and health care service uptake were calculated using Chi-Square or Fisher's exact test. For independent numerical variables, Mann Whitney U test was used to compare medians in two groups. P-values of <0.25 were set as the criterion for inclusion in the regression model. Both backward and forward stepwise logistic regression models, as well as manual entering of the variables in the model, were used to identify significant predictors of binary outcomes. All methods used gave the same final results. Entry level into the model was set to 0.1, while exit was 0.11. Crude odds ratio (OR) and adjusted odds ratios (AOR) with their 95% confidence intervals (CI) are presented. A p-value <0.05 was considered significant in the final models.

Results

General characteristics of the study sample

Sociodemographic characteristics of the study participants stratified by MSM and TG are shown in Table 1. Out of the 340 respondents, 173 (61%) identified as transgender. The mean age ($\pm$ SD) among MSM was 26.1

(± 9.2) years whereas the mean age among TG was 31.3 (± 10.5). Among the 167 respondents that identified as MSM, 20% were sex workers, whereas in the TG group 76% reported being sex workers. Out of the respondents that identified as MSM, 90% reported to have attended school, whereas only 21% of TG had received any education. Almost 1/4 of MSM were married, while among TG 40% reported being married.

HIV and syphilis prevalence among MSM and TG

The prevalence of HIV among MSM was 5%, whereas it was 13% among TG. The prevalence of active syphilis was 4% in the MSM group and 11% among TG (Figure 1).

Sexual risk behaviour, HIV knowledge, socio-structural factors and uptake of HIV services

Among MSM, 61% reported to have used condoms in their last sexual intercourse (Table 2). A relatively high proportion of MSM (81%) and TG (76%) had a high level of knowledge of HIV risk and transmission. In the MSM group, 25% reported that their family forced them to marry a female and 30% had been away from their home the past month. Among TG, 37% reported that their family forced them to marry a female, and 51% had experienced discrimination in one or more situations. In the TG group, 55% had interacted with PE/OE/CE/CM and 54% had visited outreach centre in the past year. The uptake of HIV testing and counselling (HTC) services among MSM and TG was very low, only 13% of MSM and 22% of TG had visited an HTC centre in the last 12 months.

Individual and socio-structural factors associated with sexual risk behaviour in MSM and TG

Among MSM, the multiple logistic regression model revealed a negative association between having an income above 10000 Nepalese rupees and condom use in the most recent sexual encounter (AOR: 0.36, 95% CI: 0.14 - 0.92, $p=0.032$) (Table 3). In contrast, TG that had an income between 3000-10000 rupee were more likely to have used condom in their last sexual intercourse (AOR: 2.79, 95% CI: 1.12 – 6.91, $p=0.027$) (Table 4). There was a negative association between being forced by family to marry a female and condom use during last sexual intercourse among both MSM (AOR: 0.42, 95% CI: 0.19 - 0.95, $p=0.037$) and TG (AOR: 0.42, 95% CI: 0.21 - 0.88, $p=0.021$) (Table 3 and 4). Moreover, respondents that identified as TG and reported being away from home the past month were 2.5 more likely to have used condoms in their last sexual intercourse (AOR: 2.46, 95% CI: 1.13 - 5.38, $p=0.024$) (Table 4). TG that had a moderate (AOR: 0.31, 95% CI: 0.13 - 0.74, $p=0.008$), and an active response (AOR: 0.29, 95% CI: 0.12 - 0.68, $p=0.005$) to being treated unfairly were less likely to have used condom in their last sexual encounter. In multivariable analysis, having visited an outreach centre was positively associated with condom use in the most recent sexual encounter among both MSM (AOR: 5.37, 95% CI: 2.42 - 11.94, $p<0.001$) and TG (AOR: 2.81, 95% CI: 1.30 – 6.09, $p=0.009$).

Individual and socio-structural factors associated with visiting outreach centre among MSM and TG

In the previous multivariable regression models for MSM and TG, having visited an outreach centre in the past year was significantly associated with sexual risk behaviour (Table 3 and 4). The determinants of health care service uptake among MSM and TG were thus further examined. In the final model having a high level of knowledge of HIV risk and transmission (AOR: 3.33, 95% CI: 1.22 – 9.11, $p=0.019$) was positively associated with having visited an outreach centre in the last 12 months among MSM (Supplementary Table 1). Moreover, MSM who reported that they had been forced to marry a female were 3 times more likely to have visited an outreach centre (AOR: 3.08, 95% CI: 1.49 – 6.38, $p=0.002$). Moreover, having been cheated or threatened due to sexual behaviour (AOR: 3.87, 95% CI: 1.45 – 10.36, $p=0.007$) was associated with increased health care service uptake. Multivariable regression revealed that TG with comprehensive knowledge on HIV were more likely to have visited outreach center (AOR: 6.03, 95% CI: 2.48 – 14.66, $p<0.001$) (Supplementary Table 2). Furthermore, openness to family about sexual behaviour (AOR: 2.40, 95% CI: 1.04 – 5.57, $p=0.041$) and having been cheated or threatened due to sexual behaviour (AOR: 2.36, 95% CI: 0.95 - 5.89, $p=0.065$) were positively associated with having visited outreach center.

Discussion

This study assessed the HIV and syphilis prevalence along with factors that contribute to sexual risk behaviour and health care service uptake among MSM and TG in Terai highway districts of Nepal. In this study, the prevalence of HIV among MSM was 5%, whereas it was 13% in TG. The prevalence of syphilis was also markedly higher among TG (11%) compared to MSM (4%). Overall, TG had a higher prevalence of socio-structural factors concerning discrimination and marginalisation compared to MSM. These results are consistent with previous studies which have shown that TG bear a substantial part of the HIV burden compared to MSM, and that TG faces considerable stigma and discrimination [4, 29]. Furthermore, these findings highlight the importance and need for disaggregation of data by gender identity both in terms of research, but also monitoring.

In this study, approximately $\frac{3}{4}$ of TG were involved in sex work. Due to the stigma and discrimination TG experience they often have difficulties finding employment and thus resort to sex work as a means of income [30]. Being engaged in sex work can result in additional stigma and increased HIV vulnerability. Future initiatives should address the aspect of stigma and employment among TG in Nepal. Overall, the prevalence of HIV and syphilis was higher in this survey than what was observed in the latest IBBS surveys conducted among MSM and TG in the Kathmandu valley [7, 31]. This could be due to, at least in part, its location, the Terai highway district area. The open border with India, high population numbers, as well as sex work and drug trafficking across the border makes this area particularly vulnerable to HIV and STIs spread [11].

Individual and socio-structural factors associated with sexual risk behaviour among MSM and TG

Respondents that identified as MSM and had an income above 10000 Nepalese rupees had a lower likelihood of using a condom in the last sexual encounter compared to respondents that earned less. In contrast, TG that earned between 3000-10000 rupee were more likely to have used condoms in their last sexual intercourse. Studies have shown both low and high income to be associated with more unprotected sex and the underlying reason behind the observed association is not clear [32-34]. Controlling for the other covariates in the model, being forced by family to marry a female was associated with lack of condom use during the last sexual encounter among both MSM and TG. This increased sexual risk behaviour can, on the one hand, be seen as a product of the stigmatisation and discrimination that MSM and TG in Nepal experiences and is consistent with the proposed pathway in which how structural violence such as stigma leads to sexual risk behaviour [13]. Studies from China and Cambodia have highlighted the strong societal and familial pressures on LGBT to marry and suggested it can lead to increased sexual risk behaviours [35-37]. Family rejection and pressure to marry can cause psychosocial problems and also result in MSM and TG individuals to run away from home which leads to increased sexual risk behaviour, including engaging in survival sex [17, 37]. However, this association could also be explained by that respondents who report to be forced by family to marry are in fact married and thus less likely to use condoms in sexual encounters with their wives. Moreover, among both MSM and TG, having visited an outreach centre was positively associated with condom use during the most recent sexual encounter. This indicates that the outreach services are effective in reducing sexual risk behaviour and emphasises health care service uptake as an important deterrent to sexual risk behaviour among MSM and TG in Nepal.

Multivariable regression revealed that being away from home the past month was associated with increased condom use among TG. As a considerable portion of the respondents that identified as TG was involved in sex work, this association could be explained by mobility in connection with sex work or survival sex and a subsequent increase in condom use. Sex work is often connected to mobility and sexual risk behaviour and, transgender women involved in sex work have been found to be less likely to have unprotected sex with clients compared to stable partners [38, 39]. The response to unfair treatment is thought to be a measurement of coping, i.e. how an individual adapts to a problematic situation [27, 40], and is here categorised into active, moderate, and passive. An active response was defined as talking to others and acting by trying to do something about it. A moderate response was talking to someone about it but did not act, or did not talk to anyone but acted. A passive reaction was defined as not talking to anyone and not acting [27, 40]. In the model, having a moderate and active response to unfair treatment was associated with a decreased odds of condom use compared to a passive response. This result could be interpreted as that a more accepting and less outspoken coping strategy to unfair treatment in which TG try to comply with the recommended guidelines and does not dare to speak out is associated with condom use, whereas an active response could signify empowerment and engagement in sexual risk behaviour. However, several different definitions of coping strategies have been proposed and it is not clear how these relate to unprotected sex and discrimination in different populations and contexts [41]. This is also a difficult concept to tap into and requires a well-designed culturally appropriate instrument [41].

Individual and socio-structural factors associated with visiting outreach centre among MSM and TG

Given that having visited an outreach centre was significantly associated with sexual risk behaviour among both MSM and TG, it was important to examine the factors that influenced uptake of health services. Among MSM, having been forced by family to marry a female was positively associated with visiting an outreach centre. Moreover, both MSM and TG who reported to have been cheated or threatened the past year were more likely to have visited an outreach centre. These results could mean that the outreach centres are successful in providing a safe, friendly, and judgment-free setting for the respondents who feel discriminated and marginalised. This is in line with a recent study investigating uptake of HTC services among MSM and TG in Nepal which found that MSM and TG who experienced forced sex in the past year were more likely to access HTC services [42]. For both MSM and TG, there was a positive association between having high level of knowledge on HIV and visiting an outreach centre. This is in line with previous studies which have shown that poor HIV knowledge is associated with decreased HIV testing among MSM and TG in Brazil and India [43, 44]. There was a positive association between openness to family about sexual behaviour/identity and having visited an outreach centre. This is consistent with the idea that reduced stigma removes barriers to service uptake and has been observed in a previous study of MSM in Nepal [18].

Limitations

Given that the survey had a cross-sectional design, no causal inferences can be drawn from the results. Moreover, as the interviews were performed face to face, there is the potential risk of respondents giving socially desirable answers. Since the seeds were recruited through the BDS, this might have limited the representativeness of the data as more hidden groups of MSM and TG that are not affiliated with the BDS might potentially have other characteristics. Furthermore, there were also cases of cross recruitment of participants from different areas which led to a smaller number of respondents recruited from the districts Kanchanpur and Kapilbastu than initially planned for [22].

Conclusions

The high prevalence of HIV and syphilis as well as factors related to discrimination and stigma among TG in the Terai highway districts highlights the increased need for surveillance of this population outside the Kathmandu Valley and calls for further action in terms of prevention and treatment for HIV and STI. The results further emphasise the need for disaggregation of data by gender identity, as well as interventions tailored specifically towards the different populations. Given the high HIV/STIs prevalence levels in this area efforts should be made to improve the provision of testing and antiretroviral therapy coverage. Since healthcare service uptake seems to be an important factor in determining sexual risk behaviour among MSM and TG in Nepal, outreach services targeting this key population should be scaled up. Furthermore, as the social and structural challenges MSM and TG in Nepal experiences are encountered by MSM and TG worldwide, the results presented here are not only relevant to Nepal, but also generalizable to other settings, especially MSM and TG living in rural areas in other low-income countries.

List Of Abbreviations

AOR Adjusted Odds Ratio

BDS Blue Diamond Society

CC Community Centre

CE Community Educators

CI Confidence Interval

CM Community Motivators

DIC Drop-In Centre

EOD Experience of Discrimination

FSW Female Sex Worker

HIV Human Immunodeficiency Virus

HTC HIV Testing and Counseling

IBBS Integrated Biological and Behavioural Surveillance

IC Information Centre

LGBT Lesbian, Gay, Bisexual and Transgender

MSM Men who have Sex with Men

MSW Male Sex Worker

NCASC National Centre for AIDS and STD Control

OE Outreach Educator

OR Odds Ratio

PE Peer Educator

PWID People Who Inject Drugs

RDS Respondent Driven Sampling

STI Sexually Transmitted Infection

TG Transgender

Declarations

Ethics approval and consent to participate

The survey was ethically approved by the Nepal Health Research Council and was carried out according to ethical and human rights standards [22]. All participants gave verbal and witnessed consent before taking part in the study and no information which could identify individuals were collected in the survey. Respondents were given a unique ID number to track their participation in the study. Confidentiality was ensured throughout the survey process, and quality control systems for collection of both biological and behavioural data were incorporated into the survey protocol. It was important to retrieve information about younger segments of this population since risk behaviours often start at a young age [28]. The Nepal Health Research Council granted permission to obtain information from respondents below 18 without parental consent since disclosing the sexual behaviour/identity of the respondent to their parents might compromise them and have unknown repercussions for these individuals. Respondents were given counselling before and after clinical test results. Individuals that tested positive for HIV/STIs were given available treatment or referred for further treatment.

Consent for publication

Not applicable.

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on request.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was funded by The Global Fund to Fight AIDS, Tuberculosis and Malaria . The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions

MS, KD, JD and GM designed the study, analysed and interpreted the data, in addition to writing the manuscript. KD, US, RB and BR contributed to the survey administration, managing resources and supervision of field activities. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank officials from Blue Diamond Society for coordinating and mobilizing outreach/peer educators to facilitate the field work. We also thank the individuals who agreed to participate in the IBBS

survey. In addition, we would like to thank Fahri Saatcioglu and Mari Kaarbø for proofreading the manuscript.

References

1. UNAIDS: Global HIV & AIDS statistics – 2018 fact sheet. In.; 2018.
2. World Health Organization: Consolidated guidelines on HIV prevention, diagnosis, treatment and care for key populations – 2016 update. In.; 2016.
3. Baral S, Sifakis F, Cleghorn F, Beyrer C: Elevated risk for HIV infection among men who have sex with men in low- and middle-income countries 2000-2006: A systematic review. *Plos Medicine* 2007, 4(12):1901-1911.
4. Poteat T, Scheim A, Xavier J, Reisner S, Baral S: Global Epidemiology of HIV Infection and Related Syndemics Affecting Transgender People. *JAIDS-Journal of Acquired Immune Deficiency Syndromes* 2016, 72:S210-S219.
5. World Health Organization. Regional Office for South-East Asia: Progress Report on HIV in the WHO South-East Asia Region. In.; 2016.
6. The World Factbook, Nepal [<https://www.cia.gov/library/publications/the-world-factbook/geos/np.html>]
7. Government of Nepal. Ministry of Health. National Centre for AIDS and STD Control: Country Progress Report Nepal,. In.; 2018.
8. Paudel T, Singh N, Raj Banjara M, Kafle SP, Chandra Ghimire Y, Pokharel BR, Rawal BB, Badal K, Chaulagain M, Pendse R *et al*: Epidemiology of HIV, programmatic progress and gaps in last 10 years in Nepal. *Journal of virus eradication* 2016, 2(Suppl 4):35-40.
9. UNDP, USAID: Being LGBT in Asia: Nepal Country Report. Bangkok. In.; 2014.
10. Deuba K, Ekstrom AM, Tomson G, Shrestha R, Marrone G: HIV decline associated with changes in risk behaviours among young key populations in Nepal: analysis of population-based HIV prevalence surveys between 2001 and 2012. *Int J STD AIDS* 2016, 31:0956462416678303.
11. Nepal B: Population mobility and spread of HIV across the Indo-Nepal border. *Journal of Health Population and Nutrition* 2007, 25(3):267-277.
12. Reisner SL, Poteat T, Keatley J, Cabral M, Mothopeng T, Dunham E, Holland CE, Max R, Baral SD: Global health burden and needs of transgender populations: a review. *Lancet* 2016, 388(10042):412-436.
13. Poundstone KE, Strathdee SA, Celentano DD: The social epidemiology of human immunodeficiency virus/acquired immunodeficiency syndrome. *Epidemiologic Reviews* 2004, 26:22-35.
14. Stahlman S, Beyrer C, Sullivan PS, Mayer KH, Baral SD: Engagement of Gay Men and Other Men Who Have Sex with Men (MSM) in the Response to HIV: A Critical Step in Achieving an AIDS-Free Generation. *AIDS Behav* 2016, 20(Suppl 3):330-340.

15. Regmi PR, van Teijlingen E: Importance of Health and Social Care Research into Gender and Sexual Minority Populations in Nepal. *Asia-Pacific Journal of Public Health* 2015, 27(8):806-808.
16. Singh S, Pant SB, Dhakal S, Pokhrel S, Mullany LC: Human rights violations among sexual and gender minorities in Kathmandu, Nepal: a qualitative investigation. *Bmc International Health and Human Rights* 2012, 12.
17. Wilson E, Pant SB, Comfort M, Ekstrand M: Stigma and HIV risk among Metis in Nepal. *Culture Health & Sexuality* 2011, 13(3):253-266.
18. Deuba K, Ekstrom AM, Shrestha R, Ionita G, Bhatta L, Karki DK: Psychosocial Health Problems Associated with Increased HIV Risk Behavior among Men Who Have Sex with Men in Nepal: A Cross-Sectional Survey. *Plos One* 2013, 8(3).
19. Deuba K, Karki DK, Shrestha R, Aryal UR, Bhatta L, Rai KK: Risk of HIV Infection Among Men Having Sex With Men in Kathmandu Valley, Nepal. *Asia-Pacific Journal of Public Health* 2014, 26(2):126-137.
20. Kohlbrenner V, Deuba K, Karki DK, Marrone G: Perceived Discrimination Is an Independent Risk Factor for Suicidal Ideation among Sexual and Gender Minorities in Nepal. *Plos One* 2016, 11(7).
21. White RG, Hakim AJ, Salganik MJ, Spiller MW, Johnston LG, Kerr L, Kendall C, Drake A, Wilson D, Orroth K *et al*: Strengthening the Reporting of Observational Studies in Epidemiology for respondent-driven sampling studies: "STROBE-RDS" statement. *Journal of clinical epidemiology* 2015, 68(12):1463-1471.
22. NCASC: Integrated Biological and Behavioral Surveillance (IBBS) Survey among Men who have Sex with Men (MSM) and Transgender (TG) People in Terai Highway Districts of Nepal, Round I. In.; 2016.
23. Heckathorn DD: Respondent-driven sampling: A new approach to the study of hidden populations. *Social Problems* 1997, 44(2):174-199.
24. Family Health International IAPaCP: Behavioral surveillance surveys: Guidelines for repeated behavioral surveys in populations at risk of HIV. 2000.
25. Khanal V, Adhikari M, Karkee R: Social Determinants of Poor Knowledge on HIV Among Nepalese Males: Findings from National Survey 2011. *Journal of Community Health* 2013, 38(6):1147-1156.
26. Ghimire B, Suguimoto SP, Zamani S, Ono-Kihara M, Kihara M: Vulnerability to HIV infection among female drug users in Kathmandu Valley, Nepal: a cross-sectional study. *Bmc Public Health* 2013, 13.
27. Krieger N, Smith K, Naishadham D, Hartman C, Barbeau EM: Experiences of discrimination: Validity and reliability of a self-report measure for population health research on racism and health. *Social Science & Medicine* 2005, 61(7):1576-1596.
28. Beyrer C, Baral SD, van Griensven F, Goodreau SM, Chariyalertsak S, Wirtz AL, Brookmeyer R: Global epidemiology of HIV infection in men who have sex with men. *Lancet* 2012, 380(9839):367-377.

29. Poteat T, Ackerman B, Diouf D, Ceesay N, Mothopeng T, Odette KZ, Kouanda S, Ouedraogo HG, Simplicie A, Kouame A *et al*: HIV prevalence and behavioral and psychosocial factors among transgender women and cisgender men who have sex with men in 8 African countries: A cross-sectional analysis. *PLoS Med* 2017, 14(11):e1002422.
30. Winter S, Diamond M, Green J, Karasic D, Reed T, Whittle S, Wylie K: Transgender people: health at the margins of society. *Lancet* 2016, 388(10042):390-400.
31. NCASC: Integrated Biological and Behavioral Surveillance (IBBS) Survey among Man having Sex with Man (MSM) and Transgender (TG) in Kathmandu Valley, Round 5. In.; 2015.
32. Nakamura N, Mausbach BT, Ulibarri MD, Semple SJ, Patterson TL: Methamphetamine Use, Attitudes About Condoms, and Sexual Risk Behavior Among HIV-Positive Men Who Have Sex with Men. *Archives of Sexual Behavior* 2011, 40(2):267-272.
33. Lane T, Raymond HF, Dladla S, Rasethe J, Struthers H, McFarland W, McIntyre J: High HIV Prevalence Among Men Who have Sex with Men in Soweto, South Africa: Results from the Soweto Men's Study. *Aids and Behavior* 2011, 15(3):626-634.
34. Wang Y, Wang ZZ, Jia MM, Liang AJ, Yuan D, Sun Z, Gan F, Wang YC, Cai Y, Zhang ZR: Association between a syndemic of psychosocial problems and unprotected anal intercourse among men who have sex with men in Shanghai, China. *Bmc Infectious Diseases* 2017, 17.
35. Liu JX, Choi K: Experiences of social discrimination among men who have sex with men in Shanghai, China. *Aids and Behavior* 2006, 10:S25-S33.
36. Steward WT, Miege P, Choi KH: Charting a Moral Life: The Influence of Stigma and Filial Duties on Marital Decisions among Chinese Men who Have Sex with Men. *Plos One* 2013, 8(8).
37. UNDP, USAID: Being LGBT in Asia: Cambodia Country Report. Bangkok. In.; 2014.
38. Goldenberg S, Strathdee S, Perez-Rosales M, Sued O: Mobility and HIV in Central America and Mexico: A critical review. *J Immigrant Minority Health* 2012, 14(1):48-64.
39. Poteat T, Wirtz AL, Radix A, Borquez A, Silva-Santisteban A, Deutsch MB, Khan SI, Winter S, Operario D: HIV risk and preventive interventions in transgender women sex workers. *Lancet* 2015, 385(9964):274-286.
40. Stancil TR, Hertz-Picciotto I, Schramm M, Watt-Morse M: Stress and pregnancy among African-American women. *Paediatr Perinat Epidemiol* 2000, 14(2):127-135.
41. Han CS, Ayala G, Paul JP, Boylan R, Gregorich SE, Choi KH: Stress and Coping with Racism and Their Role in Sexual Risk for HIV Among African American, Asian/Pacific Islander, and Latino Men Who Have Sex with Men. *Archives of Sexual Behavior* 2015, 44(2):411-420.
42. Shrestha R, Philip S, Shewade HD, Rawal B, Deuba K: Why don't key populations access HIV testing and counselling centres in Nepal? Findings based on national surveillance survey. *BMJ open* 2017, 7(12):e017408.

43. Brito AM, Kendall C, Kerr L, Mota RMS, Drew M, Guimaraes C, Dourado I, Pinho AA, Benzaken AS, Brignol S *et al*: Factors Associated with Low Levels of HIV Testing among Men Who Have Sex with Men (MSM) in Brazil. *Plos One* 2015, 10(6).
44. Woodford MR, Chakrapani V, Newman PA, Shunmugam M: Barriers and facilitators to voluntary HIV testing uptake among communities at high risk of HIV exposure in Chennai, India. *Global Public Health* 2016, 11(3):363-379.

44. Woodford MR, Chakrapani V, Newman PA, Shunmugam M: Barriers and facilitators to voluntary HIV testing uptake among communities at high risk of HIV exposure in Chennai, India. *Global Public Health* 2016, 11(3):363-379.

Tables

Table 1. Socio-demographic characteristics among MSM and TG in Terai highway districts of Nepal.

	MSM, n=167, n (%)		TG, n=173, n (%)		Total, n=340, n (%)	
	Unweighted	Weighted*	Unweighted	Weighted*	Unweighted	Weighted*
Age distribution						
Mean (SD) age	27.4 (9.2)	26.1	31.8 (10.5)	31.3	29.7 (10.0)	28.1
Median age	24		29		27	
Minimum	17		17		17	
Maximum	56		66		66	
Education						
Never attended school	21 (12.6)	10	42 (24.3)	78.8	63 (18.5)	14.3
Ever attended school	146 (87.4)	90	131 (75.7)	21.2	277 (81.5)	85.7
Sexual identity						
Masculine appearing	159 (95.2)	98.3	24 (13.9)	12.6	183 (53.8)	64.6
Feminine appearing	8 (4.8)	1.7	149 (86.1)	87.4	157 (46.2)	35.4
Marital status						
Unmarried	116 (69.5)	77.2	104 (60.1)	60.4	220 (64.7)	29.4
Married	51 (30.5)	22.8	69 (39.9)	39.6	120 (35.3)	70.6
Living situation						
No home/rented	41 (24.6)	23.5	45 (26.0)	26.3	86 (25.3)	24.6
Own home/resident. hotel	126 (75.4)	76.5	128 (74.0)	73.7	254 (74.7)	75.4
Sex work						
Non sex workers	118 (70.7)	79.7	44 (25.4)	23.8	162 (47.6)	57.7
Sex workers	49 (29.3)	20.3	129 (74.6)	76.2	178 (52.4)	42.3
Income level						
Less than 3000	50 (29.9)	44.5	34 (19.7)	26.1	84 (24.7)	37.3
3000 - 10000	73 (43.7)	38.5	74 (42.8)	38.0	147 (43.2)	38.3
Above 10000	44 (26.3)	17.0	65 (37.6)	35.9	109 (32.1)	24.4

*Weighted value based on RDS II Estimator.

Table 2. Sexual risk behavior, HIV knowledge, socio-structural factors, and uptake of HIV prevention services among MSM and TG in Terai highway districts.

	MSM, n=167, n (%)		TG, n=173, n (%)		Total, n=340, n (%)	
	Unweighted	Weighted*	Unweighted	Weighted*	Unweighted	Weighted*
<u>Sexual risk behavior</u>						
Condom use at last sexual intercourse						
No	69 (41.3)	38.8	76 (43.9)	58.2	145 (42.6)	46.4
Yes	96 (57.5)	61.2	96 (55.5)	41.8	192 (56.5)	53.6
No response	2 (1.2)		1 (0.6)		3 (0.9)	
<u>Knowledge level of HIV</u>						
Low knowledge	30 (18.0)	19.3	31 (17.9)	23.7	61 (17.9)	21.0
High knowledge	137 (82.0)	80.7	142 (82.1)	76.3	279 (82.1)	79.0
<u>Socio-structural factors</u>						
Forced by family to marry female						
No	112 (67.1)	75.4	98 (56.6)	63.1	210 (61.8)	70.6
Yes	55 (32.9)	24.6	73 (42.2)	36.9	128 (37.6)	29.4
No Response			2 (1.2)		2 (0.6)	
Forced to leave home due to sexual behavior						
No	148 (88.6)	91.0	142 (82.1)	87.5	290 (85.3)	89.4
Yes	19 (11.4)	9.0	31 (17.9)	12.5	50 (14.7)	10.6
Openness to family about sexual behavior						
No	142 (85.0)	93.7	115 (66.5)	22.6	257 (75.6)	87.3
Yes	25 (15.0)	6.3	58 (33.5)	77.4	83 (24.4)	12.7
Beaten due to sexual behavior past year						
No	157 (94.0)	95.7	153 (88.4)	94.9	310 (91.2)	95.4
Yes	10 (6.0)	4.3	20 (11.6)	5.1	30 (8.8)	4.6

Forced to have sex past year						
No	144 (86.2)	93.1	141 (81.5)	82.7	285 (83.8)	89.0
Yes	23 (13.8)	6.9	32 (18.5)	17.3	55 (16.2)	11.0
Cheated/threatened due to sexual behavior in past year						
No	139 (83.2)	93.9	125 (72.3)	78.9	264 (77.6)	88.0
Yes	28 (16.8)	6.1	47 (27.2)	21.1	75 (22.1)	12.0
Don't Remember/Know			1 (0.6)		1 (0.3)	
Discrimination level						
No discrimination	99 (59.3)	72.3	81 (46.8)	49.0	180 (52.9)	63.9
In one or more situations	68 (40.7)	27.7	92 (53.2)	51.0	160 (47.1)	36.1
Reaction when treated unfairly						
Accept it/keep to self	62 (37.1)	35.5	55 (31.8)	34.8	117 (34.4)	35.2
Do something/keep to self	40 (24)	26.6	60 (34.7)	30.0	100 (29.4)	28.0
Do something/talk to others	65 (38.9)	37.9	58 (33.5)	35.2	123(36.2)	36.8
Away from home last month						
No	120 (71.9)	69.6	113 (65.3)	71.1	233 (68.5)	70.3
Yes	47 (28.1)	30.4	60 (34.7)	28.9	107 (31.5)	29.7
Cross border travel for sexual activities						
No	136 (81.4)	88.9	138 (79.8)	79.7	274 (80.6)	85.3
Yes	30 (18)	11.1	35 (20.2)	20.3	65 (19.1)	14.7
No response	1 (0.6)				1 (0.3)	
Interacted with PE, OE or CM or CE last 12 months						
No	75 (44.9)	40.4	58 (33.5)	45.4	133 (39.1)	45.9
Yes	92 (55.1)	59.6	113 (65.3)	54.6	205 (60.3)	54.1

No response			2 (1.2)		2 (0.6)	
Visited outreach center in the last 12 months						
No	95 (56.9)	75.1	54 (31.2)	45.8	149 (43.8)	63.6
Yes	72 (43.1)	24.9	119 (68.8)	54.2	191 (56.2)	36.4
Visited STI clinic in the last 12 months						
No	146 (87.4)	92.6	143 (82.7)	86.6	289 (85.0)	90.3
Yes	21 (12.6)	7.4	30 (17.3)	13.4	51 (15.0)	9.7
Visited HTC center in the last 12 months						
No	128 (76.6)	87.4	128 (74.0)	78.5	256 (75.3)	83.9
Yes	39 (23.4)	12.6	45 (26.0)	21.5	84 (24.7)	16.1

*Weighted value based on RDS II Estimator.

Table 3. Logistic regression model of individual and socio-structural factors associated with condom use at last sex among MSM in Terai districts of Nepal, (n=165, missing=2).

	Crude OR	95% CI	p-value	AOR	95% CI	p-value
<u>Individual factors</u>						
Age	0.97	0.94 - 1.01	0.140			
Income level						
Less than 3000	1			1		
3000 - 10000	0.52	0.24 - 1.11	0.091	0.56	0.24 - 1.28	0.168
Above 10000	0.43	0.18 - 1.01	0.052	0.36	0.14 - 0.92	0.032
Total						
<u>Living situation</u>						
No home/rented	1			1		
Own home/resident. hotel	1.90	0.93 - 3.87	0.078	1.95	0.90 - 4.25	0.092
<u>Syphilis</u>						
Negative	1					
Positive	2.56	0.68 - 9.67	0.166			
<u>Socio-structural factors</u>						
Forced by family to marry female						
No	1			1		
Yes	0.68	0.35 - 1.31	0.251	0.42	0.19 - 0.95	0.037
Cheated/threatened due to sexual behaviour in past year						
No	1					
Yes	2.01	0.83 - 4.87	0.124			
Away from home last month						
No	1					
Yes	0.52	0.26 - 1.04	0.063			
Interacted with PE, OE or CM or CE last 12 months						

No	1					
Yes	2.40	1.28 - 4.53	0.007			
Visited outreach centre in the last 12 months						
No	1			1		
Yes	3.35	1.71 - 6.55	<0.001	5.37	2.42 - 11.94	<0.001

Table 4. Logistic regression model of individual and socio-structural factors associated with condom use at last sex among TG in Terai districts of Nepal, (n=172, missing=1).

	Crude OR	95% CI	p-value	AOR	95% CI	p-value
<u>Individual factors</u>						
Income level						
Less than 3000	1			1		
3000 - 10000	2.51	1.09 - 5.77	0.031	2.79	1.12 - 6.91	0.027
Above 10000	2.36	1.01 - 5.54	0.048	1.74	0.67 - 4.53	0.253
Active syphilis						
Negative	1			1		
Positive	0.49	0.19 - 1.26	0.136	0.40	0.14 - 1.17	0.096
Syphilis						
Negative	1					
Positive	0.55	0.22 - 1.39	0.206			
<u>Socio-structural factors</u>						
Forced by family to marry female						
No	1			1		
Yes	0.65	0.36 - 1.21	0.175	0.42	0.21 - 0.88	0.021
Reaction when treated unfairly						
Accept it/keep to self	1			1		
Do something/keep to self	0.31	0.14 - 0.68	0.004	0.31	0.13 - 0.74	0.008
Do something/talk to others	0.30	0.13 - 0.66	0.003	0.29	0.12 - 0.68	0.005
Away from home last month						
No	1			1		
Yes	1.72	0.90 - 3.28	0.103	2.46	1.13 - 5.38	0.024
Visited outreach centre in the last 12 months						
No	1			1		

Yes	1.85	0.96 - 3.56	0.065	2.81	1.30 - 6.09	0.009

Figures

Figure 1

Prevalence of HIV and syphilis among MSM and TG in eight Terai highway districts.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.docx](#)
- [supplement2.docx](#)