

Quality of Life as a predictor of mortality in the general population: a systematic review and meta-analysis

Aung Zaw Zaw Phyo

Monash University Faculty of Medicine Nursing and Health Sciences <https://orcid.org/0000-0002-8834-4072>

Rosanne Freak-Poli

Monash University Faculty of Medicine Nursing and Health Sciences

Heather Craig

Monash University Faculty of Medicine Nursing and Health Sciences

Danijela Gasevic

Monash University Faculty of Medicine Nursing and Health Sciences

Nigel Stocks

The University of Adelaide Adelaide Medical School

David A. Gonzalez-Chica

The University of Adelaide Adelaide Medical School

Joanne Ryan (✉ joanne.ryan@monash.edu)

<https://orcid.org/0000-0002-7039-6325>

Research article

Keywords: quality of life, life quality, HRQoL, mortality, meta-analysis, predictor, review

Posted Date: February 3rd, 2020

DOI: <https://doi.org/10.21203/rs.2.22428/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on November 6th, 2020. See the published version at <https://doi.org/10.1186/s12889-020-09639-9>.

Abstract

Background: Quality of life (QoL) is multi-dimensional concept of an individual' general well-being status in relation to their value, environment, cultural and social context in which they live. This study aimed to quantitatively synthesise available evidence on the association between QoL and mortality in the general population.

Methods: An electronic search was conducted using three bibliographic databases, MEDLINE, EMBASE and PsycINFO. Inclusion criteria were studies that assessed QoL using standardized tools and examined mortality risk in a non-patient population. Qualitative data synthesis and meta-analyses using a random-effects model were performed.

Results: Of 4,184 articles identified, 47 were eligible for inclusion, involving approximately 1,200,000 participants. Studies were highly heterogeneous in terms of QoL measures, population characteristics and data analysis. In total, 43 studies (91.5%) reported that better QoL was associated with lower mortality risk. The results of four meta-analyses indicated that higher health-related QoL (HRQoL) is associated with lower mortality risk, which was consistent for overall HRQoL (HR 0.633, 95% CI: 0.514 to 0.780), physical function (HR 0.987, 95% CI: 0.982 to 0.992), physical component score (OR 0.950, 95% CI: 0.935 to 0.965), and mental component score (OR 0.980, 95% CI: 0.969 to 0.992).

Conclusion: These findings provide evidence that better QoL/HRQoL was associated with lower mortality risk. The utility of these measures in predicting mortality risk indicates that they should be considered further as potential screening tools in general clinical practice, beyond the traditional objective measures such as body mass index and the results of laboratory tests.

Background

Quality of life (QoL) is a multi-dimensional concept of an individual's general well-being status in relation to the value, environment, cultural and social context in which they live [1]. Since QoL measures outcomes beyond biological functioning and morbidity [2], it is recognised as an important measure of overall [1]. The origin of the term QoL dates back to the early 1970s, as a measure of wellness with linkage to health status like diseases or disability [3, 4]. Since then, interest in QoL has increased considerably [5]. As life expectancy increases, more emphasis has been placed on the importance of better QoL, and the maintenance of good health for as long as possible [6–9]. Indeed, global leading health organizations have emphasized the importance of QoL and well-being as a goal across all life stages [10–12].

Moreover, QoL has increasingly been used in the wider context to monitor the efficacy of health services (e.g. patient reported outcome measures, PROMs), to assess intervention outcomes, and as an indicator of unmet needs [13–15]. Several studies have reported that QoL is negatively associated with rehospitalization and death in patients with diseases such as coronary disease [16, 17], and pulmonary diseases [18]. Further, QoL is also predictive of overall survival in patients affected by cancer, chronic kidney disease or after coronary bypass graft surgery [19–22]. In recent years, an increasing number of studies have investigated whether QoL is also a predictor of mortality risk in the general population [23–27].

To date, there has been only one pooled analysis of eight heterogeneous-Finnish cohorts. That study of 3,153 older adults, focused exclusively on the prognostic value of the validated 15-dimensional (15D) health-related QoL (HRQoL) measures [28] for predicting all-cause mortality [29]. However, there has been no systematic review investigating the association between QoL and all-cause mortality in population-based samples which could be used to monitor health changes in the general population. Therefore, this systematic review and meta-analysis was conducted with the aim of determining whether QoL is predictive of mortality in the general population which includes individuals with or without a range of health conditions.

Methods

Search Methods

This systematic review and meta-analysis were conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [30]. The protocol for this review was registered with the International Prospective Register of Ongoing Systematic Reviews (PROSPERO) [31], under the registration number: CRD42019139994 [32]. The electronic bibliographic databases, MEDLINE, EMBASE and PsycINFO (through OVID) were searched from database inception until June 21, 2019. The search strategy was developed in consultation with a Senior Medical Librarian. The MeSH terms and key-words were developed for MEDLINE (through OVID) and were translated to EMBASE and PsycINFO using the OVID platform (See Supplementary Tables S1-S3, Additional File 1). When the full text of an article was not available, all attempts were made to obtain it by contacting the authors directly. To identify further potentially relevant studies, another search was also developed with those specific QoL / HRQoL measures which were found in this review (See Supplementary Table S4, Additional File 1). Additionally, the bibliography lists of the included articles were also hand searched.

Inclusion and Exclusion Criteria

Articles were included if they: (a) involved adults aged 18 years and older; (b) were general population-based samples with or without a range of health conditions; (c) assessed mortality from any cause or cause-specific mortality using a longitudinal design; and (d) included a QoL / HRQoL measure using a standard tool. We excluded papers not written in English, reviews, or studies including only specific groups of patients (e.g. patients on dialysis, those with fractures, after surgery, or individuals with a terminal illness).

Study Selection

The screening of articles for eligibility according to title and abstract was undertaken independently by two reviewers (AZZP and HC). All relevant full-text articles were reviewed by two reviewers (AZZP and HC) for eligibility against inclusion criteria. Discrepancies and disagreements were resolved through discussion between two reviewers (AZZP and HC) and a consensus decision was made if required, through consultation with a third reviewer (JR).

Data Extraction

A standard data extraction form was used which included the following fields – title, authors, year of publication, setting/country, name of the study and design, sample size, follow-up period, participant characteristics (age and

sex), specific QoL measure, cause of death (if available), and results (risk estimates including 95% confidence intervals, CI) which were standardized in term of 1-unit increase or 1-SD increase for continuous risk estimate, or high vs. low for categorical risk estimates. The first reviewer (AZZP) completed the data extraction form and a second reviewer (HC) verified the extracted information. All efforts were made to contact authors when there was missing information.

Quality Appraisal

The quality of included studies was appraised using 'the Newcastle – Ottawa Quality Assessment Scale (NOS)' [33]. The NOS includes eight items, categorized into three dimensions (a) Selection, (b) Comparability, and (c) Outcome. The NOS scale uses a star system to evaluate the quality of each study, and they can be accredited a maximum of one star for each item within the Selection and Outcome dimension and two stars for the Comparability item. When considering the comparability of each study, a star was provided for studies which controlled for relevant covariates – age, sex (where appropriate), socioeconomic status or proxy (including socioeconomic position, education level or income), and some measure of co-morbidity (for example a specific health condition). An additional star was given for studies which considered other factors associated with QoL and mortality, including clinical measures, BMI, or lifestyle factors (i.e. smoking, alcohol, physical activity). The range of NOS scoring was from 0 to 9 stars, with higher scores indicating less susceptibility to bias. The methodological quality of included studies was rated by one reviewer (AZZP) and verified by a second reviewer (HC). Disagreements were resolved through discussion with a third reviewer (JR).

Data Synthesis

The clinical and methodical heterogeneity of the studies was examined, in particular considering the measure of QoL used, and the effect estimates reported (Hazard Ratio (HR), Relative Risk (RR) or Odds Ratio (OR)). Where studies were considered too methodically heterogeneous to enable pooling, the results were summarized quantitatively in tables according to related categories with risk estimates; and 95% CIs.

Meta-analysis

A meta-analysis was performed when there was a sufficient number of studies (four or more) which used the same domain of QoL measure and equivalent effect estimate parameters. In the present study, four meta-analyses were conducted for a pooled risk estimate of studies using (a) physical component score (PCS) of 36-item Short Form (SF-36) and OR / RR; (b) physical function domain of SF-36 and HR; (c) mental component score (MCS) of SF-36 and OR / RR; and (d) the 15-dimensional measure (15D) and HR. A DerSimonian-Laird random-effects model was chosen given heterogeneity in the studies in terms of population characteristics and varying health status. When more than one risk estimate was reported in the study, the fully adjusted/final regression model was included. Effect estimates were standardized where possible, so all values corresponded to a 1-unit increase in QoL. A pooled risk estimates of less than one indicates a decreased risk of mortality with higher QoL. Statistical heterogeneity was evaluated by using the I^2 statistic, and the results were interpreted based on the Cochrane guidelines (0–40% = no heterogeneity; 30–60% = moderate heterogeneity; 50–90% = substantial heterogeneity; and 75–100% = considerable heterogeneity) [34]. Funnel plots and Egger's test were used to assess publication bias. Data analysis was undertaken using Stata statistical software, version 15.0 (StataCorpLP, College Station, TX, USA).

Results

Search Result

A total of 4,175 articles were identified from the systematic database search, and six additional articles were found via searching the reference list of included articles (Figure 1). After removing duplicates, 3,140 records remained for review. After title and abstract screening, 3,058 articles were excluded and the full-text of the remaining 82 articles were evaluated for eligibility. A total of forty-four (44) articles met all inclusion criteria. Excluded articles with reasons for exclusion are presented in Supplementary Table S5, Additional File 1. Moreover, three articles from additional search were also added in this review. Therefore, a total of forty-seven (47) articles were included in this systematic review.

Description of Included Studies

Table 1 presents the characteristics of the 47 included studies. The earliest study was published in 1993 while the remaining included articles were published between 2002 and 2019, with 28% published in the past five years. All studies except the retrospective cohort study of Ul-Haq *et al.*, [73] were prospective cohort studies. The included studies were conducted in USA (34%), UK (9%), Australia (6%), Canada (6%), Spain (6%), Taiwan (6%), Belgium (4%), Finland (4%), Scotland (4%), Sweden (4%), Bangladesh (2%), China (2%), Germany (2%), South Korea (2%), Italy (2%), Norway (2%), and South Africa (2%). The sample sizes of the included studies ranged from 171 [39] to 559,985 [38]; 14 studies had a sample size of less than 1000, 17 studies between 1,000 and 10,000, 13 studies between 10,000 and 100,000, and the remaining three studies [36, 38, 51] has a sample size of more than 100,000 participants. Five studies included only males [39, 40, 52, 69, 71] and three studies only females [54, 57, 72]. The remaining 39 studies recruited between 3% to 78 % of women. The follow-up periods of the studies varied between 9 months [70] and 18 years [71].

Half of the included studies (24 studies) measured QoL using the Short Form 36 (SF-36), four studies used the shorter version SF-12 [43, 49, 60, 73], and two studies SF-20 [47, 72]. Four studies [25, 52, 61, 67] used the WHO-QOL questionnaire, two [36, 42] used the core CDC Healthy Days Measure (HRQOL-4), two [37, 51] used the EuroQoL-5 Dimension (EQ-5D), two [26, 51] used the short-form six-dimension utility index (SF-6D), two [46, 53] used the health utilities index 3 (HUI3) and one study that included five individual cohorts [29] used the 15-dimensional HRQoL measure (15D). The other studies used a variety of different QoL measures which were not used in more than one study (Table 1 and 2).

Of the 47 articles included in this review (Table 1), fourteen (14) studies involved the same cohorts and, in several cases, likely the same participants. Subsequent publications often reported effect estimates over different lengths of follow-up or using different QoL tools. Two published articles of De Buyser *et al.* reported the results of the same population-based cohort study [39, 40], three published articles by De Salvo *et al.* and Fan *et al.* were from the same study and included participants enrolled in the Veterans Affairs Ambulatory Care Quality Improvement Project [24, 41, 45], two published studies of Mold *et al.* and Lawler *et al.* used the same community-dwelling cohort [55, 59], two published studies of Higuera-Fresnillo *et al.* and Otero-Rodriguez *et al.* were from the same Spanish cohort [50, 65], two published studies of Feeny *et al.* and Kaplan *et al.* were from the same Canadian cohort [46, 53]; and Myint *et al.* published three articles [26, 62, 63] with different perspectives on the same population-based study. Additionally, Liira *et al.*'s study [29], included eight individual

cohorts, however, only five of the cohorts met the inclusion criteria for this current systematic review, and thus are shown in Table 1.

Risk of Bias Assessment

The methodological quality of included studies based on NOS ranged between five and nine stars. Among the included studies, seven were of high methodological quality, with nine stars. Across the ten studies with less than seven stars, they were scored most poorly on the items assessing how representative the cohort was in relation to the overall population being sampled and whether they adjusted for potential confounding factors in their analysis (See Supplementary Table S6 and S7, Additional File 1).

Qualitative Synthesis

Of the total 47 included studies, 43 (91.5%) studies reported for at least one of the domains examined, that better QoL was associated with lower mortality risk (Table 1). Of 33 studies which assessed physical HRQoL (nine exclusively assessed physical HRQoL), 30 studies (91%) reported better HRQoL was associated with lower mortality risk. Among the 23 studies which examined mental HRQoL (one exclusively assessed MCS), 13 studies (57%) reported that higher mental HRQoL was associated with decreased mortality risk (Table 1). The five studies [47, 50, 55, 57, 74] that measured HRQoL using SF-36 or SF-20 reported not only the physical functioning and mental health domains, but also general health perception, bodily pain, vitality, and social functioning. The findings were generally consistent in general health perception and social functioning; and it was reported that better level of general health perception and social functioning was associated with decreased mortality risk (Table 1).

The mortality risk estimates of the studies which were not included in the meta-analyses are shown in Tables 3-5. The 18 out of 20 studies which measured the PCS using the SF-36 or SF-12 or the physical functioning subscale using SF-36, RAND-36, or SF-20 reported these to be a predictor of mortality risk, with better physical health being associated with lower mortality risk (Table 3). Nine out of 16 studies which assessed the MCS or mental health subscale using SF-36 or SF-12, showed that better mental health was associated with lower mortality risk (Table 4). The 12 out of the 15 studies that measured the association between QoL and mortality risk, found that higher QoL scores were associated with lower mortality risk (Table 5).

Meta-Analyses

Four studies including 53,642 participants [23, 24, 58, 68] measured QoL using the SF-36 and examined the association between the PCS and all-cause mortality and provided estimates from logistic regression analysis (OR or RR). With an average 1.8-year follow-up, one unit increase in the SF-36 PCS was associated with a 5% decrease in all-cause mortality (pooled OR/RR = 0.950; 95% CI: 0.935 to 0.965; P-value < 0.001). There was substantial heterogeneity between studies ($I^2 = 82.1\%$) (Figure 2).

Six studies including 22,570 participants [40, 44, 55, 57, 66, 74] measured QoL using the SF-36 and investigated the association between the physical functioning and all-cause mortality using time-to-event survival analysis. With an average 8.7-year follow-up, one unit increase in the SF-36 PF was associated with a 1.3% decrease in time to death (pooled HR = 0.987; 95%CI: 0.982 to 0.992; P-value < 0.001). There was substantial heterogeneity between studies ($I^2 = 83.8\%$) (Figure 3).

Four studies including 53,642 participants [23, 24, 58, 68] measured QoL using the SF-36 and examined the association between the MCS and all-cause mortality reported estimates on logistic regression analysis (OR or RR). With an average 1.8-year follow-up, one unit increase in the SF-36 MCS was associated with a 2% decrease in all-cause mortality (pooled OR/RR = 0.980; 95% CI: 0.969 to 0.992; P-value = 0.001). There was substantial heterogeneity between studies ($I^2 = 75.9\%$) (Figure 4).

Five Finnish individual cohorts of the Liira *et al.* study including 2,377 [29] measured QoL using the 15D index and explored its association with all-cause mortality using time-to-event survival analysis. With an average 2-year follow-up, one SD (0.14) increase in the 15D index was associated with a 36.7% decrease in all-cause mortality (pooled HR = 0.633; 95%CI: 0.514 to 0.780; P-value < 0.001). There was moderate heterogeneity between studies ($I^2 = 49.4\%$) (Figure 5).

Visual inspection of the funnel plots which were used to assess for publication bias were presented in the Supplementary Figures S1-S4, Additional File 1. For three of the four meta-analyses, there was no strong evidence of publication bias, however for the meta-analysis of MCS, this test was statistically significant (P = 0.04).

Discussion

This systematic review is the first to investigate the association between QoL and mortality in community-dwelling individuals with or without health conditions. It summarizes the findings from 47 studies including approximately 1,200,000 individuals aged predominantly 65 years and older (age range 18-101 years), with 46 studies (98%) conducted in high-income or upper-middle-income countries. Overall thirteen different instruments were used to assess the association between QoL or more specifically HRQoL and mortality risk after nine months to 18 years of follow-up, with the SF-36 or its derivatives (RAND-36, SF-20, SF-6D) most commonly used. Overall, 43 (91.5%) studies of the 47 included studies reported for at least one of the domains examined, that better QoL was associated lower mortality risk, which was also supported by the results of four meta-analyses (11 studies, n=78,589) of PCS, physical function and MCS domains of the SF-36, and 15D HRQoL.

Our findings are in line with a previous study that used pooled analysis [29] of eight heterogeneous Finnish cohorts using the 15D HRQoL measure and included a wide range of both community-dwelling participants with or without morbidity, such as cardiovascular disease, dementia, and hospitalized patients with delirium. They also found that the 15D HRQoL measure was associated with two-year survival, with a slightly higher hazard ratio than that found in our study (HR per 1-SD = 0.44, 95% CI 0.40 to 0.48) [29]. These differences may relate to their inclusion of patient groups in generally poorer health, while our systematic review focused on the community dwelling population. Moreover, our findings in the general non-patient population are also comparable with studies investigating people with specific diseases such as cancer and chronic kidney disease, which reported QoL to be a predictor of mortality risk [19-21].

The findings of the present study are also consistent with those of recent population-based systematic review which investigated on the association between QoL and multimorbidity [76]. In their recent study, Makovski *et al.* (2019) systematically reviewed the evidence on the relationship between QoL and multimorbidity. They observed a stronger relationship between the PCS of QoL and multimorbidity (overall decline in QoL per additional disease = -4.37%, 95%CI -7.13% to -1.61% for WHOQoL-BREF physical domain and -1.57%, 95%CI -2.70% to -0.44% for

WHOQoL-BREF mental domain) [76]. These findings also align with the results of the present study, where the meta-analysis indicated a stronger effect size for PCS compared to MCS using the SF-36 tool (pooled OR/RR = 0.950; 95% CI: 0.935 to 0.965 for PCS; and pooled OR/RR = 0.980; 95%CI: 0.969 to 0.992 for MCS). Since physical health is generally recognised as a strong risk factor for comorbidity, hospitalisations and mortality [77-80], our findings add further support to the predictive capacity of physical HRQoL for mortality risk. Like other objective health measures such as body mass index, glycaemia, and blood pressure, these findings highlight the utility of assessing physical HRQoL in general clinical practice to help identify individuals at greatest risk of death [81].

Given the evidence regarding the longitudinal relationship between QoL and mortality risk, the finding of this review can help to increase the efficacy of disease prevention strategies in older people through identifying individuals at higher risk for adverse health outcomes in general practice / primary health settings. There is a need for further studies however, in particular to better understand the influence of gender on these associations, and whether differences could be observed for males and females. Understanding these specific relationships could help identify which particular groups are most at risk and enable specific targeting of interventions to these individuals.

Strengths of the Review

Strengths of this systematic review are that it was performed in a rigorous manner, adhering to strict systematic review guidelines. The protocol was registered with the International prospective register of systematic reviews (PROSPERO), and the review was undertaken in accordance with the preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement. A reproducible and rigorous search strategy using three electronic databases was used, which helped ensure that all relevant articles were included. The literature screening was independently performed by two reviewers, who were also involved in the process of data extraction and methodological quality assessment of the included studies in accordance with NOS. Based on the NOS, all studies received greater than or equal to five out of nine stars, which indicates that there was generally a low risk of bias. Similarly, most studies provided risk estimates that controlled for important factors including current health and socio-economic status. Since our review criteria were not limited to articles with the commonly used QoL (or HRQoL) tools such as the SF-36, this has increased the generalisability of the findings. Therefore, this review has a broad and comprehensive perspective, with results that are rigorous and can be reproduced.

Limitations of the Review

Among included articles, large heterogeneity was observed in terms of country-of-origin, participant characteristics, and evaluation of QoL. The majority of the included articles were conducted in English speaking countries, and restriction to English language articles as part of our inclusion criteria, may impact the generalisability of these findings. Since the different QoL standard tools examine different aspects [82, 83] and are not directly comparable, this made comparison of included studies in data synthesis difficult. There were also some differences in the way the data analysis was performed and the results were presented, reporting OR versus HR for example. In addition, some articles reported the risk estimates by comparing categorical QoL groups while others provided the risk estimates per 1 or more units change in the continuous scale. Hence, the different nature of each QoL scale and inconsistency in risk comparison precluded us from including some

articles in the meta-analyses. As such, only 11 studies were included across the four meta-analyses of this systematic review, and the meta-analyses still showed substantial heterogeneity. Therefore, caution should be taken with the interpretation of the overall effect estimates. Moreover, since the numbers of studies included in each meta-analysis were fewer than 10 studies, the results of funnel plots or Egger's test should also be interpreted with caution. Of particular interest here, it has commonly been reported that gender differences exist in QoL and women of all age groups have lower QoL than their male counterparts [84-88]. However, in this review, it was not possible to perform statistical pooling by gender and age groups due to the different reporting strategies of the reviewed studies.

Conclusion

This is the first systematic review and meta-analysis that has determined whether QoL is associated with mortality in the general non-patient population. In summary, the findings provide evidence that better QoL or HRQoL measured by different tools were associated with lower mortality risk in the general population. Furthermore, this study also adds further support to the predictive capacity of physical HRQoL for mortality risk. Additional research is needed to determine whether these associations differ across gender, age-group and other populations in low- and lower-middle-income countries, who have suffered of a double burden of infectious and chronic diseases, with having difficulties for accessing quality health services. Ultimately these findings suggest the utility of QoL measures to help identify populations at greatest risk of mortality and who might benefit most from routine screening in general practice and possible interventions.

Supplemental Information Note

Additional File 1: The supplementary material includes Figures S1 – S4 and Tables S1 – S7. **Figure S1.** Funnel plot of all-cause mortality risk per one unit increase in SF-36 PCS. **Figure S2.** Funnel plot of all-cause mortality risk per one unit increase in SF-36 Physical-Functioning. **Figure S3.** Funnel plot of all-cause mortality risk per one unit increase in SF-36 MCS. **Figure S4.** Funnel plot of all-cause mortality risk per one-SD (0.14) increase in 15D index. **Table S1.** Search Strategy using Ovid MEDLINE 1946 to June 21 2019. **Table S2.** Search Strategy using Embase Classic 1947 to June 21 2019. **Table S3.** Search Strategy using PsycINFO 1806 to June Week 3 2019. **Table S4.** Additional Search Strategy up to June Week 3 2019. **Table S5.** The list of excluded articles and reasons for exclusion (n = 38). **Table S6.** Appraisal Standard of Newcastle/Ottawa Scale. **Table S7.** Quality appraisal of included studies based on the Newcastle–Ottawa Quality Assessment Scale.

Abbreviations

15D = 15-dimensional

CI = confidence intervals

EQ-5D = EuroQoL-5 dimension

HR = hazard ratio

HRQoL = health-related quality of life

HUI3 = health utilities index 3

MCS = mental component score

NOS = Newcastle-Ottawa quality assessment scale

OR = odds ratio

PCS = physical component score

PRISMA = preferred reporting items for systematic reviews and meta-analyses

PROMs = patient reported outcome measures

PROSPERO = international prospective register of systematic reviews

QoL = quality of life

RR = relative risk

SD = standard deviation

SF-12 = 12-items short form

SF-20 = 20-item short form

SF-36 = 36-item short form

SF-6D = six-dimension utility index

Declarations

Ethics approval and consent to participate

This is a systematic review and meta-analysis of publicly available studies. No ethical approval was required.

Consent for publication

Not applicable

Availability of data and materials

All data analysed during this study were taken directly from published manuscripts (as referenced). All data generated during this study, are included in this published article [and its supplementary information files].

Competing interests

The authors declare no conflicts of interest.

Funding

This work was supported by Monash International Tuition Scholarship and Monash Graduate Scholarship. JR is supported by a National Health and Medical Research Council Dementia Research Leader Fellowship (APP1135727). None of the funders were involved in the design of the study, in the collection, analysis, and interpretation of data and in the writing of the manuscript.

Authors' contributions

RFP conceived the study. JR and AZPP designed the study. AZPP undertook the literature searches, screened the articles, extracted the data, performed quality assessment and data analysis. HC was the independent assessor, also completing all data screening, extraction and quality assessment. AZPP and JR interpreted the data, with input from DAGC, DG, and NS. AZPP wrote the initial manuscript draft. All authors provided critical comments and approved the final version.

Acknowledgements

We would like to thank Lorena Romero, the Senior Medical Librarian, Alfred Health, and Cassandra Freeman, the Subject Librarian, Faculty of Medicine, Nursing and Health Sciences, Monash University Library for technical support involved in developing the search strategy.

References

1. Kuyken W, Orley J, Power M, Herrman H, Schofield H, Murphy B, et al. The World Health Organization quality of life assessment (WHOQOL): Position paper from the World Health Organization. *Social Science & Medicine*. 1995;41(10):1403-9.
2. Ware JE. The Status of Health Assessment 1994. *Annu Rev Publ Health*. 1995;16(1):327-54.
3. Elkinon JR. Medicine and the quality of life. *Annals of internal medicine*. 1966;64(3):711.
4. Spitzer WO. State of science 1986: Quality of life and functional status as target variables for research. *Journal of Chronic Diseases*. 1987;40(6):465-71.
5. World Health Organization. World report on ageing and health. Geneva: World Health Organization 2015 [Available from: <http://www.who.int/ageing/publications/world-report-2015/en/>].
6. Brown GC. Living too long. EMBO reports. 2015. Report No.: 1469-221X Contract No.: 2.
7. World Health Organization. Global Health Observatory (GHO) data [cited 2019 August 27]. Available from: https://www.who.int/gho/mortality_burden_disease/life_tables/situation_trends_text/en/.
8. Centers for Disease Control and Prevention. Health-Related Quality of Life (HRQOL) Concept [cited 2019 August 27]. Available from: <https://www.cdc.gov/hrqol/concept.htm>.
9. Salomon JA, Wang H, Freeman MK, Vos T, Flaxman AD, Lopez AD, et al. Healthy life expectancy for 187 countries, 1990–2010: a systematic analysis for the Global Burden Disease Study 2010. *The Lancet*. 2012;380(9859):2144-62.
10. Centers for Disease Control and Prevention. Measuring Healthy Days: Population Assessment of Health-Related Quality of Life. Atlanta, Georgia: CDC; 2000.
11. Centers for Disease Control and Prevention. Healthy People 2020 2019 [cited 2019 August 14]. Available from: https://www.cdc.gov/nchs/healthy_people/hp2020.htm.

12. World Health Organization: International Classification of Functioning, disability, and Health: Children and Youth Version: ICF-CY. Geneva World Health Organization 2007.
13. Sintonen H. The 15-D measure of health-related quality of life : reliability, validity and sensitivity of its health state descriptive system. National Centre for Health Program E, editor. Fairfield, Vic.: Fairfield, Vic. : National Centre for Health Program Evaluation; 1994.
14. Gross RC, Limwattananon LC, Matthees LB, Zehrer LJ, Savik LK. Impact of transplantation on quality of life in patients with diabetes and renal dysfunction. *Transplantation*. 2000;70(12):1736-46.
15. Moinpour CM, Savage MJ, Troxel A, Lovato LC, Eisenberger M, Veith RW, et al. Quality of Life in Advanced Prostate Cancer: Results of a Randomized Therapeutic Trial. *The Journal of Urology*. 1999;161(4):1394-5.
16. Rodríguez-Artalejo F, Guallar-Castillón P, Otero CM. Health-Related Quality of Life as a Predictor of Hospital Readmission and Death Among Patients With Heart Failure. *ACC Current Journal Review*. 2005;14(10):28-.
17. Rumsfeld JS, Mawhinney S, McCarthy M, Shroyer AL, Villanueva CB, O'Brien M, et al. Health-related quality of life as a predictor of mortality following coronary artery bypass graft surgery. Participants of the Department of Veterans Affairs Cooperative Study Group on Processes, Structures, and Outcomes of Care in Cardiac Surgery. *JAMA*. 1999;281(14):1298.
18. Domingo-Salvany A, Lamarca R, Ferrer M, Garcia-Aymerich J, Alonso J, Félez M, et al. Health-related quality of life and mortality in male patients with chronic obstructive pulmonary disease. *American journal of respiratory and critical care medicine*. 2002;166(5):680.
19. Mehanna HM, Morton RP. Does Quality of Life Predict Long-term Survival in Patients With Head and Neck Cancer? *Archives of Otolaryngology–Head & Neck Surgery*. 2006;132(1):27-31.
20. Maisey NR, Norman A, Watson M, Allen MJ, Hill ME, Cunningham D. Baseline quality of life predicts survival in patients with advanced colorectal cancer. *European Journal of Cancer*. 2002;38(10):1351-7.
21. Tsai Y-C, Hung C-C, Hwang S-J, Wang S-L, Hsiao S-M, Lin M-Y, et al. Quality of life predicts risks of end-stage renal disease and mortality in patients with chronic kidney disease. *Nephrology Dialysis Transplantation*. 2010;25(5):1621-6.
22. Rumsfeld SJ, Mawhinney AWS, McCarthy BM, Shroyer EL, Villanueva GC, O'brien LM, et al. Health-Related Quality of Life As a Predictor of Mortality Following Coronary Artery Bypass Graft Surgery. *Survey of Anesthesiology*. 2000;44(6):326-.
23. Tsai S-Y, Chi L-Y, Lee C-H, Chou P. Health-related quality of life as a predictor of mortality among community-dwelling older persons. *European journal of epidemiology*. 2007;22(1):19-26.
24. Fan VS, Au DH, McDonnell MB, Fihn SD. Intraindividual change in SF-36 in ambulatory clinic primary care patients predicted mortality and hospitalizations. *Journal of Clinical Epidemiology*. 2004;57(3):277-83.
25. Gomez-Olive FX, Thorogood M, Bocquier P, Mee P, Kahn K, Berkman L, et al. Social conditions and disability related to the mortality of older people in rural south africa. *International Journal of Epidemiology*. 2014;43(5):1531-41.
26. Myint PK, Smith RD, Luben RN, Surtees PG, Wainwright NWJ, Wareham NJ, et al. The Short-Form Six-Dimension utility index predicted mortality in the European Prospective Investigation into Cancer-Norfolk prospective population-based study. *Journal of clinical epidemiology*. 2010;63(2):192-8.
27. Forsyth SJ, Carroll M, Lennox N, Kinner SA. Incidence and risk factors for mortality after release from prison in Australia: A prospective cohort study. *Addiction*. 2018;113(5):937-45.

28. Linde L, Sørensen J, Ostergaard M, Hørslev-Petersen K, Hetland ML. Health-related quality of life: validity, reliability, and responsiveness of SF-36, 15D, EQ-5D corrected RAQoL, and HAQ in patients with rheumatoid arthritis. *The Journal of rheumatology*. 2008;35(8):1528.
29. Liira H, Mavaddat N, Eineluoto M, Kautiainen H, Strandberg T, Suominen M, et al. Health-related quality of life as a predictor of mortality in heterogeneous samples of older adults. *European Geriatric Medicine*. 2018;9(2):227-34.
30. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009;339(jul21 1).
31. Booth A, Clarke M, Ghera D, Moher D, Petticrew M, Stewart L. An international registry of systematic-review protocols. *The Lancet*. 2011;377(9760):108-9.
32. Phyo. AZZ, Craig. H, Gonzalez-Chica. DA, Stocks. N, Freak-Poli. R, Ryan. J, et al. Quality of life as a predictor of mortality: a systematic review and meta-analysis PROSPERO 2019 CRD42019139994 [cited 2019 September 8,]. Available from: https://www.crd.york.ac.uk/prospéro/display_record.php?ID=CRD42019139994.
33. The Ottawa Hospital Research Institute. The Newcastle-Ottawa Scale (NOS) for assessing the quality if nonrandomized studies in meta-analyses [cited 2019 July 10]. Available from: http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.
34. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0. The Cochrane Collaboration 2011 [cited 2019 August 8]. Available from: <http://training.cochrane.org/handbook>.
35. Bjorkman MP, Pitkala KH, Jyvakorpi S, Strandberg TE, Tilvis RS. Bioimpedance analysis and physical functioning as mortality indicators among older sarcopenic people. *Experimental Gerontology*. 2019;122:42-6.
36. Brown D, Thompson W, Zack M, Arnold S, Barile J. Associations Between Health-Related Quality of Life and Mortality in Older Adults. *Prevention Science*. 2015;16(1):21-30.
37. Cavrini G, Broccoli S, Puccini A, Zoli M. EQ-5D as a predictor of mortality and hospitalization in elderly people. *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation*. 2012;21(2):269-80.
38. Chwastiak LA, Rosenheck RA, Desai R, Kazis LE. Association of psychiatric illness and all-cause mortality in the national Department of Veterans Affairs health care system. *Psychosomatic Medicine*. 2010;72(8):817-22.
39. De Buyser S, Petrovic M, Taes Y, Toye K, Kaufman JM, Goemaere S, et al. Three year functional changes and long-term mortality hazard in community-dwelling older men. *European Journal of Internal Medicine*. 2016;35:66-72.
40. De Buyser SL, Petrovic M, Taes YE, Toye KRC, Kaufman JM, Goemaere S. Physical function measurements predict mortality in ambulatory older men. *European Journal of Clinical Investigation*. 2013;43(4):379-86.
41. DeSalvo KB, Fan VS, McDonnell MB, Fihn SD. Predicting mortality and healthcare utilization with a single question. *Health services research*. 2005;40(4):1234-46.
42. Dominick KL, Ahern FM, Gold CH, Heller DA. Relationship of health-related quality of life to health care utilization and mortality among older adults. *Aging clinical and experimental research*. 2002;14(6):499-508.

43. Dorr DA, Jones SS, Burns L, Donnelly SM, Brunner CP, Wilcox A, et al. Use of health-related, quality-of-life metrics to predict mortality and hospitalizations in community-dwelling seniors. *Journal of the American Geriatrics Society*. 2006;54(4):667-73.
44. Drageset J, Eide GE, Ranhoff AH. Mortality in nursing home residents without cognitive impairment and its relation to self-reported health-related quality of life, sociodemographic factors, illness variables and cancer diagnosis: a 5-year follow-up study. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*. 2013;22(2):317-25.
45. Fan VS, Maciejewski ML, Liu CF, McDonell MB, Fihn SD. Comparison of risk adjustment measures based on self-report, administrative data, and pharmacy records to predict clinical outcomes. *Health Services and Outcomes Research Methodology*. 2006;6(1-2):21-36.
46. Feeny D, Huguot N, McFarland BH, Kaplan MS, Orpana H, Eckstrom E. Hearing, mobility, and pain predict mortality: a longitudinal population-based study. *Journal of clinical epidemiology*. 2012;65(7):764-77.
47. Franks P, Gold MR, Fiscella K. Sociodemographics, self-rated health, and mortality in the US. *Social Science and Medicine*. 2003;56(12):2505-14.
48. Han SS, Kim KW, Na KY, Chae DW, Kim YS, Kim S, et al. Quality of life and mortality from a nephrologist's view: A prospective observational study. *BMC Nephrology*. 2009;10(1):39.
49. Haring R, Feng Y-S, Moock J, Völzke H, Dörr M, Nauck M, et al. Self-perceived quality of life predicts mortality risk better than a multi-biomarker panel, but the combination of both does best. *BMC medical research methodology*. 2011;11(1):103.
50. Higuera-Fresnillo S, Cabanas-Sanchez V, Garcia-Esquinas E, Rodriguez-Artalejo F, Martinez-Gomez D. Physical activity attenuates the impact of poor physical, mental, and social health on total and cardiovascular mortality in older adults: a population-based prospective cohort study. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*. 2018;27(12):3293-302.
51. Jia H, Lubetkin EI, Demichele K, Stark DS, Zack MM, Thompson WW. Comparing the Performance of 2 Health Utility Measures in the Medicare Health Outcome Survey (HOS). *Medical Decision Making*. 2018;38(8):983-93.
52. Kao S, Lai K-L, Lin H-C, Lee H-S, Wen H-C. WHOQOL-BREF as predictors of mortality: A two-year follow-up study at veteran homes. *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation*. 2005;14(6):1443-54.
53. Kaplan MS, Berthelot JM, Feeny D, McFarland BH, Khan S, Orpana H. The predictive validity of health-related quality of life measures: Mortality in a longitudinal population-based study. *Quality of Life Research*. 2007;16(9):1539-46.
54. Kroenke CH, Kubzansky LD, Adler N, Kawachi I. Prospective change in health-related quality of life and subsequent mortality among middle-aged and older women. *American journal of public health*. 2008;98(11):2085-91.
55. Lawler FH, Mold JW, McCarthy LH. Do older people benefit from having a confidant? An Oklahoma Physicians Resource/Research Network (OKPRN) study. *Journal of the American Board of Family Medicine : JABFM*. 2013;26(1):9-15.

56. Lee MS, Chen RCY, Chang YH, Huang YC, Wahlqvist ML. Physical function mitigates the adverse effects of being thin on mortality in a free-living older Taiwanese cohort. *The journal of nutrition, health & aging*. 2012;16(9):776-83.
57. Leigh L, Hudson IL, Byles JE. Sleeping difficulty, disease and mortality in older women: A latent class analysis and distal survival analysis. *Journal of Sleep Research*. 2015;24(6):648-57.
58. Masel MC, Ostir GV, Ottenbacher KJ. Frailty, mortality, and health-related quality of life in older Mexican Americans. *Journal of the American Geriatrics Society*. 2010;58(11):2149-53.
59. Mold JW, Lawler F, Roberts M. The health consequences of peripheral neurological deficits in an elderly cohort: An Oklahoma physicians resource/research network study. *Journal of the American Geriatrics Society*. 2008;56(7):1259-64.
60. Munoz MA, Subirana I, Elosua R, Covas MI, Baena-Diez JM, Ramos R, et al. Utility of a short quality of life questionnaire to predict cardiovascular events. *International Journal of Cardiology*. 2011;151(3):392-4.
61. Murray C, Brett CE, Starr JM, Deary IJ. Which aspects of subjectively reported quality of life are important in predicting mortality beyond known risk factors? The Lothian Birth Cohort 1921 Study. *Quality of Life Research: An International Journal of Quality of Life Aspects of Treatment, Care & Rehabilitation*. 2011;20(1):81-90.
62. Myint PK, Luben RN, Surtees PG, Wainwright NWJ, Welch AA, Bingham SA, et al. Relation Between Self-Reported Physical Functional Health and Chronic Disease Mortality in Men And Women in the European Prospective Investigation Into Cancer (EPIC-Norfolk): A Prospective Population Study. *Annals of Epidemiology*. 2006;16(6):492-500.
63. Myint PK, Luben RN, Surtees PG, Wainwright NWJ, Welch AA, Bingham SA, et al. Self-reported mental health-related quality of life and mortality in men and women in the European Prospective Investigation into Cancer (EPIC-Norfolk): a prospective population study. *Psychosomatic medicine*. 2007;69(5):410-4.
64. Nilsson G, Ohrvik J, Lonnberg I, Hedberg P. Low psychological General Well-Being (PGWB) is associated with deteriorated 10-year survival in men but not in women among the elderly. *Archives of Gerontology and Geriatrics*. 2011;52(2):167-71.
65. Otero-Rodriguez A, Leon-Munoz LM, Balboa-Castillo T, Banegas JR, Rodriguez-Artalejo F, Guallar-Castillon P. Change in health-related quality of life as a predictor of mortality in the older adults. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*. 2010;19(1):15-23.
66. Perera S, Studenski S, Chandler JM, Guralnik JM. Magnitude and Patterns of Decline in Health and Function in 1 Year Affect Subsequent 5-Year Survival. *The Journals of Gerontology: Series A: Biological Sciences and Medical Sciences*. 2005;60(7):894-900.
67. Razzaque A, Mustafa AHMG, Streatfield PK. Do self-reported health indicators predict mortality? Evidence from Matlab, Bangladesh. *Journal of Biosocial Science*. 2014;46(5):621-34.
68. Singh JA, Borowsky SJ, Nugent S, Murdoch M, Zhao Y, Nelson DB, et al. Health-related quality of life, functional impairment, and healthcare utilization by veterans: veterans' quality of life study. *Journal of the American Geriatrics Society*. 2005;53(1):108-13.
69. St. John PD, Jiang D, Tate RB. Quality of life trajectories predict mortality in older men: The Manitoba Follow-Up Study. *Journal of Aging and Health*. 2018;30(2):247-61.

70. Sutcliffe C, Burns A, Challis D, Mozley CG, Cordingley L, Bagley H, et al. Depressed mood, cognitive impairment, and survival in older people admitted to care homes in England. *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry*. 2007;15(8):708-15.
71. Tibblin G, Svärdsudd K, Welin L, Erikson H, Larsson B. Quality of life as an outcome variable and a risk factor for total mortality and cardiovascular disease: a study of men born in 1913. *Journal of hypertension Supplement : official journal of the International Society of Hypertension*. 1993;11(4):S81.
72. Tice JA, Kanaya A, Hue T, Rubin S, Buist DSM, Lacroix A, et al. Risk Factors for Mortality in Middle-aged Women. *Archives of Internal Medicine*. 2006;166(22):2469-77.
73. Ul-Haq Z, Mackay DF, Pell JP. Association between physical and mental health-related quality of life and adverse outcomes; a retrospective cohort study of 5,272 Scottish adults. *BMC public health*. 2014;14:1197.
74. Williams ED, Rawal L, Oldenburg BF, Renwick C, Shaw JE, Tapp RJ. Risk of cardiovascular and all-cause mortality: Impact of impaired health-related functioning and diabetes - The Australian Diabetes, Obesity and Lifestyle (AusDiab) study. *Diabetes Care*. 2012;35(5):1067-73.
75. Xie G, Laskowitz DT, Turner EL, Egger JR, Shi P, Ren F, et al. Baseline health-related quality of life and 10-year all-cause mortality among 1739 Chinese adults. *PloS one*. 2014;9(7):e101527.
76. Makovski TT, Schmitz S, Zeegers MP, Stranges S, van Den Akker M. Multimorbidity and quality of life: Systematic literature review and meta-analysis. *Ageing Research Reviews*. 2019;53:100903.
77. Macnee W, Rabinovich RA, Choudhury G. Ageing and the border between health and disease. *The European respiratory journal*. 2014;44(5):1332.
78. Donat Tuna H, Ozcan Edeer A, Malkoc M, Aksakoglu G. Effect of age and physical activity level on functional fitness in older adults.(Report). *European Review of Aging and Physical Activity*. 2009;6(2):99.
79. Goldspink DF. Ageing and activity: their effects on the functional reserve capacities of the heart and vascular smooth and skeletal muscles. *Ergonomics*. 2005;48(11-14):1334.
80. Brach JS, Fitzgerald S, Newman AB, Kelsey S, Kuller L, Vanswearingen JM, et al. Physical Activity and Functional Status in Community-Dwelling Older Women: A 14-Year Prospective Study. *Archives of Internal Medicine*. 2003;163(21):2565-71.
81. Ju S-Y, Lee J-Y, Kim D-H. Association of metabolic syndrome and its components with all-cause and cardiovascular mortality in the elderly: A meta-analysis of prospective cohort studies. *Medicine*. 2017;96(45):e8491.
82. Bakas T, McLennon S, Carpenter J, Buelow J, Otte J, Hanna K, et al. Systematic review of health-related quality of life models. *Health and Quality of Life Outcomes*. 2012;10(1):134.
83. Karimi M, Brazier J. Health, Health-Related Quality of Life, and Quality of Life: What is the Difference? *Pharmacoeconomics*. 2016;34(7):645-9.
84. Hajian-Tilaki K, Heidari B, Hajian-Tilaki A. Are Gender Differences in Health-related Quality of Life Attributable to Sociodemographic Characteristics and Chronic Disease Conditions in Elderly People? *International journal of preventive medicine*. 2017;8:95.
85. Fryback GD, Dunham CN, Palta AM, Hanmer DJ, Buechner MJ, Cherepanov GD, et al. US Norms for Six Generic Health-Related Quality-of-Life Indexes From the National Health Measurement Study. *Medical Care*. 2007;45(12):1162-70.

86. González-Chica D, Adams R, Dal Grande E, Avery J, Hay P, Stocks N. Lower educational level and unemployment increase the impact of cardiometabolic conditions on the quality of life: results of a population-based study in South Australia. *An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation - Official Journal of the International Society of Quality of Life Research*. 2017;26(6):1521-30.
87. Frieling MA, Davis WR, Chiang G. The SF-36v2 and SF-12v2 health surveys in New Zealand: norms, scoring coefficients and cross-country comparisons. *Australian and New Zealand Journal of Public Health*. 2013;37(1):24-31.
88. Mishra G, Schofield M. Norms for the Physical and Mental Health Component Summary Scores of the SF-36 for Young, Middle-Aged and Older Australian Women. *An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation - Official Journal of the International Society of Quality of Life Res*. 1998;7(3):215-20.

Tables

Table 1. Characteristics of the 47 included studies

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|---------------------------------|-------------------|--|-------------|--------------------|--|-------------------|---------------|---|--|--|
| Orkman <i>et al.</i> 2019 [5] | Finland | Porvoo Sarcopenia and Nutrition Trial, Prospective | 428 | 4 yrs | 75 yrs and + 66.59 % | RAND-36 PF | all-cause | HR, 1-unit increase | PF: 0.988 (0.979 - 0.997) | age, sex, comorbidity and CRI-SMI |
| Johnson <i>et al.</i> 2015 [3]* | USA | Medicare HOS (Cohort 6-8), Prospective | 191,001 | 2.5 yrs | 65 yrs and + 58.30 % | CDC HRQOL-4 | all-cause | HR, Excellent vs. Poor HR, 0 days vs. 21-30 days | GH: 0.24 (0.21 - 0.27) Days of not good in Physical Health 0.82 (0.77 - 0.88) Days of not good in Mental Health 1.12 (1.04 - 1.22) Days of activity limitation 0.74 (0.68 - 0.79) | age, sex, race/ethnicity, education, income, range of other health and lifestyle factors |
| Verini <i>et al.</i> 2012 [7] | Italy | Pianoro Study, Prospective | 5,256 | 2 yrs | 65 yrs and + 55.3 % | EQ-5D | all-cause | HR, 1-unit increase | 0.42 (0.35 - 0.50) | sex, age, BMI, education, health and lifestyle factors |
| Wastiak <i>et al.</i> 2010 [3] | USA | 1999 Large Health Survey of Veteran Enrollees, Prospective | 559,985 | 9 yrs | 64.1 (12.9) yrs 4.1 % | SF-36 PCS | all-cause | HR, 1-unit increase | PCS: 0.97 (0.96 - 0.98) | age, race, sex, education, disability, comorbidity, BMI, lifestyle factors |
| Buyser <i>et al.</i> 2016 [9]* | Belgium | Prospective cohort | 171 | 15 yrs | 71 yrs and + 0 % | SF-36 PFI | all-cause | HR, 1-unit increase | PF: 1.01 (0.99 - 1.02) | age, polypharmacy, depression, and disability |
| Buyser <i>et al.</i> 2013 [9]* | Belgium | Prospective cohort | 352 | 15 yrs | 71 to 86 yrs 0 % | SF-36 PFI | all-cause | HR, 1-unit increase | PF: 0.992 (0.986 - 0.999) | BMI and smoking |
| Salvo <i>et al.</i> 2005 [1] | USA | VAAC Quality Improvement Project, Prospective | 21,732 | 1 yr | 64 (12) yrs 3.6 % | SF-36 PCS and MCS | all-cause | AUC | PCS: 0.73 (0.71 - 0.75) MCS: 0.68 (0.66 - 0.70) | age |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|--------------------------------|-------------------|---|-------------|--------------------|--|----------------------|---------------|---|--|--|
| Minick <i>et al.</i> 2002 [2]* | USA | Pennsylvania's Pharmaceutical Assistance Contract for the Elderly, Prospective | 84,065 | 1 yr | 78.7 (6.9) yrs 78.0 % | Core CDC HRQOL items | all-cause | RR, Excellent vs. Poor RR, 0 days vs. 21-30 days | GH: 0.24 (0.17 - 0.33) Days of not good in Physical Health 0.42 (0.38 - 0.45) Days of not good in Mental Health 0.53 (0.50 - 0.59) Days of activity limitation 0.40 (0.37 - 0.42) | age, sex, race, marital and residential status, income and comorbidity |
| Worr <i>et al.</i> 2006 [43]* | USA | Intermountain Health Care Network, Prospective | 2,166 | 2.3 yrs | 77.9 (6.8) yrs 54.9 % | SF-12 PCS and MCS | all-cause | OR, Quartile 4 (Highest) vs. Quartile 1 (Lowest) | PCS: 0.16 MCS: 0.40 | age, sex, and comorbidity |
| Ageiset <i>et al.</i> 2013 [4] | Norway | Study of Nursing Home Residents without cognitive impairment (2004-2005), Prospective | 227 | 5 yrs | 65 to 95 yrs and + 72.25 % | SF-36 PCS and MCS | all-cause | HR, 1-unit increase | PF: 0.99 (0.98 - 0.99) | age, sex, marital status, education and comorbidity |
| Worr <i>et al.</i> 2004 [24]* | USA | VAAC Quality Improvement Project, Prospective | 7,702 | 1 yr | 65.4 (10.6) yrs 3.4 % | SF-36 PCS and MCS | all-cause | OR, 1-unit increase | PCS: 0.956 (0.943 - 0.969) MCS: 0.981 (0.971 - 0.990) | age, site, distance to the VA, and comorbidity |
| Worr <i>et al.</i> 2006 [5] | USA | VAAC Quality Improvement Project, Prospective | 14,192 | 3 yrs | 64.4 (11.3) yrs 3.5 % | SF-36 PCS and MCs | all-cause | AUC | PCS: 0.721 (0.708 - 0.733) MCS: 0.689 (0.675 - 0.702) | age and sex |
| Worr <i>et al.</i> 2012 [5] | Canada | 1994/95 Canadian National Population Health Longitudinal Survey, Prospective | 12,375 | 12 yrs | 18 - 80 yrs + 52 % | HUI3 | all-cause | HR, 1-level increase | Hearing: 0.18 (0.06 - 0.57) Ambulation: 0.10 (0.04 - 0.23) Pain: 0.53 (0.29 - 0.96) | age, sex, socioeconomic, disease condition, and lifestyle factors |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|--------------------------------------|-------------------|---|-------------|--------------------|--|-------------------------------|---------------|--|---|---|
| rsyth <i>et al.</i> 2018 [7]* | Australia | RCT of a case Management Intervention for Adult transitioning from prison to the community, Prospective | 1,320 | 4.7 yrs | 32.7 (11.1) yrs 21.10% | SF-36 PCS and MCS | all-cause | HR, High vs. Low | PCS: 0.48 (0.18 - 1.20) MCS: 0.38 (0.16 - 0.91) *(CI is 99%CI) | age, sex and indigenous status |
| anks <i>et al.</i> 2003 [7]* | USA | NMES, Prospective | 21,363 | 5 yrs | 21 yrs + 55.39% | SF-20 | All-cause | HR, 1-point increase | HP: 0.993 (0.990 - 0.996) PF: 0.995 (0.992 - 0.997) RF: 0.996 (0.994 - 0.998) MH: 1.00 (0.996 - 1.003) | age, sex, race, ethnicity, education and income |
| meze-ive <i>et al.</i> 2014 [5]* | South Africa | Population under the Agincourt Health and Demographic Surveillance System, Prospective | 4,047 | 3 yrs | 50 yrs + 75.8 % | WHO QOL | all-cause | HR, Highest vs. Lowest | Overall: 0.61 | age, sex, education and union status, HH assets, and Disability Assessment |
| in <i>et al.</i> 09 [48] | South Korea | Korea Longitudinal Study on Health and Aging, Prospective | 944 | 3.25 yrs (median) | 76.0 (8.6) yrs 54.9 % | SF-36 PCS and MCS (K.V) | all-cause | HR, Tertile 3 (High) vs. Tertile 1 (Low) | PCS: 0.35 (0.19 - 0.64) MCS: 0.39 (0.22 - 0.70) | age, sex, smoking, range of serum measures |
| uring <i>et al.</i> 2011 [3]* | Germany | Population-based Study of Health in Pomerania, Prospective | 4,261 | 9.7 yrs (mean) | 20-79 yrs 50.93 % | SF-12 PCS and MCS | all-cause | HR, Highest Quartile vs. Lowest Quartile | PCS: 0.56 (0.42 - 0.75)* PCS: 0.63 (0.47 - 0.84)# MCS: 0.94 (0.73 - 1.22)* MCS: 1.04 (0.81 - 1.35)# | age, sex, * behavioural factors, # comorbidities |
| guas-esnillo <i>et al.</i> 2018 [1]* | Spain | UAM Cohort, Prospective | 3,922 | 14 yrs (median) | 71.82 (7.94) yrs 56.38 % | SF-36 PCS and MCS | all-cause | HR, Good vs. Poor | Physical: 0.74 (0.65 - 0.85) Mental: 0.85 (0.74 - 0.98) Social: 0.73 (0.63 - 0.85) | age, sex, education, lifestyle factors, BMI, waist circumference, comorbidity |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|-----------------------------|-------------------|--|-------------|--------------------|--|-------------------------------|---------------|---|--|---|
| Chen <i>et al.</i> 18 [51]* | USA | Medicare HOS Cohort 15, Prospective | 105,473 | 2 yrs | 65 yrs + 58.30 % | SF-6D and dEQ-5D | all-cause | HR, 1 st Quintile vs. 5th Quintile | SF-6D: 0.77 (0.71 - 0.80) dEQ-5D: 0.45 (0.43 - 0.49) | age, sex, socioeconomic, marital status, smoking, BMI, chronic conditions |
| Chen <i>et al.</i> 19 [52] | Taiwan | Prospective Cohort | 689 | 2 yrs | 65 yrs + 0 % | WHOQOL-(BREF) | all-cause | RR, 1-point change | Overall: 0.99 (0.77 - 1.26) | unadjusted RR |
| Chen <i>et al.</i> 20 [53] | Canada | 1994/95 Canadian National Population Health Longitudinal Survey, Prospective | 12,375 | 8 yrs | 18 - 80 yrs + 52 % | HUI3 | all-cause | HR, 1-unit increase | 0.61 (0.42 - 0.89) | age, sex, socioeconomic, other social/health, lifestyle factors |
| Chen <i>et al.</i> 21 [54] | USA | Nurses' Health Study, Prospective | 40,337 | 2.8 to 12 yrs | 46 - 71 yrs 100 % | SF-36 PCS and MCS | all-cause | RR ^a , Severe Decline vs. No Change RR ^b , Improve vs. No Change | Change in PCS 3.32 ^a (2.45 - 4.50) 0.72 ^b (0.56 - 0.91) Change in MCS 1.86 ^a (1.17 - 2.97) 0.77 ^b (0.63 - 0.95) | age, baseline HRQoL, menopausal status, social integration, BMI, educational, husbands' education, lifestyle factors, PCS/MCS |
| Chen <i>et al.</i> 22 [55] | USA | Oklahoma Longitudinal Assessment of HOMAS, Prospective | 852 | 5 yrs | 65 yrs + 56.81 % | SF-36 PCS and MCS | all-cause | HR, 1-unit increase | PF: 0.98 (0.97 - 0.98) Bodily Pain: 1.01 (1.00 - 1.01) | age, sex, socioeconomic, BMI, morbidity, functional status, having a confidant |
| Chen <i>et al.</i> 23 [56]* | Taiwan | Elderly Nutrition and Health Survey, Prospective | 1,435 | 7.9 yrs | 65 - 97 yrs 48.50 % | SF-36 PCS (T.V 1.0) | all-cause | HR, Highest PF vs. Lowest PF | PF: 0.29 (0.19 - 0.45) | age |
| Chen <i>et al.</i> 24 [57] | Australia | Australian Longitudinal Study on Women's Health, Prospective | 10,721 | 15 yrs | 70 - 75 yrs 100 % | SF-36 Vitality, Mental and PF | all-cause | HR, 1-unit increase | PF: 0.992 (0.990 - 0.994) Mental: 1.0 (0.997 - 1.002) Vitality: 1.0 (0.998 - 1.002) | age, socioeconomic, BMI, sleep, disease count, and other health factors |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|-------------------------------|-------------------|---|---|--------------------|--|--------------------------|---------------|---|---|--|
| Luoma <i>et al.</i> 2018 [29] | Finland | a. The Helsinki Businessmen Study (HBS) b. Spousal caregivers of people with dementia c. Nursing home residents d. Older persons suffering from loneliness e. Population Sample | a = 733 b = 209 c = 326 d = 208 e = 901 | 2 yrs | a. 77 (4) yrs 0 % b. 75 (7) yrs 64.6 % c. 84 (7) yrs 69.9 % d. 80 (4) yrs 75 % e. 85 (5) yrs 75.1 % | The 15D | all-cause | HR, 1SD (0.14) increase | a. 0.43 (0.31 - 0.63) b. 1.06 (0.43 - 2.63) c. 0.69 (0.58 - 0.85) d. 0.94 (0.47 - 1.87) e. 0.62 (0.49 - 0.72) | age and sex |
| Wong <i>et al.</i> 2010 [3] | USA | Hispanic Established Population for Epidemiologic Study of the Elderly, Prospective | 1,008 | 2 yrs | 74 - 101 yrs 63.2 % | SF-36 PCS and MCS | all-cause | OR, 1-point increase | PCS: 0.962 (0.941 - 0.984) MCS: 0.996 (0.974 - 1.018) | age, sex, education, marital status, financial strain, chronic illness, smoking, BMI, and frailty |
| Wong <i>et al.</i> 2008 [59] | USA | Oklahoma Longitudinal Assessment of HOMAS, Prospective | 604 | 5 yrs | 65 yrs + 56 % | SF-36 PF and bodily pain | all-cause | HR, 1-unit increase | PF: 0.98 (0.97 - 0.99) | education, income, smoking, initial and instrumental activity of daily living, health utilities / conditions |
| Wong <i>et al.</i> 2011 [6] | Spain | Prospective Cohort | 3,724 | 6.3 yrs (median) | 35 - 74 yrs 51.9 % | SF-12 PCS and MCS | all-cause | HR, 3 rd Tertile vs. 1st Tertile (Low) | PCS: 0.58 (0.39 - 0.87) MCS: 0.99 (0.69 - 1.42) | age, sex, marital status, education and cardiovascular risk factors |
| Wong <i>et al.</i> 2011 [1] | Scotland | Lothian Birth Cohort 1921, Prospective | 448 | 9 yrs | 79 yrs 56.70% | 26-item WHOQOL-BREF | all-cause | HR, 1 tertile increase / 1-point increase | Overall: 0.84 (0.67 - 1.05) GH: 0.75 (0.64 - 0.89) Physical: 0.90 (0.86 - 0.95) Psychological: 0.98 (0.91 - 1.06) Social: 0.97 (0.91 - 1.04) Environment: 0.96 (0.89 - 1.03) | age and sex |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|-------------------------|-------------------|--|-------------|--------------------|--|----------------------------------|---------------|---|--|--|
| Wright et al. 2006 [2]* | UK | EPIC-Norfolk, Prospective | 17,777 | 6.5 yrs (mean) | 41-80 yrs 56.25 % | SF-36 PCS (UK.V) | all-cause | RR, Quintiles 5 (Highest) vs. Quintiles 1 | PCS Men: 0.47 (0.33 - 0.65) Women: 0.41 (0.27 - 0.64) | age, BMI, SBP, blood cholesterol, smoking, diabetes and social class |
| Wright et al. 2007 [3]* | UK | EPIC-Norfolk, Prospective | 17,777 | 6.5 yrs (mean) | 40-79 yrs 56.25 % | SF-36 MCS (UK.V) | all-cause | HR, 1-point increase | MCS: 0.987 (0.981 - 0.993) | age, sex, PCS, lifestyle, BMI, SBP, blood cholesterol, diabetes, and social class |
| Wright et al. 2010 [5]* | UK | EPIC-Norfolk, Prospective | 17,736 | 6.5 yrs (mean) | 40-79 yrs 56.23 % | SF-6D (UK.V) | all-cause | HR, 1 SD (0.12-point) increase | 0.74 (0.69 - 0.79) | age, sex, BMI, SBP, blood cholesterol, diabetes, smoking, and social class |
| Wright et al. 2011 [1]* | Sweden | Inhabitants in the Swedish city of Vasteras, Prospective | 417 | 10 yrs | 75 yrs 51.08 % | PGWB | all-cause | RR, 1-unit change | Global Score Men: 0.984 (0.969 - 0.998) Women: 0.994 (0.978 - 1.010) | for men: smoking, obesity, living alone and other health conditions |
| Wright et al. 2010 [5]* | Spain | Spanish Population-Based Cohort, Prospective | 2,373 | 6 yrs | 60 yrs + 57.5 % | SF-36 PCS and MCS | all-cause | HR, 1-point increase | PCS: 0.952 (0.935 - 0.969) MCS: 0.990 (0.976 - 1.006) | sex, age, HRQOL, education, marital status, BMI, other health and lifestyle factors, PCS/MCS |
| Wright et al. 2005 [3]* | USA | Prospective cohort | 439 | 5 yrs | 65 yrs + 44.40 % | SF-36 PF | all-cause | HR, 1-point increase | PF: 0.991 (0.945 - 1.036) | age, sex, measure of change, number of comorbid domains, hospitalization |
| Wright et al. 2014 [7]* | Bangladesh | Matlab HDSS, Prospective | 4,037 | 2 yrs | 50 yrs + 50.06% | WHOQOL | all-cause | RR, Good/Very Good vs. Bad/Very Bad | Men: 0.26 (0.16 - 0.41) Women: 0.30 (0.10 - 0.86) | age and socio-demographic variables |
| Wright et al. 2005 [3]* | USA | Prospective | 40,508 | 1 yr | 64.5 (13.7) yrs 4.2 % | SF-36 PCS and MCS (V.V) | all-cause | OR, 1-point increase | PCS: 0.933 (0.926 - 0.941) MCS: 0.968 (0.962 - 0.973) | age, sex, socioeconomic, smoking, VA eligibility status, and prior healthcare utilization |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|---------------------------|-------------------|---|-------------|--------------------|--|-------------------|---------------|------------------------|---|--|
| Johnston et al. 2018 [1]* | Canada | Manitoba Follow-up Study, Prospective | 734 | 9 yrs | 85.5 (3.0) yrs 0 % | SF-36 PCS and MCS | all-cause | RR, High vs. Low | PCS: 0.50 (0.38 - 0.64) MCS: 0.55 (0.40 - 0.76) | age |
| McCliff et al. 2007 [2] | UK | Prospective | 308 | 0.75 yrs | 60 - 90 yrs + 68.8 % | LQOLP-R - Spitzer | all-cause | HR, increased score | 0.9805 (0.9704 - 0.9907) | unadjusted |
| Wahlblom et al. 1993 [3] | Sweden | Study of men born in 1913, Prospective | 787 | 18 yrs | 50 yrs + 0 % | Goteborg QoL | all-cause | No Data | Only Health variable was significantly related to mortality | health, physical fitness, and appetite |
| Wong et al. 2006 [72] | USA | B-FIT, Prospective | 17,748 | 9 yrs | 55 - 80 yrs + 100 % | SF-20 PF | all-cause | HR, Highest vs. Lowest | PF: 0.70 (0.60 - 0.90) | age, other health and lifestyle factors |
| Chen et al. 2007 [23]* | Taiwan | A 2000 Population-based survey in Taiwan, Prospective | 4,424 | 3 yrs | 65 yrs + | SF-36 PCS and MCS | all-cause | RR, 1-point increase | PCS: 0.954 (0.941 - 0.968) MCS: 0.985 (0.971 - 0.999) | age, sex, feel tired, other health and lifestyle factors |
| McHaugh et al. 2014 [3]* | Scotland | Scottish Health Survey 2003, Retrospective | 5,272 | 7.6 yrs (mean) | 20 - 65 yrs + 54.80 % | SF-12 PCS and MCS | all-cause | HR, Best vs. Worst | PCS: 0.36 (0.22 - 0.57) MCS: 0.80 (0.61 - 1.05) | age, sex, SIMD, education, BMI, other health and lifestyle factors |
| Williams et al. 2012 [4]* | Australia | AusDiab study, Prospective | 9,979 | 7.4 yrs | 25 yrs + 55.00 % | SF-36 PCS and MCS | all-cause | HR, 1-point change | PF: 0.983 (0.979 - 0.987) RP: 0.995 (0.993 - 0.997) Bodily Pain: 0.996 (0.992 - 0.999) GH: 0.985 (0.980 - 0.990) Vitality: 0.992 (0.987 - 0.996) Social F: 0.993 (0.990 - 0.996) RE: 0.999 (0.996 - 1.001) MH: 0.999 (0.994 - 1.004) | age, sex, BMI, smoking, health conditions, serum measures |

| Authors and Year | Setting - Country | Study Name and Design | Sample Size | Follow-up in years | Participants (Age in Range or Mean (SD), Female %) | QoL Measure | Type of Death | Comparison | Risk estimate (95% CI) | Adjustment |
|-------------------------------|-------------------|----------------------------|-------------|--------------------|--|------------------|---------------|-----------------------------|------------------------|---|
| Wang <i>et al.</i> 2014 [75]* | China | PRC-USA Study, Prospective | 1,739 | 10.1 yrs (median) | 57.7 (8.4) yrs 64.2 % | Chinese (QOL-35) | all-cause | HR, Upper 50% vs. Lower 50% | 0.69 (0.49 - 1.00) | age, sex, social-economic, other health and lifestyle factors |

AUC = Area under curve; **BMI** = Body Mass Index; **CDC HRQOL-4** = Core CDC Healthy Days Measures HRQOL-4; **Chinese (QOL-35)** = Chinese 35-item Quality of Life Instrument; **CRi-SMI** = Calf Intracellular Resistance Skeletal Muscle Index; **EQ-5D** = the EuroQoL-5 Dimension; **GH** = General Health; **HUI3** = The Health Utilities Index Mark 3 Version; **HH** = Household; **HP** = Health Perceptions; **HR** = Hazard Ratio; **K.V** = Korea Version; **LQOLP-R - Spitzer** = Lancashire Quality-of-Life Profile-Residential incorporated the Spitzer Uniscale; **MCS** = Mental Component Score; **MH** = Mental Health; **OR** = Odds Ratio; **PCS** = Physical Component Score; **PF** = Physical Functioning; **PGWB** = Psychological General Well-Being; **QoL** = Quality of Life; **RE** = Role-Emotional; **RF** = Role Function; **RP** = Role Physical; **RR** = Relative Risk; **SF-36** = Short Form 36; **SF-20** = Short Form 20; **SF-12** = Short Form 12; **SF-6D** = Short-Form Six Dimension Utility Index; **SBP** = Systolic Blood Pressure; **Social F** = Social Functioning; **SIMd** = Scottish Index of Multiple deprivation; **The 15D** = The 15 dimensional instrument; **T.V** = Taiwan Version; **UK** = United Kingdom; **UK.V** = UK Version; **USA** = United States of America; **VA** = Veterans Affairs; **V.V** = Veterans Version;

*where studies report reverse association or risk estimate per more than 1-unit increase, the risk estimates were standardised per 1-unit increase or 1-SD increase or high vs. low for the purpose of consistency across the table

Study Abbreviation

AusDiab = Australian Diabetes, Obesity and Lifestyle; **B-FIT** = Breast and Bone Follow-up Study of the Fracture Intervention Trial; **EPIC-Norfolk** = European Prospective Investigation into Cancer - Norfolk; **Matlab HDSS** = Matlab Health and Demographic Surveillance System of the International Centre for Diarrhoeal Disease Research; **Medicare HOS** = Medicare Health Outcomes Survey; **NMES** = Household Survey component of the National Medical Expenditure; **Oklahoma Longitudinal Assessment of HOMAS** = Oklahoma Longitudinal Assessment of Health Outcomes of Mature Adults Studies; **PRC-USA Study** = People's Republic of China-United States of America Chinese Collaborative Study of Cardiovascular and Cardiopulmonary Epidemiology; **VAAC** = Veterans Affairs Ambulatory Care;

Table 2. Quality of life scale included in the systematic review

| QoL Scale | | Study |
|--|---|---|
| Short Form Health Survey scales | SF-36, SF-20, SF-12, RAND-36 | Study [23, 24, 27, 35, 38-41, 43-45, 47-50, 54-60, 62, 63, 65, 66, 68, 69, 72-74] |
| World Health Organization questionnaires | WHOQOL, WHOQOL-BREF | Study [25, 52, 61, 67] |
| Centre for Diseases Control and Prevention Health Related Quality of Life scale | CDC HRQOL | Study [36, 42] |
| Six Dimensions Short Form Scale | SF-6D | Study [26, 51] |
| Euro Quality of Life scale | EQ-5D | Study [37, 51] |
| Health Utilities Index 3 | HUI3 | Study [46, 53] |
| Psychological General Well-Being Index | PGWB | Study [64] |
| 15-dimensional index | 15D | Study [29] |
| Goteborg Quality of Life Instrument | Goteborg QoL | Study [71] |
| Lancashire Quality of Life Profile-Residential incorporated the Spitzer Uniscale | LQOLP-Residential incorporated the Spitzer Uniscale | Study [70] |
| Chinese 35-Item Quality of Life Instrument | Chinese QOL-35 | Study [75] |

Table 3. Physical component score / physical functioning as predictors of all-cause mortality

| Author (Year) | Comparison | Effect estimate (95% CI) |
|---|---|--|
| SF - 36 Physical Component Score (continuous) | | |
| Chwastiak <i>et al.</i> 2010 | HR, 1-unit increase | 0.97 (0.96 - 0.98) |
| DeSalvo <i>et al.</i> 2005 | AUC | 0.73 (0.71 - 0.75) |
| Fan <i>et al.</i> 2006 | AUC | 0.721 (0.708 - 0.733) |
| Otero-Rodriguez <i>et al.</i> 2010* | HR, 1-unit increase | 0.952 (0.935 - 0.969) |
| SF-36 Physical Function Scale (continuous) | | |
| De Buyser <i>et al.</i> 2016 a* | HR, 1-unit increase | 1.01 (0.99 - 1.02) |
| Mold <i>et al.</i> 2008 b | HR, 1-unit increase | 0.98 (0.97 - 0.99) |
| RAND-36 Physical Function Scale (continuous) | | |
| Bjorkman <i>et al.</i> 2019 | HR, 1-unit increase | 0.988 (0.979 - 0.997) |
| SF - 36 Physical Component Score (categorised) | | |
| Forsyth <i>et al.</i> 2018* | HR, High vs. Low | 0.48 (0.18 - 1.20) # |
| Han <i>et al.</i> 2009 | HR, Tertile 3 High vs. Tertile 1Low | 0.35 (0.19 - 0.64) |
| Higuera-Fresnillo <i>et al.</i> 2018* | HR, Good vs. Poor | 0.74 (0.65 - 0.85) |
| Myint <i>et al.</i> 2006* | RR, Quintile 5 Highest vs. Quintile 1 Lowest | 0.47 (0.33 - 0.65) Men 0.41 (0.27 - 0.64) Women |
| St.John <i>et al.</i> 2018* | RR, High vs. Low | 0.50 (0.38 - 0.64) |
| SF - 36 Physical Functioning (categorised) | | |
| Lee <i>et al.</i> 2012* | HR, Highest vs. Lowest | 0.29 (0.19 - 0.45) |
| SF - 36 Change in Physical Component Score (categorised) | | |
| Kroenke <i>et al.</i> 2008 | RR, Severe Decline vs. No Change | 3.32 (2.45 - 4.50) |
| | RR, Improvement vs. No Change | 0.72 (0.56 - 0.91) |
| SF - 20 Physical Function Scale (continuous) | | |
| Franks <i>et al.</i> 2003* | HR, 1-point increase | 0.995 (0.992 - 0.997) |
| SF - 20 Physical Function Scale (categorised) | | |
| Tice <i>et al.</i> 2006 | HR, Highest vs. Lowest | 0.70 (0.60 - 0.90) |
| SF - 12 Physical Component Score (categorised) | | |
| Dorr <i>et al.</i> 2006* | OR, Highest Quartile vs. Lowest Quartile | 0.16 |
| Haring <i>et al.</i> 2011* | HR, Highest Quartile vs. Lowest Quartile | 0.56 (0.42 - 0.75) c |
| | | 0.63 (0.47 - 0.84) d |
| Munoz <i>et al.</i> 2011 | HR, 3 rd Tertile vs. 1 st Tertile | 0.58 (0.39 - 0.87) |
| UI-Haq <i>et al.</i> 2014* | HR, Best Quintile vs. Worst Quintile | 0.36 (0.22 - 0.57) |

a. De Buyser *et al.* (2016) and De Buyser *et al.* (2013) were from the same study. De Buyser *et al.* (2013) was included in meta-analysis

b. Lawler *et al.* (2013) and Mold *et al.* (2008) were from the same study. Lawler *et al.* (2013) was included in meta-analysis

c. behavioural factors adjusted

d. comorbidities adjusted

CI is 99% CI

*where studies report reverse association or risk estimate per more than 1-unit increase, the risk estimates were standardised per 1-unit increase or 1-SD increase or high vs. low for the purpose of consistency across the table

AUC = Area under curve

Table 4. Mental component score / mental health as predictors of all-cause mortality

| Author (Year) | Comparison | Effect estimate (95% CI) |
|---|---|--------------------------|
| SF - 36 Mental Component Score (continuous) | | |
| DeSalvo <i>et al.</i> 2005 | AUC | 0.68 (0.66 - 0.70) |
| Fan <i>et al.</i> 2006 | AUC | 0.689 (0.675 - 0.702) |
| Myint <i>et al.</i> 2007* | HR, 1-unit increase | 0.987 (0.981 - 0.993) |
| Otero-Rodriguez <i>et al.</i> 2010* | HR, 1-unit increase | 0.990 (0.976 - 1.006) |
| SF - 36 Mental Health (continuous) | | |
| Leigh <i>et al.</i> 2015 | HR, 1-unit increase | 1.00 (0.997 - 1.002) |
| Williams <i>et al.</i> 2012* | HR, 1-point-change | 0.999 (0.994 - 1.004) |
| SF - 36 Mental Component Score (categorised) | | |
| Forsyth <i>et al.</i> 2018* | HR, High vs. Low | 0.38 (0.16 - 0.91) ## |
| Han <i>et al.</i> 2009 | HR, Tertile 3 High vs. Tertile 1Low | 0.39 (0.22 - 0.70) |
| Higuera-Fresnillo <i>et al.</i> 2018* | HR, Good vs. Poor | 0.85 (0.74 - 0.98) |
| St.John <i>et al.</i> 2018* | RR, High vs. Low | 0.55 (0.40 - 0.76) |
| SF - 36 Change in Mental Component Score (categorised) | | |
| Kroenke <i>et al.</i> 2008 | RR, Severe Decline vs. No Change | 1.86 (1.17 - 2.97) |
| | RR, Improvement vs. No Change | 0.77 (0.63 - 0.95) |
| SF - 20 Physical Function Scale (continuous) | | |
| Franks <i>et al.</i> 2003* | HR, 1-point increase | 1.00 (0.996 - 1.003) |
| SF - 12 Mental Component Score (categorised) | | |
| Dorr <i>et al.</i> 2006* | OR, Highest Quartile vs. Lowest Quartile | 0.40 |
| Haring <i>et al.</i> 2011* | HR, Highest Quartile vs. Lowest Quartile | 0.94 (0.73 - 1.22) a |
| | | 1.04 (0.81 - 1.35) b |
| Munoz <i>et al.</i> 2011 | HR, 3 rd Tertile vs. 1st Tertile | 0.99 (0.69 - 1.42) |
| UI-Haq <i>et al.</i> 2014* | HR, Best Quintile vs. Worst Quintile | 0.80 (0.61 - 1.05) |

a. behavioural factors adjusted

b. comorbidities adjusted

99% CI

*where studies report reverse association or risk estimate per more than 1-unit increase, the risk estimates were standardised per 1-unit increase or 1-SD increase or high vs. low for the purpose of consistency across the table

AUC = Area under curve

Table 5. Other QoL measures rather than SF / RAND, as predictor of all-cause mortality

| Author (Year) | Comparison | Effect estimate (95% CI) |
|---|---|---|
| Core CDC Healthy Days Measures (HRQOL-4) (General Health) categorised | | |
| Rowan <i>et al.</i> 2015* | HR, Excellent vs. Poor | 0.24 (0.21 - 0.27) |
| Dominiack <i>et al.</i> 2002* | RR, Excellent vs. Poor | 0.24 (0.17 - 0.33) |
| WHO QOL - BREF (Overall) | | |
| Woo <i>et al.</i> 2005 | RR, 1-point change | 0.99 (0.77 - 1.26) |
| Murray <i>et al.</i> 2011 | HR, 1-tertile increase | 0.84 (0.67 - 1.05) |
| WHO QOL (Categorised) | | |
| Combez-Olive <i>et al.</i> 2014* | HR, Highest vs. Lowest | 0.61 |
| Alzazque <i>et al.</i> 2014* | RR, Good vs. Bad | 0.26 (0.16 - 0.41) men 0.30 (0.10 - 0.86) women |
| Psychological General Well-being (PGWB) (Global Score) continuous | | |
| Nilsson <i>et al.</i> 2011* | RR, 1-unit change | 0.984 (0.969 - 0.998) men 0.994 (0.978 - 1.010) women |
| Manchester Quality-of-life Profile-Residential (LQOLP-R) incorporated the Spitzer Uniscale | | |
| Atcliffe <i>et al.</i> 2007 | HR, increased score | 0.9805 (0.9704 - 0.9907) |
| Chinese 35-item Quality of Life (QOL-35) categorised | | |
| Ye <i>et al.</i> 2014* | HR, Upper 50% vs. Lower 50% | 0.69 (0.49 - 1.00) |
| The Health Utilities Index Mark 3 Version (HUI3) continuous | | |
| Keeney <i>et al.</i> 2012 | HR, 1-level increase | Hearing: 0.18 (0.06 - 0.57) Ambulation: 0.10 (0.04 - 0.23) Pain: 0.53 (0.29 - 0.96) |
| Waplan <i>et al.</i> 2007 | HR, 1-unit increase | Overall: 0.61 (0.42 - 0.89) |
| The EuroQoL-5 Dimension (EQ-5D) continuous | | |
| Avrini <i>et al.</i> 2012 | HR, 1-unit increase | 0.42 (0.35 - 0.50) |
| The EuroQoL-5 Dimension EQ-5D categorised | | |
| Chen <i>et al.</i> 2018* | HR, 5 th Quintile vs. 1 st Quintile | 0.45 (0.43 - 0.49) |
| Short Form Six Dimension Utility Index (SF-6D) continuous | | |
| Wright <i>et al.</i> 2010 | HR, 1SD 0.12-point increase | 0.74 (0.69 - 0.79) |
| Short Form Six Dimension Health Utility Measure (SF-6D) categorised | | |
| Chen <i>et al.</i> 2018* | HR, 5th Quintile vs. 1st Quintile | 0.77 (0.71 - 0.80) |
| Stockholm Quality of Life Assessment | | |
| Wahlbin <i>et al.</i> 1993 | Only Health variable was significantly related to mortality (No data available) | |

*where studies report reverse association or risk estimate per more than 1-unit increase, the risk estimates were standardised per 1-unit increase or 1-SD increase or high vs. low for the purpose of consistency across the table

Figures




Figure 1

Flow Diagram of Review Process




Figure 2

Forest plot of all-cause mortality risk (odds ratio/relative risk) per one unit increase in SF-36 PCS. N: sample size; FU (yrs): follow-up in years; CI: confidence interval




Figure 3

Forest plot of all-cause mortality risk (hazard ratio) per one unit increase in Physical Functioning of SF-36. N: sample size; FU (yrs): follow-up in years; CI: confidence interval




Figure 4

Forest plot of all-cause mortality risk (odds ratio/relative risk) per one unit increase in SF-36 MCS. N: sample size; FU (yrs): follow-up in years; CI: confidence interval




Figure 5

Forest plot of all-cause mortality risk (hazard ratio) per one SD (0.14) increase in the 15D index score among five individual cohorts included in the Liira et al. study 2018. N: sample size; FU (yrs): follow-up in years; CI: confidence interval

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile1.docx](#)