β -Conglycinin regulates distal intestinal immunity through the CIITA-mediated MHC II-PI3K/Akt/mTOR signaling pathway in hybrid grouper (*Epinephelus fuscoguttatus* $\times$ *E. lanceolatus*): The ameliorative effects of sodium butyrate (NaB)

Bin Yin

Guangdong Ocean University

Hongyu Liu

Guangdong Ocean University

Beiping Tan (✉ bptan@126.com)

Fisheries College, Guangdong Ocean University

Xiaohui Dong

Guangdong Ocean University

Shuyan Chi

Guangdong Ocean University

Qihui Yang

Guangdong Ocean University

Shuang Zhang

Guangdong Ocean University

Research

Keywords: hybrid grouper (*Epinephelus fuscoguttatus* $\times$ *E. lanceolatus*), β -conglycinin, sodium butyrate, serum immunity, intestinal histopathology, intestinal inflammation, MHC II PI3K/Akt/mTOR

Posted Date: December 18th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-130234/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

β -Conglycinin (7S) is a strong immunogenic protein that modulates immune responses in the intestines of aquatic animals, whereas sodium butyrate (NaB) may promote growth by protecting the intestinal tract of fish. However, the mechanisms of action of 7S and NaB in these regards have yet to be determined.

Results

In this study, we investigated the effects of low and high doses of 7S and the ameliorative effects of NaB (based on high-dose 7S) on the growth performance, serum immunity, distal histopathology, and CIITA-mediated MHC II-PI3K/Akt/mTOR pathway in hybrid groupers (*Epinephelus fuscoguttatus* $\times$ *E. lanceolatus*). The results revealed that the specific growth rate of groupers significantly increased, decreased, and increased in the low-level 7S (bL), high-level 7S (bH) high-level 7S plus NaB (bH-NaB) groups, respectively. The feed coefficient ratio was significantly increased in the bH and bH-NaB groups, whereas serum levels of IgE, IFN- γ , IL-1 β , and TNF- α were upregulated in the bH group, and IgE was upregulated in the bH-NaB group. With respect to distal intestine histopathology, the intestinal diameter/plica height ratio was significantly increased in the bH group. Furthermore, there were increases in nitric oxide, nitric oxide synthase (NOS), and peroxynitrite anion (ONOO) in the bH group, and decreases in NOS and ONOO in the bH-NaB group. In the distal intestinal tract, the mRNA levels of TSC1, mTOR C2, CIITA, and CREB1 were significantly upregulated in all three treatment groups, whereas those of IKK α , Rheb, mTOR C1, mLST8, EIF4B, NFY, GILT, and AEP were upregulated and downregulated in the bH and bH-NaB groups, respectively.

Conclusions

Collectively, these results indicate that 7S has a regulatory effect on serum immunity and can also affect distal intestinal tract development in hybrid groupers by modulating hindgut injury-related parameters. Within the distal intestinal tract, 7S can induce intestinal inflammation by activating the CIITA-mediated MHC II-PI3K/Akt/mTOR pathway, which eventually manifests as a reduction in growth performance. Supplementing feed with NaB represents an effective approach for enhancing serum immunity, and also protects the intestines from damage caused by high doses of 7S.

Background

The hybrid grouper (*Epinephelus fuscoguttatus* $\times$ *E. lanceolatus*) is characterized by a rapid growth rate and desirable meat quality, which is of high economic value (Wu, et al., 2017). As a typical carnivorous marine fish, hybrid grouper typically require up to 50–55% protein in their feed (Jiang, et al., 2016), which is generally provided by fishmeal containing high levels of nitrogen and phosphorus

(Schneider, et al., 2004). In addition, soybean meal is also a widely available and inexpensive source of protein for hybrid grouper diets (Murray, et al., 2010). However, hybrid grouper have a low tolerance to soybean meal, which can make up approximately 40–60% of fishmeal diets (Firdaus, et al., 2016). Although a high intake of soybean meal can lead to intestinal inflammation in hybrid groupers (Zhang, et al., 2019), the provision of appropriate amounts of this meal can enhance immunity to a certain extent, thereby improving growth performance. These effects are widely believed to be attributable to the immunogenicity of the soy glycinin (11S) and β -conglycinin (7S) contents of soybean meal (Li, et al., 2017a), which are the main components of soybean storage proteins (Han, et al., 2018). β -conglycinin has a molecular weight of 140–180 kD, and accounts for approximately 30% of the soy protein content. Most of the 7S ingested as a nutrient is absorbed by the intestinal tract, and contributes little to the antigenic effect. In piglets, 7S allergic reactions mainly induce a reduction in the height of the intestinal villi and thereby reduces the efficiency of nutrient absorption (Hao, et al., 2009a), whereas in grass carp, 7S can promote an increase in reactive oxygen species (ROS) production in the distal intestine, leading to disruption of the structural integrity of epithelial cells (Xu, et al., 2019). However, the mechanisms underlying the intestinal tolerance to 7S and the intestinal inflammation caused by this compound in hybrid grouper have yet to be determined.

Short-chain fatty acids include acetic acid, propionic acid, and butyric acid, with the latter playing a major role in regulating the organism metabolism (Hamer, et al., 2007). It has been established that the hydrolysis of sodium butyrate (NaB: the sodium salt form of butyric acid) to butyric acid has a wide range of physiological effects, providing an energy source for intestinal cells, promoting intestinal cell differentiation and proliferation (Liu, et al., 2019b), and maintaining intestinal morphology to facilitate digestion and absorption (Liu, et al., 2014). Accordingly, given its unique nutritional and physiological functions and lack of polluting and harmful residues, NaB has become a valuable additive that is used to supplement the diets of cultured aquatic animals.

The intestinal tract of fish is well supplied by nerves and microvasculature, and is often referred to as the second brain of the fish. Moreover, it also contributes to the provision of 70% of fish immunity. Accordingly, as the most fully functional immune organ in fish, the intestines play a dual role in providing tolerance to feed antigens and immune defense. Unlike mammals, fish epitopes are primarily distributed in the hindgut, and consequently, compared with the foregut and midgut, the hindgut is more susceptible to attack by antigens present in feed, which can potentially cause intestinal inflammation. In this regard, studies have shown that the major histocompatibility complex MHC II plays an important role in identifying and presenting foreign antigens, and research in our laboratory has revealed that intestinal inflammation in hybrid grouper caused by ingesting feed containing high levels of soybean meal is related to an abnormal activation of the PI3K/Akt/mTOR signaling pathway. However, it has yet to be established whether this activation is due to the recognition of foreign antigens by MHC II. Furthermore, it has also been found that NaB can significantly inhibit the Akt/mTOR signaling pathway, and thus we speculate that by inhibiting PI3K/Akt/mTOR, NaB could be used to relieve the intestinal inflammation caused by 7S.

In this study, we therefore used hybrid groupers as experimental animals to evaluate the hypothesis that dietary β -conglycinin might affect growth by regulating serum immunity, distal intestinal histopathology, epithelial injury, and the PI3K/Akt/mTOR signaling pathway. In addition, we also investigated the potential protective effects of appropriate doses of NaB against high levels of dietary 7S. We anticipate that the findings of this study will provide a reference for research on improving the tolerance of hybrid grouper to soybean meal and reveal the mechanisms underlying the protective effects of NaB on the distal intestine.

Materials And Methods

Animals

The juvenile hybrid grouper used in this study were purchased from a commercial hatchery on East Island (Zhanjiang, China), and were derived from the same batch of artificially hatched fish. Prior to experimentation, the fish were acclimated to the local environment by placing in an outdoor pond of the Biological Research Base of Guangdong Ocean University for 1 week, during which time they were provisioned with commercial feed.

Feeding trial and challenge test

Having acclimated, 480 healthy uniformly sized hybrid groupers were selected and randomly divided into four groups, each with four replicates of 30 fish, and the fish were cultured in 300-L glass fiber buckets. During the culture period, the fish were fed twice daily (at 08:00 and 16:00) to apparent satiety with a water change of approximately 70%. Dissolved oxygen levels in the containers were maintained at ≥ 7.00 mg/L via continuous aeration using air stones, whereas water temperature, pH, and ammonia nitrogen content were maintained at $29.00 \pm 1.30^\circ\text{C}$, 7.8–8.1, and <0.09 mg/L, respectively. At the end of the 8-week culture period, 40 fish per treatment were randomly selected for a challenge test (10 fish per replicate). Each fish was injected intraperitoneally with 200 μL of *Vibrio parahaemolyticus* at a concentration of 7.41×10^8 CFU/ml, and thereafter the fish were continuously observed for 1 week, with mortality being recorded daily.

Diet formulations

Experimental fish were provided with one four diets containing protein and fat at levels of approximately 48% and 10%, respectively, which were prepared using redfish meal, casein, and gelatin as the main protein sources, and fish oil and soy lecithin as the main lipid sources. As treatments, the diet was supplanted as follows: 0% 7S, 0% NaB (fishmeal group, FM); 1.50% 7S (low-level 7S, bL), 0% NaB; 6.00% 7S, 0% NaB (high-level 7S, bH); 6.00% 7S, 0.13% NaB (high-level 7S with NaB, bH-NaB) (Table S1). The three experimental diets were also supplemented with methionine and lysine to same level as that in the FM diet. The essential amino acid profiles of the diets used in this experiment are shown in Table S2. Purified 7S was purchased from the China Agricultural University (Patent No. 200410029589.4, China). For use in the experimental diets, water was added to the purified β -conglycinin several times in small

amounts, stirred thoroughly, and the pH of the solution was adjusted to 7.2. The 7S solution was then placed in a glass petri dish and dried for 72 h in a low-temperature freeze dryer. After crushing the different raw materials of the diet through 380-µm meshes, the components mixed in a stepwise manner, adding fish oil, soybean lecithin, and 30% water and mixing evenly. The resulting mixture was then placed into a feed pelletizer, which produced feed pellets of 2.5 mm in diameter. The pellets were air-dried at room temperature until the moisture content was approximately 10% (approx. 48 h), sealed in a self-sealing bag, and stored at -20°C until required. The 30% microencapsulated NaB used in this study was kindly provided by Shanghai Menon Animal Nutrition Technology Co., Ltd.

Sample collection

After the end of the 8-week breeding experiment, the fish in each bucket were fasted for 24 h and then counted and weighed. For each replicate, two fish were randomly selected, from which blood was drawn using a 1 mL syringe. After being left to stand for 12 h, the blood samples were centrifuged for 10 min at 5000 g/min and 4°C. The serum thus obtained was collected and stored at -80°C for the determination of serum immune indicators. A further six fish were randomly selected for each replicate, from which the intestines were gently removed, and the segment of intestine near the cloacal hole was cut as the distal intestine (DI). Two of the DI segments for each replicate were placed in 4% formaldehyde solution for the preparation of Alcian Blue-Periodic acid-Shiff (AB-PAS) stained intestinal sections, and a further two DI segments were placed in liquid nitrogen for the determination of hindgut-related enzyme activities. After the DI tissue had been thawed on ice, dried with qualitative filter paper, and weighed, a sample was homogenized in homogenization medium (0.8% sodium chloride solution) in an ice bath at a sample: medium ratio of 1:9 (mass to volume ratio), centrifuged at 5000 g/min for 10 min at 4°C, and the supernatant was collected for analysis. The remaining two DI samples from each replicate were placed in RNA lysis buffer, and after being left to stand for 12 h, were stored in a freezer at -80°C for subsequent determination of related gene expression.

Determination of growth and DI evaluation indicators

Parameters of interest were determined as follows:

WGR (weight gain rate, %) = $100 \times (\text{final weight} - \text{initial weight}) / \text{initial weight}$

SGR (specific growth rate, %) = $100 \times [\ln(\text{final weight}) - \ln(\text{initial weight})] / \text{days of experiment}$

SR (survival rate, %) = $100 \times \text{final fish number} / \text{initial fish number}$

FCR (feed coefficient ratio) = $\text{dry feed consumed} / \text{weight gain}$

Id/Ph = $\text{intestinal diameter} / \text{plica height}$

Immune index determinations

Serum immune and DI injury indices were determined using Shanghai Enzyme-Linked ELISA immunoassay kits (Full-wavelength microplate reader-1510). AB-PAS sections of DI were prepared by Wuhan Servicebio Biological Technology Co., Ltd. Observations were made using a Leica DM 6000 optical microscope, and measurements and photographs were taken using cellSens Standard 1.8 and LAS 3.8 software, respectively, with 10 randomly selected plicae and muscle layers being measured in each section.

Total RNA from hybrid grouper DI was extracted using a TranZol Up Plus RNA kit (Beijing TransGen Biotechnology Co., Ltd) in accordance with the manufacturer's instructions. The integrity of the extracted RNA was assessed by 1% agarose gel electrophoresis, and a NanoDrop ND-2000 spectrophotometer (Thermo Scientific, USA) was used to determine the purity and concentration of RNA. RNA samples of sufficient quality should run as a clear and complete band, and have an OD₂₆₀/OD₂₈₀ ratio of between 1.8 and 2.0. A PrimerScriptTM RT-PCR Kit (TaKaRa, Kusatsu, Japan) was used to reverse transcribe total RNA to cDNA, which was stored at -20°C prior to RT-PCR analysis. Specific primers for RT-PCR (Table S3) were designed using Primer Premier 5.0 software based on full-length transcriptome sequences of hybrid grouper in the NCBI Sequence Read Archive with accession number PRJNA664416. All the real-time PCR reactions were run in an Applied Biosystems 7500 Real-Time PCR System (Life Technologies, Carlsbad, CA, USA) using a SYBR® Premix Ex TaqTM Kit (Takara). The β-actin gene was used as an internal reference to normalize the amplification efficiency of each target gene, and the mRNA expression levels of genes were compared and analyzed using 2^{-ΔΔCT} methods according to Livak et al. (Livak, Schmittgen, 2001).

Statistical analyses

All data were subjected to one-way analysis of variance followed by Tukey multiple range tests to determine significant differences among treatment groups using SPSS v. 22 (IBM, USA) at a significance level of $P < 0.05$, as described by Guo et al. (Guo, et al., 2017). The results are presented as the means ± SE.

Results

Growth performance and challenge test

The results obtained for the growth performance of fish in different groups are presented in Table 1. These results indicated that compared with the FM group, the final body weight (FBW), weight gain rate (WGR), and specific growth rate (SGR) were all significantly higher in the bL group and lower in the bH group, whereas no significant differences were observed between the FM and bH-NaB groups. The feed coefficient ratio (FCR) in the bH group was found to differ significantly from that in the other three groups, although there was no significant difference between the FM and bL groups. In contrast, we detected no significant differences among the four groups with respect to survival rate (SR).

The cumulative mortality of fish in response to injecting with *V. parahaemolyticus* is shown in Figure S1, which indicates that fish in the bH and bH-NaB groups suffered heavier mortalities than those in the FM and bL groups, with all groups reaching an asymptote at approximately 5 days post-injection. Among the different groups, the bH group showed a sustained and rapid increase in cumulative mortality from the first day post-injection, whereas in the bH group, mortality increased slowly over the first 3 days after injection, although thereafter showed a rapid increase.

Serum immune indices

The serum immune indices determined for the different groups are presented in Figure. 1. Compared with the FM group, we detected significant increases in serum IFN- γ in the bL, bH, and bH-NaB groups, among which the bH-NaB group showed the most significant difference. There were no significant difference between the FM and bL groups with respect to IgE levels, and the same trend was observed between the bH and bH-NaB groups, whereas IgE levels in these two groups were found to be significantly upregulated compared with the FM and bL groups. Furthermore, the levels of IL-1 β were found to be significantly increased in the bH group and significantly reduced in the bL and bH-NaB groups, although no significant differences were detected between the bL and bH-NaB groups. With respect to TNF- α , we observed significantly downregulated and upregulated levels in the bL and bH groups, respectively, compared with the FM group, whereas in contrast, there was no significant difference between the bH-NaB and FM groups.

Distal intestinal histopathological evaluation and damage indicators

The results obtained for the different groups with respect to distal intestinal histopathology evaluated by AB-PAS staining, developmental status assessment, and damage indicators determined by ELISA are presented in Figures 2, 3, and 4, respectively. As shown in Figure 3, the intestinal diameter/plica height ratios (Id/Ph) obtained for the bL and bH-NaB groups were significantly higher than that for the FM group, and significantly lower than that for the bH group. Among the four groups, the bH group was found to have the lowest density of mucus cells, whereas we detected no significant differences among the other three groups.

Compared with the FM group, the levels of nitric oxide (NO) were significantly upregulated in the bH and bH-NaB groups (Figure 4). Similarly, although the levels of NO in the bL groups were somewhat lower, they were still significantly higher than those in the FM group. Whereas we observed that levels of nitric oxide synthase (NOS) in the bH group were significantly higher than those in the FM, bL, and bH-NaB groups, we detected no significant difference among the FM, bL, and bH-NaB groups. Moreover, levels of the peroxynitrite anion (ONOO) were found to be lowest in the bL group, whereas no significant differences were observed among the FM, bH, and bH-NaB groups.

Distal intestinal immune-related gene expression and protein expression

The effects of 7S and NaB on the distal intestine of hybrid grouper are presented in Figure 5. Compared with the FM group, the mRNA levels of PI3K, RS5, PRAS40, p70 S6K, and PRR5 were upregulated in the bL and bH groups, and were significantly upregulated in the bH-NaB group. Similarly, we observed that the mRNA levels of 3-PDK1, mTOR C1, TEL2, EIF4E, EIF4B, Sin1, NFY, MHC I, GILT, AEP, and TCR were all upregulated in the bL and bH-NaB groups, and were significantly upregulated in the bH group. Although Rheb, Raptor, mLST8, and 4EBP1 mRNA levels were also significantly upregulated in the bH group, no changes were detected between the bL and bH-NaB groups and the FM group.

In addition, the bL, bH and bH-NaB groups all showed significantly higher levels of TSC1, mTOR C2, 4EBP1, CIITA and CREB1 mRNA compared with the FM group, whereas no significant differences were detected between the bH and bH-NaB groups. Although we found that the mRNA levels of dectator, RICTOR, SGK1, RFX5, and MHC II in the bH and bH-NaB group were significantly higher than those in the FM and bL groups, no significant differences were observed between the FM and bL groups or between the bH and bH-NaB groups. PKC mRNA levels were upregulated in the bH-NaB group and downregulated in the bL group, and those of RhoA were upregulated in the bL and bH-NaB groups compared with the FM group. Moreover, the mRNA expression of CD4 was significantly downregulated in the bL, bH, and bH-NaB groups, whereas we detected no significant differences among the four groups with respect to TSC2 and S6.

For western blot analysis of PI3K-Akt-mTOR (Figure 6), the expressions of P-PI3K/GAPDH and P-Akt/GAPDH and P-Akt/mTOR were significantly up-regulated in bL group, meanwhile, the PI3K/GAPDH, P-PI3K/GAPDH and P-PI3K/PI3K were significantly up-regulated in bH group. Compared with bH group, the PI3K/GAPDH, Akt/GAPDH and P-PI3K/PI3K were all significantly up-regulated after supplementing NaB in bH-NaB group.

Discussion

Growth performance and challenge test

In general, due to the incomplete development of the immune system in juvenile animals, the intestines of individuals often show an allergic reaction to 7S, resulting in metabolic disorders and adverse physiological and biochemical reactions (Han, et al., 2018), which can have the effect of disrupting the digestion and absorption of nutrients in the intestines (Zhang, et al., 2014). Interestingly, a low dose of 7S can often induce a response similar to that obtained with immunostimulants (Zhou, et al., 2015), which is often manifested in terms of modified growth performance. For example, in turbot (*Scophthalmus maximus* L.), a reduction in feed utilization efficiency following ingestion of feeds containing 4%–8% 7S has been observed to result in a significant reduction in specific growth rate (SGR) (Li, et al., 2017b). These findings were similar to those obtained in the present study, in which we observed that hybrid grouper receiving a diet containing 8% 7S showed a significant reduction in SGR and weight gain (WG). However, whereas in turbot, the effect of ingesting of a diet containing 2% 7S did not differ significantly from that of a control diet with respect to SGR, we observed an increase in SGR in hybrid grouper fed a

diet supplemented with low-dose 7S. Moreover, in turbot, it was found that the level of soybean supplantation and WG showed a first-degree relationship (Gu, et al., 2016). Hybrid groupers can, however, tolerate up to 30%–50% replacement of fishmeal with soybean meal in their feed, and experience no adverse effects on growth, even at soybean meal substitution levels of 10% and 30%, WG increased while did not show a significant difference (He, et al., 2020). These observations thus indicate that hybrid grouper may be more tolerant to soybean meal than are turbot, and therefore we speculate that for turbot, the addition of 2% soybean globulin is higher than the dose used for immune enhancement.

In the present study, we found that the addition of sodium butyrate (NaB) significantly increased SGR in the bH-NaB group compared with that in the bH group, indicating that NaB can be effective in ameliorating the growth inhibition effects attributable to high doses of 7S. We also found that the feed coefficient ratio of fish in the bH and bH-NaB groups was significantly higher than that in the FM and bL groups, whereas the addition of NaB significantly reduced the FCR, indicating that NaB may enhance the growth performance by improving the feed utilization of hybrid grouper, which is similar to results obtained for grass carp (Liu, et al., 2017b; Liu, et al., 2014). We speculate that this effect of NaB is associated with an increase intestinal antioxidant capacity and enhancement of intestinal immunity (Mirghaed, et al., 2019).

To further investigate the effects of 7S and NaB on the disease resistance of the hybrid grouper, we conducted a 7-day challenge test using *Vibrio parahaemolyticus* injected into fish at a concentration of 7.41×10^8 CFU/mL at the end of the breeding experiments. We accordingly observed the highest cumulative mortality in the bH group, thereby indicating that high doses of 7S can significantly reduce disease resistance in hybrid groupers, whereas resistance was enhanced to some extent in groupers fed a diet supplemented with NaB. Interestingly, whereas the trend in mortality in the bH group showed a consistent and rapid increase during the first 4 days post-injection, the upward trend only became apparent at day 4 post-injection in groupers fed the NaB-supplemented diet. These observations thus tend to indicate that although NaB may not be effective in ameliorating the immunosuppression caused by high doses of 7S, it can, to a certain extent, delay the onset of disease caused by *V. parahaemolyticus*, thereby gaining valuable time for treatment. However, given that it was unclear how this resistance is regulated, we conducted the research described in the following section.

Serum immunity

It is widely believed that the allergic responses to specific antibody-mediated humoral immunity and T-cell-mediated cellular immunity, which induce the production of serum-specific antibodies, occur primarily in the intestinal tract of fish ingesting 7S (Mclean, Ash, 1986; O'Donnell, et al., 1996). In this regard, cytokines play an important role in regulating immune responses and the functions of IgE as an important mediator of allergic responses. As an antigenic protein, 7S is processed by antigen-presenting cells and is bound by IgE antibodies produced by plasma cells (Krishnan, et al., 2009; Zheng, et al., 2012). When the antigen re-enters the body, binding to antibodies causes mast cells to release allergic substances such as histamine, thereby inducing an adverse reaction (Amigo-Benavent, et al., 2011;

Ogawa, et al., 1995). With respect to the findings of the present study, we found that low-dose 7S did not appear to stimulate an increase in IgE production in hybrid grouper, whereas in contrast, supplementation with high-dose 7S significantly increased the levels of IgE, thereby indicating that ingestion of high doses of 7S may cause intestinal damage by activating B cells or T cells, which thus produce higher amounts of IgE, and consequently generate higher levels of toxic substances (Sun, et al., 2013). Similar results have also been obtained in piglets (Hao, et al., 2009b), and in several previous studies, NaB has been shown to enhance IL-4-induced IgE production (Yamamoto, et al., 1996), and can also improve clinical symptoms and nasal mucosal epithelial morphology by reducing serum IgE levels in mice (Wang, et al., 2016). Thus, on the basis of the findings of previous studies and the study reported herein, it would appear that NaB does not alleviate high-dose 7S-induced allergic reactions by lowering serum IgE.

As a typical pro-inflammatory cytokine, IL-1 β is primarily produced by mononuclear macrophages, and can induce the activation of neutrophils and macrophages, stimulate the release of thromboxane and platelet-activating factor, increase the permeability of epithelial and endothelial cells, and induce intestinal inflammation (Al-Sadi, et al., 2012). In the present study, we found that levels of IL-1 β in bL and bH group fish were significantly down- and upregulated, respectively. The lower levels of IL-1 β in the bL group conceivably indicate that low-dose 7S can maintain normal serum immune responses by modifying IL-1 β secretion, whereas the significantly elevated levels of IL-1 β in the bH group, could indicate that at higher doses, 7S may promote CD4⁺ T-cell activation, proliferation, and differentiation by enhancing the abnormal secretion of IL-1 β , thereby exacerbating the inflammatory response (Sun, et al., 2013). We also observed that NaB supplementation of the grouper diet had the effect of significantly reducing serum IL-1 β , thereby indicating that NaB may modulate serum immunity by inhibiting the abnormal secretion of IL-1 β . In this regard, studies on rats have found that NaB can induce the overexpression of IL-1 β by inhibiting the binding of p65 to the IL-1 β promoter (Zhong, et al., 2017). We accordingly speculate that a similar mechanism operates in the hybrid grouper.

TNF- α is a further key cytokine that plays role in mediating intestinal inflammatory processes (Parker, et al., 2019), and has been shown to exacerbate the progression of enteritis by inducing the release of pro-inflammatory factors (Lang, et al., 2004). Consequently, determining changes in the levels of TNF- α is important with respect to monitoring inflammatory responses in the body. We found that a low dose of 7S significantly reduced serum TNF- α levels in the hybrid grouper, whereas in contrast, a high dose had the effect of promoting the secretion of TNF- α . Similarly, in turbot, it has been found that supplementing feed with 4% to 8% 7S increased the expression of TNF- α , which ultimately inhibited growth (Li, et al., 2017b). These observations indicate that 7S has a regulatory effect on TNF- α secretion in both hybrid grouper and other fish. Indeed, it has previously been found that the percentage of CD4⁺ lymphocyte sub-populations in mice increases linearly with increasing 7S levels, and thus we hypothesized that the elevation of TNF- α in the high-dose 7S group may be associated with an increase in CD4⁺ lymphocyte sub-populations in groupers (Fei, et al.). We also found that supplementation of the grouper diet with NaB had the effect of reducing serum TNF- α to levels comparable with those observed in FM group fish, which we suspect could be attributable to the fact that NaB can inhibit TNF- α secretion by down-regulating

LITAF expression (Han, et al., 2018). This effect may also be associated with the fact that NaB inhibits mast cell activation and suppresses the release of mast cell inflammatory mediators (Assem, et al., 2008; Wang, et al., 2018).

Similarly to TNF- α , the type II interferon IFN- γ play roles in immune and inflammatory responses. 7S is treated by APC cells and the 7S soymetide is exposed, which binds to B- or T-cell-transformed immunoglobulins (Tsuruki, et al., 2003). On the basis of the findings of the present study, it would appear that both low and high doses of 7S can promote the secretion of IFN- γ , and we accordingly speculate that this may be closely related to the activation of T cells by 7S as an antigenic protein. This assumption tends to be supported by our observations indicating that dietary supplementation with NaB was associated with a significant elevation in the levels of serum IFN- γ . NaB was also found to inhibit certain IFN- γ functions by inhibiting the downstream signaling of IFN- γ and thus the production of IFN- γ -induced factors (Kitamura, et al., 2010). We accordingly conjecture that, similar to humans, NaB can reduce the production of toxic substances in groupers by blocking the induction of IFN- γ on inflammatory cells. To examine this possibility, we conducted the experiments described in the following section.

Distal intestinal damage and histopathology

Nitric oxide (NO), a free radical gas produced from L-arginine via the catalytic activity of nitric oxide synthase (NOS), plays an important role in host defense and inflammatory responses, and can interact with cytokines such as TNF- α and IFN- γ , thereby influencing the course of inflammatory responses (Kim, et al., 2019; Kubes, Mccafferty, 2000; Nakanishi, et al., 2018). Currently, however, there is a degree of controversy as to the mechanism whereby NO is regulated in the intestines (Anand, et al., 2006; Kubes, Mccafferty, 2000). In the present study, we found that NOS showed an elevated trend in the bL group and NO was significantly elevated in the fish in this group. A similar pattern was also observed in groupers in the bH group. Notably in this regard, we also detected significant increases and decreases in the levels of IFN- γ and TNF- α in the bL group, respectively, the former of which can enhance NO secretion by stimulating macrophages and inducing the production of nitric oxide synthase (Nieto-Patlán, et al., 2019). However, given that TNF- α showed an increase and decrease in the bL and bH groups, respectively, we were unable to determine whether TNF- α is involved in the regulation of NO production. Nevertheless, we speculate that either TNF- α is not involved at low doses of 7S but is involved at high doses, or, alternatively, that TNF- α does not play a role in regulating the macrophage production of NO in the hybrid grouper. In addition, we observed that dietary supplementation with NaB maintained the significant upregulated production of IFN- γ , and resulted in the recovery of TNF- α to FM group levels.

Correspondingly, NOS levels were significantly reduced, and NO showed a non-significant downward trend. Thus, taking into consideration the results obtained for both IFN- γ and TNF- α , we are more inclined to believe that NaB downregulates NOS activity by inhibiting the release of TNF- α (Liu, et al., 2019a), thereby reducing NO levels. The peroxynitrite anion (ONOO⁻) is produced by the rapid binding of NO to oxygen radicals, which may be an important factor in cell damage, energy consumption, and cell death (Ródenas, et al., 1995). We found that levels of ONOO⁻ in the bL group were significantly downregulated, which contrasts with the elevated levels of NO in this group. Our observations tended to indicate that both

IFN- γ and IL-1 β and TNF- α are involved in the regulation of the NO-binding peroxide-generating step of ONOO production. And from the analysis just presented, TNF- α and IL-1 β are very important in the regulation of ONOO generation (Hou, et al., 2010; Neumann, et al., 2006), particularly at high doses of 7S. We also found that NaB supplementation significantly reduced NOS levels, which may be related to the inhibition of NOS promoter-dependent transcriptional activity by NaB (Sasahara, et al., 2002). Excess ONOO can adversely affect tissues primarily by damaging DNA, inactivating enzymes, and degrading mitochondria (Gerdes, et al., 2020), and thus the downregulation of ONOO in the bH-NaB group may be related to the fact that NaB can enhance the provision of energy to cells (Fang, et al., 2019; Liu, et al., 2017a), thereby preventing the generation of ONOO and thus preventing its negative effects.

As previously mentioned, the strong oxidizing properties of ONOO $^-$ and its interaction with cytokines such as TNF- α and IL-1 β can induce cellular damage and thus disrupt the normal morphology of tissues. To better visualize the effects of NO and ONOO on the intestinal tract of hybrid grouper, we prepared histomorphological sections of the hindgut, which is particularly susceptible to 7S-related disturbance (Periodic acid Schiff and Alcian Blue staining, AB-PAS). Intestinal development in fish can often reflect digestive absorption, and in this regard, plica height is typically used as an indicator of intestinal health. However, we would argue that plica height alone does not truly reflect the status of intestinal health. Therefore, as an alternative, we used the ratio of intestinal diameter to plica height (Id/Ph), which we believe to be a more reliable indicator of intestinal integrity. The larger this ratio, the greater is the space occupied by folds in the intestines, and thus the greater is the efficiency of absorption (Wang, et al., 2017). On the basis of the findings of the present study, we established that a high dose of 7S can significantly inhibited the development of plicae in the distal intestine of hybrid grouper, thereby reducing the absorptive area of the intestinal fold and thus the efficiency with which feed nutrients are absorbed. However, supplementing the grouper diet with NaB reversed the detrimental effects of high-dose 7, resulting in a reduction in the Id/Ph ratio, and significantly enhancing intestinal development. Similar results have also been obtained for the yellow drum (*Nibea albiflora*, Richardson) (Wu, et al., 2020a).

The mucus secreted by the mucous cells of fish contains non-specific immunochemical substances, including lysozyme, transfer factors, chitin, and complementary substances, that are resistant to pathogenic microorganisms (Bosi, et al., 2017). On the basis of AB-PAS staining, mucous cells can be classified into four types: I (pure red), II (pure blue), III (purple reddish), and IV (blue purple) (Sun, et al., 2019). From Figure 2, it can be seen that most of the mucous cells in the distal intestine of hybrid grouper are type II cells. As an acidic mucous cell, the acidic mucus secreted by type II mucous cells plays an important role in regulating the transport of proteins and their residues, lubricating coarse feed, and increasing the viscosity of mucus and immune protection (Gui, et al., 2007). In the present study, we observed that the numbers of type II mucus cells in the intestines of bL and bH-NaB group fish were not significantly different from those in the FM group, whereas in contrast, numbers were significantly reduced in the bH group, thereby indicating that a high dose of 7S can significantly inhibit the formation of type II mucus cells and thus result in the reduced secretion of antimicrobial active substances. Conversely, supplementing the grouper diet with NaB was found to effectively increase the abundance of

type II mucous cells, thereby enhancing the ability of the distal intestine of hybrid groupers to resist pathogenic microorganisms, and thus protecting the distal intestine.

Distal intestinal transduction of the MHC II-PI3K/Akt/mTOR signaling pathway induced by CIITA

Subsequent to the ingestion of antigenic proteins in feed, most is converted to amino acids and other small molecules. However, owing to the high molecular weight of 7S, a small fraction is not readily broken down and directly enters the lymphatic system via the intestinal mucosal spaces to induce an immune response (Chen, et al., 2011). When the antigen re-enters the body from the intestines, it will cause intestinal mucosa damage and edema, leading to inflammation. In this regard, although numerous studies have examined the effects of antigenic proteins on the intestines of fish, few have investigated the root cause of the inflammatory effects.

The major histocompatibility complex (MHC) processes foreign antigens, generating antigenic peptide-MHC molecular complexes, which are present in T cells and play an important role in inducing immune responses (Bremberg, 2018). In this process, MHC II and MHC I present exogenous and endogenous antigens, respectively, and in the present study, we focused on the mechanisms associated with the former. In mammals, MHC II encapsulates foreign antigens within vesicular primary endosomes in response to an antigen signal. The principal genes involved in the MHC II antigen presentation process are GILT, AEP, and CTSB (Ganesan, et al., 2015; Maehr, et al., 2005), which function cooperatively to transport the primary endosomes to lysosomes for hydrolysis and antigen degradation. Thereafter, the complexes enter the Golgi apparatus for processing, and are thereby converted to antigenic peptide-MHC II molecular complexes, which activate CD4 T cells and produce corresponding immune responses (Lvaro-Benito, et al., 2018). In this regard, rather than binding directly to the promoter of the MHC II gene, CIITA has been shown to act by controlling MHC II gene transcription via a synergistic interaction with the essential transcription factors RFX5, CREB, and NFY (Gobin, et al., 1998; Wong, et al., 2014). In the present study, we found that the addition of a low dose of 7S promoted the synergistic association between CIITA and CREB1 and NFY to initiate the transcription of MHC II, thus presenting the foreign 7S antigenic protein (Figure 7(a)). In contrast, we detected no significant changes in RFX5, and thus MHC I was not upregulated. Studies in patients with type III naked lymphocyte syndrome (BLS) have shown that the RFX complex is required for CIITA-mediated MHC I activity (Gobin, et al., 1998), and it is speculated that this is also the case in hybrid grouper. In response to receiving the antigenic signal from 7S, MHC II forms an intranuclear body regulated by GILT, AEP, and CTSB, and is subsequent to processing within lysosomes and the Golgi system to form an antigenic peptide-MHC II molecular complex, which then transmits the signal to CD4 T cells to activate the immune response. In the distal intestine of hybrid grouper, AEP does not appear to be involved in transduction to nucleosomes, but rather promotes the transport of nucleosomes to lysosomes by upregulating GILT and downregulating CTSB, as has also been observed in mice (Testa, 2010). However, the involvement of AEP appears to be related to the type of antigen detected by an organism, as AEP preferentially deals with viral antigens, as opposed to macromolecular protein antigens (Watts, et al., 2005). CD4 T cells are activated upon receiving signals from the molecular complex, which in turn secrete cytokines with immunomodulatory properties. In combination with the

results obtained from our examination of serum-related immunity, these findings indicate that CD4 T cells and macrophages can enhance intestinal immunity in hybrid groupers by inhibiting the secretion of the pro-inflammatory factors IL-1 β and TNF- α and promoting secretion of the anti-inflammatory factor IFN- γ .

The development of intestinal inflammation is generally accompanied by the abnormal expression of constituents of the PI3K/Akt/mTOR pathway (Kim, et al., 2016), the regulation of which by anti-nutritional factor-induced intestinal inflammation is currently incompletely understood. PI3K is a heterodimer composed of the regulatory subunit p85 and the catalytic subunit p110, the activities of which provide a signal for Akt, using PIP3 as a second messenger (Thapa, et al., 2015). mTOR is an atypical serine/threonine protein kinase comprising two complexes, mTOR C1 and mTOR C2. The former consists of three core components, namely, mTOR, Raptor, and mLST8. mTOR is the catalytic subunit of the complex, whereas mLST8 is associated with the catalytic domain of mTOR C1, which stabilizes kinase activation, although is not the catalytic subunit of mTOR C1. mLST8 however, is associated with the catalytic domain of mTOR C1 that stabilizes kinase activation. Rictor does not seem to affect mTOR C1 activity (Lang, et al., 2010). mTOR C2 consists of three core components, namely, mTOR, Rictor, and mLST8, among which mTOR is a component of the catalytic subunit of the complex, as in mTOR C1. mTOR is a rapamycin-insensitive partner that can interact with Protor, although the physiological function of this interaction remains to be determined. mLST8 is a complex catalytic subunit of mTOR C2 (Rosner, Hengstschlager, 2008).

Akt also receives signals from mTOR C2, thereby promoting the expression of the core protein Rheb via the TSC1/TSC2 complex, which regulates mTOR C1 (Hsieh, et al., 2014). On the basis of the low-dose 7S results obtained in the present study, we established that in response to this treatment, Akt is not regulated by mTOR C2 or PI3K, and does not promote Rheb expression via TSC2, thereby disrupting the regulation of mTOR C1. Moreover, in mTOR C1, TEL2 promotes the expression of 4E-BP2, although EIF4E is unaffected, whereas in mTOR C2, TEL2, and Sin1 are likely to be the key components responsible for the upward modulation of RhoA. Furthermore, although the expression of TSC1 is elevated, in combination with Rheb, the regulation of mTOR C1 does not appear to be controlled by Akt. Given the aforementioned observations, we speculate that activation of the PI3K/Akt/mTOR pathway in response to low doses of 7S is superficial and does not have a fundamental effect on key downstream proteins, with CD4 T-cell activation playing a major role.

In response to supplementation with a high dose of 7S (Figure 7(b)), MHC I synthesis is also upregulated via RFX5 activation, and a comparison with the response to low doses indicates that RFX5 is essential for MHC I regulation (Gobin, et al., 1998). Subsequent to the formation of nuclear endosomes, it appears that GILT and AEP, although not GILT and CTSB, are involved in regulation. This observation indicates that the expression of AEP may be dependent not to the type of antigen, but rather the dose of antigen. In this case, following the activation of CD4 T cells by the antigenic peptide-MHC II molecular complex, the secretion of IFN- γ remains at high levels, and there are also significant increases in the expression of the pro-inflammatory factors IL-1 β and TNF- α , thereby contributing to the development of intestinal inflammation. Interestingly in this regard, we also detected an upregulation of endogenous antigen

presentation by MHC I, which we suspect may be associated with the elevated levels of body endogenous antigens induced by high doses of 7S. Likewise, we found that the PI3K/Akt/mTOR pathway was also fully activated in response to stimulation with high doses of 7S. Commencing with PI3K, which activates Akt by upregulating 3-PDK1, we found that IKK α was also upregulated, and although we did not examine the role played by NF- κ B in the present study, we speculate that Akt may induce the transfer of NF- κ B to the nucleus via IKK α . With respect to mTOR C1, the components depressor, Raptor, and mLST8 are activated, resulting in an mTOR C1-induced upregulation of 4E-BP1 and 4E-BP2, which in turn results in an increase in EIF4E secretion. Moreover, p70 S6K also increased the level of EIF4B via mTOR C1. Upregulation of eIF-4E is often considered a marker of early colon cancer, and it has also been reported that 4E-BP1 and 4E-BP2 limit the anti-inflammatory response of macrophages by inhibiting IL-10 (William, et al., 2018). Thus, high doses of 7S-induced intestinal inflammation in the hybrid grouper may be associated with elevated levels of 4E-BP1 and 4E-BP2 and a suppression of the anti-inflammatory effects of macrophages. In addition, the expression of TSC1 is elevated, thereby in turn promoting the expression of Rheb, which relays signals from Akt to mTOR C1. Furthermore, with respect to mTOR C2, mLST8, depressor, and RICTOR are also activated, thereby elevating the levels of SGK1, which is involved extensively in the regulation of inflammation, and not only induces cardiac inflammation by activating the NLRP3 inflammasome (Gan, et al., 2017) but can also exacerbate inflammation by stimulating NF- κ B (Wu, et al., 2020b).

As a common feed additive used to supplement the diets of livestock, poultry, and aquatic animals, NaB can be hydrolyzed in the gastrointestinal tract to form butyric acid, which provides a source of energy to the intestinal epithelial cells, and contributes to the repair of damaged intestinal mucosa and strengthening intestinal immunity. However, the mechanisms whereby NaB influences intestinal repair have yet to be determined. In the present study, we found that supplementation of feed containing a high dose of 7S with appropriate levels of NaB facilitated the transcription of MHC I and MHC II via the upregulated expression of CREB1 and downregulation of NFY, and that MHC II expression was maintained at a high level (Figure(c)). These observation thus tend to indicate that dietary supplementation with NaB has the effect of enhancing the efficiency of antigenic protein presentation and processing in hybrid groupers. In addition, a concomitant reduction in the expression of MHC I may be indicative of a reduction in the severity of the inflammatory response compared with that seen in groupers receiving a high dose of 7S in the absence of NaB (Wu, et al., 1994). In addition GILT and AEP are inhibited during the development of nucleosomes, which subsequently migrate to the lysosomes and Golgi apparatus. There is also a reduction in expression of the pro-inflammatory factors IL-1 β and TNF- α , thereby indicating a weakening of the inflammatory response. Although PI3K expression was also upregulated, the expression of 3-PDK1 and Akt remained at levels similar to those observed in groupers receiving a high dose of 7S. However, we found that the key transcription factor IKK α , which is transduced by Akt to NF- κ B, was suppressed. On the basis of these observations, we thus speculate that NaB supplementation could suppress the activation of NF- κ B (Si, et al., 2015), although we did not verify this assumption in the present study. With regards to the mTOR C1 complex, we noted an inhibition of mLST8 and Raptor, which in turn downregulated the expression and secretion of 4E-BP1 and EIF4B, and although

the expression of p70 S6K was maintained at a high level, the formation of the downstream product EIF4B was inhibited.

Collectively, these findings indicate that the protective effect of NaB on the hindgut of hybrid grouper is closely associated with small molecule proteins downstream of mTOR C1, such as EIF4E. Although TSC1 showed high expression, that of Rheb was significantly reduced, and taken together with the results obtained for groupers receiving feed supplemented with high and low doses of 7S, we speculate that there are probably alternative mechanisms regulating the pathway from Akt to mTOR C1. With respect to mTOR C2, we established that the downregulated expression of mLST8 and Sin1 and upregulation of PRR5 promoted an increase in the expression of RhoA and PKC, and that SGK1 concomitantly remained at high levels. RhoA and PKC are generally considered to function as signal molecules associated with the inflammatory response (Cataisson, et al., 2006; Xie, et al., 2013; Yu, et al., 2019), and are accordingly also commonly used as targets to suppress inflammation (Fard, et al., 2013; Kong, et al., 2013). Thus, NaB does not appear to alleviate intestinal inflammation via mTOR C2, but rather exacerbates the progression of intestinal inflammation.

Conclusion

In the present study, we demonstrated that low and high doses of β -conglycinin (7S) significantly enhanced and reduced the weight gain rate, specific growth rate, and feed utilization of hybrid grouper, respectively. However, we found that supplementation of the grouper diet with sodium butyrate was effective in relieving the growth inhibition caused by high doses of 7S. High-dose 7S was shown to disrupt the development of distal intestinal plica, suppress the density of distal intestinal type II mucus cells, and enhance the synthesis of NO, NOS, and ONOO associated with intestinal injury, thereby exacerbating the process of distal intestinal inflammation. The enhancement of immune responses in the distal intestine of hybrid grouper receiving a low dose of 7S appears to be associated primarily with the activation of CD4 T cells, which promotes the release of the anti-inflammatory factor IFN- γ and reduces that of TNF- α and IL-1 β . However, this does not appear to be mediated via by the PI3K/Akt/mTOR pathway. In contrast, this pathway was found to be fully activated upon stimulation with high doses of 7S. Within this pathway, Akt activates mTOR C1 via the promotion of Rheb expression, which in turn enhances the expression of the downstream pro-inflammatory small molecule proteins EIF4E and 4E-BP2, whereas Akt also upregulates SGK1 expression via mTOR C2. These dual effects synergistically contribute to exacerbating inflammation in the distal intestine of hybrid grouper. Furthermore, we established that sodium butyrate relieves intestinal inflammation by inhibiting the expression of the inflammatory-related small molecule proteins 4E-BP2 and EIF4B downstream of mTOR C1. However, in the case of mTOR C2, we found that sodium butyrate does not protect the distal intestine by inhibiting RhoA, PKC, and SGK1, but instead further intensified intestinal inflammation by upregulating RhoA and PKC. In combination with the results obtained for growth performance and histopathological analyses, our findings indicate PI3K/Akt/mTOR C1 and PI3K/Akt/mTOR C2 play major roles in mediating 7S-induced immune responses in the distal intestine of hybrid grouper, and with respect to the mechanisms

underlying the ameliorative effect of sodium butyrate, we suspect that mTOR C1 plays a more important role than mTOR C2.

Abbreviations

NaB: sodium butyrate; 7S:β-conglycinin; FM group:fishmeal group; bL group:low conglycinin group; bH group:high beta conglycinin group; bH-NaB group:high beta conglycinin with sodium butyrate group; kD:Kilo Dalton; ROS:reactive oxygen species; DI:distal intestine; AB-PAS:Alcian Blue-Periodic acid-Shiff; IBW:Initial body weight; FBW:Final body weight; WGR:weight gain rate; SGR:specific growth rate; SR:survival rate; FCR:feed coefficient ratio; Id/Ph:intestinal diameter/plica height; IFN-γ:interferon γ; IgE:Immunoglobulin E; IL-1β:interleukin-1β; TNF-α:tumor necrosis factor-α; NO:nitric oxide; NOS:nitric oxide synthase; ONOO:peroxynitriteanion; PI3K RS5, phosphatidylinositol 3-kinase regulatory subunit 5; 3-PDK1, 3-phosphoinositide dependent kinase-1; TSC1, tuberous 1; TSC2, tuberous 2; Akt, serine/threonine-protein kinase; IKKα, inhibitor of nuclear factor kappa-B kinase subunit α; Rheb, Ras homolog enriched in brain; Raptor, regulatory associated protein of mTOR; PRAS40, proline-rich Akt1 substrate 1; mTOR, mammalian target of rapamycin; mTOR C1, mammalian target of rapamycin complex 1; mTOR C2, mammalian target of rapamycin complex 2; deptor, DEP domain-containing mTOR-interacting protein; mLST8, target of rapamycin complex subunit Ist8; TEL2, telomere length regulation protein; 4EBP1, eukaryotic translation initiation factor 4E binding protein 1; 4EBP2, eukaryotic translation initiation factor 4E binding protein 2; p70 S6K, ribosomal protein S6 kinase β1; EIF4E, translation initiation factor 4E; EIF4B, translation initiation factor 4B; S6, small subunit ribosomal protein S6; Sin1, target of rapamycin complex 2 subunit; RICTOR, rapamycin-insensitive companion of mTOR; PRR5, proline-rich protein 5; RhoA, Ras homolog gene family, member A; PKCα, protein kinase C α; SGK1, serum/glucocorticoid-regulated kinase 1; CIITA, major histocompatibility complex class II trans-activator; RFX5, regulatory factor X5; CREB1, cyclic AMP-responsive element-binding protein 1; NFY, nuclear transcription factor Y subunit α; MHC I, major histocompatibility complex class I antigen; MHC II, major histocompatibility complex class II antigen; GILT, gamma-interferon-inducible-lysosomal thiol reductase; AEP, asparaginyl endopeptidase; CTSB, cathepsin B; TCR, T cell receptor;

Declarations

Ethics approval and consent to participate

The animal protocol used in the present study was approved by the ethics review board of Guangdong Ocean University, and all procedures were performed in accordance with the standards of the National Institutes of Health Guidelines for the Care and Use of Laboratory Animals (NIH Publication No. 8023, revised 1978) and relevant Chinese policies.

Consent for publication

Not applicable.

Availability of data and material

All datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Key R&D Program of China (2019YFD0900200) and the National Natural Science Foundation of China (No.31772864) and the Natural Science Foundation of Guangdong Province (2018A030313154&2020A1515011129).

Author contributions

All indicated authors have actively contributed to this study. The contributions of the co-authors are as follows: H. Y. Liu and B. P. Tan designed the study; B. Yin conducted the study and analyzed the data; X. H. Dong participated in the interpretation of results; B. Yin wrote the manuscript; Q. H. Yang, S. Y. Chi, and S. Zhang purchased the reagent supplies; H. Y. Liu revised the manuscript. All authors have read and approved the final version of the manuscript.

Acknowledgments

We would like to thank the Key Laboratory of Control for Disease of Aquatic Economic Animals of Guangdong Higher Education Institutes (Zhanjiang, China) for kindly providing the *Vibrio parahaemolyticus* strain used in this study.

References

- Al-Sadi, R., Guo, S.H., Dokladny, K., Smith, M.A., Ye, D.M., Kaza, A., Watterson, D.M., Ma, T.Y., 2012. Mechanism of interleukin-1 β induced-increase in mouse intestinal permeability in vivo. J. Interferon Cytokine Res. 32, 474-484. doi: 10.1089/jir.2012.0031
- Amigo-Benavent, M., Clemente, A., Ferranti, P., Caira, S., Castillo, M.D.d., 2011. Digestibility and immunoreactivity of soybean β -conglycinin and its deglycosylated form. Food Chem. 129, 1598-1605. doi: 10.1016/J.FOODCHEM.2011.06.015
- Anand, R.J., Rippel, C.A., Leapheart, C.L., Li, J., Hackam, D.J., 2006. Activated macrophages inhibit enterocyte gap junctions during intestinal inflammation through the release of nitric oxide. J. Surg. Res. 130, 227. doi: 10.1016/J.JSS.2005.11.183

- Assem, E.S.K., Peh, K.H., Wan, B.Y.C., Middleton, B.J., Dines, J., Marson, C.M., 2008. Effects of a selection of histone deacetylase inhibitors on mast cell activation and airway and colonic smooth muscle contraction. *Int. Immunopharmacol.* 8, 1793-1801. doi: 10.1016/J.INTIMP.2008.08.017
- Bosi, G., DePasquale, J.A., Carosi, A., Lorenzoni, M., Dezfuli, B.S., 2017. Protective responses of intestinal mucous cells in a range of fish–helminth systems. *J. Fish Dis.* 40, 1001-1014. doi: 10.1111/JFD.12576
- Bremberg, N., 2018. Exploring the external dimension of european cooperation in the area of internal security: the case of euro-mediterranean civil protection. *J. Immunol. Methods.* 169, 273-285. doi: 10.1016/0022-1759(94)90271-2
- Cataisson, C., Pearson, A.J., Tsien, M.Z., Mascia, F., Yuspa, S.H., 2006. CXCR2 ligands and G-CSF mediate PKC α -induced intraepidermal inflammation. *J. Clin. Invest.* 116, 2757-2766. doi: 10.1172/JCI27514
- Chen, F., Hao, Y., Piao, X.S., Ma, X., Wu, G.Y., Qiao, S.Y., Li, D.F., Wang, J.J., 2011. Soybean-derived [Beta]-conglycinin affects proteome expression in pig intestinal cells in vivo and in vitro. *J. Anim. Sci.* 89, p.743-753. doi: 10.2527/jas.2010-3146
- Fang, W.J., Xue, H.L., Chen, X., Chen, K., Ling, W.H., 2019. Supplementation with sodium butyrate modulates the composition of the gut microbiota and ameliorates high-fat diet-induced obesity in mice. *J. Nutr.* 149, 747-754. doi: 10.1093/JN/NXY324
- Fard, M.A., Ebrahimi, K.B., Miller, N.R., 2013. RhoA activity and post-ischemic inflammation in an experimental model of adult rodent anterior ischemic optic neuropathy. *Brain Res.* 1534, 76-86. doi: 10.1016/J.BRAINRES.2013.07.053
- Fei, G.P., Piao, X.S., Ou, D.Y., Li, D.F., Hao, Y., Characterization of the antigenic specificity of soybean protein β -conglycinin and its effects on growth and immune function in rats. *Arch. Anim. Nutr.* 61, 189-200. doi: 10.1080/17450390701318358
- Firdaus, R.F., Lim, L.S., Kawamura, G., Shapawi, R., 2016. Assessment on the acceptability of hybrid grouper, *Epinephelus fuscoguttatus* $\times$ *Epinephelus lanceolatus*, to soybean meal-based diets. *AACL Bioflux.* 9, 284-290.
- Gan, W.Q., Ren, J.Y., Li, T.G., Lv, S.L., Li, C.H., Liu, Z.L., Yang, M., 2017. The SGK1 inhibitor EMD638683, prevents Angiotensin II-induced cardiac inflammation and fibrosis by blocking NLRP3 inflammasome activation. *BBA-Biomembranes.* 1864, 1-10. doi: 10.1016/J.BBADIS.2017.10.001
- Ganesan, A.P.V., Miller, M.A., Eisenlohr, L.C., 2015. Endogenously-processed influenza epitopes intersect with the classical MHCII presentation pathway via the GILT endosomal reductase (APP2P.104). *J. Immunol.* 192.
- Gerdes, H.J., Yang, M., Heisner, J.S., Camara, A.K.S., Stowe, D.F., 2020. Modulation of peroxynitrite in cardiac mitochondria. *BBA-Bioenergetics.* 1861, 148290. doi: 10.1016/j.bbabbio.2020.148290

- Gobin, S.J.P., Peijnenburg, A., Eggermond, M.V., Zutphen, M.V., Elsen, P.J.V.D., 1998. The RFX complex is crucial for the constitutive and CIITA-mediated transactivation of MHC class I and beta2-microglobulin genes. *Immunity*. 9, 531-541. doi: 10.1016/S1074-7613(00)80636-6
- Gu, M., Bai, N., Zhang, Y.Q., Krogh, S., 2016. Soybean meal induces enteritis in turbot *Scophthalmus maximus* at high supplementation levels. *Aquaculture*. 464, 286-295. doi: 10.1016/j.aquaculture.2016.06.035
- Gui, K.J., Park, J.J., Lee, J.W., Lee, J.S., 2007. Microscopic anatomy of foot of the spiny top shell, *Batillus cornutus* (lightfoot, 1786) (Gastropoda: Turbinidae). *J. Shellfish Res.* 25, 1071-1077. doi: 10.2983/0730-8000(2006)25[1071:MAOFOT]2.0.CO;2
- Guo, X.Z., Ran, C., Zhang, Z., He, S.X., Jin, M., Zhou, Z.G., 2017. The growth-promoting effect of dietary nucleotides in fish is associated with an intestinal microbiota-mediated reduction in energy expenditure. *J. Nutr.* 147, 781-788. doi: 10.3945/jn.116.245506
- Hamer, H.M.H., Jonkers, D., Venema, K., Vanhoutvin, S., Troost, F.J., Brummer, R.J.M., 2007. Review article: the role of butyrate on colonic function. *Aliment. Pharm. Ther.* 27, 104-119. doi: 10.1111/J.1365-2036.2007.03562.X
- Han, F.L., Wang, X.D., Guo, J.L., Qi, C.L., Xu, C., Luo, Y., Li, E.C., Qin, J.G., Chen, L.Q., 2018. Effects of glycinin and β -conglycinin on growth performance and intestinal health in juvenile Chinese mitten crabs (*Eriocheir sinensis*). *Fish Shellfish Immun.* 84, 269-279. doi: 10.1016/j.fsi.2018.10.013
- Hao, Y., Zhan, Z.F., Guo, P.F., Piao, X.S., Li, D.F., 2009a. Soybean beta-conglycinin-induced gut hypersensitivity reaction in a piglet model. *Arch. Anim. Nutr.* 63, 188-202. doi: 10.1080/17450390902860026
- Hao, Y., Zhan, Z., Guo, P., Piao, X., Li, D., 2009b. Soybean beta -conglycinin-induced gut hypersensitivity reaction in a piglet model. *Archives of Animal Nutrition*. 63, 188-202.
- He, Y.F., Ye, G.L., Chi, S.Y., Tan, B.P., Dong, X.H., Yang, Q.H., Liu, H.Y., Zhang, S., 2020. Integrative transcriptomic and small RNA sequencing reveals immune-related miRNA-mRNA regulation network for soybean meal-induced enteritis in hybrid grouper, *Epinephelus fuscoguttatus* × *Epinephelus lanceolatus*. *Front. Immunol.* 11, 1-15. doi: 10.3389/FIMMU.2020.01502
- Hou, R.R., Zhang, J., Yin, T., Cao, H.W., Zhang, N.Y., 2010. Upregulation of PTEN by peroxynitrite contributes to cytokine-induced apoptosis in pancreatic β -cells. *Apoptosis*. 15, 877-886. doi: 10.1007/S10495-010-0510-Z
- Hsieh, C.T., Chuang, J.H., Yang, W.C., Yin, Y., Lin, Y.-S., 2014. Ceramide inhibits insulin-stimulated Akt phosphorylation through activation of Rheb/mTORC1/S6K signaling in skeletal muscle. *Cell. Signal.* 26, 1400-1408. doi: 10.1016/J.CELLSIG.2014.03.004

- Jiang, S.T., Wu, X.Y., Luo, Y., Wu, M.J., Lu, S.D., Jin, Z.B., Yao, W., 2016. Optimal dietary protein level and protein to energy ratio for hybrid grouper (*Epinephelus fuscoguttatus* × *Epinephelus lanceolatus*) juveniles. *Aquaculture*. 465, 28-36. doi: 10.1016/j.aquaculture.2016.08.030
- Kim, H., Banerjee, N., Barnes, R.C., Pfent, C.M., Talcott, S.T., Dashwood, R.H., Mertens-Talcott, S.U., 2016. Mango polyphenolics reduce inflammation in intestinal colitis-involvement of the miR-126/PI3K/AKT/mTOR axis in vitro and in vivo. *Mol. Carcinogen*. 56, 197-207. doi: 10.1002/MC.22484
- Kim, M.K., Kim, K.B., Kim, H.S., Lee, B.M., 2019. Alternative skin sensitization prediction and risk assessment using proinflammatory biomarkers, interleukin-1 beta (IL-1 β) and inducible nitric oxide synthase (iNOS). *J. Toxicol. Env. Heal. A*. 82, 361-378. doi: 10.1080/15287394.2019.1609183
- Kitamura, K., Andoh, A., Inoue, T., Amakata, Y., Hodohara, K., Fujiyama, Y., Bamba, T., 2010. Sodium butyrate blocks interferon-gamma (IFN-gamma)-induced biosynthesis of MHC class III gene products (complement C4 and factor B) in human fetal intestinal epithelial cells. *Clin. Exp. Immunol*. 118, 16-22. doi: 10.1046/J.1365-2249.1999.01004.X
- Kong, L., Shen, X., Lin, L., Leitges, M., Rosario, R., Zou, Y.S., Yan, S.F., 2013. PKC β promotes vascular inflammation and acceleration of atherosclerosis in diabetic ApoE null mice. *Arterioscl. Throm. Vas*. 33, 1779-1787. doi: 10.1161/ATVBAHA.112.301113
- Krishnan, H.B., Kim, W.S., Jang, S.C., Kerley, M.S., 2009. All three subunits of soybean β -conglycinin are potential food allergens. *J. Agric. Food. Chem*. 57, 938-943. doi: 10.1021/JF802451G
- Kubes, P., Mccafferty, D.M., 2000. Nitric oxide and intestinal inflammation. *Am. J. Med*. 109, 150-158. doi: 10.1016/S0002-9343(00)00480-0
- Lang, A., Lahav, M., Sakhnini, E., Barshack, I., Fidder, H.H., Avidan, B., Bardan, E., 2004. Allicin inhibits spontaneous and TNF- α induced secretion of proinflammatory cytokines and chemokines from intestinal epithelial cells. *Clin. Nutr*. 23, 1199-1208. doi: 10.1016/j.clnu.2004.03.011
- Lang, S.A., Moser, C., Hackl, C., Fichtner-Feigl, S., Stoeltzing, O., 2010. Abstract 299: Blockade of Rictor impairs mTORC1 inhibitor-induced upregulation of IGF-IR and Her2 expression in human gastrointestinal cancer cells. *Cancer Res*. 70, 299-299. doi: 10.1158/1538-7445.AM10-299
- Li, Y.X., Yang, P., Zhang, Y.J., Ai, Q.H., Xu, W., 2017a. Effects of dietary glycinin on the growth performance, digestion, intestinal morphology and bacterial community of juvenile turbot, *Scophthalmus maximus* L. *Aquaculture*. 479, 125-133. doi: 10.1016/j.aquaculture.2017.05.008
- Li, Y.X., Hu, H.B., Liu, J.T., Yang, P., Zhang, Y.J., Ai, Q.H., Xu, W., Zhang, W.B., Mai, K.S., 2017b. Dietary soya allergen β -conglycinin induces intestinal inflammatory reactions, serum-specific antibody response and growth reduction in a carnivorous fish species, turbot *Scophthalmus maximus* L. *Aquac. Res*. 48, 4022-4037. doi: 10.1111/ARE.13224

- Liu, J., Chang, G.J., Huang, J., Wang, Y., Ma, N.N., 2019a. Sodium butyrate inhibits the inflammation of lipopolysaccharide-induced acute lung injury in mice by regulating the toll-like receptor 4/nuclear factor κ B signaling pathway. *J. Agr. Food Chem.* 67, 1674-1682. doi: 10.1021/ACS.JAFC.8B06359
- Liu, J.D., Bayir, H.O., Cosby, D.E., Cox, N.A., Williams, S.M., Fowler, J., 2017a. Evaluation of encapsulated sodium butyrate on growth performance, energy digestibility, gut development, and *Salmonella* colonization in broilers. *J. Poult. Sci.* 96, 3638-3644. doi: 10.3382/ps/pex174
- Liu, M.M., Guo, W., Wu, F., Qu, Q.C., Tan, Q.S., Gong, W.B., 2017b. Dietary supplementation of sodium butyrate may benefit growth performance and intestinal function in juvenile grass carp (*Ctenopharyngodon idellus*). *Aquac. Res.* 48, 4102-4111. doi: 10.1111/ARE.13230
- Liu, W.S., Yang, Y.O., Zhang, J.L., Gatlin, D.M., Ring?, E., Zhou, Z.G., 2014. Effects of dietary microencapsulated sodium butyrate on growth, intestinal mucosal morphology, immune response and adhesive bacteria in juvenile common carp (*Cyprinus carpio*) pre-fed with or without oxidised oil. *Br. J. Nutr.* 112, 15-29. doi: 10.1017/S0007114514000610
- Liu, Y., Chen, Z.C., Dai, J.H., Yang, P., Xu, W.Q., Ai, Q.H., Zhang, W.B., Zhang, Y.A., Zhang, Y.J., Mai, K.S., 2019b. Sodium butyrate supplementation in high-soybean meal diets for turbot (*Scophthalmus maximus* L.): Effects on inflammatory status, mucosal barriers and microbiota in the intestine. *Fish Shellfish Immun.* 88, 65-75. doi: 10.1016/J.FSI.2019.02.064
- Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods.* 25, 402-408. doi: 10.1006/meth.2001
- Lvaro-Benito, M., Morrison, E., Ebner, F., Abualrous, E.T., Freund, C., 2018. Distinct editing functions of natural HLA-DM allotypes impact antigen presentation and CD4 + T cell activation. *Cell. Mol. Immunol.* 17, 133-142. doi: 10.1038/s41423-018-0181-1
- Maehr, R., Hang, H.C., Minter, J.D., Kim, Y.M., Cuvillier, A., Nishimura, M., Yamada, K., Shirahama-Noda, K., Hara-Nishimura, I., Ploegh, H.L., 2005. Asparagine endopeptidase is not essential for Class II MHC antigen presentation but is required for processing of Cathepsin L in mice. *J. Immunol.* 174, 7066-7074. doi: 10.4049/jimmunol.174.11.7066
- McLean, E., Ash, R., 1986. The time-course of appearance and net accumulation of horseradish peroxidase (HRP) presented orally to juvenile carp *Cyprinus carpio* (L.). *Comp Biochem. Phys. A.* 84, 687-690. doi: 10.1016/0300-9629(86)90388-9
- Mirghaed, A.T., Yarahmadi, P., Soltani, M., Paknejad, H., Hoseini, S.M., 2019. Dietary sodium butyrate (Butirex® C4) supplementation modulates intestinal transcriptomic responses and augments disease resistance of rainbow trout (*Oncorhynchus mykiss*). *Fish Shellfish Immunol.* 92, 621-628. doi: 10.1016/J.FSI.2019.06.046

- Murray, H.M., Lall, S.P., Rajaselvam, R., Boutilier, L.A., Blanchard, B., Flight, R.M., Colombo, S., Mohindra, V., Douglas, S.E., 2010. A nutrigenomic analysis of intestinal response to partial soybean meal replacement in diets for juvenile Atlantic halibut, *Hippoglossus hippoglossus*, L. *Aquaculture*. 298, 282-293. doi: 10.1016/j.aquaculture.2009.11.001
- Nakanishi, Y., Sato, T., Takahashi, K., Ohteki, T., 2018. IFN- γ -dependent epigenetic regulation instructs colitogenic monocyte/macrophage lineage differentiation in vivo. *Mucosal Immunol.* 11, 871-880. doi: 10.1038/mi.2017.104
- Neumann, P., Gertzberg, N., Vaughan, E., Weisbrot, J., Woodburn, R., Lambert, W., Johnson, A., 2006. Peroxynitrite mediates TNF-induced endothelial barrier dysfunction and nitration of actin. *AM. J. Physiol-Lung C.* 290, L674-L684. doi: 10.1152/AJPLUNG.00391.2005
- Nieto-Patlán, E., Serafín-López, J., Wong-Baeza, I., Pérez-Tapia, M.S., Cobos-Marin, L., Estrada-Parra, S., Estrada-Garcia, I., 2019. Valproic acid promotes a decrease in mycobacterial survival by enhancing nitric oxide production in macrophages stimulated with IFN- γ . *Tuberculosis*. 114, 123-126. doi: 10.1016/J.TUBE.2018.12.007
- O'Donnell, G.B., Reilly, P., Davidson, G.A., Ellis, A.E., 1996. The uptake of human gamma globulin incorporated into poly (D,L-lactide-co-glycolide) microparticles following oral intubation in Atlantic salmon, *Salmo salar* L. *Fish Shellfish Immunol.* 6, 507–520. doi: 10.1006/fsim.1996.0048
- Ogawa, T., Bando, N., Tsuji, H., Nishikawa, K., Kitamura, K., 1995. Alpha-subunit of beta-conglycinin, an allergenic protein recognized by IgE antibodies of soybean-sensitive patients with atopic dermatitis. *J. Agric. Chem. Soc. Japan.* 59, 831-833. doi: 10.1271/bbb.59.831
- Parker, A., Vaux, L., Patterson, A.M., Modasia, A., Muraro, D., Fletcher, A.G., Byrne, H.M., Maini, P.K., Watson, A.J.M., Pin, C., 2019. Elevated apoptosis impairs epithelial cell turnover and shortens villi in TNF-driven intestinal inflammation. *Cell Death Dis.* 10, 108-108. doi: 10.1038/S41419-018-1275-5
- Ródenas, J., Mitjavila, M.T., Carbonell, T., 1995. Simultaneous generation of nitric oxide and superoxide by inflammatory cells in rats. *Free Radical Bio. Med.* 18, 869-875. doi: 10.1016/0891-5849(94)00215-6
- Rosner, M., Hengstschlager, M., 2008. Cytoplasmic and nuclear distribution of the protein complexes mTORC1 and mTORC2: rapamycin triggers dephosphorylation and delocalization of the mTORC2 components rictor and sin1. *Hum. Mol. Genet.* 17, 2934-2948. doi: 10.1093/HMG/DDN192
- Sasahara, Y., Mutoh, M., Takahashi, M., Fukuda, K., Tanaka, N., Sugimura, T., Wakabayashi, K., 2002. Suppression of promoter-dependent transcriptional activity of inducible nitric oxide synthase by sodium butyrate in colon cancer cells. *Cancer Lett.* 177, 155-161. doi: 10.1016/S0304-3835(01)00778-9
- Schneider, O., Amirkolaie, A.K., Vera-Cartas, J., Eding, E.H., Schrama, J.W., Verreth, J.A.J., 2004. Digestibility, faeces recovery, and related carbon, nitrogen and phosphorus balances of five feed

ingredients evaluated as fishmeal alternatives in Nile tilapia, *Oreochromis niloticus* L. Aquac. Res. 35, 1370-1379. doi: 10.1111/j.1365-2109.2004.01179.x

Si, J.J., Yang, J.L., Xue, L.F., Yang, C.H., Luo, Y.X., Shi, H.S., Lin, L., 2015. Sodium butyrate (NaB) could rescue memory reconsolidation in the presence of IKK inhibition with SSZ. Plos One. g006, 0043973. doi: <https://doi.org/10.1371/journal.pone.0043973.g006>

Sun, H., Liu, X., Wang, Y.Z., Liu, J.X., Feng, J., 2013. Allergen-specific immunoglobulin, histamine and T-cell responses induced by soybean glycinin and β -conglycinin in BALB/c mice of oral sensitisation. Food Agr. Immunol. 24, 489-501. doi: 10.1080/09540105.2012.730501

Sun, J.F., Wang, Y.Z., Lv, A.J., Xian, J.A., Wang, Q.K., Zhang, S.L., Guo, Y.J., Xing, K.Z., 2019. Histochemical distribution of four types of enzymes and mucous cells in the intestine of koi carp (*Cyprinus carpio* var. koi). Fish Physiol. Biochem. 45, 1367-1376. doi: 10.1007/S10695-019-00673-Y

Testa, J.S., 2010. Elucidation of MHC class II-restricted antigen processing pathways using influenza virus epitopes, Dissertations & Theses - Gradworks. Thomas Jefferson University.

Thapa, N., Choi, S., Tan, X., Wise, T., Anderson, R.A., 2015. Phosphatidylinositol phosphate 5-kinase I γ and PI3K/Akt signaling couple to promote oncogenic growth. J. Biol. Chem. 290, 1-23. doi: 10.1074/jbc.M114.596742

Tsuruki, T., Kishi, K., Takahashi, M., Tanaka, M., Matsukawa, T., Yoshikawa, M., 2003. Soymetide, an immunostimulating peptide derived from soybean β -conglycinin, is an fMLP agonist. Febs Lett. 540, 206-210. doi: 10.1016/S0014-5793(03)00265-5

Wang, C.C., Wu, H., Lin, F.H., Gong, R., Hu, C.H., 2018. Sodium butyrate enhances intestinal integrity, inhibits mast cell activation, inflammatory mediator production and JNK signaling pathway in weaned pigs. Innate Immun-London. 24, 40-46. doi: 10.1177/1753425917741970

Wang, J., Wen, L.T., Wang, Y., Chen, F.Q., 2016. Therapeutic effect of histone deacetylase inhibitor, sodium butyrate, on allergic rhinitis in vivo. Dna Cell Biol. 35, 203-208. doi: 10.1089/DNA.2015.3037

Wang, Y.R., Wang, L., Zhang, C.X., Song, K., 2017. Effects of substituting fishmeal with soybean meal on growth performance and intestinal morphology in orange-spotted grouper (*Epinephelus coioides*). Aquacult. Rep. 5, 52-57. doi: 10.1016/J.AQREP.2016.12.005

Watts, C., Matthews, S.P., Mazzeo, D., Manoury, B., Moss, C.X., 2005. Asparaginyl endopeptidase: case history of a class II MHC compartment protease. Immunol. Rev. 207, 218-228. doi: 10.1111/J.0105-2896.2005.00312.X

William, M., Leroux, L.P., Chaparro, V., Lorent, J., Graber, T.E., M'Boutchou, M.-N.I., Charpentier, T., Fabié, A., Dozois, C.M., St?Ger, S., 2018. eIF4E-Binding Proteins 1 and 2 Limit Macrophage Anti-Inflammatory

- Responses through Translational Repression of IL-10 and Cyclooxygenase-2. J. Immunol. 200, 4102-4116. doi: 10.4049/JIMMUNOL.1701670
- Wong, D., Lee, W., Humburg, P., Makino, S., Lau, E., Naranbhai, V., Fairfax, B.P., Chan, K., Plant, K., Knight, J.C., 2014. Genomic mapping of the MHC transactivator CIITA using an integrated ChIP-seq and genetical genomics approach. Genome Biol. 15, 494-494. doi: 10.1186/S13059-014-0494-Z
- Wu, M.J., Lu, S.D., Wu, X.Y., Jiang, S.T., Luo, Y., Yao, W., Jin, Z.B., 2017. Effects of dietary amino acid patterns on growth, feed utilization and hepatic IGF-I, TOR gene expression levels of hybrid grouper (*Epinephelus fuscoguttatus* × *Epinephelus lanceolatus*) juveniles. Aquaculture. 468, 508-514. doi: 10.1016/J.AQUACULTURE.2016.11.019
- Wu, Q.H., Trymbulak, W., Tatake, R.J., Forman, S.J., Zeff, R.A., Shanley, J.D., 1994. Replication of human cytomegalovirus in cells deficient in β 2-microglobulin gene expression. J. Gen. Virology. 75, 2755-2759. doi: 10.1099/0022-1317-75-10-2755
- Wu, X., Wang, L.G., Xie, Q.P., Tan, P., 2020a. Effects of dietary sodium butyrate on growth, diet conversion, body chemical compositions and distal intestinal health in yellow drum (*Nibea albiflora*, Richardson). Aquac. Res. 51, 69-79. doi: 10.1111/ARE.14348
- Wu, X.Q., Jiang, W., Wang, X.L., Zhang, C., Tian, Z.L., 2020b. SGK1 enhances Th9 cell differentiation and airway inflammation through NF- κ B signaling pathway in asthma. Cell Tissue Res., 1-12. doi: 10.1007/S00441-020-03252-3
- Xie, X., Chang, X.T., Chen, L., Huang, K.P., Huang, J., Wang, S.G., Shen, X.Y., Liu, P.Q., Huang, H.Q., 2013. Berberine ameliorates experimental diabetes-induced renal inflammation and fibronectin by inhibiting the activation of RhoA/ROCK signaling. Mol. Cell. Endocrinol. 381, 56-65. doi: 10.1016/J.MCE.2013.07.019
- Xu, D.D., Jiang, W.D., Wu, P., Liu, Y., Jiang, J., Tan, B.P., Yang, Q.H., Kuang, S.Y., Tang, L., Zhou, X.Q., 2019. Soybean β -conglycinin caused intestinal inflammation and oxidative damage in association with NF- κ B, TOR and Nrf2 in juvenile grass carp (*Ctenopharyngodon idella*): varying among different intestinal segments. Fish Shellfish Immunol. 95, 105-116. doi: 10.1016/J.FSI.2019.10.021
- Yamamoto, I., Matsunaga, T., Sakata, K., Nakamura, Y., Doi, S., Hanmyou, F., 1996. Histone hyperacetylation plays a role in augmentation of IL-4-induced IgE production in LPS-stimulated murine B-lymphocytes by sodium butyrate. J. Biochem. 119, 1056-1061. doi: 10.1093/OXFORDJOURNALS.JBCHEM.A021347
- Yu, X.F., Wang, D., Wang, X.H., Sun, S.Y., Zhang, Y.H., Wang, S.Q., Miao, R.R., Xu, X.X., Qu, X.J., 2019. CXCL12/CXCR4 promotes inflammation-driven colorectal cancer progression through activation of RhoA signaling by sponging miR-133a-3p. J. Exp. Clin. Canc. Res. 38. doi: 10.1186/s13046-018-1014-x

- Zhang, J.X., Guo, L.Y., Feng, L., Jiang, W.D., Kuang, S.Y., Liu, Y., Hu, K., Jiang, J., Li, S.H., Tang, L., 2014. Soybean β -conglycinin induces inflammation and oxidation and causes dysfunction of intestinal digestion and absorption in fish. *Plos One*. 8, e58115. doi: 10.1371/journal.pone.0058115
- Zhang, W., Tan, B., Ye, G., Wang, J., Zhang, H., 2019. Identification of potential biomarkers for soybean meal-induced enteritis in juvenile pearl gentian grouper, *Epinephelus lanceolatus* $\times$ *Epinephelus fuscoguttatus*. *Aquaculture*. 512, 734337. doi: 10.1016/j.aquaculture.2017.05.008
- Zheng, H.G., Tian, H., Ma, N., Qin, G.X., Sun, Z.W., Yu, C.Y., 2012. Purification and IgE-binding properties of soybean β -conglycinin subunit. *Process Biochem*. 47, 2531-2537. doi: 10.1016/J.PROCBIO.2012.07.003
- Zhong, X., Savidge, T., Cong, Y., Li, Q., 2017. PTH-073 Severe neonatal inflammation epigenetically sensitises the colonic epithelium for exacerbated il-1B activation. *Gut*. 66, A242-. doi: 10.1136/gutjnl-2017-314472.472
- Zhou, Q.C., Mai, K.S., Tan, B.P., Liu, Y.J., 2015. Partial replacement of fishmeal by soybean meal in diets for juvenile cobia (*Rachycentron canadum*). *Aquacult. Nutr*. 11, 175-182. doi: 10.1111/J.1365-2095.2005.00335.X

Tables

Table 1 Growth parameters and feed utilization of juvenile hybrid grouper (*Epinephelus fuscoguttatus* $\times$ *E. lanceolatus*) fed the experimental diets for 8 weeks.

	FM	bL	bH	bH-NaB
IBW (g)	7.70 $\pm$ 0.05	7.72 $\pm$ 0.06	7.70 $\pm$ 0.04	7.69 $\pm$ 0.05
FBW (g)	50.60 $\pm$ 0.26 ^b	56.21 $\pm$ 2.71 ^c	42.09 $\pm$ 6.52 ^a	51.11 $\pm$ 1.31 ^b
WGR (%)	557.22 $\pm$ 3.33 ^b	636.50 $\pm$ 11.37 ^c	464.18 $\pm$ 6.76 ^a	608.87 $\pm$ 32.51 ^{bc}
SGR (%/day)	3.36 $\pm$ 0.01 ^b	3.57 $\pm$ 0.03 ^c	3.09 $\pm$ 0.02 ^a	3.49 $\pm$ 0.16 ^{bc}
SR (%)	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00	99.17 $\pm$ 0.83	100.00 $\pm$ 0.00
FCR	0.81 $\pm$ 0.05 ^a	0.77 $\pm$ 0.05 ^a	1.52 $\pm$ 0.21 ^c	1.04 $\pm$ 0.08 ^b

Value show means $\pm$ SE (n=4); Significance was evaluated by one-way ANOVA followed by Tukey's multiple range tests. FM, control diet; bL, containing 1.5% 7S diet, bH, containing 6% 7S diet, bH-NaB, containing 6% 7S and 0.13% SB diet. IBW, initial body weight; FBW, final body weight; WGR, weight gain rate; SGR, specific growth rate; SGR, survival rate; FCR, feed coefficient ratio. ^{a, b, c, d}Mean values among all treatments with different letters were significantly different when the interaction was significant ($P < 0.05$).

Figures

Figure 1

Serum immune indexes of juvenile hybrid groupers (*Epinephelus fuscoguttatus* × *E. lanceolatus*) fed experimental diets for 8 weeks. Values are the means ± SE (n = 4); Significance was evaluated by one-way ANOVA followed by Tukey multiple range tests. FM, control diet; bL, containing 1.5% 7S diet, bH, containing 6% 7S diet, bH-NaB, containing 6% 7S and 0.13% NaB diet. IFN-γ, interferon-gamma; IgE, immunoglobulin E; IL-1β, interleukin-1 beta; TNF-α, tumor necrosis factor-alpha. a, b, c, d Mean values denoted by different superscripted letters differ significantly at the P < 0.05 level.

Figure 1

Distal intestinal histological evaluation based Alcian Blue-Periodic acid-Shiff staining of distal intestine sections obtained from juvenile hybrid groupers (*Epinephelus fuscoguttatus* × *E. lanceolatus*) fed experimental diets for 8 weeks. Green arrow: type II mucous cell; red arrow: muscle thickness; yellow arrow: plica height; blue arrow: plica width.

Figure 1

Distal intestinal damage indicators of juvenile hybrid groupers (*Epinephelus fuscoguttatus* × *E. lanceolatus*) fed the experimental diets for 8 weeks ($P < 0.05$).

Figure 1

Distal intestinal antigen processing and presentation-related gene expression in juvenile hybrid groupers (Epinephelus fuscoguttatus × E. lanceolatus) fed the experimental diets for 8 weeks. Results are represented as the means ± SE (n = 4). PI3K RS5, phosphatidylinositol 3-kinase regulatory subunit 5; 3-PDK1, 3-phosphoinositide dependent kinase-1; TSC1, tuberous 1; TSC2, tuberous 2; Akt, serine/threonine-protein kinase; IKKα, inhibitor of nuclear factor kappa-B kinase subunit α; Rheb, Ras homolog enriched in the brain; Raptor, regulatory associated protein of mTOR; PRAS40, proline-rich Akt1 substrate 1; mTOR, mammalian target of rapamycin; mTOR C1, mammalian target of rapamycin complex 1; mTOR C2, mammalian target of rapamycin complex 2; deutor, DEP domain-containing mTOR-interacting protein; mLST8, target of rapamycin complex subunit 1st8; TEL2, telomere length regulation protein; 4EBP1, eukaryotic translation initiation factor 4E binding protein 1; 4EBP2, eukaryotic translation initiation factor 4E binding protein 2; p70 S6K, ribosomal protein S6 kinase β1; EIF4E, translation initiation factor 4E; EIF4B, translation initiation factor 4B; S6, small subunit ribosomal protein S6; Sin1, target of rapamycin complex 2 subunit; RICTOR, rapamycin-insensitive companion of mTOR; PRR5, proline-rich protein 5; RhoA, Ras homolog gene family, member A; PKCα, protein kinase C α; SGK1, serum/glucocorticoid-regulated kinase 1; CIITA, major histocompatibility complex class II trans-activator; RFX5, regulatory factor X5; CREB1, cyclic AMP-responsive element-binding protein 1; NFY, nuclear transcription factor Y subunit α; MHC I, major histocompatibility complex class I antigen; MHC II, major histocompatibility complex class II antigen; GILT, gamma-interferon-inducible-lysosomal thiol reductase; AEP, asparaginyl endopeptidase; CTSB, cathepsin B; TCR, T cell receptor; CD4, T-cell surface glycoprotein. Significance was evaluated by one-way ANOVA followed by Tukey multiple range tests. FM, control diet; bL, containing

1.5% 7S diet, bH, containing 6% 7S diet, bH-NaB, containing 6% 7S and 0.13% NaB diet. a, b, c, d Mean values denoted by different superscripted letters differ significantly at the $P < 0.05$ level ($P < 0.05$).

Figure 1

Summary of the potential pathways associated with exogenous antigen β -conglycinin (7S) processing, presentation, and CIITA-mediated MHC II-PI3K/Akt/mTOR in the distal intestine of juvenile hybrid groupers (*Epinephelus fuscoguttatus* $\times$ *E. lanceolatus*). (a): the bL group (diet containing 1.5% 7S); (b): the bH group (diet containing 6% 7S); (c): the bH-NaB group (diet containing 6% 7S and 0.13% NaB).

Figure 3

Distal intestinal histopathological observation of juvenile hybrid groupers (*Epinephelus fuscoguttatus* × *E. lanceolatus*) fed experimental diets for 8 weeks ($P<0.05$).

Figure 6

Dietary modulations of distal intestinal responses in the target of PI3K/Akt/mTOR signaling pathway. Results are represented as mean ± SE (n=4). Significance was evaluated by one-way ANOVA followed by Tukey’s multiple range tests. FM, control diet; bL, containing 1.5% 7S diet, bH, containing 6% 7S diet, bH-NaB, containing 6% 7S and 0.13% NaB diet. a, b, c, dMean values among all treatments with different letters were significantly different when the interaction was significant ($P<0.05$).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterial.docx](#)