Knowledge, Practice and Associated Factors of Catheter-Associated Urinary Tract Infection Prevention Among Nurses Working at University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia, 2021

Alebachew Ferede Zegeye (✉ alexferede24@gmail.com)

University of Gondar

Chanyalew Worku Kassahun

University of Gondar

Yemataw Zewdu Temechu

University of Gondar

Research Article

Keywords: Catheter-associated Urinary Tract Infection, Gondar, Ethiopia, Knowledge, Nurses, Practice

Posted Date: February 2nd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1306083/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Catheter-associated urinary tract infection is the source of about 20% of episodes of health-care acquired infections in acute care facilities and it is over 50% in long-term care facilities. To date, knowledge and practice of nurses towards catheter-associated urinary tract infection is still lower. However, in Ethiopia, there is no published scientific research describing the knowledge and practice of nurses on catheter-associated urinary tract infections prevention. Therefore, this study aimed to assess knowledge, practice and associated factors of catheter-associated urinary tract infection prevention among nurses working at university of Gondar comprehensive specialized hospital, northwest Ethiopia, 2021

Methods: Institutional based cross sectional study was conducted at University of Gondar Comprehensive Specialized Referral Hospital from April 01 to May 01, 2021 among 423 nurses. Simple random sampling technique was used. Data were collected by using self-administered questionnaire then coded and entered into EPI-Data version 4.6 and then exported to SPSS version 23. All variables with P-value <0.25 in the binary logistic regression analyses were included in the multivariable regression analysis. The degree of association was interpreted by using the adjusted odds ratio with 95% confidence intervals, and the significance level was declared at P-value <0.05 . The Hosmer–Lemeshow test was used to check the fitness of the model.

Results: The knowledge and practice of nurses towards CAUTI prevention was 53% and 50% respectively. Good knowledge was associated with sex [AOR=1.84, 95% CI (1.09, 3.11)], work experience [AOR=2.36, 95% CI (1.09, 5.10)], working unit [AOR=6.9, 95% CI (2.11, 22.52)], received training [AOR=2.33, 95% CI (1.17, 4.65)] and had guideline [AOR=2.78, 95% CI (1.59, 4.88)]. Good practice was significantly associated with sex, work experience, good attitude and good knowledge towards CAUTI prevention [AOR=2.19, 95% CI (1.23, 3.88)], [AOR=5.44, 95% CI (1.66, 17.84)], [AOR=3.06, 95% CI (1.65, 5.67)], and [AOR=5.28, 95% CI (2.86, 9.76)] respectively.

Conclusions: In this study nearly half of nurses had good knowledge and practice towards CAUTI prevention. Sex, work experience, work unit, presence of guideline, and training were significantly associated with knowledge; while sex, work experience, good attitude and, knowledge were significantly associated with practice of nurses towards CAUTI prevention.

Background

Urinary tract infection (UTI) is one of the most common healthcare acquired infection caused by disease-causing organisms in the ordinarily sterile urine or tissues of the genitourinary tract involving the urethra, bladder, and kidneys [1, 2]. Catheter associated urinary tract infection is resulted from introduction of indwelling catheter into the bladder and it is one of the most common health care acquired infections encountered in clinical practice which has been performed by unsterile technique, and by not taking

adequate measures to maintain cleanliness of the urinary catheter during insertion, maintenance, and removal[3].

Globally urinary tract infection affecting more than 150 million people each year [4] and 70-80% of these urinary tract infections are attributed to catheter associated urinary tract infection[5]. Catheter-associated urinary tract infection is the source of about 20% of episodes of health-care acquired infections in acute care facilities and it is over 50% in long-term care facilities [6]. Additionally, the use of indwelling urethral catheters increases the risk of UTI occurrence in health-care settings by 14-fold [7]. Catheter-associated urinary tract infections are responsible for prolonged hospital lengths of stay, bacterial resistance, urosepsis, and even cause death [8].

In United states CAUTI accounted for 32% of all hospital acquired infections, making it the most frequent type of infection experienced in the hospital with approximately 449,000 CAUTI incidences with an estimated cost of 450 million dollar per year[9]. In developing countries including Africa, the prevalence and extent of catheter-associated urinary tract infection remains high due to limited resources and deficiency in health care system. In Uganda, one-third of UTI is accounted for CAUTI[10]. In Ethiopia, catheter associated urinary tract infection (CAUTI) is attributed 60.19 cases per 1000 persons-days[11].

Nurses play an important role in urinary catheter insertion, maintenance, and removal [12]. Catheter-associated urinary tract infection is considered as the most frequent and preventable infections, if nurses take into consideration the recommended catheter placement indications and evidenced-based practice of catheter maintenance [13]. Therefore, nurses should have adequate knowledge regarding infection control in the use of urethral catheters and their practice must be adhered to guidelines of catheter associated urinary tract infection prevention [14].

Even though 17% - 69% of catheter-associated urinary tract infections can be prevented by applying infection prevention methods, the knowledge and practice level of many nurses are not well known regarding catheter insertion, maintenance and removal [15]. According to the study conducted in Pakistan the adequate knowledge and practice level of nurses towards CAUTI prevention was 3.13% and 49% respectively[16]. Another study in Saudi Arabia shows that only 0.73% of nurses have adequate knowledge and 55.7% of nurses have good practice regarding CAUTI prevention[17].

Factors such as advanced level of education and long years of work experience are found to have a positive impact on the level of nurse's knowledge and practice towards catheter-associated urinary tract infections prevention [15, 18, 19].

In recent years, certain measures have been taken to control CAUTI. Center for disease control (CDC), Infectious Diseases Society of America (IDSA), world health organization (WHO), and department of health in Britain have published their efforts towards CAUTI prevention including reducing inappropriate use of urinary catheters, Perform proper techniques for indwelling catheter insertion, implement proper urinary catheter maintenance procedures, and removing urinary catheters in a timely manner[20-23]. Ethiopia has been taken certain measures to tackle CAUTI through applying recommendations such as

limiting use of indwelling urinary catheters and applying practices guidelines for insertion, maintenance, and removal of catheter[24].

Even though many efforts have been made to prevent CAUTI, the burden continues to rise. As my best searching in Ethiopia there is no study conducted regarding nurse's knowledge and practice towards CAUTI prevention. Thus the current research is aimed to assess knowledge, practice and its associated factors towards CAUTI prevention among nurses at University of Gondar comprehensive specialized hospital.

Methods

Study period and area

This study was conducted from April 01 to May 01, 2021 in University of Gondar comprehensive specialized hospital. The hospital is found in Gondar town which is 727 km from Addis Ababa, the capital of Ethiopia and 180 km from Bahir Dar, the capital of Amhara region. The hospital is used as the referral center for more than 5 million catchment populations. It has more than 1000-bed capacity and provides service to over 200,000 patients annually[25]. It provides both specialty and subspecialty services including pediatrics, surgery, gynecology and obstetrics, internal medicine, psychiatry, and ophthalmology in its in-patient and outpatient services. There are around 495 nurses who are working in the hospital. The nurse's patient-ratio varied within working units and nurses are performing comprehensive cares in each working unit

Study design

Institutional based cross sectional study design was employed in University of Gondar Comprehensive Specialized Hospital

Source and study population

Source population

All nurses working at University of Gondar Comprehensive Specialized Hospital

Study population

All nurses working at University of Gondar Comprehensive Specialized Hospital who were available at the time of data collection period

Inclusion and exclusion criteria

Inclusion criteria

All nurses who were working in university of Gondar Comprehensive Specialized Hospital

Sample size determination

Sample size was determined by using single population proportion formula by taking the proportion of knowledge and practice (p) 50%(because there was no study conducted in our country regarding this title)with expected margin of error (d) 5% just to maximize the sample size and improve the precision of study, 95% confidence interval and non-response rate 10%.

$$n = (Z_{\alpha/2})^2 p (1-p) / d^2$$

$$n = (1.96)^2 \cdot 0.5(1-0.5) / (0.05)^2$$

$$n = 384$$

$$\text{Non response rate} = (384) (10\%) = 38.4$$

$$\text{Total sample (n)} = 384 + 38.4 = \underline{423}$$

Where n= minimum sample size required to the study

d= margin of error

p= prevalence of knowledge and practice

$Z_{\alpha/2}$ = value of standard normal distribution

Sampling technique and procedure

The total number of sample size was obtained from university of Gondar comprehensive specialized hospital in all inpatient and outpatient services. Study participants were selected by using simple random sampling method and were assigned randomly by lottery method. There are a total 495 of nurses in University of Gondar comprehensive specialized hospital. 70 nurses from medical units, 59 from pediatric units, 83 from surgical units, 27 from ICUs, 13 from dialysis unit, 34 from OR units, 31 from recovery units, 20 from fistula unit, 38 from emergency OPD, and 48 from outpatient department nurses were selected by using proportional allocation. And the total sample size was 423.

Data collection tool and procedure

A structured English version self-administered questionnaire was used as instrument to collect data which is adopted from different literatures. The data were collected by three trained BSc unemployed nurses for duration of one month and supervised by one midwifery professional [13, 17, 26]. The questionnaire had five sections. The first section dealing with demographic characteristics of respondents, the second deals with work related characteristics, and the third section was about personal factors towards CAUTI prevention.

The fourth section was about 10 questions related to catheter indications, maintenance, care, removal, risk factors for CAUTI, and complications of CAUTI to assess nurses' knowledge towards CAUTI prevention and each correct answer was scored one and each wrong answer was scored zero. The total score for correct answers from all of the items was computed and classified as follows; good knowledge when study participants respond equal and above from the mean score of knowledge questions towards CAUTI prevention and poor knowledge when study participants respond below the mean score from knowledge questions towards CAUTI prevention

The fifth section deals the extent of nurses' practices toward prevention of CAUTI. It included 15 questions; five items for practices of nurses before catheter insertion, two items for practices of nurses during catheter insertion, and eight items for practices of nurses after catheter insertion. Study participants were asked to choose the appropriate answer for each item in the questionnaire. Every correct answer was given one point and wrong answer was given zero. The total score for correct answers from all of the items was computed and classified as follows; good practice when study participants respond equal and above from the mean score of practice questions towards CAUTI prevention and poor practice when study participants respond below the mean score from practice questions towards CAUTI prevention

The questionnaire was validated by following face validity method. To test the reliability of the tool, Cronbach alpha were calculated after a pre-test by taking 5% of the sample size at Tibebe Ghion specialized hospital and the value was 0.79, 0.83 and 0.89 for knowledge, attitude and practice tools respectively that were in acceptable ranges. The participants involved in the pre-test study were not included in the present study.

Study variables

Dependent variables

- Knowledge of nurses towards CAUTI prevention
- Practice of nurses towards CAUTI prevention

Independent variables

- Socio-demographic variables
- ▢ Age, sex, religion, ethnicity, level of education, marital status, monthly income
- Work related variables
- ▢ Work experience, working unit, training on CAUTI prevention, number of attending training on CAUTI prevention, availability of CAUTI prevention guidelines, usage of CAUTI prevention guidelines
- Personal related variables
- ▢ Attitude of nurses towards CAUTI prevention

Operational definition

Knowledge: is awareness and understanding of nurses regarding CAUTI prevention

Good Knowledge: Refers to those study participants who scored point greater than or equal to the mean score of knowledge questions towardsCAUTI prevention[13].

Poor knowledge: Refers to those study participants who scored point less than the mean score of knowledge questions towardsCAUTI prevention[13].

Practice: In this study it refers to the nurses actions regarding CAUTI prevention

Good Practice:Refers to those study participants who scored point greater than or equal to the mean score of practice questions towardsCAUTI prevention [17].

Poor Practice:Refers to those study participants who scored point less than the mean score of practice questions towardsCAUTI prevention [17].

Favorable attitude:Refers to those study participants who scored point greater than or equal to the mean score of attitude questions about CAUTI prevention[26].

Unfavorableattitude:Refers to those study participants who scored point less than the mean score of attitude questions about CAUTI prevention[26].

Data processing and analysis

The data were coded and entered into Epi-data version 4.6 and then exported to statistical package for social science (SPSS) version 23 for analysis. The data were cleaned then frequencies and cross-tabulation were computed before regression analysis. Descriptive statistics was computed, and the result

was summarized by texts, tables, and charts and presented in terms of mean and standard deviation. The Hosmer–Lemeshow test was used to check the model fitness. Binary logistic regression model was used to test the association between dependent and each independent variable. All variables with P-value <0.25 in the bivariable logistic regression analyses were included in the multivariable regression analysis. In the multivariable logistic regression analysis the degree of association was interpreted by using the adjusted odds ratio with 95% confidence intervals, and the significance level was declared at P-value <0.05.

Data quality assurance

Data quality was ensured by providing half day training to data collectors about the research objective, participant selection method, eligible study subjects, data collection tools, and procedures. The data collection tool was adopted from different literatures which was developed in English language and was checked for its consistency in the meaning of words. A pretest was done on five percent (22) from the total sample size at Tibebe Ghion specialized hospital two weeks before the actual data collection period. Based on the result of the pretest, necessary corrections and amendments were taken on the data collection tools (i.e. knowledge related question). Regular supervision was held during the data collection period and the collected data were checked on daily basis for completeness.

Results

Socio-demographic characteristics of nurses

A total of 406 participants were included in this study with a response rate of 96%. Among respondents 221 (54.4%) were female. In this study the mean age of the participants was 31.36 years $\pm$ 4.03 SD. Most of the participants 306(75.4%) were orthodox Christian and 239(58.8%) were married and 380(93.6%) had more than 5000 birr income and 307(75.6%) nurses had a bachelor's degree (Table 1).

Table 1: Socio-demographic characteristics of nurses to assess knowledge, practice and associated factors towards CAUTI prevention among nurses at UoGCSH, Northwest Ethiopia, 2021 (N = 406).

Socio-demographic variables	Frequency(n)	Percent (%)
Age		
20 -29	143	35.2%
30 -39	243	59.9%
≥40	20	4.9%
Sex		
Male	185	45.6%
Female	221	54.4%
Marital status		
Single	162	39.9%
Married	244	60.1%
Level of education		
Diploma	9	2.2%
BSc degree	307	75.6%
Masters and above	90	22.2%
Monthly income		
<5000 ETB	26	6.4%
5000 – 7000 ETB	201	49.5%
7001 – 9000 ETB	156	38.4%
>9000 ETB	23	5.7%

NB. BSc.....Bachelor of Science, ETB.....Ethiopian birr

Work related characteristics of nurses on prevention of CAUTI

More than half of study participants 204(50.2%) had years of experience ranged from 5 to 10 years. Out of 406 nurses who participated in the study, 89(21.9%) nurses were working in major and minor OR, and while 36(9.1%) of nurses were working in Adult emergency OPD. More than half of study participants 223(54.9%) attended training program regarding catheter associated UTI and 26(6.8%) nurses attended training program on catheter associated UTI prevention more than once. More than half of study participants 218(53.7%) had CAUTI prevention guidelines in their working unit (Table 2).

Table 2: work related factors to assess knowledge, practice and associated factors towards CAUTI prevention among nurses at UoGCSH, Northwest Ethiopia, 2021 (N= 406).

Work related variables	Frequency (n)	Percent (%)
Work experience		
<5 years	116	28.6%
5 to 10 years	204	50.2%
>10 years	86	21.2%
Attend training on CAUTI prevention		
Yes	223	54.9%
No	183	45.1%
Number of attending training		
Only once	197	48.5%
More than once	26	6.4%
Working unit		
Medical units	67	16.5%
Surgical units	82	20.2%
OR	89	21.9%
Pediatric units	58	14.3%
Fistula unit	33	8.1%
EOPD	37	9.1%
Others	40	9.9%
Presence of CAUTI prevention guidelines		
Yes	218	53.7%
No	188	46.3%
Following of CAUTI prevention guidelines		
Yes	204	50.2%
No	202	49.8%

NB. Others.....*Obstetrics ward*Malnutrition ward*Oncology unit*Chronic OPD

OR..... Operating room, EOPD..... Emergency Out Patient Department

Attitude of nurses towards prevention of CAUTI

From a total of 406 nurses 53.4% had good attitude and 46.6% had poor attitude towards CAUTI prevention (Table 3).

Table 3: Frequency distribution of participants response of Attitude towards CAUTI prevention among nurses at UoGCSH, Northwest Ethiopia, 2021 (N = 406).

Attitude	Frequency(n)	Percent (%)
Poor	189	46.6
Good	217	53.4

Nurses' Knowledge regarding to prevention of CAUTI

Participants were asked 10 questions to assess their knowledge on CAUTI prevention, and they were categorized into two groups based on their score in relation to the mean. The mean score was 5.64 (SD = 2.49). More than half 215(53 %) of the respondents were found to have good knowledge, while 191(47 %) of the respondents were found to have poor knowledge towards CAUTI prevention (Table 4).

Nurses' practice regarding to prevention of CAUTI

By using 15 practice based questions, the mean practice score of the respondents was found to be 10.22 (SD = 3.27). Half 203(50 %) of the respondents had good practice; whereas the remaining 203(50 %) respondents had poor practice of CAUTI prevention (Table 5).

Table 4: Frequency distribution of participants response of knowledge towards CAUTI prevention among nurses at UoGCSH, Northwest Ethiopia, 2021 (N = 406).

Questions	Yes		No	
	Freq.	%	Freq.	%
Among the following what is an inappropriate indication for indwelling urinary catheterization	210	51.7	196	48.3
Which is an appropriate indication of urinary catheterization among the following	198	48.8	208	51.2
Proper technique used for indwelling urinary catheter Insertion	296	72.9	110	27.1
Based on CDC 2009 Guidelines for prevention of CAUTI, it is advised to remove the catheter as soon as possible post operatively, preferably with in	221	54.4	185	45.6
If catheter is obstructed during you are assess patient, what are you going to do	254	62.6	152	37.4
One of the following is not a nursing action to prevent infections from urinary catheter	209	51.5	197	48.5
Which one among the following is a risk factor for CAUTI	238	58.6	168	41.4
Which patient is at high risk of mortality or developing CAUTI	214	52.7	192	47.3
Which action is perform before catheter insertion from the following except	203	50.0	203	50.0
All the following are complications of CAUTI except	247	60.8	159	39.2

Table 5: Frequency distribution of participants response of practice towards CAUTI prevention among nurses at UoGCSH, Northwest Ethiopia, 2021 (N = 406).

Questions	Yes		No	
	Freq.	%	Freq.	%
Do you wash your hands before you insert the catheter	380	93.6	26	6.6
What type of gloves do you use when you insert the catheter	297	73.2	109	26.8
Do you use sterile lubricant	376	92.6	30	7.4
What type of solution do you use to clean the urethra before you insert the catheter	251	61.8	155	38.2
How do you open and handle the indwelling catheter from its packaging	223	54.9	183	45.1
What type of solution do you inject in the needleless port to inflate the balloon	333	82.0	73	18.0
Do you secure indwelling catheter after insertion to prevent movement of urinary catheter	301	74.1	105	25.9
Do you keep the catheter and collecting tube free from kinking	301	74.1	105	25.9
How do you keep the urine collecting bag	215	53	191	47.0
Do you empty the collecting bag regularly and use a separate, clean urine collecting jug for each patient	295	72.7	111	27.3
How do you maintain the system	198	48.8	208	51.2
Do you keep the urine collecting bag out off the floor	308	75.9	98	24.1
Do you avoid contact of the drainage spigot with the collecting container	258	63.5	148	36.5
Do you cleaning the perianal area during daily bathing or showering	220	54.2	186	45.8
What do you use during any manipulation of the catheter or collecting system	195	48.0	211	52.0

Factors Associated with the Knowledge of Nurses towards CAUTI prevention

Sex, work experience, working unit, presence of guideline, and training on prevention of catheter associated urinary tract infection were found to have significant association with nurses' knowledge regarding to prevention of CAUTI.

For males the odds of good knowledge towards catheter associated urinary tract infection prevention was 1.84 times that of females [AOR = 1.84, 95% CI (1.09, 3.11)]. The odds of good knowledge towards catheter associated UTI prevention among nurses with 5 to 10 years work experience was 2.36 times that of nurses with <5 years of work experience [AOR = 2.36, 95 % CI (1.09, 5.10)]. The odds of good knowledge towards catheter associated UTI prevention among nurses working in emergency OPD was 6.9 times that of working in medical units [AOR 6.9, 95% CI (2.11, 22.52)]. The odds of good knowledge

towards catheter associated UTI prevention among nurses who took training related to catheter associated UTI was 2.33 times that of nurses who had not taken [AOR = 2.33, 95 % CI (1.17, 4.65)]. For nurses who had catheter associated UTI prevention guidelines the odds of good knowledge towards CAUTI prevention was 2.78 times that of nurses who had not[AOR=2.78; 95%CI (1.59, 4.88)] (Table 6).

Table 6: Factors Associated with the Knowledge of Nurses towards CAUTI prevention at UoGCSH, Northwest Ethiopia, 2021 (N = 406).

Variables	Knowledge		COR(95% CI)	AOR(95% CI)
	Good	Poor		
	N	N		
Age (in years)				
20 -29	107	36	1	1
30 -39	76	167	6.53(4.10, 10.40)	2.04(0.99 , 4.17)
>40	8	72	4.46(1.69, 11.77)	0.52(0.13, 2.05)
Sex				
Male	115	70	1.99(1.335, 2.96)	1.84(1.09, 3.11)
Female	100	121	1	1
Monthly income				
<5000 ETB	18	8	1	1
5001-7000 ETB	130	71	1.23(0.51, 2.97)	0.49(0.18, 1.32)
7001-9000 ETB	40	116	6.53(2.63, 16.16)	1.12(0.38, 3.32)
>9000 ETB	3	20	15.00(3.44, 65.36)	4.99(0.83, 29.98)
Work experience				
<5 years	93	23	1	1
5 to 10 years	74	130	7.10(4.15, 12.17)	2.36(1.09, 5.10)
>10 years	24	62	10.45(5.42, 20.13)	1.25(0.44, 3.57)
Attend training on CAUTI prevention				
Yes				
No	168	55	8.84(5.64, 13.87)	2.33(1.17, 4.65)
	47	136	1	1
Working unit				
Medical units	35	32	1	1

Surgical units	53	29	0.60(0.31, 1.16)	0.49(0.21, 1.14)
OR	39	50	1.40(0.74, 2.65)	1.51(0.67, 3.40)
Pediatric units	28	30	1.17(0.58, 2.37)	1.13(0.45, 2.83)
Fistula unit	6	27	4.92(1.80, 13.46)	2.73(0.81, 9.20)
EOPD	7	30	4.69(1.809, 12.15)	0.81(0.37, 1.78)
Others	23	17		6.90(2.11, 22.52)
				0.76(0.29, 2.00)
Presence of CAUTI prevention guidelines				
Yes				
No	156(38.4)	62(15.3)	5.50(3.59, 8.42)	2.78(1.59, 4.88)
	59(14.5)	129(31.8)	1	1

Key: Bold numbers indicate significantly associated variables with nurses' knowledge

Hosmer and Lemeshow test= 0.194

Factors Associated with the practice of Nurses towards CAUTI prevention

Sex, work experience, good attitude and good knowledge were found to have significant association with practice of nurses' towards prevention of catheter associated UTI, while presence of guidelines and working unit were not significantly associated at p-value of < 0.05.

For males the odds of good practice towards catheter associated UTI prevention was 2.19 times that of females [AOR = 2.19, 95% CI (1.23, 3.88)]. The odds of good practice towards catheter associated UTI prevention among nurses with >10 years work experience was 5.44 times that of nurses with <5 years of work experience [AOR = 5.44, 95 % CI (1.66, 17.84)]. Moreover the odds of good practice towards catheter associated UTI prevention among nurses with 5 to 10 years work experience was 3.59 times that of nurses with <5 years of work experience [AOR = 3.59, 95 % CI (1.46, 8.87)]. The odds of good practice towards catheter associated UTI prevention among nurses having favorable attitude was 3.06 times that of nurses having unfavorable attitude [AOR = 3.06, 95 % CI (1.65, 5.67)]. The odds of good practice towards catheter associated UTI prevention among nurses having good knowledge was 5.28 times that of nurses having poor knowledge [AOR = 5.28, 95 % CI (2.86, 9.76)] (Table 7).

Table 7: Factors Associated with the practice of Nurses towards CAUTI prevention at UoGCSH, Northwest Ethiopia, 2021 (N = 406).

Variables	Practice		COR(95% CI)	AOR(95% CI)
	Good	Poor		
	N (%)	N (%)		
Age (in years)				
20 -29	106	37	1	1
30 -39	92	151	4.70(2.98, 7.41)	0.68(0.29, 1.58)
>40	5	15	8.60(2.92, 25.29)	0.92(0.21, 4.08)
Sex				
Male	115	70	2.48(1.66, 3.71)	2.19(1.23, 3.88)
Female	88	133	1	1
Monthly income				
<5000 ETB	13	13	1	1
5000-7000 ETB	148	53	0.36(0.16, 0.82)	0.24(0.08, 1.74)
7001-9000 ETB	37	119	3.22(1.37, 7.55)	1.04(0.31, 3.50)
>9000 ETB	5	18	3.60(1.03, 12.62)	0.51(0.09, 2.84)
Work experience				
< 5 years	98	18	1	1
5 to 10 years	86	118	7.47(4.21, 13.27)	3.59 (1.46, 8.87)
>10 years	19	67	19.20(9.39, 39.27)	5.44(1.66, 17.84)
Attend training on CAUTI prevention				
Yes				
No	154	69	6.10(3.96, 9.41)	0.76(0.35, 1.66)
	49	134	1	1
Working unit				
Medical units	37	30	1	1
Surgical units	51	31	0.75(0.40, 1.45)	0.65(0.25, 1.70)
OR	34	55	1.99(1.05, 3.80)	1.26(0.51, 3.14)
Pediatric units	34	24	0.87(0.43, 1.77)	0.60(0.22, 1.68)
Fistula unit	11	22	2.47(1.03, 5.88)	0.89(0.28, 2.74)
EOPD	10	27	3.33(1.39, 7.96)	2.43(0.73, 8.03)

Others	26	14	0.66 (0.30, 1.49)	0.43(0.15, 1.27)
Presence of CAUTI prevention guidelines				
Yes				
No	142	76	3.89(2.57, 5.88)	1.69(0.88, 3.25)
	61	127	1	1
Attitude				
Poor	139	50	1	1
Good	64	153	6.65(4.30, 10.27)	3.06(1.65, 5.67)
Knowledge				
Poor	151	40	1	1
Good	52	163	11.83(7.41, 18.89)	5.28(2.86, 9.76)

Key: Bold numbers indicate significantly associated variables with nurses' Practice Hosmer and Lemeshow test= 0.461

Discussion

Catheter associated urinary tract infection (CAUTI) is one of major global health concern, which places a huge burden on the health care system. Nurses are expected to play a significant role in the prevention of CAUTIs[27].

The result of this study showed that 53%with 95%CI (48.3, 57.6) of study participants had good knowledge about catheter associated urinary tract infection prevention. This finding is in line with study conducted in Pakistan in which (51.8%) of the respondents had good knowledge[28]. Even though, there is difference in socioeconomic status and level of health sector development, the possible reason of similarity between the current study and study in Pakistan might be using a similar study population (staff nurse), study unit and study design.

The result of this study also lower than the study conducted in Sri Lanka (63.9%)[29], in India (56.7%) [30]. and in Rwanda (64.52%)[13]. The possible reason for this difference might be due to difference in study unit and study settings. On this study nurses working in inpatient and outpatient services were included in only one hospital; whereas in study done in Rwanda the study subject was nurses working only in ICUs at two different referral hospital.

However, the result of this study was higher than the study conducted in India 16.7%[31], in Saudi Arabia (37.23%)[17], and in another study in India (46.7%) [32]. The possible justification might be due to differences in study design, differences in study setting, and the differences in result rating;for example, in Saudi Arabia study,the knowledge was rated as Good, moderate, and poor. As well, the result rating in

India's study is quite different; it was rate as high, moderate and low. This may differ from the figurative interpretation of this study as good and poor knowledge.

For males the odds of good knowledge towards catheter associated urinary tract infection prevention was 1.84 times that of females [AOR = 1.84, 95% CI (1.09, 3.11)]. This finding was inconsistent with the study conducted in Turkey[33]. The possible explanation might be because males are more exposed to different sociocultural and social activities than females in the Ethiopian context. Furthermore, females have extraordinary activities as compared with males such as cooking, washing, and preparing for daily home consumption. Therefore, they might have less time to read and scale up their knowledge[34].

The odds of good knowledge towards catheter associated UTI prevention among nurses with 5 to 10 years work experience was 2.36 times that of nurses with <5 years of work experience [AOR = 2.36, 95 % CI (1.09, 5.10)]. Similar finding was reported in study conducted in India; where years of experience were significantly associated with knowledge[32]. Another study has conducted in Rwanda revealed that, the greater the working experience the higher the knowledge gained[13]. The reason might be nurses with more years of work experience have more chance to work with different professionals so that they can learn from their coworker's experience. Moreover, in the fact that the participants working for extended period of time in their professions, they could obtain various exposures that increase their knowledge towards CAUTI prevention related to their day to day activities.

The odds of good knowledge towards catheter associated UTI prevention among nurses who took training related to catheter associated UTI was 2.33 times that of nurses who had not taken [AOR = 2.33, 95 % CI (1.17, 4.65)]. This finding was consistent with the study conducted in India[32], Egypt [35]. The possible explanation might be due to that training can enhance an individual's body of knowledge regarding CAUTI prevention.

The odds of good knowledge towards catheter associated UTI prevention among nurses working in emergency OPD was 6.9 times that of working in medical units [AOR 6.9, 95% CI (2.11, 22.52)]. This finding was consistent with the study conducted in Saudi Arabia[17]. The possible explanation might be because of the prevalence of violence and none violence causality emergency cases occurred more in emergency units than the medical units. Therefore, nurses who are working in emergency ward face patients who need immediate intervention and treatments. They may lead them to read and update themselves.

For nurses who had catheter associated UTI prevention guidelines the odds of good knowledge towards CAUTI prevention was 2.78 times that of nurses who had not [AOR=2.78; 95%CI (1.59, 4.88)]. This finding was supported by a study conducted in KP Pakistan [36]. This can be justified by the fact that the presence of guideline will advocate the art and science of CAUTI prevention. Availability of CAUTI prevention guideline is likely improving nurses' knowledge and providing continuity of care.

Half of 203(50%) with 95% CI (45.3, 54.7) of study participants had good practice towards prevention of catheter associated urinary tract infection. This result was lower than the studies conducted in India

(86.5%)[30], Sri Lanka (79.5%)[29], Saudi Arabia (55.7%),[17], and Rwanda (79.9%) [13]. The possible justification might be due to difference in socioeconomic status, level of health sector development, study setting, study design, and difference in tool used to rating the outcome.

On the other hand, this study finding was higher than the study conducted in Pakistan (49%)[19]. The reason behind this difference might be difference in study setting and the difference in instrument (tool) used to assess nurses' practice towards prevention of catheter associated urinary tract infection.

In a multivariable logistic regression analysis, variables like sex, work experience, having good attitude and, good knowledge were found to have significantly associated with the practice of nurses towards catheter associated UTI prevention

For males the odds of good practice towards catheter associated UTI prevention was 2.19 times that of females [AOR = 2.19, 95% CI (1.23, 3.88)]. This finding was inconsistent with the study conducted in Rwanda [13]. The possible explanation might be in Ethiopia, most of the home-based activities such as food preparation and food serving, child feeding, cloth hygiene, and home-based sanitation are left for women. Therefore, women may not get time to practice CAUTI prevention because of being busy taking care of the family members compared with men.

The odds of good practice towards catheter associated UTI prevention among nurses with >10 years work experience was 5.44 times that of nurses with <5 years of work experience [AOR = 5.44, 95 % CI (1.66, 17.84)], Moreover the odds of good practice towards catheter associated UTI prevention among nurses with 5 to 10 years work experience was 3.59 times that of nurses with <5 years of work experience [AOR = 3.59, 95 % CI (1.46, 8.87)]. Similar finding was reported in studies conducted in India [37], Pakistan [16]. The finding might be justified as nurses' experience increased; they become familiar with the subject matter thereby acquire good practice to carry out the procedure with safety and precision.

The odds of good practice towards catheter associated UTI prevention among nurses having good knowledge was 5.28 times that of nurses having poor knowledge [AOR = 5.28, 95 % CI (2.86, 9.76)]. This finding was consistent with the study conducted in India [38], Iran [39] and Saudi Arabia[17]. The possible explanation could be because of the more knowledgeable to a particular subject matter the more skillful to a particular activity.

The odds of good practice towards catheter associated UTI prevention among nurses having favorable attitude was 3.06 times that of nurses having unfavorable attitude [AOR = 3.06, 95 % CI (1.65, 5.67)]. This finding was in line with the study conducted in Tamil Nadu, India[40]. The possible explanation might be individuals with good attitude towards CAUTI prevention have good practice level towards CAUTI prevention.

Limitation of the study

Despite, the high response rate in this study, social desirability bias and recall bias are potential limitations of these self-reported results.

I.e. study participants might not give true and genuine responses on the self-administered questionnaire, preferring to provide more socially acceptable responses than their actual day to day practice.

Conclusion

This study revealed that more than half of nurses who working in university of Gondar comprehensive specialized hospital had good knowledge regarding CAUTI prevention. Factors associated with good knowledge towards CAUTI prevention were sex, length of work experience, working unit, presence of guideline, and having training.

Half of study participants had good practice towards CAUTI prevention. Factors associated with good practice towards CAUTI prevention were sex, work experience, having good attitude and, good knowledge were found to have significantly associated with the practice of nurses towards catheter associated UTI prevention.

Abbreviations

CAUTI: Catheter Associated Urinary Tract Infection; CDC: Center for Disease Control; DC: data collector; ERC: Ethical Review Committee; HAI: Health Care Acquired Infection; IDSA: Infectious Diseases Society of America; IUC: Indwelling Urinary Catheter; KM: Kilometer; KP: Knowledge, Practice; NHSN: National Healthcare Safety Network; PI: Principal Investigator; SPSS: Statistical Package for Social Science; UoGCSH: University of Gondar Comprehensive Specialized Hospital; UTI: Urinary Tract Infection; WHO: World Health Organization

Declarations

Ethical approval and consent to participate

Ethical clearance and support letters were obtained from the Ethical Review Committee (ERC) of the University of Gondar College of Medicine School of nursing and it was submitted and approved by the Ethical Review Committee (S/N/164/7/2013). The support letter was submitted to University of Gondar comprehensive specialized hospital. Then, permission was obtained from the hospital director and departments or section heads. Study participants were informed about the purpose and objective of the study through verbal and written consent before the data collection. In addition, participants who were unwilling to take part in the study and those who needed to quit their participation at any stage were informed to do so without any restriction. Study participants privacy, secrecy, and wellbeing were protected. The data collected for the purpose of this study were not including study participants identification such as the name of the study participants.

Consent to publication

Not applicable.

Availability of data and materials

Data will be available upon request from the corresponding author.

Competing interest

The authors declared that there is no competing interest.

Funding

No funding

Author's contributions

AFZ: conceptualized the study and was involved in design, analysis, interpretation, report and manuscript writing.

CWK: involved in analysis, interpretation, report, and manuscript writing.

YZT: involved in analysis, interpretation, report, and manuscript writing.

Acknowledgments

The authors acknowledge University of Gondar for its financial support, University of Gondar Specialized Comprehensive Hospitals found in the northwest Ethiopia, and data collectors.

References

1. Monegro AF, Muppidi V, Regunath H: Hospital acquired infections. *StatPearls [Internet]* 2020.
2. Gudino C: Eliminating Catheter-Associated Urinary Tract Infections. 2015.
3. Benny AM, Idiculla AS, Kunjumon A, George A, Sequera SK: Nurses' Knowledge on Prevention of Catheter-Associated Urinary Tract Infection in a Selected Hospital of Mangaluru. *Journal of Health and Allied Sciences NU* 2020, 10(03):128-131.
4. Öztürk R, Murt A: Epidemiology of urological infections: A global burden. *World journal of urology* 2020:1-11.
5. Weber DJ, Sickbert-Bennett EE, Gould CV, Brown VM, Huslage K, Rutala WA: Incidence of catheter-associated and non-catheter-associated urinary tract infections in a healthcare system. *Infection Control & Hospital Epidemiology* 2011, 32(8):822-823.
6. Nicolle LE: Catheter associated urinary tract infections. *Antimicrobial resistance and infection control* 2014, 3(1):1-8.

7. Doughton TV, Bankole HS, Johnson RC, Hounmanou G, Moussa Toure I, Houessou C, Boko M, Baba-Moussa L: Catheter-associated urinary tract infections at a hospital in Zinvie, Benin (West Africa). *International Journal of Infection* 2016, 3(2).
8. Navoa-Ng JA, Berba R, Rosenthal VD, Villanueva VD, Tolentino MCV, Genuino GAS, Consunji RJ, Mantaring III JBV: Impact of an International Nosocomial Infection Control Consortium multidimensional approach on catheter-associated urinary tract infections in adult intensive care units in the Philippines: International Nosocomial Infection Control Consortium (INICC) findings. *Journal of infection and public health* 2013, 6(5):389-399.
9. Abiodun KO: Catheter-Associated Urinary Tract Infection in New York and North Carolina. 2018.
10. Odoki M, Almustapha Aliero A, Tibyangye J, Nyabayo Maniga J, Wampande E, Drago Kato C, Agwu E, Bazira J: Prevalence of bacterial urinary tract infections and associated factors among patients attending hospitals in Bushenyi district, Uganda. *International journal of microbiology* 2019, 2019.
11. Zewdu A: Healthcare Associated Infection at Adama Hospital Medical College, Adama, Oromia, Ethiopia. Addis Ababa University; 2017.
12. Sobeih H, Nasr M: Indwelling urinary catheter management: Effect of an interactive workshop on nurses' practice and perception. *New York Science Journal* 2015, 8(5):117-124.
13. Mukakamanzi J: Knowledge, attitude and practices of nurses towards the prevention of catheter-associated urinary tract infection in selected Referral Hospitals in Rwanda. University of Rwanda; 2017.
14. Opina MLF, Oducado RMF: Infection control in the use of urethral catheters: Knowledge and practices of nurses in a private hospital in Iloilo city. *Asia Pacific Journal of Education, Arts and Sciences* 2014, 1(5):93-100.
15. Köse Y, Leblebici Y, Akdere SŞ, Çakmakçı H, Ötünçtemur S, Egici MT, Bektemür G: Level of knowledge of the nurses work in a public hospital about the prevention of catheter associated urinary tract infections. *The Medical Bulletin of Sisli Etfal Hospital* 2016, 50(1):70-79.
16. Anwar G, Nawaz G, Afzal M, Majeed I, Waqas A: Assessment of Perceptions and Practices of the Nurses to Prevent Indwelling Catheter Associated Infection; Jinnah Hospital Lahore, Pakistan. *International Journal of Applied Sciences and Biotechnology* 2017, 5(2):150-158.
17. Algarni SS, Sofar SSS, Wazqar DY: NURSES'KNOWLEDGE AND PRACTICES TOWARD PREVENTION OF CATHETER-ASSOCIATED URINARY TRACT INFECTION AT KING ABDULAZIZ UNIVERSITY HOSPITAL. *Journal of Health, Medicine and Nursing* 2019, 4(1):50-73.
18. Sessa A, Di Giuseppe G, Albano L, Angelillo IF: An investigation of nurses' knowledge, attitudes, and practices regarding disinfection procedures in Italy. *BMC infectious diseases* 2011, 11(1):1-7.
19. Wicks K: Investigation of nursing knowledge of catheter selection following the introduction of a catheter decision support tool. ACU Research Bank; 2015.
20. Chenoweth C, Saint S: Preventing catheter-associated urinary tract infections in the intensive care unit. *Critical care clinics* 2013, 29(1):19-32.

21. Barbadoro P, Labricciosa FM, Recanatini C, Gori G, Tirabassi F, Martini E, Gioia MG, D'Errico MM, Prospero E: Catheter-associated urinary tract infection: Role of the setting of catheter insertion. *American Journal of infection control* 2015, 43(7):707-710.
22. Lo E, Nicolle LE, Coffin SE, Gould C, Maragakis LL, Meddings J, Pegues DA, Pettis AM, Saint S, Yokoe DS: Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infection Control & Hospital Epidemiology* 2014, 35(S2):S32-S47.
23. Schneider MA: Prevention of catheter-associated urinary tract infections in patients with hip fractures through education of nurses to specific catheter protocols. *Orthopaedic Nursing* 2012, 31(1):12-18.
24. NATIONAL INFECTION PREVENTION AND CONTROL (IPC) REFERENCE MANUAL FOR HEALTH CARE SERVICE PROVIDERS AND MANAGERS *Ministry of health- Ethiopia* 2019.
25. Teni FS, Gebresillassie BM, Birru EM, Belachew SA, Tefera YG, Wubishet BL, Tekleyes BH, Yimer BT: Costs incurred by outpatients at a university hospital in northwestern Ethiopia: a cross-sectional study. *BMC health services research* 2018, 18(1):1-10.
26. Latief NAK, Kurniawati ND, Pratiwi IN: Nurses' Compliance with the Catheter Associated Urinary Tract Infection Prevention Bundle. *International Journal of Psychosocial Rehabilitation*, 24(7).
27. Withanagamage A: Nursing interventions for the prevention of catheter associated urinary tract infections. 2020.
28. Shehzadi A, Ali A, Bhatti MR: Saudi Journal of Nursing and Health Care ISSN: 2616-7921 (Print). 2018.
29. Rohan P, Amarasekara TD: Knowledge and Practices among Female Nurses on Insertion and Care of Indwelling Urinary Catheter to Female Patients in a Selected Teaching Hospital in Sri Lanka. *The Open University of Sri Lanka* 2019, 14(01):101-118.
30. Parker V, Giles M, Graham L, Suthers B, Watts W, O'Brien T, Searles A: Avoiding inappropriate urinary catheter use and catheter-associated urinary tract infection (CAUTI): a pre-post control intervention study. *BMC health services research* 2017, 17(1):1-9.
31. Cutinho MC, Sheilini M, Harish B: Knowledge on practice of urinary catheter care and compliance to urinary catheter care guidelines-A hospital based study. *Indian Journal of Public Health Research and Development* 2018, 9(11):218-224.
32. Prasanna K, Radhika M: Knowledge regarding catheter care among staff nurses. *International Journal of Applied Research* 2015, 1(8):182-186.
33. Kose Y, Leblebici Y, Akdere SS, Cakmakci H, Otunctemur S, Egici MT, Bektemur G: Level of knowledge of the nurses work in a public hospital about the prevention of catheter associated urinary tract infections. *mortality* 2016, 4:5.
34. Dessalegn M, Ayele M, Hailu Y, Addisu G, Abebe S, Solomon H, Mogess G, Stulz V: Gender inequality and the sexual and reproductive health status of young and older women in the Afar Region of Ethiopia. *International Journal of Environmental Research and Public Health* 2020, 17(12):4592.
35. M Abdelmoaty A, A Sabry H, H ElSebaie E, M Kenawy A: Indwelling Urinary Catheter: Effect of Training on Nurses Knowledge and Skills. *The Egyptian Family Medicine Journal* 2020, 4(1):144-157.

36. Shah M, Wahab F, Ullah F, Gul U, Aziz A, Ullah Z: Infection control in the use of urethral catheter: Knowledge and practises of nurses. *Am J Adv Drug Deliv [Internet]* 2017, 5(01).
37. Jagdeep Kaur CNK: Adoption of Guidelines on Knowledge and Practice Regarding Care of Urinary Catheter in Situ among Staff Nurses. *International Journal of Health Sciences & Research* 2017.
38. Zachariah CA: Effectiveness of SIM on knowledge and practice among nurses regarding prevention of UTI in patients with indwelling catheter. 2016.
39. Kalantarzadeh M, Mohammadnejad E, Ehsani SR, Tamizi Z: Knowledge and practice of nurses about the control and prevention of nosocomial infections in emergency departments. *Archives of clinical infectious diseases* 2014, 9(4).
40. Balu P, Ravikumar D, Somasunder V: Assessment of Knowledge, Attitude and Practice on Prevention of Catheter-associated Urinary Tract Infection (CAUTI) among Health Care Professionals Working in a Tertiary Care Teaching Hospital. *J Pure Appl Microbiol* 2021, 15(1):335-345.