Maternal Mental Health, Handwashing Practices, and Infant Illness in Rural China

Qi Jiang

Stanford University <https://orcid.org/0000-0001-8054-5390>

Nourya Cohen

Stanford University

Mika Ohtori

Phillips Exeter Academy

Jie Gao

Sichuan University West China School of Public Health

Qingzhi Wang (✉ wangqz@stu.scu.edu.cn)

Sichuan University West China School of Public Health

Evelyn Zhang

Stanford University

Sabrina Zhu

The Harker School

Hannah Johnstone

Stanford University

Yian Guo

Stanford University

Sarah-Eve Dill

Stanford University

Huan Zhou

Sichuan University West China School of Public Health

Scott Rozelle

Stanford University

Research

Keywords: maternal mental health, depression, anxiety, stress, hygiene practices, handwashing practices, infant illness, rural China

Posted Date: December 21st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-130715/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Maternal Mental Health, Handwashing Practices, and Infant Illness in Rural China

Qi Jiang¹, Nourya Cohen¹, Mika Ohtori², Jie Gao³, Qingzhi Wang^{3*}, Evelyn Zhang¹, Sabrina Zhu⁴, Hannah Johnstone¹, Yian Guo¹, Sarah-Eve Dill¹, Huan Zhou³, Scott Rozelle¹

¹ Freeman Spogli Institute for International Studies, Stanford University, Stanford, California, USA

² Phillips Exeter Academy, Exeter, New Hampshire, USA

³ West China School of Public Health and West China Forth Hospital, Sichuan University, Chengdu, China.

⁴ Harker School, San Jose, California, USA

* Correspondence:

Qingzhi Wang

wangqz@stu.scu.edu.cn

Abstract

Background: Maternal mental health problems are a major burden to global public health and play an important role in infant wellbeing. While western countries have extensively studied the associations between maternal mental disorders, infant health, and hygiene practices, little is known in developing settings. This study investigates the correlations between maternal mental health problems and infant illness as well as handwashing practices of mothers in rural western China.

Methods: The research team followed a multi-stage random cluster sampling method to select a total of 720 mothers of infants aged 0-6 months from poverty counties in rural western China. Surveys questions about mental health were based on the Depression, Anxiety, and Stress Scale-21 (DASS-21), and questions about infant illness and handwashing practices followed evaluative surveys of prior studies. Adjusted ordinary least squares (OLS) regressions were run to examine how maternal mental health problems (including depression, anxiety, and stress symptoms) are associated with infant illnesses and maternal handwashing practices.

Results: We found significant associations between all three symptoms of maternal mental health and whether the infant had shown symptoms of illness twice in the past two weeks; depression and anxiety symptoms are significantly associated with whether the infant had been taken to the doctor in the past two weeks. Additionally, depression symptoms had a significant negative association with all categories of handwashing practices, and symptoms of anxiety and stress were significantly associated with handwashing count and frequently washing hands after cleaning infant's bottom.

Conclusion: These results indicate that maternal mental health problems are significantly associated with infant illness and suggest that postnatal mental health interventions may be effective in mitigating poor infant health outcomes.

Keywords: maternal mental health, depression, anxiety, stress, hygiene practices, handwashing practices, infant illness, rural China

Background

Studies have found that a child's health in the earliest years of development is essential for lifelong health and development [1]. However, as many as 5.2 million children under 5 years old die from preventable illnesses each year, the majority of which reside in low and middle income countries (LMICs), including China [2]. In LMICs, the chance of child and infant illness is much higher due to a number of risk factors such as poor access to clean water and fewer cleaning agents. Additionally, the rate of recovery from preventable illness is much lower due to fewer resources and a lack of adequate healthcare [2]. Similarly, in rural China, as many as 22% of all infant deaths were found to be associated with diarrheal illness and as many as 71% of rural infant deaths were associated with acute respiratory infections in 2015 [3]. Such high rates of fatality due to preventable infant illness require further study and intervention.

Furthermore, the international literature has established that mother and caregiver hygiene practices are a key factor in child and infant health outcomes in LMICs. For example, the World Health Organization (WHO) has predicted that better hygiene practices could improve over half of diarrheal illnesses worldwide [4]. International studies similarly show that poor hygiene practices, particularly among caregivers, are associated with infant and child illnesses [5]. A study in Ethiopia found that children under 5 whose mothers had poor handwashing practices were more likely to be admitted to the hospital for episodes of diarrhea [6]. Conversely, a study in South Africa found that infants whose caregivers frequently engaged in handwashing practices had reduced incidences of diarrhea and pneumonia [7].

Beyond the relationship between caregiver hygiene practices and child health, maternal mental health has been linked to infant health outcomes in LMICs. Because infants are most dependent on mothers for care [8], maternal mental health issues which may

manifest in poor hygiene practices, have been associated with higher rates of infections, diarrheal diseases and hospital visits among infants [9–13]. In addition, there is evidence that maternal mental health issues are associated with higher rates of admission to neonatal care units [14,15].

While there is not definitive evidence connecting maternal mental health and the hygiene practices of mothers, studies conducted LMICs did find significant relationships between adolescent mental health and hygiene practices [16,17]. However, to our knowledge, only one study has examined the association between maternal mental health and hygiene practices. This study, conducted in Ethiopia, Bangladesh, and Vietnam, found that maternal mental health issues were significantly associated with poor hygiene practices in the household; however, hygiene was not a significant focus of the paper [18]. With this in mind, the hygiene scores presented in the study were limited in their scope and the hygiene standards listed were vague. Therefore, more thorough research in developing settings, is necessary to confirm the correlations between maternal mental health and hygiene practices.

Like many LMICs, research has found high rates of infant illnesses and poor maternal mental health in rural areas of China, thus making it an ideal site to examine maternal mental health and hygiene practices. A study of child. illnesses in rural areas of China found that among rural children under five, 48% had experienced diarrhea, 50% had experienced fevers, and 60% had spells of coughing in the two-week period before the survey [19]. These rates are comparable to, and in some cases even higher than, other developing countries, where rates of illness in any given two-week period tend to be around 40–50% for diarrhea [20,21], 10-50% for fever, and 10-20% for coughing spells [22,23]. Research on maternal mental health in China has also discovered rates of depression to be around 20-25% [24–26], which is higher than both the 13% global rate of maternal mental disorders and similar to the rate of mental disorders among women in other LMICs (20%) [27].

In this study, the research team aims to investigate the association between maternal mental health problems, handwashing practices, and infant illness in rural China. More specifically, the study seeks to achieve three objectives. First, we describe the prevalence of maternal mental health problems, maternal handwashing practices, and infant illness in rural western China. Second, we investigate the correlations between maternal mental health problems and infant illness. Finally, we examine the correlations between maternal mental health problems and handwashing practices.

Methods

This study was conducted among low-income families residing in rural areas of Nanchong prefecture in Sichuan Province. According to the National Bureau of Statistics of China, 48% of Sichuan's population are rural residents [28]. The average per capita disposable annual income in rural areas of Sichuan is RMB 13,331 (\$1906), far lower than the national average of RMB 28,228 (\$4,033) (CNBS, 2019). These factors allow us to define our study sample as being rural, low-income participants.

Sampling

The research team followed a multi-stage random cluster sampling method. First, within Nanchong prefecture, four nationally designated “poverty counties” were randomly selected. Next, sample townships were selected from within each county. The research team excluded all non-rural townships and rural townships with populations less than 10,000. Of the remaining townships, 20 townships per county were randomly selected to reach a total of 80 townships.

Third, sample households were selected for participation in the study. The research team obtained a list of eligible households (households with pregnant women beyond the second trimester or new mothers with infants under six months) in each township from the

local county-level Maternal and Child Hospital. The research team aimed to recruit 25 such households from each township. If a township had more than 25 eligible households, 25 households were randomly selected for inclusions. In townships that did not have enough eligible households, the research team included all eligible households in the township and expanded the sampling frame to include villages surrounding the township. Because this study aims to identify the role of the mother's mental health in handwashing practices and infant health outcomes, we excluded 353 households whose respondents were pregnant women and 41 households whose respondents were not the mothers of the infants. After dropping mothers with missing key data, our analytical sample totalled 720 mothers and their infants in 80 rural townships.

Data collection

The research team surveyed sample households from November to December 2019. One-on-one survey interviews were conducted with each participant by survey enumerators who were trained to implement the survey instruments for each of the study's main components. The survey collected four blocks of data: maternal mental health (depression, anxiety, and stress symptoms), infant illness outcomes, caregiver handwashing practices, and demographic variables.

Maternal mental health

To measure the mental health of sampled mothers, enumerators administered the Depression, Anxiety, and Stress Scale-21 (DASS-21). DASS-21 is a 21-item, short-form version of the DASS-42 created by Lovibond and Lovibond [29] and validated in China [30,31]. Participants were read a total of 21 statements about their emotions in the past week: with seven items corresponding to depression, seven to anxiety and seven to stress. Participants were then asked to rank each statement on a scale from 0 to 3 (where 0=did not apply to me at all and 3=applied to me most of the time). Scores for depression, anxiety and

stress were calculated by totalling the item scores for each subscale and multiplying by two. Hence, the scores for each subscale could range from 0 to 42. According to the DASS manual, individuals are considered symptomatic of a given mental health issue if they scored above 9 for depression, 7 for anxiety, and 14 for stress. It is important to note, however, that the DASS-21 only evaluates the symptoms of mental illness and cannot be used as a clinical diagnosis.

Infant illness outcomes

To assess infant illness outcomes mothers were asked to report whether their infant had exhibited specific symptoms twice in the two weeks prior to the survey. Symptoms included fever, cough, runny nose, blood in the stool, diarrhoea, vomiting, difficulty breathing, skin rash, scrapes, bruising, burning, lethargy, ear or eye irritation or persistent crying in the past two weeks. Infants were considered symptomatic if they answered yes to any of the items. For infants who did have symptoms of illness, enumerators then asked the mother whether the infant had visited a doctor or for the symptoms. Questions were derived from China Health and Nutrition Survey and the International Food Policy Research Institute (IFPRI) Alive & Thrive baseline survey [32,33].

Handwashing practices

To assess handwashing practices, enumerators asked caregivers questions related to handwashing. The enumerator's questions were based on measures used in other studies to evaluate hygiene practices [16,17,34]. In the past, studies have looked at hygiene by checking for frequency of handwashing at critical times and by asking for a recall of handwashing times. In this study, participants were similarly asked to recall when they had washed their hands in the previous day (e.g. before or after going to the bathroom, before or after cooking, before or after handling the baby). The survey then counted the number of times they remembered washing their hands in the previous day to create an overall handwashing count

for each mother. Additionally, mothers were asked how often (always, often, rarely, or never) they washed their hands when feeding the baby and after cleaning their infant's bottom. The research team then converted these into binary variables, assigning participants who answered "often" or "always" as "frequent" and any other answer as "infrequent".

Enumerators also collected observational data on whether a cleansing agent (such as soap or hand sanitizer) was available in the place of handwashing.

Demographic characteristics

The final survey block collected data on the infant and family demographic characteristics. Infant characteristics were obtained from each infant's birth certificate and included the infant's gender, age in months, premature birth status, and whether the infant was born with a low birth weight. Family characteristics collected included mother's age, whether the infant's mother and father had graduated from high school, and the value of family assets. To provide a quantifiable estimate of family assets, enumerators asked whether the family owned or had access to certain household items, such as tap water, computer, internet, or a car. A standardized family asset index was then calculated using principal component analysis [35].

Statistical analysis

Our statistical analysis is composed of three parts. First, the research team calculated the overall prevalence of maternal depression, anxiety, and stress symptoms, as well as the overall prevalence of infant illness and maternal handwashing practices. Second, an adjusted ordinary least squares (OLS) regression was run to examine the associations between mental health issues and infant illness, while controlling for other potential confounders. The specification of the regression model is:

$$Infant_illness_i = \beta_0 + \beta_1 Mental_health_i + Control_i + \epsilon_i \quad (1)$$

where *Infant_illness_i* refers to infant illness outcomes, including whether infant *i* has shown symptoms of illness at least twice in the last two weeks, or whether infant *i* has been taken to the doctor for said symptoms in the last two weeks. *Mental_health_i* is a dummy variable for whether the mother of infant *i* shows symptoms of mental health issues (including depression, anxiety or stress). *Control_i* represents a set of control variables for infant and household characteristics, including infant gender and age, whether the infant was premature, whether the infant had low birth weight, maternal age, maternal and paternal and education level, and family asset index.

Finally, an adjusted OLS regression was used to identify the correlations between mental health problems and hygiene practices:

$$Handwashing_practice_i = \beta_0 + \beta_1 Mental_health_i + Control_i + \epsilon_i \quad (2)$$

where *Handwashing_practice_i* refers to the hygiene practices of mother *i*, which includes the handwashing count, whether mother *i* frequently washed her hands before feeding the infant, whether mother *i* frequently washed her hands before cleaning the infant's bottom, and whether the household of mother *i* has soap, detergent, or other cleaning agents at the place of handwashing. *Mental_health_i* and *Control_i* are the same as described in model (1).

Results

Table 1 reports the demographic characteristics of our sample. Among 720 infants, 393 infants (55.0%) were boys, and the average age of the sample was 2.7 months at the time of the survey. Only 27 infants (3.8%) were born prematurely, and 22 (3.1%) were born with low birth weight. Among sample mothers, the average age was 28 years old. Only 283 mothers (39.3%), and 325 fathers (45.1%) had graduated from high school.

Table 1. Descriptive statistics of demographic characteristics (N=720).

	N (%) or Mean (SD)
Boys, N (%)	393 (55.0)
Infant age (months), Mean (SD)	2.7 (2.0)
Premature, N (%)	27 (3.8)
Low birth weight, N (%)	22 (3.1)
Maternal age (years), Mean (SD)	28.02 (4.95)
Mother graduated high school, N (%)	283 (39.3)
Father graduated high school, N (%)	325 (45.1)
Family assets (factor scores), Mean (SD)	-0.02 (0.99)

The prevalence of mental health problems is presented in Panel A of Table 2. Within our sample, 93 mothers (12.9%) scored above the cut-off, indicating symptoms of depression. There were 110 mothers (15.3%) who scored above the cut-off for symptoms of anxiety, and 68 mothers (9.4%) who scored above the cut-off for symptoms of stress.

The health outcomes of infants and the hygiene practices of their mothers in our sample are presented in Panel B and C of Table 2. Panel B presents infant health outcomes. The results show that 282 infants (39.1%) showed symptoms of illness at least twice in the last two weeks. Among infants with symptoms of illness, 217 infants (30.1%) had been taken to a doctor for their symptoms. Panel C presents the handwashing practices of mothers. The data show that, on average, mothers washed their hands about 6.2 times in the past day. Additionally, 571 mothers (79.3%) stated that they frequently washed their hands before feeding their infants, and 663 mothers (92.1%) reported that they frequently washed their hands after cleaning their infant's bottom. Enumerator observations also show that 565 households (78.5%) had some type of cleaning agent at the place of handwashing.

Table 2. Descriptive statistics of mental health, infant illness and handwashing practices (N=720).

	N (%) or Mean (SD)
<i>Panel A. Mental health problems</i>	
Mother has symptoms of depression, N (%)	93 (12.9)
Mother has symptoms of anxiety, N (%)	110 (15.3)
Mother has symptoms of stress, N (%)	68 (9.4)
<i>Panel B. Infant illness outcomes</i>	
The infant has shown different symptoms at least twice in the last two weeks, N (%)	282 (39.1)
The infant has visited the doctor for said symptoms in the last two weeks, N (%)	217 (30.1)
<i>Panel C. Handwashing practices</i>	
Handwashing yesterday (times)	6.2 (3.5)
Frequent handwashing before feeding the infant N (%)	571 (79.3)
Frequent handwashing after cleaning the infant's bottom, N (%)	663 (92.1)
Has cleaning agent at the place of handwashing, N (%)	565 (78.5)

Table 3 presents the results of the OLS regression analysis of correlations between maternal mental health problems (depression, anxiety, and stress symptoms) and infant illness outcomes (symptoms of illness and doctor visits). A mother showing symptoms of depression in the past week was shown to have a significant positive association with the infant showing symptoms of illness at least twice in past two weeks ($p=0.004$), and the infant visiting the doctor in the past two weeks ($p=0.041$). Similarly, mothers with symptoms of anxiety were significantly more likely to have an infant who showed symptoms of illness or required doctor visits in the past two weeks ($p=0.011$, $p=0.047$, respectively). Finally, mothers who displayed symptoms of stress were significantly associated with infants displaying symptoms of illness at least twice in the past two weeks ($p=0.042$).

Table 3. Associations between mental health problems and infant illness (N=720).

	Infant had illness at least twice in past two weeks	Infant visited doctors in the past two weeks
Symptoms of depression	0.190*** (0.053)	0.104* (0.051)
Symptoms of anxiety	0.163** (0.050)	0.130** (0.047)
Symptoms of stress	0.125* (0.061)	0.044 (0.059)

Table 4 reports the OLS correlations between maternal mental health problems and handwashing practices. Depressive symptoms were significantly negatively associated with handwashing count the previous day ($p < 0.001$), frequently washing hands before feeding the infant ($p = 0.020$), frequently washing hands after cleaning infant's bottom ($p = 0.006$) and having a cleaning agent at the place of handwashing ($p = 0.030$). Symptoms of anxiety were negatively associated with handwashing count the previous day ($p = 0.006$) and frequently washing hands after cleaning infant's bottom ($p = 0.017$). In addition, symptoms of stress were negatively associated with handwashing count the previous day ($p < 0.001$) and frequently washing hands after cleaning infant's bottom ($p = 0.024$).

Table 4. Associations between mental health problems and handwashing practices (N=720).

	Handwashing count the previous day	Frequently washing hands before feeding the infant	Frequently washing hands after cleaning infant's bottom	Has cleaning agent at the place of handwashing
Symptoms of depression	-1.582*** (0.384)	-0.103* (0.044)	-0.081** (0.030)	-0.086* (0.040)
Symptoms of anxiety	-0.989** (0.361)	-0.025 (0.041)	-0.067* (0.028)	-0.020 (0.036)
Symptoms of stress	-1.595*** (0.433)	-0.072 (0.051)	-0.078* (0.034)	-0.031 [0.045]

Discussion

This study examined the associations between maternal mental health, handwashing practices, and child illness outcomes using extensive survey data from 720 new mothers with infants aged 0-6 months. We found that there were significant negative associations between most of the maternal mental health problems and infant illness and doctor visits. We also found maternal mental health problems were significantly correlated to maternal handwashing practices, especially handwashing count and frequently washing hands after cleaning infant's bottom. Collectively, these results suggest that maternal mental health is significantly related to infant care and infant health outcomes. In order to improve maternal hygiene practices, policymakers should consider implementing postnatal mental health interventions.

When looking at the prevalence of infant illness, our analyses showed that more than one-third (39%) of infants had displayed at least two different symptoms of illnesses in the last two weeks. Given this high rate of illness, it is unsurprising that the observed hygiene practices in our sample of rural China were generally poorer than those of high-income countries. Of the fourteen possible handwashing counts, the data show that on average mothers washed their hands 6.2 times. In contrast, when Americans were asked to give an average handwashing count on any given day, the average answer was 8.6 [36]. Moreover, and more specifically, our results show that mothers in our sample washed their hands infrequently, with a substantial percentage not washing their hands before feeding the infant or after cleaning their infant's bottom, and some mothers not having any cleaning agent near their sinks. These infrequent hygiene practices are likely contributing to the spread of disease. These findings are consistent with a study conducted in rural China that found that only 23.8% of rural adults practice proper handwashing behaviours, in that they frequently

handwashed at key times, used soap, and didn't share a towel with family members after handwashing [37].

When investigating the associations between maternal mental health problems and infant health, we found that mothers showing symptoms of depression, anxiety, and stress were more likely to have an infant who had an illness at least twice in the past two weeks. We also found that that mothers showing symptoms of depression, anxiety, and stress were more likely to have taken their infants to the doctor for symptoms of illness. This is consistent with other studies that show an inverse relationship between mental health and general early childhood outcomes, with depressed mothers more likely to have children who are ill, malnourished, and have low birth weight [9–15]. Patel et al. found that depressed postpartum mothers spent about twice the number of days in the previous 30 days unable to complete their daily activities [38]. It is likely that maternal mental health issues are associated with lower maternal health-seeking behaviors and practices, thus infants are more likely to experience symptoms of illness. Similarly, higher rates of maternal mental health problems may be leading to an increase in infant doctor visits. These results confirm the inverse relationship between maternal mental health and infant health outcomes.

Additionally, when examining the correlations between maternal mental health and handwashing practices, we found significant overall negative associations between symptoms of mental health problems and handwashing practices. Specifically, mothers at risk of depression is significantly less likely to have a higher handwashing count, frequently wash their hands before feeding the infant or cleaning infant's bottom, or having cleaning agent at the place of handwashing; mother with anxiety or stress symptoms is less likely to have a higher handwashing account or frequently wash their hands after cleaning infant's bottom. These findings are consistent with the few studies that exist on mental health and hygiene, which have similarly found that the two are negatively associated [16–18]. Given these

findings, it is likely that mental health issues are associated with mothers paying less attention to their infant's hygiene—an association which is common with other child care practices, such as infant nutrition [39–41] and immunizations [9].

Our findings show that maternal mental health in rural China has far reaching implications for both hygiene and infant illnesses, and thus requires more attention from policymakers and researchers. For instance, proactively including mental health care into perinatal care may mitigate the associations found between poor maternal mental health and hygiene practices [16–18]. This type of mental health support is likely to result in positive behavioural outcomes such as increased engagement and greater attentiveness towards infant health [25,42–46]. Additional interventions which teach mothers about the importance of hygiene practices are another good step in preventing poor infant health outcomes. While depressed mothers displayed the lowest adherence to hygiene practices, general hygiene practices were low across the entire sample population, indicating that additional interventions focusing on health and hygiene education may be useful in preventing poor infant health outcomes. Emphasis on when and how to properly wash their hands (such as before feeding their infants or after cleaning the bottom of their infants) is likely to result in better hygiene practices across the community and lower rates of infant illness.

This study has many strengths. It is one of the first to examine the correlations between maternal mental health, handwashing practices, and infant illness outcomes, and the first paper to do so in China. This study also fills a gap in the literature on maternal mental health and infant care in rural China, by at women with infants aged 0-6 months, as opposed to previous studies which have rarely sampled women with young children.

We acknowledge that this study has several limitations. For one, since our analysis utilizes cross sectional data, we are not able to draw causal relationships between mental health and hygiene practices. Second, the hygiene measures could be more extensive as they

only focused on handwashing and soap ownership. In addition, handwashing was measured based solely on recall which can be inaccurate. Future research should examine hygiene practices in detail and over time to understand the causal relationships between maternal mental health, hygiene, and infant health in rural China.

Conclusion

This study found that maternal mental health is significantly associated with hygiene practices and infant health outcomes. The results show a significantly negative association between symptoms of mental health problems and handwashing practices. Infants who have mothers with poor mental health are also more likely to have higher rates of illness. The study suggests that postnatal mental health interventions may be effective in mitigating poor infant health outcomes.

Abbreviations

LMICs: Low- and Middle-Income Countries

DASS-21: Depression, Anxiety, and Stress Scale (short form)

International Food Policy Research Institute: IFPRI

Declarations

Ethics approval and consent to participate

This study received ethical approval from the Stanford University Institutional Review Board (Protocol # 44312). All participants provided oral consent for participation in this study.

Individuals found to have severe mental health problems were referred to the local hospital for diagnosis and treatment.

Availability of data and materials

Datasets used in the current study are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was funded by National Natural Science Foundation of China (No.71874114).

Funding sources played no role in the design of the study, data collection, analysis, or interpretation, or in writing the manuscript.

Authors' contributions

QJ, NC, MO, EZ, SZ, YG, SD and SR contributed to the conceptualization of the study. QJ, QW, EZ, YG, SD and SR contributed to the acquisition, analysis or interpretation of data. QJ, NC, MO, EZ, SZ, YG drafted the manuscript, and QJ, JG, QW, HJ, SD, HZ, and SR made substantial revisions to the manuscript.

References

1. Connecting the Brain to the Rest of the Body: Early Childhood Development and Lifelong Health Are Deeply Intertwined Working Paper No.15. [Internet]. Center on the Developing Child Harvard University. 2020 [cited 2020 Dec 10]. Available from: <https://developingchild.harvard.edu/resources/connecting-the-brain-to-the-rest-of-the-body-early-childhood-development-and-lifelong-health-are-deeply-intertwined/>
2. Children: improving survival and well-being [Internet]. World Health Organization. 2020 [cited 2020 Dec 10]. Available from: <https://www.who.int/news-room/fact-sheets/detail/children-reducing-mortality>
3. Wang K, Xiang L, Kang L, Miao L, Li Q, Li X, et al. Communicable disease mortality trends and characteristics of infants in rural China, 1996-2015. Vol. 20, BMC Public Health. BioMed Central Ltd.; 2020. p. 455.
4. Diarrhoeal disease [Internet]. World Health Organization. 2017 [cited 2020 Dec 10]. Available from: <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease>
5. Edward A, Jung Y, Chhorvann C, Ghee AE, Chege J. Association of mother's handwashing practices and pediatric diarrhea: Evidence from a multi-country study on community oriented interventions. Journal of Preventive Medicine and Hygiene. 2019;60(2):E93–102.
6. Taddese AA, Dagne B, Dagne H, Andualem Z. <p>Mother's Handwashing Practices and Health Outcomes of Under-Five Children in Northwest Ethiopia</p>. Pediatric Health, Medicine and Therapeutics. 2020 Mar;Volume 11:101–8.
7. Nsele NM, McKerrow NH. A study of self-reported handwashing practices of caregivers in relation to acute respiratory infections and gastroenteritis in infants in a

peri-urban community in kwazulu-natal province, south africa. SAJCH South African Journal of Child Health. 2019;13(1):23–6.

8. Maternal, Newborn & Child Health [Internet]. Bill and Melinda Gates Foundation. 2015 [cited 2020 Dec 10]. p. d. Available from:
https://www.mendeley.com/catalogue/3d58003b-039c-3558-8e4e-1680ad759d4e/?utm_source=desktop&utm_medium=1.19.4&utm_campaign=open_catalog&userDocumentId=%7B56ce60c5-c1cc-3923-ad1a-343664fd29de%7D
9. Rahman A, Iqbal Z, Bunn J, Lovel H, Harrington R. Impact of maternal depression on infant nutritional status and illness: A cohort study. Archives of General Psychiatry. 2004 Sep 1;61(9):946–52.
10. Rahman A, Bunn J, Lovel H, Creed F. Maternal depression increases infant risk of diarrhoeal illness: - A cohort study. Archives of Disease in Childhood. 2007 Jan 1;92(1):24–8.
11. Adewuya AO, Ola BO, Aloba OO, Mapayi BM, Okeniyi JAO. Impact of postnatal depression on infants' growth in Nigeria. Journal of Affective Disorders. 2008 May 1;108(1–2):191–3.
12. Ross J, Hanlon C, Medhin G, Alem A, Tesfaye F, Worku B, et al. Perinatal mental distress and infant morbidity in Ethiopia: A cohort study. Archives of Disease in Childhood: Fetal and Neonatal Edition. 2011 Jan 1;96(1):F59–64.
13. Ban L, Gibson JE, West J, Tata LJ. Association between perinatal depression in mothers and the risk of childhood infections in offspring: A population-based cohort study. BMC Public Health. 2010;10:799.

14. Chung TKHMFLTKMMYASKMCHFkMFrcpLDTSMrcp. Antepartum Depressive Symptomatology Is Associated With Adverse Obstetric and Neonatal Outcomes. *Psychosomatic Medicine*. 2001 Oct;63(5):830–4.
15. Misri S, Oberlander TF, Fairbrother N, Carter D, Ryan D, Kuan AJ, et al. Relation between prenatal maternal mood and anxiety and neonatal health. *Canadian Journal of Psychiatry*. 2004 Oct 29;49(10):684–9.
16. Ranasinghe S, Ramesh S, Jacobsen KH. Hygiene and mental health among middle school students in India and 11 other countries. *Journal of Infection and Public Health*. 2016 Jul 1;9(4):429–35.
17. Slekiene J, Mosler HJ. Does depression moderate handwashing in children? *BMC Public Health*. 2018 Aug 1;18(1):1–9.
18. Nguyen PH, Saha KK, Ali D, Menon P, Manohar S, Mai LT, et al. Maternal mental health is associated with child undernutrition and illness in Bangladesh, Vietnam and Ethiopia. *Public Health Nutrition*. 2013;17(6):1318–27.
19. Zhang J, Shi L, Wang J, Wang Y. An infant and child feeding index is associated with child nutritional status in rural China. *Early Human Development*. 2009 Apr 1;85(4):247–52.
20. Shah SM, Yousafzai M, Lakhani NB, Chotani RA, Nowshad G. Prevalence and correlates of diarrhea. *Indian Journal of Pediatrics*. 2003 Mar 1;70(3):207–11.
21. Mbonye AK. Prevalence of childhood illnesses and care-seeking practices in rural Uganda. *The Scientific World Journal*. 2003;3:721–30.

22. Bhandari N, Mazumder S, Bahl R, Martines J, Black RE, Bhan MK. An educational intervention to promote appropriate complementary feeding practices and physical growth in infants and young children in Rural Haryana, India. *Journal of Nutrition*. 2004;134(9):2342–8.
23. Yarnoff BO, Allaire BT, Detzel P. Associations between infant feeding practices and length, weight, and disease in developing countries. *Frontiers in Pediatrics*. 2013 Sep 1;1(SEP).
24. Jiang W, Mo M, Li M, Wang S, Muyiduli X, Shao B, et al. The relationship of dietary diversity score with depression and anxiety among prenatal and post-partum women. *Journal of Obstetrics and Gynaecology Research*. 2018 Jul 26;44(10):1929–36.
25. Zhang Y, Muyiduli X, Wang S, Jiang W, Wu J, Li M, et al. Prevalence and relevant factors of anxiety and depression among pregnant women in a cohort study from south-east China. *Journal of Reproductive and Infant Psychology*. 2018 Oct 20;36(5):519–29.
26. Yue A, Gao J, Yang M, Swinnen L, Medina A, Rozelle S. Caregiver depression and early child development: A mixed-methods study from rural China. *Frontiers in Psychology*. 2018 Dec 10;9(DEC):2500.
27. Mental Health and Substance Use [Internet]. World Health Organization. 2016 [cited 2020 Dec 10]. Available from: <https://www.who.int/teams/mental-health-and-substance-use/maternal-mental-health>
28. National Bureau of Statistics of China [Internet]. 2019 [cited 2020 Dec 10]. Available from: <http://www.stats.gov.cn/english/>

29. Lovibond PF, Lovibond SH. The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour Research and Therapy*. 1995;33(3):335–43.
30. Wang K, Shi HS, Geng FL, Zou LQ, Tan SP, Wang Y, et al. Research on translations of tests: Cross-cultural validation of the depression anxiety stress scale-21 in China. *Psychological Assessment*. 2016 May 1;28(5):e88–100.
31. Chan RCK, Xu T, Huang J, Wang Y, Zhao Q, Shum DHK, et al. Extending the utility of the Depression Anxiety Stress scale by examining its psychometric properties in Chinese settings. *Psychiatry Research*. 2012 Dec 30;200(2–3):879–83.
32. Bangladesh Alive & Thrive Baseline Survey 2010: Shasthya Shebika (SS) | IFPRI : International Food Policy Research Institute [Internet]. 2010 [cited 2020 Dec 10]. Available from: <https://www.ifpri.org/publication/bangladesh-alive-thrive-baseline-survey-2010-shasthya-shebika-ss>
33. China Health and Nutrition Survey 1989-2011 | GHDx [Internet]. 2011 [cited 2020 Dec 10]. Available from: <http://ghdx.healthdata.org/record/china-health-and-nutrition-survey-1989-2011>
34. Song IH, Kim SA, Park WS. Family factors associated with children's handwashing hygiene behavior. *Journal of Child Health Care*. 2013 Jun 28;17(2):164–73.
35. Kolenikov S, Angeles G. Socioeconomic status measurement with discrete proxy variables: Is principal component analysis a reliable answer? *Review of Income and Wealth*. 2009 Mar 1;55(1):128–65.

36. 9/21/09 Summary | The American Cleaning Institute (ACI) [Internet]. 2009 [cited 2020 Dec 10]. Available from: <https://www.cleaninginstitute.org/newsroom/surveys/92109-summary>
37. Tao SY, Cheng YL, Lu Y, Hu YH, Chen DF. Handwashing behaviour among Chinese adults: A cross-sectional study in five provinces. *Public Health*. 2013 Jul;127(7):620–8.
38. Patel V, Rodrigues M, DeSouza N. Gender, poverty, and postnatal depression: A study of mothers in Goa, India. *American Journal of Psychiatry*. 2002 Jan;159(1):43–7.
39. Harpham T, Huttly S, De Silva MJ, Abramsky T, Harpham T. Maternal mental health and child nutritional status in four developing countries. *J Epidemiol Community Health*. 2005;59:1060–4.
40. Barker ED, Kirkham N, Ng J, Jensen SKG. Prenatal maternal depression symptoms and nutrition, and child cognitive function. *British Journal of Psychiatry*. 2013 Dec;203(6):417–21.
41. Haithar S, Kuria MW, Sheikh A, Kumar M, Vander Stoep A. Maternal depression and child severe acute malnutrition: A case-control study from Kenya. *BMC Pediatrics*. 2018 Sep 3;18(1):1–9.
42. Whiffen VE, Gotlib IH. Infants of Postpartum Depressed Mothers: Temperament and Cognitive Status. *Journal of Abnormal Psychology*. 1989;98(3):274–9.
43. Downey G, Coyne JC. Children of depressed parents: An integrative review. *Psychological Bulletin*. 1990;108(1):50–76.

44. Lovejoy MC, Graczyk PA, O'Hare E, Neuman G. Maternal depression and parenting behavior: A meta-analytic review. *Clinical Psychology Review*. 2000 Aug 1;20(5):561–92.
45. Cummings EM, Cd K, Cummings EM, Kouros CD. Maternal Depression and its Relation to Children's Development and Adjustment. In: *Encyclopedia on Early Childhood Development*. 2009.
46. Goodman JH, Santangelo G. Group treatment for postpartum depression: A systematic review. Vol. 14, *Archives of Women's Mental Health*. Springer; 2011. p. 277–93.