

Using the Risk Perception Attitude Framework to understand Nasopharyngeal Cancer risk reducing behaviors

Su-Hie Ting (✉ suhieting@gmail.com)

Universiti Malaysia Sarawak Fakulti Sains dan Teknologi Sumber <https://orcid.org/0000-0002-2479-6395>

Rayenda Khresna BRAHMANA

Universiti Malaysia Sarawak Fakulti Ekonomi dan Bisnis

Collin Jerome

Universiti Malaysia Sarawak

Yuwana Podin

Universiti Malaysia Sarawak

Research article

Keywords: Nasopharyngeal cancer, Risk Perception Attitude framework, risk reducing behaviours, Malaysians

Posted Date: February 3rd, 2020

DOI: <https://doi.org/10.21203/rs.2.22503/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: To reduce deaths due to nasopharyngeal cancer (NPC), it is important to understand factors that predict NPC risk reducing behaviors. This study employed the Risk Perception Attitude (RPA) framework to predict NPC risk reducing behaviors among a group of Malaysians.

Methods: A sample of Malaysians (n=215) completed a questionnaire about perceived susceptibility, perceived severity, self-efficacy, response efficacy and intention to enact self-protective actions to reduce NPC risk.

Results: A majority of the participants had responsive (high risk, high efficacy) and proactive attitudes (low risk, high efficacy). Hierarchical regression of mediation effect under structural equation model (SEM) approach was used to test the theory. Response efficacy and self-efficacy were negatively associated with perceived risk ($p < 0.01$). Intention was negatively associated with perceived risk and positively associated with response efficacy and self-efficacy ($p < 0.01$). The testing of the indirect effect from perceived risk effect yielded different conclusions for each efficacy measure. For the relationship between perceived risk and intention, there was no mediation effect of response efficacy but the mediation effect of self-efficacy was significant ($p < 0.1$). Heightened perceived risk weakens efficacy beliefs and intention to enact self-protective behavior, suggesting that low risk messages may work better to avert fatalistic thinking for this group. Perceived risk and response efficacy explained 26.5% of the variance in self-efficacy, suggesting the importance of framing NPC risk messages to heighten the audience's confidence to enact self-protective health behaviors.

Background

Nasopharyngeal cancer (NPC) when diagnosed before Stage 3 has good prognostic outcomes. Considering that NPC is sensitive to chemo-radiotherapy and results in a two- and three-year survival rate of 84% and 78% respectively in cases of early detection [1], regular screening is important to reduce mortality. Worldwide, deaths due to NPC number 50,000 out of 86,000 cases [2] and 71% of new NPC cases are from East and Southeast Asia [3]. In Malaysia NPC is the fifth most common cancer, and the lifetime risk for males was 1 in 143 and for females it was 1 in 417 in the 2007–2011 period [4]. Most of the NPC cases were detected at Stages 3 and 4 (63% for males, 60% for females) [5], suggesting lack of awareness towards NPC symptoms.

Minimizing risk factors can reduce NPC deaths. It is widely reported among NPC researchers that Epstein-Barr virus infection, smoking, and frequent consumption of preserved food and salted fish are associated with a high incidence of NPC [6]. Non-environmental risk factors of NPC include gender, ethnicity, and family history [7]. Certain ethnic groups such as the Bidayuh [8], Cantonese [9] and Malaysian residents in Sarawak, Penang and Labuan have higher NPC incidences [5]. Exposure to formaldehyde, wood dust, smoke and chemicals may also be involved in the pathogenesis of NPC [10]. Awareness of NPC risk factors can lead to adoption of risk reducing behaviors such as screening and reduced intake of NPC-

causing foods. However, taboos surrounding cancer discourage screening for early cancer detection [11, 12, 13]. However, little is known about NPC risk reducing behaviors and factors that predict these behaviors among Malaysians. Thus far, research on NPC in Malaysia are mostly clinical studies on the epidemiology of NPC [14, 15, 16, 17, 18, 19, 20, 21]. Some studies examined the knowledge of the primary care doctors on NPC [22] and social impact of NPC [23]. The present study is the first to investigate individuals' reasons for enacting preventive or self-protective measures to reduce risk in a setting where incidence of NPC deaths is high: NPC ranks number five in cancer incidences in Malaysia period [4]. This study aimed to employ the Risk Perception Attitude (RPA) framework to predict NPC risk reducing behaviors among a group of Malaysians.

Risk Perception Attitude (RPA) Framework

The RPA framework was developed by Rimal and Real [24] based on Witte's Extended Parallel Process Model [25] to understand the relationship between health risk perceptions and health behaviors, moderated by their efficacy beliefs. Risk perception is related to beliefs about perceived severity and perceived susceptibility, while risk prevention behaviour is related to efficacy beliefs, operationalised as the product of self-efficacy and response efficacy. Health behaviors refer to self-protective measures that individuals can take or intend to take to avoid certain diseases, including information-seeking behaviours. Based on their risk perceptions and efficacy beliefs, individuals are categorized into one of the four attitudinal groups: responsive (high risk, high efficacy), avoidance (high risk, low efficacy), proactive (low risk, high efficacy) and indifference (low risk, low efficacy). Their non-intervention study on skin cancer-related behaviours showed that the responsive group reported healthier outcomes than the avoidance group, and the proactive group reported healthier outcomes than the indifferent group [24].

The use of RPA has identified diverse risk factors influencing health behavioral intentions. For example, people who reported high levels of perceived cancer risk and strong self-efficacy were more motivated and able to engage in various health actions such as cancer-prevention diets [26], information seeking on cancer, and cancer screening [27, 28, 29]. Besides efficacy beliefs, other studies found that non-environmental risk factors moderated breast cancer risk perceptions and intention to undergo mammography screening such as rural-urban locality, religion, social-cultural beliefs [30], family history [31], age [32], and health literacy [33]. Spiritual health locus of control, for example, influenced American women's perceived risk and efficacy, which resulted in several health outcomes: message acceptance, talking about breast cancer, information seeking and behavioural intentions [34]. In addition, cancer fatalism, mistrust of health care providers, and previous negative experiences with the medical system have also been found to be associated with people's invariable attitudes towards cancer precautionary measures such as cancer screening and adherence to health care recommendations [35]. In their review of the field concluded that the causal relationship between people's health risk perceptions and their health behaviours is "more tenuous" than expected, given the "discrepanc(ies) in (research) findings" while some studies found a positive connection between people's perceptions of health risks and their health-related precautionary behavior, others discovered otherwise [27]. They further argued that the

causes for these discrepancies could be attributed to researchers' own failure to take into account the various moderators that may influence the causal link between risk perception and health behavior. Consistent with the RPA and the findings of studies that have employed this framework, we hypothesise that – efficacy beliefs, in addition to various modifying factors, will predict NPC risk reducing behaviors.

Methods

Participants and Procedure

Eligible participants were Malaysian residents ages 15 or older from various ethnic and socio-economic status groups. Given that the focus of the larger study was to investigate NPC risk reducing behaviors and factors that predict these behaviors, Malaysians with and without experience of NPC were eligible to be enrolled into the study. Based on the general rule, the minimum number of respondents or sample size is five-to-one ratio of the number of independent variables to be tested. However, Hair et al. [36] proposed that the acceptable ratio is ten-to-one. In the end, our sample size was 215 respondents. Since we could not get a list of all the elements of the population, we used a non-probability sampling of purposive sampling. The characteristics of 215 participants are reported in Table 1.

Table 1
Demographic characteristics of participants (n = 215)

Demographic Variables	n	%	Demographic Variables	n	%
Gender			Education		
Male	99	46%	Primary	5	2%
Female	116	54%	Form 3	8	4%
Age (Years)			Form 5/Certificate	77	36%
< 20	37	17%	Form 6/Diploma/Matriculation	39	18%
21–30	62	29%	Bachelor Degree	64	30%
31–40	58	27%	Postgraduate Degree	18	8%
41–50	22	10%	Professional Qualification	4	2%
51–60	15	7%	Income		
> 60	21	10%	No Income	45	21%
Marriage Status			<RM20,000	58	27%
Single	107	50%	RM2,000-RM3,999	66	31%
Married	99	46%	RM4,000-RM5,999	35	16%
Divorced	6	3%	>RM6,000	11	5%
Widowed	3	1%	Knowledge of NPC		
Ethnic Group			Some knowledge of NPC	123	57%
Malay	38	18%	Experienced NPC	28	13%
Chinese	83	39%	Family experienced NPC	14	7%
Sarawakian	88	41%	Work deals with NPC	12	6%
Others	6	3%	Friends and colleagues experienced NPC	43	20%
			Undertaken medical tests for NPC	12	6%

The objective of the study was explained to eligible Malaysian residents who agreed to participate. Enumerators administered the written informed consent and the questionnaire face-to-face. The study was conducted in Kuching, the capital city of Sarawak, a Malaysian state located on Borneo Kalimantan Island. Because this was a low-risk survey study, ethical approval was not needed at the university. However, to ensure the participants' safety, enumerators were trained to answer questions about NPC, and contact details of the researchers were provided to participants. The participants received RM5 (equivalent of USD 1.20) as a token of appreciation upon completion of the questionnaire.

Measures

The survey comprised a questionnaire eliciting self-reports of independent measures: perceived risk (three items, [37]); response efficacy (four items, [38]); and self-efficacy (six items, [39]). The dependent measures consisted of items on intended behaviour to reduce NPC risk (six items, [39, 40]). The questionnaire also contained demographic and health status questions.

All the items used 7-point Likert-type scales with a neutral option. The questionnaire items are shown in Table 2. The instrument was developed by first conducting a pilot study in which the researchers surveyed their contacts for appropriateness and clarity of the items, and then refining the items based on responses received.

Table 2
Results of measurement model

Model Construct	Items	Loading	CR	AVE	Cronbach Alpha
Perceived Risk	1. Nose and throat cancer could happen to me.	0.6243	0.8802	0.7163	0.8241
	2. How likely are you to get nose and throat cancer?	0.9334			
	3. How likely are you to get nose and throat cancer compared with other people your age?	0.9423			
Response Efficacy	2. How confident are you that changing to a healthier lifestyle (e.g., diet, exercise) can reduce the risk of nose and throat cancer?	0.7874	0.8088	0.5852	0.6494
	3. How confident are you that avoiding certain food	0.7605			
	4. How confident are you that avoiding environmental pollutants can reduce the risk of nose and throat cancer?	0.7464			
Self-Efficacy	1. How confident are you that you can change to a healthier lifestyle	0.6374	0.8630	0.6134	0.8140
	2. How easy is it for you to change to a healthier lifestyle to reduce risk to nose and throat cancer?	0.6640			
	3. How confident are you that you can avoid certain food said to cause nose and throat cancer to reduce risk to the disease?	0.7920			
	4. How easy is it for you to avoid certain food believed to cause nose and throat cancer to reduce risk to the disease?	0.7283			
	5. How confident are you that you can go for medical tests for nose and throat cancer for early detection/treatment of nose and throat cancer?	0.7502			
	6. How easy is it for you to go for medical tests for nose and throat cancer for early detection/treatment of nose and throat cancer?	0.7163			
Intention	1. I intend to lead a healthier lifestyle after today.	0.8279	0.9100	0.6280	0.8819

Model Construct	Items	Loading	CR	AVE	Cronbach Alpha
	2. I intend to lead a healthier lifestyle after today to reduce the risk of getting nose and throat cancer.	0.8358			
	3. I intend to avoid environmental pollutants after today.	0.8008			
	4. I intend to avoid environmental pollutants after today to reduce the risk of getting nose and throat cancer.	0.8018			
	5. I intend to do medical tests to monitor my health after today.	0.7571			
	6. I intend to do medical tests after today for early detection/treatment of nose and throat cancer.	0.7259			

Goodness of Measures: Reliability Test. This current research takes the coefficient of Cronbach's alpha to assess the reliability of the items. Table 2 reports the alpha values (0.8241 for perceived risk, 0.6494 for response efficacy, 0.8140 for self-efficacy, and 0.8819 for intention). Those alpha values are above 0.6, implying the items for each construct is reliable [41].

Goodness of Measures: Construct Validity. We tested the construct validity to obtain the measure fit [42]. It can be assessed from the respective loadings and cross loadings. We followed Hair et al.'s [36] threshold of 0.5 to infer the validation of an item. This means that an item needs to have value of 0.5 for its dimension only. As such, if an item has a loading value higher than 0.5 on two or more dimensions then it can be concluded there is an issue of cross loading. All items measuring a particular dimension loaded highly on that construct (higher than 0.5) and loaded lower on other construct (lower than 0.5) thus confirming construct validity.

Goodness of Measures: Convergent Validity. We tested further the items convergent validity by utilizing composite reliability and average variance extracted as the assessment. This test is important to ensure the degree to which multiple items that measure the same concept are in agreement. Table 2 reports composite reliability values of 0.8802, 0.8088, 0.8630, and 0.9100 for constructing perceived risk, response efficacy, self-efficacy, and intention respectively. Those values exceed the recommended threshold of 0.7 from Hair et al. [36]) implying the items have good convergent validity. Table 2 also reports the AVE values, whereas the values are 0.7163, 0.5852, 0.6134, and 0.6280 for constructing perceived risk, response efficacy, self-efficacy, and intention respectively. Those values exceed the recommended threshold of 0.5 from Barclays et al. [43]. Hence, we conclude that all four constructs (perceived risk, response efficacy, self-efficacy, and intention) are all valid measures of their respective dimensions based on the convergent validity test. All constructs (perceived risk, response efficacy, self-

efficacy, and intention) have valid measures based on their parameter estimates and statistical significance [44].

Goodness of Measures: Discriminant Validity. Lastly, we test discriminant validity of the measures to assess the degree to which items differentiate among constructs. This examination is important to ensure there is no potentially overlapping constructs. Table 3 shows that the squared correlation for each construct is less than the AVE values implying adequate discriminant validity.

Table 3
Discriminant validity of constructs

	Intention	Perceived Risk	Response Efficacy	Self-Efficacy
Intention	0.6280			
Perceived Risk	0.0325	0.7163		
Response Efficacy	0.1672	0.0384	0.5852	
Self-Efficacy	0.2307	0.0388	0.2756	0.6134

To test the hypotheses, we used hierarchical regression of mediation effect under Structural Equation Model approach. The dependent variable is intention to adopt NPC risk reducing behaviors while the main independent variables are perceived risk, response efficacy and self-efficacy. We added a mediator in our model due to the argument that the prediction of intention is intervened by response efficacy and self-efficacy.

Results

Based on Rimal and Real’s RPA framework, the participants were categorized into one of the four attitudinal groups using their means scores in perceived risk, response efficacy and self-efficacy beliefs [24]. Over half of the participants had a proactive attitude (118 low risk, high response efficacy; 114 low risk, high self-efficacy); “they are not motivated by their perceived risk status, but rather by their desire to remain disease free” [24]. Table 4 shows that about 44% of the participants had a responsive attitude (96 high risk and high response efficacy; 92 high risk and high self-efficacy), characterised as “being aware of their risk status and believing they have the requisite skills to avert the threat of the disease” and “motivated in enacting self-protective behaviour” [24]. Very few participants had avoidant (high risk, low self-efficacy), and indifferent (low risk, low self-efficacy) attitudes towards averting the NPC threat.

Table 4
Level of intention of participants for each RPA attitudes group

Scenario	Low		Medium		High		Total
	n	%	n	%	N	%	
Response Efficacy							
Indifferent (Low Risk/Low Response Efficacy)	0	0%	0	0%	0	0%	0
Proactive (Low Risk/High Response Efficacy)	1	1%	78	66%	39	33%	118
Avoidant (High Risk/Low Response Efficacy)	0	0%	0	0%	0	0%	0
Responsive (High Risk/High Response Efficacy)	1	1%	76	79%	19	20%	96
Self-Efficacy							
Indifferent (Low Risk/Low Self-Efficacy)	1	25%	3	75%	0	0%	4
Proactive (Low Risk/High Self-Efficacy)	0	0%	75	66%	39	34%	114
Avoidant (High Risk/Low Self-Efficacy)	1	20%	3	60%	1	20%	5
*Responsive (High Risk/High Self-Efficacy)	0	0%	74	80%	18	20%	92

Path analysis was employed to analyze the seven hypotheses. Figure 1 presents the results. In term of goodness of fit, our R2 value is 0.265 suggesting that 26.5% of the variance in the extent of self-efficacy can be explained by the hierarchical mediation of perceived risk and response-efficacy. Hence, we can conclude that our model has good explanation power.

We divided the results into three stages, namely, 1) direct effect between independent variables and mediator including the dependent variable; 2) direct effect between mediator and dependent variable; and 3) indirect effect between independent variable and dependent variable.

Hypothesis 1 stated that perceived risk has an effect on response efficacy. Our results show the negative effect of perceived risk on response efficacy with the coefficient value of -0.1960. The effect is significant at 1% level ($p < 0.01$) supporting the H1. Hence, higher perceived risk leads to lower response efficacy. In RPA framework, this behavior is categorized as avoidance. The same conclusion is found for the relationship between perceived risk and self-efficacy.

Hypothesis 2 stated that perceived risk has an effect on self-efficacy. We found that perceived risk has a negative effect on self-efficacy at the significance level of 1% with the coefficient value of -0.1968 supporting the H2. This implies that higher perceived risk leads to lower value of self-efficacy which is categorized as avoidance attitudes in the RPA framework. Thus, heightened risk perceptions may weaken efficacy beliefs. Perceived vulnerability to NPC seems to weaken confidence in taking precautionary measures and confidence in the measures.

Figure 1 depicts the direct effect between the independent variables and dependent variable. Hypothesis 3 stated that perceived risk has an effect on intention. Perceived risk has a significant effect on intention at 1% significance level supporting H3. The coefficient value is -0.1146 implying that higher perceived risk worsens the intention. It supports our H3. Heightened perceived risk may result in lower motivations to enact self-protective behavior to avert the threat of NPC, reflecting a fatalistic view of their vulnerability to the cancer (Rimal, 2000).

Hypothesis 4 stated that response efficacy has an effect on intention and Hypothesis 5 stated that self-efficacy has an effect on intention. In terms of the effect of efficacy effects on intention, Fig. 1 shows that both efficacy measures have significant effects on intention at 1% significance level. First, response efficacy has a significant effect on intention with a coefficient value of 0.214 implying higher response efficacy strengthens the intention. Self-efficacy is also found to have a significant effect on intention with the coefficient value of 0.3667 implying that a high level of self-efficacy strengthens intention to enacting NPC risk reducing behaviors. These findings provide support for H4 and H5. Participants who believe they can take effective measures to avert the threat of NPC may be motivated to engage in more extensive precautionary behavior.

We ran the indirect effect for testing the path analysis of response efficacy. The findings show that there is a significant effect of response efficacy as the mediator in linking perceived risk and self-efficacy. Therefore, confidence in the effectiveness of precautionary behavior induces self-confidence to adopt NPC risk reducing behaviors.

We also tested the indirect effect from perceived risk effect. Hypothesis 6 stated that response efficacy mediates the relationship between perceived risk and intention whereas Hypothesis 7 stated that self-efficacy mediates the relationship between perceived risk and intention. Our estimations have different conclusions for each efficacy measure. When we take response efficacy as the mediator, the result depicts that there is no mediation effect of response efficacy for the relationship between perceived risk and intention. Meanwhile, when self-efficacy is used as the mediator, there is a significant mediation effect to the extent that perceived risk influences intention. Yet, the relationship is arguably significant due to its 10% significance level. The mediation effect is negative implying that self-efficacy significantly carries the influence of perceived risk to reduce the intention of participants to enact NPC risk reducing behaviors. It seems that while efficacy behaviors play a limited role in mediating the effect of perceived risk upon intention, self-efficacy or confidence in adopting precautionary measures have some mediation effect.

Discussion

The current study examined the viability of RPA in predicting intention to enact NPC risk reducing behaviors in a sample of Malaysians. To our knowledge, this is the first study to use the RPA framework to understand factors influencing individuals' motivation to enact preventive or self-protective measures to reduce NPC risk in a setting of high NPC incidence. There were no similar surveys in Malaysia or

neighboring Asian countries with high incidence of NPC such as Indonesia and Hong Kong as the studies are also on epidemiology of NPC [45, 46, 47, 48, 49]. Hence, one strength of our study is that it has obtained information that is hitherto not available on perceptions of the severity of NPC, susceptibility to NPC and efficacy beliefs pertaining to reducing risk of NPC. Despite National Cancer Registry records showing high incidence of NPC among Malaysians (NPC ranking number five in cancer incidences [4]), the participants in our study perceived themselves to be at low risk to NPC and rated the severity of NPC as marginally low. However, their efficacy beliefs and intentions to enact self-protective behaviors were moderately high. Their perceptions of low threat of NPC could be due to lack of awareness of NPC incidences. The latest National Cancer Registry showed that age-standardised rates for nose and throat cancer incidences among men and women in Malaysia are 6.4 and 2.2 respectively in the 2007–2011 period [4]. The results on the lack of awareness of NPC are consistent with findings on the insufficient knowledge on breast cancer [50, 51], colorectal cancer [52], oral cancer [53], and cervical cancer [54, 55] in Malaysia. In fact, Samat et al.'s [55] survey showed that only 68% of the participants have some knowledge of cancer risk factors, and the study involved urban and educated Malaysians. In our study, 57% of the participants reported having some knowledge of NPC (Table 1). It is expected that the level of awareness on NPC would be even lower among the rural population, indicating the need for education.

Our study yielded somewhat unexpected results on the negative relationship between perceived risk with intention, response efficacy, and self-efficacy. In the RPA framework, Rimal and Real [24] predicted that high risk would increase motivation to enact self-protective health behaviours but in our study, it had the opposite effect. Two possible reasons are discussed here. First, there could be a strong sense of fatalism because the participants believed that there was not much they could do to avert NPC since cancer is unlike infectious diseases whereby people can take measures to minimize their risk. Ting et al. [13] found taboos surrounding cancer. The participants avoided the topic of NPC and were apprehensive when talking about it, for fear of inviting cancer into their lives [12]. They were mostly resigned to the fact that chance determined whether they got cancer. Elsewhere, taboos have been found to hinder women from taking up breast cancer screening [11]. Second, the participants could be unaware of their risk. As a group, the participants had a low level of perceived risk. Few had a family history of cancer (only 14 out of 215, Table 1), and they believed they led a generally healthy lifestyle, and were not exposed to potential risk factors such as preserved food and environmental pollutants.

Furthermore, these risk factors of NPC are not established because of mixed results in past studies. Our study indicated that a high level of perceived risk weakens efficacy beliefs and intention to enact self-protective measures. However, in information campaigns, it is still necessary to impart information on the risk so that the public is informed of the correct facts on NPC risk and severity but the framing should take account their psychological profile, determined using the RPA framework. In addition to health risk, the framing of the NPC risk messages can take into account financial risk because cancer treatment and management is costly [57], and the awareness may motivate the public to consider screening and lifestyle changes (e.g., preserved food, exposure to environmental pollutants) although the latter may not be confirmed risk factors of NPC.

Conclusions

The study provides important information on how risk and efficacy perceptions of a sample of the Malaysian population with respect to intention to enact self-protective behaviors. The most striking finding from our study is the stronger mediating influence of self-efficacy on perceived risk and intention, compared to response efficacy. In fact, response efficacy does not mediate between perceived risk and intention. Previous studies examined the relationship between perceived cancer risk and self-efficacy for intention to engage in cancer prevention diets [26], information seeking and cancer screening [27, 28, 29]. However, these studies treated efficacy beliefs as one construct. By treating response efficacy and self-efficacy as distinct constructs, our study has indicated that it is self-confidence to enact self-protective health behaviors which motivated participants to act on perceived risk of NPC. Surprisingly, response efficacy did not influence participants to alleviate their risk of NPC. In our study, the self-protective measures examined were screening, reducing preserved food (e.g., preserved vegetables, salted eggs), reducing exposure to environmental pollutants, and adopting a healthy lifestyle (e.g., exercise). The selection of these measures was based on research which has shown the association of NPC with preserved food [15, 58] and environmental pollutants [23, 59]. In our study, the participants had confidence that these self-protective measures were somewhat effective but this was not a factor in heightening their intention to adopt these measures. Instead their intention to enact these measures was influenced by their self-efficacy or their confidence to exert personal control over their health behaviors. The group of participants had high response and self-efficacy, with almost equal numbers having proactive and responsive attitudes. They had different motivations to enact self-protective behavior; the proactive group seeks to remain disease free whereas the response group seeks to avert the disease threat [24]. Hence, taking into view the mediating effect of self-efficacy between perceived risk and intention, education to increase awareness of NPC needs to focus on elevating the confidence of the target audience to take self-protective measures, and perhaps convincing them of the ease of performing those measures.

However, since it is the first of such studies to examine factors influencing individuals' motivation to enact self-protective measures to reduce NPC risk using the RPA, further studies should be conducted to test the model with other samples within the Malaysian population to find out whether it can explain intention to enact NPC risk reducing behaviors. While the current model has identified self-efficacy as an important construct to target in NPC risk message design, one limitation of the study is that the findings were based on self-reports of perceived risk of NPC which is variable and subject to an individual's prior beliefs. This was also clouded by the participants' taboos in assessing their own risk for fear that it would invite cancer incidences. In view of this, an important angle of investigation to undertake in future is message effects on perceptions of risk and efficacy to obtain evidence-based information on individual's conceptualization of perceived threat of NPC as a property of the message. The findings will inform the design of group-specific risk messages, taking into consideration the audience segmentation afforded by the survey findings on the influence of risk and efficacy beliefs on motivation to enact self-protective health behaviors.

Abbreviations

NPC

Nasopharyngeal cancer

RPA

Risk Perception Attitude

Declarations

Ethics approval and consent to participate

The enumerators explained the study to participants and asked them to sign a consent form indicating their willingness to participate in the study. They were told that no personal information would be revealed in any reports on the study. Written consent was obtained from participants who agreed to participate in the study. Because this was a low-risk survey study, ethical approval was not needed at the university. However, to ensure the participants' safety, enumerators were trained to answer questions about NPC, and contact details of the researchers were provided to participants. There were no participants under 16 years old in the study.

Consent for publication

Not applicable

Availability of data and materials

The data used to support the findings of this study are available from the corresponding author upon request.

Competing interests

The authors declare that they have no competing interests

Funding

This work was funded by the Ministry of Education, Malaysia via the Transdisciplinary Research Grant Scheme (FRGS) [Grant no. C09/TRGS/1519/2016].

Author's contributions

SHT conceived the study, reviewed the literature, designed the instruments, contributed to the analysis and interpretation of the data, and wrote the manuscript. RKB designed the statistical model, and analysed and interpreted the data. CJ reviewed the literature and co-wrote the manuscript. YP assisted with the interpretation of the results. The final version of the manuscript was read and approved by each contributing author.

Acknowledgements

Not applicable

References

1. Fles R, Bos ACRK, Rachmawati D, Waliyanti E, Tan IB, Haryana SM, ... Dewi FST. *Evaluation of diagnosis and treatment of nasopharyngeal carcinoma in Indonesia*. Amsterdam: Box Press; 2016.
2. Parkin DM, Bray F, Ferlay J. Global cancer statistics: 2002. *CA: A Cancer Journal for Clinicians*, 2005; 55(2):74-108.
3. Chang ET, Adami HO. The enigmatic epidemiology of nasopharyngeal carcinoma. *Cancer Epidemiology and Prevention Biomarkers*, 2006; 15(10):1765-77.
4. Ministry of Health, Malaysia. *Malaysian national cancer registry report: 2007-2011 Malaysia cancer statistics, data and figure*. Putrajaya, Malaysia: National Cancer Institute; 2015.
5. Ministry of Health, Malaysia. *Summary of Malaysian national cancer registry report 2007-2011*. Putrajaya, Malaysia: National Cancer Registry Department and National Cancer Institute; 2017.
6. Chua MLK, Wee JTS, Hui EP, Chan ATC. *Nasopharyngeal carcinoma*. *Lancet*, 2016; 5, 387(10022):1012-24.
7. Fles R, Wildeman MA, Sulistiono B. Knowledge of general practitioners about nasopharyngeal cancer at the Puskesmas in Yogyakarta, Indonesia. *BMC Med Educ*, 2010; 10(1):81.
8. Devi BC, Pisani P, Tang TS, Parkin DM. High incidence of nasopharyngeal carcinoma in native people of Sarawak, Borneo Island. *Cancer Epidemiol Biomarkers Prev*. 2005; 13(3):482-6.
9. Wee J, Ha TC, Loong S, Qian C. Is nasopharyngeal cancer really a "Cantonese cancer"? *Chin J Cancer*, 2010; 29(5):517-26.
10. Mahdavifar N, Ghoncheh M, Mohammadian-Hafshejani A, Khosravi B, Salehiniya H. Epidemiology and inequality in the incidence and mortality of nasopharynx cancer in Asia. *Osong Public Health Res Perspect*, 2016; 7(6):360-72.
11. Banning M, Hafeez H. A two-center study of Muslim women's views of breast cancer and breast health practices in Pakistan and the UK. *Journal of Cancer Education*, 2010; 25, 349-53.
12. O'Callaghan C, Schofield P, Butow P, Nolte L, Price M, Tsintziras S, ... Goldstein D. "I might not have cancer if you didn't mention it": a qualitative study on information needed by culturally diverse cancer survivors. *Supportive Care in Cancer*, 2016; 24(1):409-18.
13. Ting SH, Jerome C, Podin Y, Wan Ahmad SS. *Evaluating nose and throat cancer risk: Number sense and taboos on ill-health*. Proceedings of 2nd International Conference on Social and Economic Development (ICSED2) 2018, Kuala Terengganu, Malaysia; 2018.
14. Armstrong RW, Kutty MK, Dharmalingam MK, Ponnudurai JR. Incidence of nasopharyngeal carcinoma in Malaysia, 1968-1977. *British Journal of Cancer*, 1979; 40(4):557-67.

15. Armstrong RW, Armstrong MJ, Mimi CY, Henderson BE. Salted fish and inhalants as risk factors for nasopharyngeal carcinoma in Malaysian Chinese. *Cancer Research*, 1983; 43(6):2967-70.
16. Aziz A, Ramli RR, Mohamad I, Bhavaraju VMK. Young nasopharyngeal carcinoma: a review of an 8-year experience in the East Coast Malaysia Hospital. *The Egyptian Journal of Otolaryngology*, 2017; 33(2):490-4.
17. Prasad U, Rampal L. Descriptive epidemiology of nasopharyngeal carcinoma in Peninsular Malaysia. *Cancer Causes & Control*, 1992; 3(2):179-82.
18. Tarone RE, Levine PH, Yadav M, Pandey JP. Relationship between immunoglobulin allotypes and susceptibility to nasopharyngeal carcinoma in Malaysia. *Cancer Research*, 1990; 50(11):3186-8.
19. Teoh JW, Yunus RM, Hassan F, Ghazali N, Abidin ZA. Nasopharyngeal carcinoma in dermatomyositis patients: A 10-year retrospective review in Hospital Selayang, Malaysia. *Reports of Practical Oncology & Radiotherapy*, 2014; 19(5):332-6.
20. Tiong TS, Selva KS. Clinical presentation of nasopharyngeal carcinoma in Sarawak Malaysia. *Medical Journal of Malaysia*, 2005; 60(5): 624-8.
21. Yu ZH, Xu GZ, Hu YH, Su XG, Gu XZ. Value of computed tomography in staging the primary lesion (T-staging) of nasopharyngeal carcinoma (NPC): an analysis of 54 patients with special reference to the parapharyngeal space. *International Journal of Radiation Oncology, Biology, Physics*, 1985; 11(12): 2143-7.
22. Balachandran R, Philip R, Avatar S, Simon R, Mann GS, Benedict CTW, ... Ch'ng ML. Exploring the knowledge of nasopharyngeal carcinoma among medical doctors at primary health care level in Perak state, Malaysia. *European Archives of Oto-Rhino-Laryngology*, 2012; 269(2):649-58.
23. Armstrong RW, Armstrong MJ, Lye MS. Social impact of nasopharyngeal carcinoma on Chinese households in Selangor, Malaysia. *Singapore Medical Journal*, 2000; 41(12):582-7.
24. Rimal RN, Real K. Perceived risk and efficacy beliefs as motivators of change: Use of the risk perception attitude (RPA) framework to understand health behaviors. *Human Communication Research*, 2003; 29(3):370-99.
25. Witte K. Fear control and danger control: A test of the extended parallel process model (EPPM). *Communication Monographs*, 1994; 61:113-4.
26. Sullivan HW, Beckjord EB, Rutten LJF, Hesse BW. Nutrition-related cancer prevention, cognitions and behavioral intentions: Testing the risk perception attitude framework. *Health Education & Behavior*, 2008; 35(6):866-79.
27. Rimal RN, Juon H-S. Use of the risk perception attitude framework for promoting breast cancer prevention. *Journal of Applied Social Psychology*, 2010; 40(2):287-310.
28. Wong CH. Investigating the effects of cancer risk and efficacy perceptions on cancer prevention adherence and intentions. *Health Communication*, 2009; 24(2):95-105.
29. Wong CH. Interaction of comparative cancer risk and cancer efficacy perceptions on cancer-related information seeking and scanning behaviors. *Communication Research Reports*, 2012; 29(3):193-203.

30. Allo TA, Edewor PA, & Imhonopi D. Assessment of perceived risks of breast cancer and breast cancer screening among women in five selected local governments in Ogun State, Nigeria. *SAGE Open*, 2019; 9(2):1-11.
31. Chen LS, Kaphingst KA. Risk perceptions and family history of lung cancer: differences by smoking status. *Public Health Genomics*, 2011; 14(1):26-34.
32. Hanoch Y, Rolison JJ, Freund AM. Does medical risk perception and risk taking with age? *Society for Risk Analysis*, 2016; 38(5):917-28.
33. Peters E, Dieckmann N, Dixon A, Hibbard JH, Mertz CK. Less is more in presenting quality information to consumers. *Medical Care Research and Review*, 2007; 64(2): 169-90.
34. Leshner G, I-Huei C, Hyun JS, Yoon YC, Frisby C. The role of spiritual health locus of control in breast cancer information processing between African American and Caucasian women. *Integrative Medicine Insights*, 2006; 2(2):35-44.
35. Morris NS, Field TS, Wagner JL, Cutrona SL, Roblin DW, Gaglio B, Williams AE, Han PJK, Costanza ME, Mazor KM. The association between health literacy and cancer-related attitudes, behaviors, and knowledge. *Journal of Health Communication*, 2013; 18(Sup1):223-41.
36. Hair JF, Black WC, Babin BJ, Anderson. RE, 2010. *Multivariate Data Analysis*. New Jersey, Pearson Prentice Hall; 2010.
37. Katapodi MC, Lee KA, Facione NC, Dodd M. Predictors of perceived breast cancer risk and the relation between perceived risk and breast cancer screening: A meta-analytic review. *Preventive Medicine*, 2004; 38:388-402.
38. Grasso KL, Bell RA. Understanding health information seeking: A test of the risk perception attitude framework. *Journal of Health Communication*, 2015; 20(12):1406–14.
39. Bell RA, McGlone MS, Dragojevic M. Bacteria as bullies: Effects of linguistic agency assignment in health message. *Journal of Health Communication*, 2014; 19(3):340-58.
40. Rimal RN, Böse K, Brown J, Mkandawire G, Folda L. Extending the purview of the risk perception attitude framework: findings from HIV/AIDS prevention research in Malawi. *Health Communication*, 2009; 24(3):210-18.
41. Nunnally JC., Bernstein IH. *Psychometric theory*. New York: McGraw-Hill; 1994.
42. Sekaran U, Bougie R. *Research methods for business: A skill building approach*. United Kingdom: Wiley; 2010.
43. Barclay D, Higgins C, Thompson R. The Partial Least Squares (PLS) approach to causal modeling: Personal computer adoption and use as an illustration. *Technology Studies*, 1995; 2(2):285
44. Chow WS, Chan LS. Social network, social trust and shared goals in organizational knowledge sharing. *Information & Management*, 2008; 45(7):458-65.
45. Adham M, Kurniawan AN, Muhtadi AI, Roezin A, Hermani B, Gondhowiardjo S, ... & Middeldorp JM. Nasopharyngeal carcinoma in Indonesia: epidemiology, incidence, signs, and symptoms at presentation. *Chinese Journal of Cancer*, 2012; 31(4), 185.

46. Cheung ST, Lo KW, Leung SF, Chan WY, Choi PH, Johnson PJ ... Huang DP. Prevalence of LMP1 deletion variant of Epstein-Barr virus in nasopharyngeal carcinoma and gastric tumors in Hong Kong. *International Journal of Cancer*, 1996; 66(5):711-2.
47. Tse LA, Yu IT, Mang OW, Wong SL. Incidence rate trends of histological subtypes of nasopharyngeal carcinoma in Hong Kong. *British Journal of Cancer*, 2006; 95(9):1269-73.
48. Yu MC, Ho JH, Lai SH, Henderson BE. Cantonese-style salted fish as a cause of nasopharyngeal carcinoma: report of a case-control study in Hong Kong. *Cancer Research*, 1986; 46(2):956-61.
49. Rahman S, Subroto H, Novianti D. Clinical presentation of nasopharyngeal carcinoma in West Sumatra, Indonesia. *Proceeding of the 20th World Congress of IFOS*.
50. Hadi MA, Hassali MA, Shafie AA, Awaisu A. Evaluation of breast cancer awareness among female university students in Malaysia. *Pharmacy Practice*, 2010; 8(1):29-34.
51. Norlaili AA, Fatimah MA, Daliana NFN, Maznah D. Breast cancer awareness of rural women in Malaysia: Is it the same as in the cities? *Asian Pacific Journal of Cancer Prevention*, 2013; 14(12):7161-4.
52. Su TT, Goh JY, Tan J et al. Level of colorectal cancer awareness: A cross sectional exploratory study among multi-ethnic rural population in Malaysia. *BMC Cancer*, 2010; 13(1):1-8.
53. Ghani WMN, Doss JG, Jamaluddin M, Kamaruzaman D, Zain RB. Oral cancer awareness and its determinants among a selected Malaysian population. *Asian Pacific Journal of Cancer Prevention*, 2013; 14(3):1957-63.
54. Khoo C L, Teoh S, Rashid AK. Awareness of cervical cancer and HPV vaccination and its affordability among rural folks in Penang Malaysia. *Asian Pac J Cancer Prev*, 2011; 12(6):1429-33.
55. Wong LP, Wong YL, Low WY, Khoo EM, Shuib R. Knowledge and awareness of cervical cancer and screening among Malaysian women who have never had a Pap smear: a qualitative study. *Singapore Medical Journal*, 2009; 50(1):49-53.
56. Samat N, Ghazali S, Atang C. Awareness and knowledge of cancer: a community survey in Kedah and Perlis. *Asian Social Science*, 2014; 10(21):10-8.
57. Zafar SY, Peppercorn JM, Schrag D, Taylor DH, Goetzinger AM, Zhong X, Abernethy AP. The financial toxicity of cancer treatment: a pilot study assessing out-of-pocket expenses and the insured cancer patient's experience. *The Oncologist*, 2013; 18(4):381-90.
58. Armstrong RW, Imrey PB, Lye MS, Armstrong MJ, Yu MC, Sani S. Nasopharyngeal carcinoma in Malaysian Chinese: Salted fish and other dietary exposures. *International Journal of Cancer*, 1998; 77(2):228-35.
59. Armstrong RW, Kutty MK, Armstrong MJ. Self-specific environments associated with nasopharyngeal carcinoma in Selangor, Malaysia. *Social Science & Medicine. Part D: Medical Geography*, 1978; 12(3-4):149-156.

Figures




Figure 1

Structured model of the RPA framework predicting intention to adopt NPC risk reducing behaviors among Malaysians.