Insecticidal features displayed by the beneficial rhizobacterium *Pseudomonas chlororaphis* PCL1606

Eva Arrebola (✉ ead@uma.es)

Universidad de Málaga <https://orcid.org/0000-0002-2308-7829>

Francesca R. Aprile

Universidad de Málaga: Universidad de Málaga

Claudia E. Calderón

Universidad de Málaga

Antonio de Vicente

Universidad de Málaga: Universidad de Málaga

Francisco M. Cazorla

Universidad de Málaga: Universidad de Málaga

Research Article

Keywords: Insecticidal, *Galleria mellonella*, *Pseudomonas chlororaphis*, avocado, biocontrol.

Posted Date: February 16th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1311663/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The biocontrol rhizobacterium *Pseudomonas chlororaphis* is one of the bacterial species of the *P. fluorescens* group where insecticide *fit* genes have been found. Fit toxin, supported with other antimicrobial compounds, gives the bacterial capacity to repel and fight against predators, such as nematodes and insect larvae, thus protecting the plant host and itself. *Pseudomonas chlororaphis* PCL1606 is an antagonistic rhizobacterium isolated from avocado roots with efficient biocontrol against fungal soil-borne disease. The main antimicrobial compound produced by *P. chlororaphis* PCL606 is 2-hexyl,5-propyl resorcinol (HPR), which plays a crucial role in effective biocontrol against fungal pathogens. Further analysis of the *P. chlororaphis* PCL1606 genome showed the presence of hydrogen cyanide (HCN), pyrrolnitrin (PRN) and homologous *fit* genes. To test the insecticidal activity and to determine the bases for such activity, single and double mutants on the biosynthetic genes of these four compounds were tested in a *Galleria mellonella* larval model using inoculation by injection. The results revealed that fit toxin and HPR are the main compounds responsible for the insecticide phenotype in *P. chlororaphis* PCL1606, and HCN and PRN could be considered supporting compounds.

Introduction

The greatest variety of living beings is concentrated in the soil, and most of them gather around the roots of plants. Beneficial microorganisms coexist in this ecological niche along with pathogens and predators. Among the beneficial rhizobacteria associated with plant roots, the genus *Pseudomonas* is one of the most studied. The principal interactions of these rhizobacteria with the plants include cooperation with the plant host by plant growth promotion (PGPR), induction of systemic acquisition resistance (SAR; Haas and Défago 2005; Bakker et al. 2007), and competition or antagonism to soil-borne phytopathogens. In addition, these bacteria can show insecticidal activity and use insects as vectors for dispersal (Kupferschmied et al. 2013).

The main group of rhizobacteria of the genus *Pseudomonas* is the *Pseudomonas fluorescens* group, where insecticidal activity is not a common trait. Moreover, according to genomic diversity analysis, the *P. fluorescens* group is divided into three subclades (Loper et al. 2012). Interestingly, the strains belonging to subclade number 2 neither harbour the Fit toxin nor have the ability to kill *Galleria mellonella* larvae (Ruffner et al. 2015). However, all strains belonging to *P. protegens* and *P. chlororaphis* species tested in that study (which represent subclade 1) have both entomotoxicities (Ruffner et al. 2015). There are some examples of *P. protegens* and *P. chlororaphis* strains that are able to infect and efficiently kill insect larvae after oral uptake (Kupferschmied et al. 2013), and, this trait could also be related to the rhizobacterial ability to resist the deleterious effects of grazing predators, resulting in better host protection and competition against surrounding organisms (Nandi et al. 2015).

A close association with insecticidal activity has been demonstrated in the model bacterium *P. protegens* Pf-5 for a set of genes named the *fit*ABCDEFGH cluster (*fit*, *fluorescens* insecticidal toxin; Péchy-Tarr et al. 2008, 2013; Ruffner et al. 2013). The unique virulence cassette harbours the *fitD* gene encoding the

protein Fit toxin as well as regulatory and secretion protein functions (Péchy-Tarr et al. 2008). However, *fitD* deletion mutants retain substantial toxicity, indicating the presence of additional virulence factors (Péchy-Tarr et al. 2008; Ruffner et al. 2013). Many antagonistic pseudomonads are able to produce several antimicrobial compounds, such as phenazines (PHZ), 2,4-diacetylphloroglucinol (DAPG), pyoluteorin (PLT), pyrrolnitrin (PRN), hydrogen cyanide (HCN), or cyclic lipopeptides, that form a cocktail able to repel plant pathogens (Haas and Keel 2003; Gross and Loper 2009; Rochat et al. 2010; Kupferschmied et al. 2013). It has been reported that some of these primary antifungal compounds can also be used for self-defence against predators, such as protozoa and nematodes (Bjørnlund et al. 2009; Jousset et al. 2009; Raaijmakers and Mazzola 2012). Specifically, HCN has shown nematocidal activity against *Meloidogyne hapla*, supported by PHZ and PRN production (Lee et al. 2011). Moreover, Nandi et al. (2015) reported that HCN and PRN were able to act as powerful fungal repellents; in addition, these two compounds are the key compounds that affect the interaction of *P. chlororaphis* PA23 and *Caenorhabditis elegans*. In this sense, *Galleria mellonella* larvae are being used as an insect model for antibiotic susceptibility testing (Tsai et al. 2016; Ignasiak and Maxwell 2017; Andrea et al. 2019), toxicity of chemicals (Allegra et al. 2018) and virulence factors (Durieux et al. 2021), revealing their usefulness for insecticidal activity studies of several rhizobacteria.

The model rhizobacterium *P. chlororaphis* PCL1606 was isolated from the root of healthy avocado trees (*Persea Americana* Mill.) in a crop infected by *Rosellinia necatrix*, the causal agent of white rot root. This bacterium is characterized by being a highly efficient antagonist against many soil-borne phytopathogenic fungi (Cazorla et al. 2006; Calderón et al. 2014). Analysis of antifungal compounds produced by PCL1606 has shown the production of HCN and PRN; however, its main antifungal antibiotic produced by PCL1606 is 2-hexyl,5-propyl resorcinol (HPR; Cazorla et al. 2006; Calderón et al. 2015). Previous studies using random insertional derivative Tn5 mutants and selected insertional defective mutants lacking HPR have indicated a direct correlation between resorcinol and the biocontrol of soil-borne fungal diseases (Cazorla et al. 2006; Calderón et al. 2013). Furthermore, the locus encoding a putative cytotoxin homologous to FitD has been allocated to the PCL1606 genome (Calderon et al. 2015). These circumstances suggest that *P. chlororaphis* PCL1606 could be a good candidate to have insecticidal functions.

The main goal of this study was to unravel the roles of HCN, PRN, HPR and FitD compounds in the insecticidal features of *P. chlororaphis* PCL1606. For this, the insecticidal activity of PCL1606 was tested in a *Galleria mellonella* (greater wax moth) model.

Materials And Methods

Bacterial strains, plasmids and growth conditions

The plasmids and bacterial strains are described in Table 1. *Escherichia coli* strains were grown on LB medium (Ausubel et al. 1995) at 37°C for 24 h and supplemented with antibiotics according to the plasmid requirements. *Pseudomonas* spp. were grown and maintained on LB broth, KB medium (King et

al. 1954) and/or tryptone-peptone-glycerol medium (TPG, Calderón et al. 2013), supplemented with antibiotics as necessary (Table 1), and incubated at 25°C for 48 h. The antibiotic concentrations used in this study were 50 µg/mL kanamycin (km), 80 µg/mL gentamycin (Gm) for defective *P. chlororaphis* PCL1606 mutants and 40 µg/mL for the *E. coli* strain. *Galleria mellonella* larvae were obtained by commercial production (Animal-Center S.C., Spain), used immediately and maintained at 25°C in the dark during the experiments, and eventually preserved for one week at 4-10°C.

Table 1
Bacteria and plasmids used in the current study.

Strain	Relevant characteristics ^a	Reference
Bacteria		
<i>Pseudomonas</i> spp		
AVO110	<i>P. alcaligenes</i> ; efficient colonizer of avocado roots; antagonistic to <i>Rosellinia necatrix</i> .	Pliego et al. 2007 Pintado et al. 2021
BL915	<i>P. chlororaphis</i> subsp aurantiaca; antagonistic to <i>Rhizoctonia solani</i> .	Hill et al. 1994 Nowak-Thompson et al. 2003.
Pf-5	<i>P. protegens</i> ; antagonistic to <i>Rhizoctonia solani</i> ; insecticidal activity on <i>Galleria mellonella</i>	Howell and Stipanovic 1979; Péchy-Tarr et al. 2008; Lim et al. 2013
PCL1606	<i>P. chlororaphis</i> ; isolated from avocado rhizosphere; biocontrol, efficient root colonizer and antagonistic to <i>Rosellinia necatrix</i> and <i>Fusarium oxysporum</i>	Cazorla et al. 2006
PCL1606:: <i>darB</i>	PCL1606 derivative insertional mutant in <i>darB</i> gene; HPR-; Km ^r (former name Δ <i>darB</i>).	Calderon et al. 2013, 2019
ComB	PCL1606:: <i>darB</i> transformed with pCOMB; HPR+; Gm ^r	Calderon et al. 2013
PCL1606:: <i>prnC</i>	PCL1606 derivative insertional mutant in <i>prnC</i> gene; PRN-	Calderon et al. 2015
PCL1606:: <i>hcnB</i>	PCL1606 derivative insertional mutant in <i>hcnB</i> gene; HCN-	Calderon et al. 2015
PCL1606:: <i>fitD</i>	PCL1606 derivative insertional mutant in locus PCL1606_RS12180.	This study
PCL1606:: <i>darBprnC</i>	PCL1606 derivative double insertional mutant in <i>darB</i> and <i>prnC</i> genes; HPR-; PRN-	Calderón et al. 2015
PCL1606:: <i>darBhcnB</i>	PCL1606 derivative double insertional mutant in <i>darB</i> and <i>hcnB</i> genes; HPR-; HCN-	Calderón et al. 2015
PCL1606:: <i>prnC hcnB</i>	PCL1606 derivative double insertional mutant in <i>prnC</i> and <i>hcnB</i> genes; PRN-; HCN-	Calderón et al. 2015
PCL1606:: <i>darBfitD</i>	PCL1606 derivative double insertional mutant in <i>darB</i> and <i>fitD</i> genes; HPR-; FIT-	This study
^a HCN: production of hydrogen cyanide; HPR: production of 2-hexyl, 5-propyl resorcinol; PRN: production of pyrrolnitrin; FIT: production of FitD protein; Km ^r : kanamycin resistant; Gm ^r : gentamycin resistant; Amp ^r : ampicillin resistant		

Strain	Relevant characteristics ^a	Reference
PCL1606:: <i>gacS</i>	PCL1606 derivative insertional mutant in <i>gacS</i> gene, involved in secondary metabolism regulation.	Martín-Pérez et al. 2007
<i>Escherichia coli</i>		
DH5α	General cloning and sub-cloning applications; dlacZ Delta M15 Delta(lacZYA-argF) U169 recA1 endA1 hsdR17(rK-mK+) supE44 thi-1 gyrA96 relA1	Taylor et al. 1993
Plasmids		
pCR®2.1-TOPO®	PCR products cloning vector <i>lacZ</i> , Km ^r , Amp ^r	Invitrogen, California, USA
pJQ200SK	Suicide vector, P15A <i>oriV sacB mob</i> , Gm ^r	Quandt and Hynes 1993
pCRfitD	A fragment of PCL1606_RS12180 [PCL1606_24850] sequence cloned into pCR2.1, Km ^r for integrative mutation	This study
pJQfitD	A fragment of PCL1606_RS12180 [PCL1606_24850] sequence cloned into pJQ200SK, Km ^r for integrative mutation in PCL1606:: <i>darB</i>	This study
^a HCN: production of hydrogen cyanide; HPR: production of 2-hexyl, 5-propyl resorcinol; PRN: production of pyrrolnitrin; FIT: production of FitD protein; Km ^r : kanamycin resistant; Gm ^r : gentamycin resistant; Amp ^r : ampicillin resistant		

Table 2

Pseudomonas chlororaphis PCL1606 genes description used in the current study and tested by directed mutagenesis. Information obtained from National Center for Biotechnology Information (NCBI).

Locus	Gene	Product description	% Identity ncl-ncl	Reference Microorganism
PCL1606_RS08425	<i>gacS</i>	Response regulator	94	<i>Pseudomonas chlororaphis</i> 189
PCL1606_RS10175	<i>darB</i>	β -ketoacyl synthase DarB	92	<i>Pseudomonas aurantiaca</i> BL915
PCL1606_RS12180	<i>fitD</i>	Cytotoxin	84	<i>Pseudomonas protegens</i> Pf-5
PCL1606_RS13730	<i>prnC</i>	FAD-dependent oxidoreductase	94	<i>Pseudomonas aurantiaca</i> BL915
PCL1606_RS17610	<i>hcnB</i>	Cyanide-forming glycine dehydrogenase subunit HcnB	85	<i>Pseudomonas protegens</i> CHA0

Insecticide experiment in *Galleria mellonella*

The insecticide capacity of *Pseudomonas* spp. strains was performed in a *Galleria mellonella* larval model (Burgess 1976). The *Pseudomonas* spp. strains were grown in 30 mL of LB broth without antibiotics at 25°C overnight and 200 rpm of orbital agitation. Afterwards, the cultures were adjusted to 3×10^5 cfu/mL or 3×10^3 cfu/mL according to the assay. Ten millilitres from cultures was centrifuged at 4000 rpm for 10 min, and the pellets were resuspended in the same volume using 10 mM MgSO_4 buffer.

Commercial *Galleria mellonella* larvae approximately 1.5-cm long and 0.5-cm wide were selected for assays. The larvae were inoculated with approximately 10 μL of bacterial suspension injected into a 1-mL syringe with a needle 13-mm long and a 0.3-mm internal diameter (Becton Dickinson, Ireland). The needle and inoculation point were disinfected by 96% ethanol every time. Injections with strain Pf-5 (Table 1) and MgSO_4 buffer were performed as positive and negative controls, respectively. The experiments were performed by three independent assays with 15 larvae each. Once inoculated, the larvae were kept in the dark at 25°C for 3.5 days. Insecticidal activity was monitored after 17, 24, 30, 40, 60 and 80 hours by checking the absence of motility and melanisation of the larval body.

Strains Manipulation And Molecular Assays

Insertion mutagenesis for gene inactivation in *P. chlororaphis* PCL1606 was used to test the insecticide characteristics. Specifically, for this study, a disruptive vector was inserted into the putative *fitD* gene located on the PCL1606 chromosome via single-crossover homologous recombination (Arrebola et al. 2012). The cloning of vector pCR2.1 (Invitrogen Life Technologies USA) and plasmid purification were performed using standard procedures. The plasmids obtained were transformed into wild-type PCL1606 by electroporation (Choi et al. 2006). The double insertional mutant PCL1606::*darBfitD* was constructed using the previously obtained single insertional mutant PCL1606::*darB* (Calderón et al. 2013) in which the pCR2.1 derivative plasmid was already present in the mutant. Consequently, the suicide vector pJQ200SK (Table 1) was used to clone a *fitD* gene fragment and mutate the *fitD* gene by insertion of the chromosome via single-crossover homologous recombination.

The correct insertions of the disruption vectors were verified by PCR amplification. Bacterial growth curves were obtained in LB broth culture media to confirm similarities among the constructed defective mutants and the wild-type *P. chlororaphis* PCL1606 (data not shown).

Statistical analysis

The data were statistically analysed using an analysis of variance ANOVA (Sokal and Rohlf 1986), followed by Fisher's least significant difference test (LSD, $P = 0.05$) using IBM SPSS Statistics 22 software (SPSS Inc., Chicago, IL, United States). All experiments were performed at least three times independently.

Results

Genotypic and phenotypic study of the insecticidal capacity of *P. chlororaphis* PCL1606

An *in silico* analysis of putatively *fit* genes found in *P. chlororaphis* PCL1606 revealed a high similarity with *fit* gene operons detected in the insecticidal bacterium *Pseudomonas protegens* Pf-5, in which putative products are equivalents (Figure 1). Loci PCL1606_RS12165, PCL1606_RS12170 and PCL1606_RS12175 are related to the transport of cytotoxin, PCL1606_RS12180 is the largest locus and corresponds to the insect toxin FitD, PCL1606_12185 is annotated as an outer membrane protein, and PCL1606_RS12190, PCL1606_RS12195 and PCL1606_RS12200 are related to the regulation of toxin production. Equivalent genes were described in *P. protegens* Pf-5, where *fitABC* is related to transport by the type I secretion system, *fitD* has been described as a cytotoxin coding gene, *fitE* as a type I secretion outer membrane protein and *fitFGH* as a regulatory gene (Péchy-Tarr et al. 2008). Furthermore, the distribution and arrangement of these loci in the genome of PCL1606 are homologous to the *fit* cluster in *P. protegens* Pf-5. Likewise, the protein size and percentage of identity of the majority of loci with its equivalent *fit* gene showed that it was highly similar (Figure 1). Once the putative *fitD* gene was located in PCL1606, an insertional defective mutant was constructed to analyse its involvement in insecticide activity. Insecticidal activity using commercial *Galleria mellonella* larvae as a model by derivatising defective mutants of *P. chlororaphis* PCL1606 in antibiotic compounds, such as pirrolnitrine (PRN),

hydrogen cyanide (HCN) and 2-hexyl,5-propyl resorcinol (HPR), was studied, as well as in the mutant defective in the global regulator GacS (Table 1). The results have been compared with the wild-type strains *P. protegens* Pf-5, where a *fit* cluster was described (Péchy-Tarr et al. 2008), *P. chlororaphis* subsp. *aurantiaca* BL915, a producer of HPR (Nowak-Thompson et al., 2003), and *P. alcaligenes* AVO110, whose biocontrol activity was not related to antibiotic production (Pliego et al. 2007; Pintado et al. 2021). Twenty-four hours postinoculation with 3×10^5 cfu/mL as a dose (Figure 2), the control wild-type strains Pf-5, BL915 and PCL1606 caused a high insect mortality percentage (100% of dead larvae with intense larval melanization), contrary to AVO110, in which inoculation produced a comparable response to inoculation with the control buffer. Single mutants of *P. chlororaphis* PCL1606 in *darB* and *fitD* resulted in a slight decrease in mortality of approximately 15%, but they still had above 85% mortality in both cases. This drop in larval mortality shown by the PCL1606::*darB* mutant (defective in HPR) recovered to wild-type levels when this mutation was complemented (strain ComB). A decrease of approximately 40% in mortality occurred when single and double mutants in *hcnB* and *prnC* genes were tested. However, differences were observed when *darB* was involved in a double mutation with *hcnB* and *prnC*. Thus, in PCL1606 double mutants lacking HPR and PRN (PCL1606::*darBprnC*) or HPR and HCN (PCL1606::*darBhcnB*), the mortality levels observed increased to those shown by the wild-type strain values, which had higher mortality values than those displayed by the single mutant in *darB*, *prnC* or *hcnB*. A great difference was observed with the double mutant in the *darB* and *fitD* genes (PCL1606::*darBfitD*). The inoculation of this double mutant PCL1606::*darBfitD* decreased larval mortality to no inoculated control levels (Figure 2).

Involvement of HPR and Fit toxin in the toxicity of *G. mellonella*

A mortality survey eighty hours after inoculation with two different bacterial doses (3×10^3 and 3×10^5 cfu/mL; Figure 3 and S1) was performed. The results showed that 40 h after inoculation, the wild-type *P. chlororaphis* PCL1606 and single mutants in *darB*, *fitD* and *gacS* retained the highest mortality; however, the double mutant PCL1606::*darBfitD* reduced the *G. mellonella* mortality level to the buffer control values. This double mutant reached 60% *G. mellonella* larval mortality 80 h after inoculation. Therefore, there was a delay in the insecticidal activity of PCL1606 when HPR and Fit toxin were impaired in the same strain (Figure 3a). With the higher dose, all events reached the highest mortality from 16 h to 30 h after inoculation, but the double mutant HPR-Fit toxin showed a delay in killing the larvae, reaching 100% mortality 60 h after inoculation (Figure S1).

The symptoms observed in *G. mellonella* larvae 24 h after the injection (3×10^3 cfu/mL) with *fitD*- and *darB*-defective mutants (Figure 3b) showed that several larvae started to melanise. After 30 h, nearly all the larvae inoculated with the single mutants PCL1606::*fitD*, PCL1606::*darB* and PCL1606::*gacS* and the wild-type PCL1606 showed dark melanization. However, at 30 h postinoculation with buffer or with double mutant PCL1606::*darBfitD*, the inoculated larvae did not display signs of melanization and mobility, and only 60 h after inoculation, they began to show symptoms of larval intoxication in those infected with the HPR-Fit toxin double mutant (Figure 3b). The different mortality kinetics could be better distinguished at 30, 40 and 60 h after inoculation (Figure S2), showing that the wild-type PCL1606 strain

presented progressively increasing mortality, reaching 100% at 60 h. The single mutants in *fitD* and *darB* showed significantly higher virulence than the wild-type strain at 30 h. The single mutant in *gacS* seemed to be delayed until 40 h to display higher virulence. Finally, the double mutant seemed to be innocuous after 60 h, when insecticide activity began (Figure S2).

Discussion

Pseudomonas chlororaphis PCL1606 is a rhizobacterium isolated from avocado roots characterized by 2-hexyl,5-propyl resorcinol (HPR) production, the main compound involved in biocontrol (Cazorla et al. 2006), but also in other relevant phenotypes, such as biofilm formation and colonization (Calderón et al. 2019; Arrebola et al. 2019). A previous analysis of additional antifungal antibiotic compounds produced by PCL1606 revealed the production and presence of coding genes for hydrogen cyanide (HCN) and pyrrolnitrin (PRN); however, phenazine production and the presence of coding genes were not found in PCL1606 (Calderón et al. 2015). Additionally, the presence of genes homologous to the *fit*ABCDEFGH cluster, encoding a putative cytotoxin similar to Fit toxin, was detected in the genome of *P. chlororaphis* PCL1606 (Calderón et al. 2015). According to Ruffner et al. (2015), the Fit cytotoxin is restricted to a particular group of rhizobacteria comprised of *P. protegens* and *P. chlororaphis*, and it is strongly correlated with high insect toxicity.

PCL1606 displayed insecticidal activity at the same level as the control strains *P. chlororaphis* subsp *aurantiaca* (former *P. aurantiaca*) BL915 and *P. protegens* Pf-5, where HPR and Fit production were first described, respectively (Nowak-Thompson et al. 2003; Péchy-Tarr et al. 2008). On the other hand, the nonantagonistic rhizobacterium *P. alcaligenes* AVO110 did not show insecticidal capacity, did not have the *fit* or any antifungal antibiotic genes in its genome (Pintado et al. 2021), and did not produce antifungal secondary metabolites (Pliego et al. 2007; Pintado et al. 2021). The typical symptoms produced by the wild-type strain PCL1606 correspond to mortality accompanied by intense melanization of the *G. mellonella* larvae from 24 h postinoculation. Melanization is a typical insecticidal symptom corresponding to the synthesis and deposition of melanin to encapsulate pathogens at the wound site followed by haemolymph coagulation and opsonisation (Tsai et al. 2016).

Single and double mutants impaired the production of antifungal hydrogen cyanide (HCN) or pyrrolnitrin (PRN) compounds in PCL1606, resulting in a mortality reduction of *G. mellonella* larvae, but with no differences among them. This suggests that both compounds could only contribute to the insecticidal background and were not the main insecticidal compounds produced, as previously described for other insecticidal strains (Flury et al. 2017). It has been previously described that the insecticidal virulence of *P. protegens* CHA0 and *P. chlororaphis* PCL1391 mutants lacking one or several antibiotics, such as 2,4-diacetylphloroglucinol, phenazine, pyrrolnitrin or pyoluterin, was also not reduced (Flury et al. 2017).

The role of HCN and PRN on insecticidal and nematocidal activity by antagonistic bacteria has been illustrated by other authors (Nandi et al. 2015; Kang et al. 2019), and our results indicate their involvement in the insecticidal phenotype displayed by the model strain PCL1606. In addition, this study

included the PCL1606 derivative mutant defective in *gacS*, since *gacS* is a global regulator that can interfere with activities related to the production of secondary compounds, such as antifungals and insecticides (Saraf et al. 2011; Nandi et al. 2015). Previous studies have reported that mutants deficient in the global regulator *gacS* showed no insecticide activity (Kang et al. 2019). These results do not agree with our observation where the impaired mutants in the *gacS* gene of PCL1606 still retain insecticidal activity, suggesting a *gacS*-regulated independent pathway responsible for the insecticidal phenotype.

For this, the effect of HPR and Fit products was studied. Single mutants of these genes scarcely lowered *G. mellonella* larval mortality, but the absence of both genes completely impaired the insecticidal activity of the derivative bacteria, even in the presence of *hcnB* and *pmc* genes.

The insecticidal and nematocidal effectiveness of the Fit cytotoxin in *P. protegens* and *P. chlororaphis* has been documented (Kupferschmied et al. 2013; Péchy-Tarr et al. 2008). Heterologous expression of the Fit toxin in *Escherichia coli* resulted in the capacity of the bacterium to kill the insect host upon injection (Kupferschmied et al. 2013). In addition, the deletionally defective *fitD* mutants of *P. protegens* Pf-5 and CHA0 show a decrease in insecticidal activity, demonstrating that FitD makes an important contribution to insect virulence but may not be essential for the capacity for these bacteria to multiply in the infected host. The fact that the *fitD* mutants still retained a certain level of toxicity indicated to the authors that additional factors may contribute to anti-insect activity (Péchy-Tarr et al. 2008), in agreement with our observations on PCL1606. However, the PCL1606 derivative mutant with the fit toxin gene disrupted showed the capacity to kill *G. mellonella* at almost 80% 30 h postinfection. This result is not only due to HCN and PRN, as reported by the double derivative mutant in HPR and Fit toxin, with no virulence at this time postinfection. On the other hand, the double mutant HCN and PRN still showed that 60% of *G. mellonella* larvae died. Therefore, we can conclude that HCN and PRN could contribute to the insecticidal capacity of PCL1606, but they could be considered accompanying compounds. Thus, HPR and the Fit toxin could be considered the main toxins responsible for insecticidal activity. Single derivative mutants in HPR and Fit toxin displayed the same levels of *G. mellonella* mortality, suggesting that the mutation of one of them can complement the mortality index by the other gene and vice versa. However, the mortality shown by the derivative mutant PCL1606::*darBfitD* was innocuous after approximately one day and a half, when the first sign of mortality started to appear.

HPR has been described as the main antibiotic against fungal pathogens such as *Rosellinia necatrix* or *Fusarium oxysporum* (Cazorla et al. 2006). However, HPR has been revealed as a versatile compound involved in bacterial adhesion, colony morphology and typical air-liquid interphase pellicles produced by PCL1606 (Calderón et al. 2019). Due to the chemical nature of HPR, it is possibly not an integral component of the biofilm matrix, together with the evidence of a regulatory role in *Photorhabdus asymbiotica* (Hapeshi and Waterfield 2017), which extends its putative role in PCL1606.

The combined antimicrobial and regulatory roles of HPR in PCL1606 could explain several results obtained in the present study. The toxic nature of HPR against different organisms (Kanda et al. 1975) could directly affect *G. mellonella* cells, as happens with antifungal phenazines (Wand et al. 2013),

helping the insecticidal characteristics of *P. chlororaphis* PCL1606. This insecticidal activity would rival FitD in toxicity, thus justifying the single mutants PCL1606:*darB* and PCL1606:*fitD* results in comparison with double mutant PCL1606::*darBfitD*, also revealing that because of the absence of HPR and Fit toxin, there was no mortality due to HCN and PRN. On the other hand, HPR could have additional regulatory roles on secondary metabolites since alkylresorcinols can be involved as signalling molecules in a novel quorum sensing two-component regulatory system (Brameyer et al. 2015). Belonging to these sets of secondary metabolites could be chitinase and phospholipase C, putative genes that have been found in the PCL1606 genome (PCL1606_RS13585 as chitinase and PCL1606_RS14060 as phosphatase C). Both enzymes have also been reported to be important in insect pathogenicity (Flury et al. 2016). This also helps to explain the *gacS* mutant results, whose killing ability was not affected, suggesting a regulation of all the compounds at a higher hierarchy, but this regulatory role of HPR is still a hypothesis under study.

In summary, PCL1606 could produce the Fit toxin, a described compound with insecticidal capacity, and HPR, which has been shown to have insecticidal potential, to which its fungicidal character and its possible regulatory role must be added. This confirms HPR as one the main compounds produced by *P. chlororaphis* PCL1606 involved in the beneficial phenotypes displayed by this model bacterium.

Declarations

Acknowledgements

Thank to Mrs. Irene Linares Rueda for their technical assistance with laboratory research.

Data and materials availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request

Funding

This work was supported by research project AGL2017-83368-CO2-1-R of Ministerio de Ciencias y Tecnología, and the project UMA-FEDERJA-046 of Junta de Andalucía.

Competing Interests

The authors declare they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author's contribution

Eva Arrebola: Conceptualization, Data curation, Investigation, Formal analysis, Writing original draft. Francesca R. Aprile and Claudia E. Calderón: Assays performant and results. Antonio de Vicente:

Conceptualization and writing support. Francisco M. Cazorla: Funding acquisition, Methodology, Supervision, Writing, review and editing.

Ethics approval: This article does not contain any studies with human or animal subjects performed by any of the authors

Consent to participate: No applicable

Consent for publication: No applicable

Conflict of interest: The authors declare no competing interests

References

1. Allegra E, Titball RW, Carter J, Champion OL (2018) *Galleria mellonella* larvae allow the discrimination of toxic and non-toxic chemicals. *Chemosphere* 198:469–472. <https://doi.org/10.1016/j.chemosphere.2018.01.175>
2. Andrea A, Krogfelt KA, Jenssen H (2019) Methods and challenges of using the greater wax moth (*Galleria mellonella*) as a model organism in antimicrobial compound discovery. *Microorganisms* 7(3):85. <https://doi.org/10.3390/microorganisms7030085>
3. Arrebola E, Carrión VJ, Cazorla FM, Pérez-García A, Murillo J, de Vicente A (2012) Characterization of the mgo operon in *Pseudomonas syringae* pv. *syringae* UMAF0158 that is required for mangotoxin production. *BMC Microbiol* 12:2-17. <http://www.biomedcentral.com/1471-2180/12/10>
4. Arrebola E, Tienda S, Vida C, de Vicente A, Cazorla FM (2019) Fitness features involved in the biocontrol interaction of *Pseudomonas chlororaphis* with host plants: the case study of PcPCL1606. *Front Microbiol* 10:719. <https://doi.org/10.3389/fmicb.2019.00719>
5. Ausubel FM, Brent R, Kingston RE, Moore DD, Seidman JG, Smith JA, Struhl K (1995) Current protocols in molecular biology. John Wiley and Sons (ed), New York
6. Bakker PAHM, Pieterse CMJ, van Loon LC (2007) Induced systemic resistance by fluorescent *Pseudomonas* spp. *Phytopathology* 97(2):239–243. <https://doi.org/10.1094/PHYTO-97-2-0239>
7. Bjørnlund L, Rønn R, Péchy-Tarr M, Maurhofer M, Keel C, Nybroe O (2009) Functional GacS in *Pseudomonas* DSS73 prevents digestion by *Caenorhabditis elegans* and protects the nematode from killer flagellates. *ISME J* 3:770–779. <https://doi.org/10.1038/ismej.2009.28>
8. Brameyer S, Kresovic D, Bode HB, Heermann R (2015) Dialkylresorcinols as bacterial signaling molecules. *PNAS* 112(2):572–577. <https://www.pnas.org/cgi/doi/10.1073/pnas.1417685112>
9. Burges HD (1976) Techniques for the bioassay of *Bacillus thuringiensis* with *Galleria mellonella*. *Entomol Exp Appl* 19(3):243–254. <https://doi.org/10.1111/j.1570-7458.1976.tb02604.x>
10. Calderón CE, Pérez-García A, de Vicente A, Cazorla FM (2013) The *dar* genes of *Pseudomonas chlororaphis* PCL1606 are crucial for biocontrol activity via production of the antifungal compound

- 2-hexyl, 5-propyl resorcinol. *Mol Plant-Microbe Interact* 26(5):554–565.
<https://doi.org/10.1094/MPMI-01-13-0012-R>
11. Calderón CE, Ramos C, de Vicente A, Cazorla FM (2015) Comparative genomes analysis of *Pseudomonas chlororaphis* PCL1606 reveals new insight into antifungal compounds involved in biocontrol. *Mol Plant-Microbe Interact* 28(3):249–260. <https://doi.org/10.1094/MPMI-10-14-0326-FI>
 12. Calderón CE, Tienda S, Heredia-Ponce Z, Arrebola E, Cárcamo-Oyarce G, Eberl L, Cazorla FM (2019) The compound 2-hexyl, 5-propyl resorcinol has a key role in biofilm formation by the biocontrol rhizobacterium *Pseudomonas chlororaphis* PCL1606. *Front Microbiol* 10:396.
<https://doi.org/10.3389/fmicb.2019.00396>
 13. Calderón CE, de Vicente A, Cazorla FM (2014) Role of 2-hexyl, 5-propyl resorcinol production by *Pseudomonas chlororaphis* PCL1606 in the multitrophic interaction in the avocado rhizosphere during the biocontrol process. *FEMS Microbiol Ecol* 89:20–31. <https://doi.org/10.1111/1574-6941.12319>
 14. Cazorla FM, Duckett S, Bergstrom E, Noreen S, Odijk R, Lugtenberg BJJ, Thomas-Oates J, Bloemberg GV (2006) Biocontrol of avocado *Dematophora* root rot by antagonistic *Pseudomonas fluorescens* PCL1606 correlates with the production of 2-hexyl, 5-propyl resorcinol. *Mol Plant-Microbe Interact* 19(4):418–428. <https://doi.org/10.1094/MPMI-19-0418>
 15. Choi KH, Kumar A, Schweizer HP (2006) A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: Application for DNA fragment transfer between chromosomes and plasmid transformation. *J Microbiol Meth* 64:391–397.
<https://doi.org/10.1016/j.mimet.2005.06.001>
 16. Durieux MF, Melloul É, Jemel S, Roisin L, Dardé ML, Guillot J, Dannaoui É, Botterel F (2021) *Galleria mellonella* as a screening tool to study virulence factors of *Aspergillus fumigatus*. *Virulence* 12(1):818–834. <https://doi.org/10.1080/21505594.2021.1893945>
 17. Flury P, Aellen N, Ruffner B, Péchy-Tarr M, Fataar S, Metla Z, Dominguez-Ferreras A, Bloemberg G, Frey J, Goesmann A, Raaijmakers JM, Duffy B, Höfte M, Blom J, Smits TH, Keel C, Maurhofer M (2016) Insect pathogenicity in plant-beneficial pseudomonads: phylogenetic distribution and comparative genomics. *ISME J* 10(10):2527–2542. <https://doi.org/10.1038/ismej.2016.5>
 18. Flury P, Vesga P, Péchy-Tarr M, Aellen N, Denet F, Hofer N, Kupferschmied KP, Kupferschmied P, Metla Z, Ma Z, Siegfried S, de Weert S, Bloemberg G, Höfte M, Keel CJ, Maurhofer M (2017) Antimicrobial and insecticidal: cyclic lipopeptides and hydrogen cyanide produced by plant-beneficial *Pseudomonas* strains CHA0, CMR12a, and PCL1391 contribute to insect killing. *Front Microbiol* 8:100. <https://doi.org/10.3389/fmicb.2017.00100>
 19. Gross H, Loper JE (2009) Genomics of secondary metabolite production by *Pseudomonas* spp. *Nat Prod Rep* 26:1408–1446. <https://doi.org/10.1039/b817075b>
 20. Haas D, Défago G (2005) Biological control of soil borne pathogens by fluorescent pseudomonads. *Nat Rev Microbiol* 3(4):307–319. <https://doi.org/10.1038/nrmicro1129>

21. Haas D, Keel C (2003) Regulation of antibiotic production in root colonizing *Pseudomonas* spp. and relevance for biological control of plant disease. *Annu Rev Phytopathol* 41:117–153.
<https://doi.org/10.1146/annurev.phyto.41.052002.095656>
22. Hapeshi A, Waterfield NR (2017) *Photorhabdus asymbiotica* as an insect and human pathogen. *Curr Top Microbiol Immunol* 402:159–177. https://doi.org/10.1007/82_2016_29
23. Hill DS, Stein JI, Torkewitz NR, Morse AM, Howell CR, Pachlatko JP, Becker JO, Ligon JM (1994) Cloning of genes involved in the synthesis of pyrrolnitrin from *Pseudomonas fluorescens* and role of pyrrolnitrin synthesis in biological control of plant disease. *Appl Environ Microbiol* 60(1):78–85
24. Howell CR, Stipanovic RD (1979) Control of *Rhizoctonia solani* in cotton seedlings with *Pseudomonas fluorescens* and with an antibiotic produced by the bacterium. *Phytopathology* 69:480–482
25. Ignasiak K, Maxwell A (2017) *Galleria mellonella* (greater wax moth) larvae as a model for antibiotic susceptibility testing and acute toxicity trials. *BMC Res Notes* 10(1):428.
<https://doi.org/10.1186/s13104-017-2757-8>
26. Jousset A, Rochat L, Péchy-Tarr M, Keel C, Scheu S, Bonkowski M (2009) Predators promote defense of rhizosphere bacterial population by selective feeding on non-toxic cheaters. *ISME J* 3:666–674.
<https://doi.org/10.1038/ismej.2009.26>
27. Kanda N, Ishizaki N, Inoue N, Oshima M, Handa A (1975) DB-2073, a new alkylresorcinol antibiotic. I. Taxonomy, isolation and characterization. *J Antibiot (Tokyo)* 28(12):935–942.
<https://doi.org/10.7164/antibiotics.28.935>
28. Kang BR, Anderson AJ, Kim YC (2019) Hydrogen cyanide produced by *Pseudomonas chlororaphis* O6 is a key aphicidal metabolite. *Can J Microbiol* 65:185–190. <https://doi.org/10.1139/cjm-2018-0372>
29. King EO, Ward MK, Raney DE (1954) Two simple media for the demonstration of pyocyanin and fluorescin. *J Lab Clin Med* 44(2):301–307
30. Kupferschmied P, Maurhofer M, Keel C (2013) Promise for plant pest control: root-associated pseudomonads with insecticidal activities. *Front Plant Sci* 4:287.
<https://doi.org/10.3389/fpls.2013.00287>
31. Lee JH, Ma KC (2011) Nematicidal activity of a nonpathogenic biocontrol bacterium, *Pseudomonas chlororaphis* O6. *Curr Microbiol* 62:746–751. <https://doi.org/10.1007/s00284-010-9779-y>
32. Lim CK, Hassan KA, Penesyan A, Loper JE, Paulsen IT (2013) The effect of zinc limitation on the transcriptome of *Pseudomonas protegens* Pf-5. *Environ Microbiol* 15(3):702–715.
<https://doi.org/10.1111/j.1462-2920.2012.02849.x>
33. Loh JM, Adenwalla N, Wiles S, Proft T (2013) *Galleria mellonella* larvae as an infection model for group A streptococcus. *Virulence* 4:419–428. <http://dx.doi.org/10.4161/viru.24930>
34. Loper JE, Hassan KA, Mavrodi DV 2nd, Lim EW, Shaffer CK BT, et al (2012) Comparative genomics of plants associated *Pseudomonas* spp. insights into diversity and inheritance of traits involved in multitrophic interactions *PLoS Genet* 8(7):e10027834.
<https://doi.org/10.1371/journal.pgen.1002784>

35. Martín-Pérez R, Romero DF, Bonilla N, Pérez-García A, de Vicente A, Cazorla FM (2007) Identification of genes involved in the production of antibiotic 2-hexyl, 5-propyl resorcinol and its role in biocontrol. Lorito M, Scala F, Woo S, Ruocco M, Capodilupo C, Zoina A (eds) XIII International Congress on Molecular Plant-Microbe Interaction. Proceeding Book, Edizioni Ziino, Sorrento, Italy, pp. 220
36. Nandi M, Selin C, Brassinga AKC, Belmonte ML, Fernando WGD, Loewen PC, de Kievit TR (2015) Pyrrolnitrin and hydrogencyanide production by *Pseudomonas chlororaphis* strain PA23 exhibits nematicidal and repellent activity against *Caenorhabditis elegans*. PLoS ONE 10(4):e0123184. <https://doi.org/10.1371/journal.pone.0123184>
37. Nowak-Thompson B, Hammer PE, Hill DS, Stafford J, Torkewitz N, Gaffney TD, Lam ST, Molnár I, Ligon JM (2003) 2,5-Dialkylresorcinol biosynthesis in *Pseudomonas aurantiaca*: novel head-to-head condensation of two fatty acid-derived precursors. J Bacteriol 185(3):860–869. <https://doi.org/10.1128/JB.185.3.860-869.2003>
38. Péchy-Tarr M, Borel N, Kupferschmied P, Turner V, Binggeli O, Radovanovic D, Maurhofer M, Keel C (2013) Control and host-dependent activation of insect toxin expression in a root-associated biocontrol pseudomonad. Environ Microbiol 15(3):736–750. <https://doi.org/10.1111/1462-2920.12050>
39. Péchy-Tarr M, Bruck DJ, Maurhofer M, Fischer E, Vogne C, Henkels MD, Donahue KM, Grunder J, Loper JE, Keel CJ (2008) Molecular analysis of a novel gene cluster encoding an insect toxin in plant-associated strains of *Pseudomonas fluorescens*. Environ Microbiol 10(9):2368–2386. <https://doi.org/10.1111/j.1462-2920.2008.01662.x>
40. Pintado A, Pérez-Martínez I, Aragón IM, Gutiérrez-Barranquero JA, de Vicente A, Cazorla FM, Ramos C (2021) The rhizobacterium *Pseudomonas alcaligenes* AVO110 induces the expression of biofilm-related genes in response to *Rosellinia necatrix* exudates. Microorganisms 9:1388. <https://doi.org/10.3390/microorganisms9071388>
41. Pliego C, Cazorla FM, González-Sánchez MA, Pérez-Jiménez RM, de Vicente A, Ramos C (2007) Selection for biocontrol bacteria antagonistic toward *Rosellinia necatrix* by enrichment of competitive avocado root tip colonizers. Res Microbiol 158:463–470. <https://doi.org/10.1016/j.resmic.2007.02.011>
42. Quandt J, Hynes MF (1993) Versatile suicide vectors which allow direct selection for gene replacement in gram-negative bacteria. Gene 15:15–21
43. Raaijmakers JM, Mazzola M (2012) Diversity and natural functions of antibiotics produced by beneficial and plant pathogenic bacteria. Annu Rev Phytopathol 50:403–424. <https://doi.org/10.1146/annurev-phyto-081211-172908>
44. Rochat L, Péchy-Tarr M, Baehler E, Maurhofer M, Keel C (2010) Combination of fluorescent reporters for simultaneous monitoring of root colonization and antifungal gene expression by a biocontrol pseudomonad on cereals with flow cytometry. Mol Plant-Microbe Interact 23:949–961. <https://doi.org/10.1094/MPMI-23-7-0949>

45. Ruffner B, Péchy-Tarr M, Höfte M, Bloemberg G, Grunder J, Keel C, Maurhofer M (2015) Evolutionary patchwork of an insecticidal toxin shared between plant-associated pseudomonads and the insect pathogens *Photorhabdus* and *Xenorhabdus*. BMC Genomics 16:609. <https://doi.org/10.1186/s12864-015-1763-2>
46. Saraf M, Rajkumar S, Saha T (2011) Perspectives of PGPR in agri-ecosystems. In: Maheshwari DK (ed) Bacteria in agrobiology: crop ecosystems. Springer, Berlin, Heidelberg, Germany, pp 361–385. https://doi.org/10.1007/978-3-642-18357-7_13
47. Sokal RR, Rohlf FJ (1986) Introducción a la Bioestadística, Barcelona. Dover Publications. https://www.reverte.com/libro/introduccion-a-la-bioestadistica_91504/
48. Taylor RG, Walker DC, McInnes RR (1993) *E. coli* host strains significantly affect the quality of small scale plasmid DNA preparations used for sequencing. Nucleic Acids Res 21:1677–1678
49. Tsai CJ-Y, Loh JMS, Proft T (2016) *Galleria mellonella* infection models for the study of bacterial disease and for antimicrobial drug testing. Virulence 7(3):214–229. <http://dx.doi.org/10.1080/21505594.2015.1135289>
50. Wang J, Zhi X, Yu X, Xu H (2013) Synthesis and insecticidal activity of new deoxypodophyllotoxin based phenazine analogues against *Mythimna separata* walker. J Agric Food Chem 61(26):6336–6343. <https://doi.org/10.1021/j4011033>

Figures

Figure 1

Organization of the insect toxin cluster in *Pseudomonas protegens* Pf-5, based in Péchy-Tarr et al. (2008) information, and *Pseudomonas chlororaphis* PCL1606. Protein sizes are indicated as number of amino acid (Aa size) for *P. protegens* Pf-5 and *P. chlororaphis* PCL1606. Beside, reliability of comparison (E-value), percentage of unique reads (% Coverage), and percentage of nucleotides identical (% Identity) are indicates for *P. chlororaphis* PCL1606. The mutation point is indicated by black inverted triangle. Genes with transport function, outer membrane protein (OMP), toxin and regulation putative functions are also indicated

Figure 2

Mortality percentage of *Galleria mellonella* larvae, infected by injection of dose 3×10^5 cfu/mL and 24h of incubation at 25°C. Statistical analysis by ANOVA was performed using IBM SPSS 22 software (SPSS

Inc., Chicago, IL, United States) and error bars are represented

Figure 3

Galleria mellonella mortality assay, a) Percentage of mortality of *Galleria mellonella*, infected at dose 3×10^3 cfu/mL along eighty hours. Counting made at 0, 17, 24, 30, 40, 60 and 80-hours post-inoculation. b) Symptom's development of *Galleria mellonella* infected at dose 3×10^3 cfu/mL incubated at 25°C during 17, 24, 30, 40 and 60 hours. Buffer inoculation was used as negative control. Healthy larva present beige color and black larvae represent the complete melanization and larvae death soon after

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryInformation.pdf](#)