Genome-wide identification of the peptide transporter family in rice and analysis of the PTR expression modulation in two near-isogenic lines with different nitrogen use efficiency

Xinghai Yang (✉ yangxinghai514@163.com)

Rice Research Institute, Guangxi Academy of Agricultural Sciences <https://orcid.org/0000-0002-3476-2578>

Xiuzhong Xia

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Yu Zeng

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Baoxuan Nong

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Zongqiong Zhang

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Yanyan Wu

Biotechnology Research Institute, Guangxi Academy of Agricultural Sciences

Qinglan Tian

Biotechnology Research Institute, Guangxi Academy of Agricultural Sciences

Weiyong Zeng

Cash Crops Research Institute, Guangxi Academy of Agricultural Sciences

Ju Gao

Guangxi Crops Genetic Improvement and Biotechnology, Guangxi Academy of Agricultural Sciences

Weiyong Zhou

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Haifu Liang

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Danting Li

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Guofu Deng

Rice Research Institute, Guangxi Academy of Agricultural Sciences

Keywords: Rice, PTR family, Phylogenetic analysis, Expression profile, Gene regulatory network

Posted Date: March 31st, 2020

DOI: <https://doi.org/10.21203/rs.2.22543/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Plant Biology on May 6th, 2020. See the published version at <https://doi.org/10.1186/s12870-020-02419-y>.

Abstract

Background: Nitrogen (N) is a major nutrient element for crop growth. In plants, the members of the peptide transporter (PTR) gene family are involved in nitrate uptake and transport. Here, we identified PTR gene family in rice and analyzed their expression level in near-isogenic lines. **Results:** We identified 96, 85 and 78 PTR genes in Nipponbare, R498 and *Oryza glaberrima*, and the phylogenetic were similar in Asian cultivated rice and African cultivated rice. The number of PTR genes was higher in peanuts (125) and soybeans (127). The 521 PTR genes in rice, maize, sorghum, peanut, soybean and *Arabidopsis* could be classified into 4 groups, and their distribution was different between monocots and dicots. In Nipponbare genome, the 25 PTR genes were distributed in 5 segmental duplication regions on chromosome 1, 2, 3, 4, 5, 7, 8, 9, and 10. The PTR genes in rice have 0-11 introns and 1-12 exons, and 16 of them have the NPF (NRT1/PTR family) domain. The results of RNA-Seq showed that the number of differentially expressed genes (DEGs) between NIL15 and NIL19 at three stages were 928, 1467, and 1586, respectively. Under low N conditions, the number of differentially expressed PTR genes increased significantly. The RNA-Seq data was analyzed using WGCNA to predict the potential interaction between genes. We classified the genes with similar expression pattern into one module, and obtained 25 target modules. Among these modules, three modules may be involved in rice nitrogen uptake and utilization, especially the brown module, in which hub genes were annotated as protein kinase that may regulate rice N metabolism. **Conclusions:** In this study, we comprehensively analyzed the PTR gene family in rice. 96 PTR genes were identified in Nipponbare genome and 25 of them were located on five large segmental duplication regions. The Ka/Ks ratio indicated that many PTR genes had undergone positive selection. The RNA-seq results showed that many PTR genes were involved in riceNUE, and protein kinases might play an important role in this process. These results provide a fundamental basis to improve the rice NUE via molecular breeding.

Background

Rice is one of the most important food crops in the world. As the population continues to grow, the food security issues are becoming more and more prominent^[1]. The application of N fertilizer has been the main method to improve rice production, but the excessive use of N fertilizer also increases the production costs, and causes serious damage to environment. As the concept of “green super rice” has been proposed^[2], developing the rice varieties with more efficient use of N fertilizers has become the key to improving agricultural production while preserving the environment.

NRT1s play important roles in nitrate uptake and transport in rice^[3,4]. NRT1 family mainly includes low-affinity nitrate transporters and belongs to PTR family. Thus, NRT1/PTR family is also named as NPF.

The major NUE-related PTR genes have been identified in rice^[9]. *OsNRT1* is the first PTR gene identified in rice^[10] and is located at the same locus as *OsNPF8.9*^[11]; *OsNRT1* has two transcripts, *OsNRT1.1a* and *OsNRT1.1b*. *OsNRT1.1b* encodes a PTR protein with 6 transmembrane domains, and its overexpression can increase rice N uptake under high or low N conditions, while *OsNRT1.1a* only works under high N conditions^[12]. Based on the homology with *AtNRT1.1*, Plett et al.^[13] named the three homologous genes

in rice as *OsNRT1.1A*, *OsNRT1.1B*, and *OsNRT1.1C*. The overexpression of *OsNRT1.1A* can up-regulate the genes related to nitrate and ammonium transport^[14]. Hu et al.^[15] found that *LOC_Os10g40600* encoded a nitrate transporter NRT1.1B, which could affect the NUE of *indica* and *japonica* via regulating rice root microbial population and altering the rhizosphere microenvironment^[16]. *OsNPF2.2* can unload nitrate from the rice xylem and participate in nitrate transport from root to stem, thus affecting the growth and development of vascular system, or even the whole plant^[17]. Overexpression of *OsPTR6* increases the expression of ammonium transporter gene and activates glutamine synthetase, which effects can promote rice growth, but reduces NUE under high ammonium conditions^[18]. Fang et al.^[19] showed that altering the expression of *OsPTR9* affected NUE, plant growth and rice production. *OsNPF2.4* plays a role in NO₃-absorption, long-distance transport and redistribution; also, altering its expression indirectly affects the reuse of potassium in roots and stems. Hu et al.^[20] concluded that *OsNPF7.2* played a role in the intracellular distribution of nitrate in roots, and affected rice growth under high nitrate condition. Recently, Gao et al.^[22] used the difference in NUE between *indica* and *japonica* rice to clone a gene encoding NADH / NADPH-dependent nitrate reductase OsNR2, which interacts with the nitrate transporter OsNRT1.1B to promote nitrate absorption by indica. Tang et al.^[23] cloned a dual-affinity nitrate transporter gene *OsNPF6.1* and found that the transcription factor *OsNAC42* can activate *OsNPF6.1*, which then enhances N absorption and NUE in rice. In summary, these rice PTR genes may play an important role in NUE, and it's likely that there are still other PTR genes in rice genome involved in N metabolism pathway.

In the previous study, we identified a NUE-related QTL *qNUE6* in rice, and *LOC_Os06g15370* may be the ideal candidate gene^[5]. The annotation for *LOC_Os06g15370* is a peptide transporter, and we named this gene as *OsPTR10*. Here, we use the latest Nipponbare genome to identify PTR genes, and compare them with the PTR genes in R498, *Oryza glaberrima*, maize, sorghum, peanut, soybean and *Arabidopsis*. The main aims of this study are: (i) determine the number of PTR genes in rice; (ii) understand the evolutionary relationship of PTR genes in rice and other 5 plants; (iii) analyze the regulatory network of PTR genes in rice NUE.

Results

Identification of PTR gene in cultivated rice

Among the 27 *Oryza* species, Asian cultivated rice (*Oryza sativa* L.) and African cultivated rice (*Oryza glaberrima* Steud.) are the two species that have been domesticated and utilized by humans^[24]. Using the conserved region sequence of PTR gene family, 96 PTR genes were identified in Nipponbare (*Oryza sativa* L. ssp. *japonica*) (Table S2), 85 were identified in R498 (*Oryza sativa* L. ssp. *indica*) (Table S3), and 78 were identified in *Oryza glaberrima* (Table S4).

Phylogenetic analysis of PTR genes

Multi-sequence alignment and phylogenetic analysis were performed on the protein sequences of 96 rice PTR genes using MEGA. Based on the evolutionary relationship, the 96 PTR genes were classified into 5 groups, and group I to V contained 20, 15, 26, 4 and 31 genes, respectively. The group V contained the most genes, which could be further classified into four subgroups, Va (3), Vb (4), Vc (8), and Vd (16) (Fig. 1).

To understand the evolutionary relationship of PTR genes between different *Oryza* species or subspecies, phylogenetic analysis was carried out using the PTR genes from Nipponbare, R498 and *Oryza glaberrima*. These PTR genes could be classified into 8 groups: group I contained 73 genes, which was the most, whereas group II had only 11 genes (Fig. 2a). The distribution of PTR genes in the 8 groups were similar among the three *Oryza* species (Table S5). Nipponbare genome was used for the further analysis as shown below.

Leguminous plants can form a symbiotic relationship with N-fixing bacteria, which can convert N_2 in the air into NH_3 . Thus, we analyzed the leguminous plants peanuts and soybeans. We also analyzed the dicotyledonous plant *Arabidopsis*, and the important monocotyledonous plants maize, sorghum and rice. The plant genome sizes of these six species were 2.5 Gb [25], 1.1 Gb [26], 125 Mb [27], 2.3 Gb [28], 730 Mb [29] and 466 Mb [30], respectively. Although the maize genome was large, it only contained 66 PTR genes, whereas the legume genome contained many more PTR genes: 125 for peanuts and 117 for soybeans. This might be related to the need of N transport after N fixation in legumes.

The phylogenetic tree was constructed based on the protein sequences of 543 PTR genes from the six species (Table S6), and these genes could be classified into 4 groups, I, II, III and IV (Fig. 2b). The PTR gene distribution in rice, maize and sorghum were similar, and group I contained the most monocots PTR genes. Similarly, the PTR gene distribution in peanut, soybean and *Arabidopsis* were similar, and group II had the most dicots PTR genes.

The chromosome location and segmental duplication events of rice PTR genes

The 96 PTR genes were unevenly distributed on 12 chromosomes in rice. Chromosome 1 had the most PTR genes and chromosome 9 contained the least. The details are shown in Fig. 3. 59 PTR genes, accounting for 60.4% of the total PTR genes, formed 17 gene clusters. Except chromosome 7, 8, and 9, all other chromosomes contained PTR gene clusters, and the cluster in 37.79-37.83 Mb region of chromosome 1 was the biggest, containing 9 PTR genes.

The segmental duplication events were analyzed using the 96 PTR genes. The results showed that there were 5 large genomic duplication regions, which contained 25 PTR genes: (i) 37.76 Mb -38.75 Mb on chromosome 1 corresponding to 20.31 Mb -21.21 Mb on chromosome 5; (ii) 27.74 Mb -28.95 Mb on chromosome 2 corresponding to 29.11 Mb -30.42 Mb on chromosome 4; (iii) 0.15 Mb -0.21 Mb on chromosome 3 corresponding to 21.76 Mb - 21.90 Mb on chromosome 10; (iv) 33.56 Mb -33.92 Mb on chromosome 3 corresponding to 5.31 Mb -6.14 Mb on chromosome 7; (v) 19.49 Mb -19.744 Mb on chromosome 8 corresponding to 12.13 Mb -12.45 Mb on chromosome 9 (Table 1).

Whole genome duplications were detected using the synonymous mutation rate K_s . The rice genome experienced three genome duplication events (Fig. 4). The whole genome duplication event shared by gramineous plants occurred at ~96 million years ago^[31]. After that, another two independent genomic duplications events occurred in rice^[32,33]. The K_a/K_s ratio reflects the extent to which all nucleotide sequences of a gene are positively selected during differentiation. If K_a/K_s is greater than 1, the gene is positively selected; if K_a/K_s is equal to 1, the gene is neutrally selected; if K_a/K_s is less than 1, the gene undergoes purify selection. The K_a/K_s ratios of paralogous PTR gene pairs were 0.08-1.83, and 693 gene pairs were greater than 1, indicating that many PTR genes underwent positive selection (Table S7, Fig. 5).

Gene structure and motif composition of rice PTR genes

The information related to the evolution of the gene family can be obtained from gene structure analysis. Thus, we performed a gene structural analysis on the 96 PTR genes (Fig. 6a). In terms of the intron-exon composition, the PTR genes had 0-11 introns and 1-12 exons, and 10 genes contained only 1 exon. Usually, the structurally similar genes have closer evolutionary relationship.

Previous studies have shown that there are three highly conserved domains in most PTR genes^[34]. In this study, there were five types of conserved domains in rice PTR genes, which were MFS, RNase_H_like, zf-RV, RT_like superfamily, and PBD. MFS can be further classified into MFS superfamily, MFS_NPF, MFS_NPF1_2, MFS_spinster_like, MFS_NPF5, MFS_NPF7, and MFS_NPF4. By performing conserved gene analysis on PTR genes, we found 77 genes had MFS superfamily domain and 16 genes contained NPF domain (Fig. 6b).

The expression profiles of PTR genes in near isogenic lines

The expression profile of a gene is often related to its function. Previous studies have shown that the proteins encoded by PTR genes can transport nitrate, playing an important role in plant growth and development. Therefore, we used RNA-Seq to analyze the expression profiles of 96 PTR genes in the near-isogenic rice lines NIL15 and NIL19 under HN and LN conditions.

The qualities of transcriptome sequencing and sequence alignment results are shown in Table S8. The cDNA library of 18 samples had high sequencing quality and good genome coverage, which was suitable for further analysis. Based on the gene expression levels in different samples, we performed correlation analysis on commonly expressed genes and differentially expressed genes (DEG). The averaged correlation coefficient between biological replicates of the same sample was $r = 0.9357$ (Fig. S1), indicating good reproducibility and reliable experimental results.

We used the qRT-PCR method to validate the 18 expression genes identified from RNA-Seq. The results showed that the qRT-PCR expression patterns of the 18 DEGs were consistent with RNA-Seq analysis (Fig. S2), suggesting that the RNA-Seq results were reliable for further analysis.

Previous studies have shown that the differences between NIL15 and NIL19 are on chromosomes 6, 8, 9, and 10^[5]. These regions contain 6 PTR genes, which were *LOC_Os06g13200*, *LOC_Os06g13210*, *LOC_Os06g15370*, *LOC_Os06g21900*, *LOC_Os06g38294*, and *LOC_Os10g22560*. The transcriptome sequencing results showed that the number of DEGs between the two lines at three stages were 928, 1,467 and 1,586 (Fig. S3), respectively. After shifted to LN condition, the number of DEGs significantly increased first, and then stabilized. We also performed functional annotation on DEGs using the GO database (<http://geneontology.org/>), and the classification results are shown in Fig. 7. In the biological process category, metabolic process contained the most genes; in the cellular component category, there were more genes related to membrane or membrane components; in the molecular function category, transporter activity related genes showed up.

In living organisms, different genes cooperate together to perform biological functions, and the same action across different genes forms a pathway. We classified and annotated the DEGs using KEGG database (<https://www.kegg.jp/>), and the classification results are shown in Fig. 8. The main involved pathways were: metabolism, genetic information processing, environmental information processing, cellular processes, organismal systems and human diseases, in which the metabolism pathway contained the most genes. Also, N metabolism-related genes were included in energy metabolism group.

The expression profiles of 96 PTR genes showed in Fig. 9. There were only two differentially expressed PTR genes between the two rice lines at 0 d, which were *LOC_Os06g13210* and *LOC_Os10g02100*. However, at 3 d and 6 d, the number of differentially expressed PTR genes increased to 12 and 7 (Table S9), in which the *LOC_Os04g50950*(*OsPTR6*)^[18], *LOC_Os06g49250* (*OsPTR9*)^[19] and *LOC_Os11g12740* (*SP1*)^[21] have been reported to participate in N uptake and transport in rice.

The gene regulation network of NUE in rice

The expression profile data was analyzed using weighted gene co-expression network analysis (WGCNA) to predict the potential interaction between genes. The correlation coefficient of the gene expression levels was calculated and taken n-degree power, so that the numerical distribution of the correlation coefficient gradually followed scale-free distribution. 58,176 genes were screened, and the criteria were: (i) at least 50% of the samples are expressed the gene; (ii) remove the genes with the least variance change $\geq 25\%$. Finally, 20,282 genes were left for further analysis.

Then, we clustered the samples and found that the samples from 0 d, 3 d and 9 d could be well separated (Fig. S4a), indicating that the data quality was reliable enough for subsequent analysis. Then, a soft threshold was chosen to construct gene coexpression network (Fig. S4b,c). 20,282 genes were used for WGCNA analysis, and 400 genes were randomly selected from the gene set to draw the expression cluster heatmap (Fig. S5). We classified the genes with similar expression pattern into one module, and identified 25 modules. We analyze the gene features in each module and found the modules with biological significance (Fig. S6). Subsequently, GO and KEGG databases were used for functional enrichment analysis, and the blue, brown and turquoise modules were found to participate in N metabolism. The

blue module contained six NUE-related genes: *LOC_Os01g36720*^[35], *LOC_Os03g62200*^[36], *LOC_Os04g40410*^[37], *LOC_Os04g43070*^[38], *LOC_Os05g39240*^[39] and *LOC_Os06g49250*^[19]. The brown module contained *LOC_Os01g61510*^[36], *LOC_Os01g65000*^[36], *LOC_Os02g47090*^[40] and *LOC_Os04g50950*^[18]. The turquoise module contained *LOC_Os02g02190*^[41], *LOC_Os02g40710*^[42], *LOC_Os02g40730*^[43], *LOC_Os03g13274*^[10], *LOC_Os03g48180*^[44], *LOC_Os10g40600*^[16,45], *LOC_Os11g12740*^[21] and *LOC_Os12g44100*^[17].

We performed WGCNA on these three modular genes. There were 59 DEGs in the brown module, and *LOC_Os03g29410*, *LOC_Os02g14480*, *LOC_Os04g24220*, *LOC_Os11g39370*, *LOC_Os09g30120* were hub genes in the regulatory network (Table S10, Fig. 10a). The functions of the most connected genes *LOC_Os03g29410*, *LOC_Os04g24220*, *LOC_Os02g14480*, and *LOC_Os11g39370* are related to protein kinase. Based on GO annotation, the functions of these genes were related to activity (GO: 0016301), stress (GO:0006950), biotic stimulus (GO: 0009607), signal transduction (GO: 0007165), and metabolic process (GO: 0008152). There were 129 DEGs in the turquoise module, and *LOC_Os06g11990*, *LOC_Os10g22590*, *LOC_Os08g44360*, *LOC_Os07g45060*, *LOC_Os07g06680*, *LOC_Os05g08370*, *LOC_Os04g59330* and *LOC_Os02g51710* were hub gene (Table S11, Fig. 10b). Due to the small number of DEGs in the blue module, we were not able to identify the hub genes.

Discussion

In this study, we identified 96 PTR genes in Nipponbare genome, which was different from the 84 PTR genes identified by Zhao et al. ^[34]. The discrepancy was possibly due to: (i) with the development of sequencing technology, the Nipponbare genome database has been continuously edited and improved ^[46,47]; (ii) the methods used in the two studies are different. Zhao et al. used BLASTP and BLASTN to search for PTR genes based on the conserved amino acid sequence of PTR domain. Our study used HMMER3 to search for PTR genes based on the conserved DNA sequence of PTR gene family, followed by manual verification.

The 96 PTR genes in rice can be classified into 5 groups, and the number of each group was not same. In addition, *LOC_Os03g13250* and *LOC_Os03g13274* (*OsNRT1*) ^[10], *LOC_Os03g01290* and *LOC_Os10g40600* (*OsNRT1.1B*) ^[15], *LOC_Os07g21960* and *LOC_Os12g44100* (*OsPTR2*) ^[17], *LOC_Os04g50940* and *LOC_Os04g50950* (*OsPTR6*) ^[18], *LOC_Os04g36040* and *LOC_Os11g12740* (*SP1*) ^[21] were more closely related. Therefore, these 5 genes may participate in N uptake and transport in rice.

Among the Asian cultivated rice, *japonica* rice had more PTR genes than *indica* rice; also, the Asian cultivated rice had more PTR genes than African cultivated rice. Although Nipponbare, R498, and *Oryza glaberrima* contained different numbers of PTR genes, their distributions in the 8 groups were evolutionarily consistent. When comparing rice with maize and sorghum, rice had the smallest genome but contained the most PTR genes. Leguminous plants have rhizobium on the roots, which can convert inorganic N in the air into organic N. We found that peanuts and soybeans had more PTR genes, which

may be related to the need of N transport after N fixation. In the phylogenetic relationship, the distribution of PTR genes in monocots and dicots were significantly different, which may be because the two types of plants produced their own unique PTR genes during evolution ^[48].

Gene family can be formed by whole genome duplication or polyploidization, tandem duplication, and segmental duplication. In this study, we found that many PTR genes formed by tandem duplication were often closely arranged on the same chromosome. Previous studies have also shown that the genes in the same gene cluster have similar sequences and functions ^[49]. By using the PTR gene to analyze segmental duplication events, we found that there were five duplicated blocks in rice genome, which is consistent with the previous results ^[32]. Among the 96 PTR genes in rice, 25 were contained within the segmental duplication. Moreover, rice underwent three whole genome duplication events: the first whole genome duplication event shared by all gramineous plants, and the other two chromosomal fragments doubling events occurred independently on rice genome. This result is consistent with previous studies ^[32,33,50]. In addition, the PTR genes in the duplication region were clustered into the same group, with closer phylogenetic relationship. There were many PTR gene pairs with Ka/Ks values greater than 1, indicating that non-synonymous mutations led to functional changes in PTR gene, making it more suitable for the environmental change ^[51].

Gene structure is related its function. Previous studies have shown that there are three conserved motifs in the protein sequences of rice PTR genes ^[34]; and all of them are contained in MFS family, which has 12 transmembrane domains ^[52]. In this study, we found most PTR genes contained MFS family motifs, and the known rice NUE gene, *OsNRT1*^[10], *OsNRT1.1A*^[14], *OsNRT1.1b*^[12], *OsPTR2*^[17], *OsPTR6*^[18], *OsPTR9*^[19], *OsNPF7.2*^[20] and *SP1*^[21] all contains MFS family domain. Moreover, 16 out of 96 PTR genes have NPF domain. In plants, NPF proteins transport a variety of substrates: nitrates, peptides, amino acids, etc ^[11]. Due to the long intron of *LOC_Os11g18044*, the sequence length (10.7 kb) was greater than other PTR genes in rice; moreover, it contained a longer MFS family domain, suggesting that the function of this gene might be more complex.

Transcriptome analysis of the N15 and NIL19 revealed that most of the DEGs are associated with metabolism and transport activation, which is consistent with plant N metabolism pathways. There were only 2 PTR genes that were differentially expressed under HN condition, but 12 PTR genes changed expression level under LN condition. These results indicated that the PTR genes may play an important role in N metabolism in a cooperative way. Previous studies have shown that *LOC_Os06g49250* (*OsPTR9*)^[19], *LOC_Os04g50950* (*OsPTR6*)^[18], *LOC_Os11g12740* (*SP1*)^[21] affect rice NUE, and these genes were all differentially expressed under LN condition.

In the three modules that may be associated with rice NUE, each module had different numbers of genes involved in N metabolism, but they all contained the genes related to the uptake and utilization of ammonium and nitrate, indicating the synergistic expression of ammonium and nitrate genes. This result

suggested that N uptake, transport, assimilation and signal transduction involved complex gene regulatory networks in rice [9].

In the brown module, the functional annotation of hub genes were protein kinase. Previous studies have shown that some protein kinases are related to N metabolism in *Arabidopsis*. AtCIPK8, a calcineurin B-like (CBL)-interacting protein kinase (*CIPK*), was found to participate in early nitrate signaling [53]. CIPK and mitogen-activated kinase kinase kinase (MEKK) are putative regulatory proteins involved in the early nitrate signalling [54]. Ca^{2+} -sensor protein kinases (CPKs) are master regulators that orchestrate primary nitrate responses [57]. Furthermore, brassinosteroid (BR) signal kinase BSK3 regulates root elongation under limited N conditions [58]. A number of protein kinases have been identified in rice, mainly Calcium-dependent protein kinase, mitogen-activated protein kinase, stress-activated protein kinase. They play important roles in rice growth and development, biotic stress, abiotic stress, and expression regulation. Recently, researchers have found that multiple protein genes are related to nitrogen use efficiency in rice. OsCPK12 participates in the response signal pathway of low nitrogen stress, and overexpression of OsCPK12 enhances rice growth under low nitrogen conditions [59]. Hsieh et al. [60] found that the N-regulated genes, *Os02g0120100*, *Os02g0807000* and *Os06g0692600* etc., which encode transcription factors, protein kinases and protein phosphatases and may be involved in the regulation of early -N responses in rice roots. The serine/threonine/tyrosine (STY) protein kinase, ACTPK1, enhances ammonium uptake and use, and promotes growth of rice seedlings under sufficient ammonium [61]. A *calcium-dependent protein kinase* gene, *esl4*, which may function upstream of nitrogen-metabolism genes [62]. The overexpression of *Calcineurin B-like interacting protein kinase 2* gene, *OsCIPK2*, could increase nitrogen use efficiency in rice [63]. OsSAPK8 is a counterpart of AtOST1 in the activation of OsSLAC1, which is a nitrate-selective anion channel in rice [56]. *LOC_Os11g39370* is a BRASSINOSTROID INNOVATIVE 1-associated receiver kinase 1 precursor gene. Therefore, the brown module may be related to the regulation of N uptake and utilization in rice. The candidate gene *OsPTR10* identified in the previous study was also included in the brown module, and its function might be related to N metabolism.

Conclusions

In this study, we comprehensively analyzed the PTR gene family in rice. 96 PTR genes were characterized and classified into 5 main groups in Nippobare genome. The distribution of PTR genes between monocots and dicots was different, and legumes had a greater number of PTR genes. The rice genome experienced three genomic duplication events, and 25 rice PTR genes locates on 5 large segmental duplication regions. The K_a/K_s ratio indicated that many PTR genes had undergone positive selection. The structure and motif analysis revealed five types of conserved domains in rice PTR genes. The differentially expressed PTR genes increased significantly under LN conditions. Using WGCNA, we found three gene modules associated with NUE, especially the yellow module. These results provide information for a better understanding of the biological function for the PTR gene and will contribute to the genetic improvement of NUE in rice.

Methods

Genome sequence retrieval and PTR gene identification

The genome sequences of six species were downloaded from the following database: Nipponbare (MSU_7.0, <http://rice.plantbiology.msu.edu/index.shtml>), R498^[64] (CANU, <http://www.mgbkbase.org/R498/>), *Oryza glaberrima*^[65] (Oryza_glaberrima_V1, http://peanutgr.fafu.edu.cn/Genome_Browse.php), *Arabidopsis*^[27] (TAIR10, http://plants.ensembl.org/Arabidopsis_thaliana/Info/Index), *Arachis hypogaea*^[25] (PGR, http://peanutgr.fafu.edu.cn/Genome_Browse.php), *Zea mays*^[28] (B73_RefGen_v4, http://plants.ensembl.org/Zea_mays/Info/Index), *Glycine max*^[26] (Glycine_max_v2.1, http://plants.ensembl.org/Glycine_max/Info/Index), and *Sorghum bicolor*^[29] (Sorghum_bicolor_NCBIv3, http://plants.ensembl.org/Sorghum_bicolor/Info/Index). The PTR gene family HMM (hidden markov model) file was downloaded from the Pfam (<http://pfam.xfam.org/>). We did a genome-wide search on PTR genes using HMMER3, with e-value cut off = 0.001 and alignment sequence greater than 198 (50% of 395). The protein sequences were extracted and put into the three major databases: CDD (<https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>), SMART (<http://smart.embl.de/>), and PFAM (<http://pfam.xfam.org/>). Then, the resulting homologous family genes were manually verified for domain conservation. If conserved domain was detected in one of the three databases, it was considered as reliable family gene. Finally, the conserved domains and protein sequences were extracted.

Phylogenetic tree construction

ClustalW was used to perform multi-sequence alignment on the identified PTR genes. The NJ tree was constructed using MEGA7, with the bootstrap repeat number of 1000 and other parameters as default. The phylogenetic tree was colored by iTOL (<https://itol.embl.de/>).

Chromosomal distribution and gene duplication

All vs all protein protein alignment was performed using BLAST (Basic Local Alignment Search Tool), and the protein pair with E-value < 0.00001, identity > 90%, and minimum coverage of matching region on query sequence to subject sequence > 75% were extracted as homologous protein. The homologous protein file and the protein coordinate file were input into MCScanX (<http://chibba.pgml.uga.edu/mcscan2/>) for collinear region identification. Finally, the collinear results were presented using Circos (<http://circos.ca/>).

Analysis of gene structure and conserved motif

The PTR gene structure, mainly including UTR (untranslated region), introns and exons, was analyzed through GSDS (<http://gsds.cbi.pku.edu.cn/>). The motif sequence prediction was performed via MEME (<http://meme-suite.org/tools/meme>) and the motif map was presented using TBtools.

Plant materials and hydroponics

This study used the two near-isogenic lines: NIL15 (NIL-13B4, low NUE) and NIL19 (GH998, high NUE). The information about the two lines can be found in previous studies^[5]. The evenly germinated seeds were selected and cultured in 96-well cultivation instrument with 1.4 mM NH₄NO₃ (high nitrogen, HN) nutrient solution. The nutrient solution was prepared according to the method in Yoshida et al.^[66]. The nutrient solution was changed every 3 days, and the pH was adjusted to ~5.5 using MERS-NaOH. The seeds were cultured in a 28 °C light incubator (13 h light /11 h dark) until the 3-leaf stage. Then, at 3-leaf stage, the nutrient solution was replaced with 0.14 mM NH₄NO₃ (low nitrogen, LN) solution, and samples were taken at 0 d, 3 d, and 6 d, with 3 biologically replicates for each sample.

RNA extraction and sequencing

The RNA of each sample was extracted using Trizol method. The purity and integrity of each RNA sample were examined by agarose gel electrophoresis. The purity of DNA was checked by Nanodrop (Thermo Fisher Scientific, USA) (OD 260/280 ≈ 2.0). RNA concentration was quantified by Qubit (Thermo Fisher Scientific, USA) accurately, and the minimum concentration was 50 ng/ul.

mRNA was isolated from total RNA using the Oligo (dT) coated magnetic beads. Then, the mRNA was randomly fragmented into ~300 bp fragments by fragmentation buffer. With reverse transcriptase and a six-base random primer (random hexamers), the mRNA was subsequently reverse-transcribed into single-strand cDNA, which then formed into a stable double-stranded structure via two-strand synthesis. The constructed library was sequenced using Illumina NovaSeq 6000.

Sequencing quality control and alignment

After quality-control filtering, the clean reads were compared with the reference genome (<http://rice.plantbiology.msu.edu/index.shtml>) to obtain the mapped reads for subsequent analysis. Sequence alignment was performed using TopHat2^[67]. The mapping rate was usually higher than 70% when the reference genome was completely annotated and the experiment was free of contamination.

Quantitative RT-PCR

The expression level of 6 genes were measured by qRT-PCR. The primers were designed based on target gene sequence (Table S1), and *Actin3* was used as a reference^[68].

Differentially expressed gene analysis

According to the results of feature counts alignment to reference genome and the annotation file, the gene read counts for each sample were obtained, followed by FPKM (fragments per kilobase million) conversion to obtain standardized gene expression levels. DESeq2^[69] was used to perform statistical analysis on raw counts. The default thresholds were: p-adjust < 0.05 and |log2FC| ≥ 1, which yielded the

differentially expressed genes between two groups or two samples which expression folds greater than 2 and *P* value less than 0.05.

Weighted gene co-expression network analysis

After background correction and normalization of gene expression data, the nonstandard genes and less altered genes were filtered, so that the gene correlation intensity fell into scale-free distribution. After data pre-processing, the genes were classified: the genes with similar expression patterns were classified into one type, which was called a module. Then, the relationships among inter-module genes or intra-module genes were analyzed. Key modules were obtained by correlating with phenotypic data, and the module's hub gene was obtained by visualization network analysis.

Abbreviations

N: Nitrogen; PTR: Peptide transporter; NUE: Nitrogen use efficiency; NPF: NRT1/PTR family; DEG: Differentially expressed genes; NIL: Near-isogenic line; HMM: Hidden markov model; NJ: Neighbor-joining method; UTR: untranslated region; qRT-PCR: Quantitative RT-PCR; FPKM: Fragments per kilobase million; GO: Gene ontology; KEGG: Kyoto encyclopedia of genes and genomes; WGCNA: Weighted gene co-expression network analysis

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All data analysed in this study are included in the main manuscript and its additional files. The data supporting the findings of this work are available in the paper and its supplementary Information files. The RNA-seq data that support the findings of this study have been deposited to National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) with the accession code PRJNA573824 [<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA573824>].

Funding

This work was supported by the Guangxi Natural Science Foundation of China (2015GXNSFAA139054, 2018GXNSFAA138124), Guangxi Ministry of Science and Technology (AB16380117), National Natural Science Foundation of China (31860371), State Key Laboratory of Rice Biology (160105), Guangxi Academy of Agricultural Sciences (2015JZ16, 2019Z08 and QN-02). The funding bodies had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Authors' contributions

XHY and DTL conceived this study. BXN, ZQZ, YYW and QLT performed data analysis. WYZ (Weiyong Zeng), JG, WYZ (Weiyong Zhou) and HFL performed the experiments, XZX and YZ wrote this manuscript. XHY and GFD revised this manuscript. All authors read and approve the paper.

Acknowledgements

The authors thank to Jie Zhuang and Zhiqiong Yan for help.

Authors' Information

¹Rice Research Institute, Guangxi Academy of Agricultural Sciences, Nanning, China. ²Biotechnology Research Institute, Guangxi Academy of Agricultural Sciences, Nanning, China. ³Cash Crops Research Institute, Guangxi Academy of Agricultural Sciences, Nanning, China. ⁴Guangxi Crop Genetic Improvement and Biotechnology Laboratory, Guangxi Academy of Agricultural Sciences, Nanning, China.

References

1. Godfray HC, Beddington JR, Crute IR, Haddad L, Lawrence D, Muir JF, et al. Food Security: The Challenge of Feeding 9 Billion People. *Science*. 2010; 327: 812-818.
2. Zhang Q. Strategies for developing Green Super Rice. *Proc Natl Acad Sci USA*. 2007; 104: 11640-16402.
3. Li H, Hu B, Chu C. Nitrogen use efficiency in crops: lessons from Arabidopsis and rice. *J Exp Bot*. 2017; 68: 2477-2488.
4. Xuan W, Beeckman T, Xu G. Plant nitrogen nutrition: sensing and signaling. *Curr Opin Plant Biol*. 2017; 39: 57-65.
5. Yang XH, Xia XZ, Zhang ZQ, Nong BX, Zeng Y, Xiong FQ, et al. QTL Mapping by Whole Genome Re-sequencing and Analysis of Candidate Genes for Nitrogen Use Efficiency in Rice. *Front Plant Sci*. 2017; 8: 1634.
6. Tsay Y, Schroeder J, Feldmann KA, Crawford NM. The herbicide sensitivity gene CHL1 of Arabidopsis encodes a nitrate-inducible nitrate transporter. *Cell*. 1993; 72: 705-713.
7. Lran S, Muos S, Brachet C, Tillard P, Gojon A, Lacombe B. Arabidopsis NRT1.1 is a bidirectional transporter involved in root-to-shoot nitrate translocation. *Mol Plant*. 2013; 6: 1984-1987.

8. Liu KH, Yf T. Switching between the two action modes of the dual-affinity nitrate transporter CHL1 by phosphorylation. *EMBO J.* 2003; 22: 1005-1013.
9. Wang YY, Cheng YH, Chen KE, Tsay YF. Nitrate Transport, Signaling, and Use Efficiency. *Annu Rev Plant Biol.* 2018; 69: 85-122.
10. Lin CM, Koh S, Stacey G, Yu SM, Lin TY, Yf T. Cloning and functional characterization of a constitutively expressed nitrate transporter gene, OsNRT1, from rice. *Plant Physiol.* 2000; 122: 379-388.
11. Lran S, Varala K, Boyer JC, Chiurazzi M, Crawford N, Daniel-vedele F, et al. A unified nomenclature of NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family members in plants. *Trends Plant Sci.* 2014; 19: 5-9.
12. Fan X, Feng H, Tan Y, Xu Y, Miao Q, Xu G. A putative 6-transmembrane nitrate transporter OsNRT1.1b plays a key role in rice under low nitrogen. *J Integr Plant Biol.* 2016; 58: 590-599.
13. Plett D, Toubia J, Garnett T, Tester M, Kaiser BN, Baumann U. Dichotomy in the NRT gene families of dicots and grass species. *PLoS One.* 2010; 5: e15289.
14. Wang W, Hu B, Yuan D, Liu Y, Che R, Hu Y, et al. Expression of the Nitrate Transporter Gene OsNRT1.1A/OsNPF6.3 Confers High Yield and Early Maturation in Rice. *Plant Cell.* 2018; 30: 638-651.
15. Hu B, Wang W, Ou S, Tang J, Li H, Che R, et al. Variation in NRT1.1B contributes to nitrate-use divergence between rice subspecies. *Nat Genet.* 2015; 47: 834-838.
16. Zhang J, Liu YX, Zhang N, Hu B, Jin T, Xu H, et al. NRT1.1B is associated with root microbiota composition and nitrogen use in field-grown rice. *Nat Biotechnol.* 2019; 37: 676-684.
17. Li Y, Ouyang J, Wang YY, Hu R, Xia K, Duan J, et al. Disruption of the rice nitrate transporter OsNPF2.2 hinders root-to-shoot nitrate transport and vascular development. *Sci Rep.* 2015; 5: 9635.
18. Fan X, Xie D, Chen J, Lu H, Xu Y, Ma C, et al. Over-expression of OsPTR6 in rice increased plant growth at different nitrogen supplies but decreased nitrogen use efficiency at high ammonium supply. *Plant Sci.* 2014; 227: 1-11.
19. Fang Z, Xia K, Yang X, Grotemeyer MS, Meier S, Rentsch D, et al. Altered expression of the PTR/NRT1 homologue OsPTR9 affects nitrogen utilization efficiency, growth and grain yield in rice. *Plant Biotechnol J.* 2013; 11: 446-458.
20. Hu R, Qiu D, Chen Y, Miller AJ, Fan X, Pan X, et al. Knock-Down of a Tonoplast Localized Low-Affinity Nitrate Transporter OsNPF7.2 Affects Rice Growth under High Nitrate Supply. *Front Plant Sci.* 2016; 7: 1529.
21. Li S, Qian Q, Fu Z, Zeng D, Meng X, Kyojuka J, et al. Short panicle1 encodes a putative PTR family transporter and determines rice panicle size. *Plant J.* 2009; 58: 592-605 .
22. Gao Z, Wang Y, Chen G, Zhang A, Yang S, Shang L, et al. The indica nitrate reductase gene OsNR2 allele enhances rice yield potential and nitrogen use efficiency. *Nat Commun.* 2019; 10: 5207.

23. Tang W, Ye J, Yao X, Zhao P, Xuan W, Tian Y, et al. Genome-wide associated study identifies NAC42-activated nitrate transporter conferring high nitrogen use efficiency in rice. *Nat Commun.* 2019; 10: 5279.
24. Stein JC, Yu Y, Copetti D, Zwickl DJ, Zhang L, Zhang C, et al. Genomes of 13 domesticated and wild rice relatives highlight genetic conservation, turnover and innovation across the genus *Oryza*. *Nat Genet.* 2018; 50: 285-296.
25. Zhuang W, Chen H, Yang M, Wang J, Pandey MK, Zhang C, et al. The genome of cultivated peanut provides insight into legume karyotypes, polyploid evolution and crop domestication. *Nat Genet.* 2019; 51: 865-876.
26. Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, et al. Genome sequence of the palaeopolyploid soybean. *Nature.* 2010; 463: 178-183.
27. Initiative AG. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature.* 2000; 408: 796-815.
28. Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, et al. The B73 maize genome: complexity, diversity, and dynamics. *Science.* 2009; 326: 1112-1115.
29. Paterson AH, Bowers JE, Bruggmann R, Dubchak I, Grimwood J, Gundlach H, et al. The *Sorghum bicolor* genome and the diversification of grasses. *Nature.* 2009; 457: 551-556.
30. Goff S, Ricke D, Lan TH, Presting G, Wang R, Dunn M, et al. A draft sequence of the rice genome (*Oryza sativa* L. ssp. *japonica*). *Science.* 2002; 296: 92-100.
31. Wang X, Wang J, Jin D, Guo H, Lee TH, Liu T, et al. Genome Alignment Spanning Major Poaceae Lineages Reveals Heterogeneous Evolutionary Rates and Alters Inferred Dates for Key Evolutionary Events. *Mol Plant.* 2015; 8: 885-898.
32. Wang X, Shi X, Hao B, Ge S, Luo J. Duplication and DNA segmental loss in the rice genome: implications for diploidization. *New Phytol.* 2005; 165: 937-946.
33. Wang X, Shi X, Li Z, Zhu Q, Kong L, Tang W, et al. Statistical inference of chromosomal homology based on gene colinearity and applications to *Arabidopsis* and rice. *BMC Bioinformatics.* 2006; 7: 447.
34. Zhao X, Huang J, Yu H, Wang L, Xie W. Genomic survey, characterization and expression profile analysis of the peptide transporter family in rice (*Oryza sativa* L.). *BMC Plant Biol.* 2010; 10: 92.
35. Feng H, Yan M, Fan X, Li B, Shen Q, Miller AJ, et al. Spatial expression and regulation of rice high-affinity nitrate transporters by nitrogen and carbon status. *J Exp Bot.* 2011; 62: 2319-2332.
36. Suenaga A, Moriya K, Sonoda Y, Ikeda A, Von WN, Hayakawa T, et al. Constitutive expression of a novel-type ammonium transporter OsAMT2 in rice plants. *Plant Cell Physiol.* 2003; 44: 206-211.
37. Yan M, Fan X, Feng H, Miller AJ, Shen Q, Xu G. Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges. *Plant Cell Environ.* 2011; 34: 1360-1372.

38. Yang S, Hao D, Cong Y, Jin M, Su Y. The rice OsAMT1;1 is a proton-independent feedback regulated ammonium transporter. *Plant Cell Rep.* 2015; 34: 321-330.
39. Gaur VS, Singh US, Gupta AK, Kumar A. Influence of different nitrogen inputs on the members of ammonium transporter and glutamine synthetase genes in two rice genotypes having differential responsiveness to nitrogen. *Mol Biol Rep.* 2012; 39: 8035-8044.
40. Hu R, Qiu D, Chen Y, Miller AJ, Fan X, Pan X, et al. Knock-down of a tonoplast localized low-affinity nitrate transporter OsNPF7.2 affects rice growth under high nitrate supply. *Front Plant Sci.* 2016; 7: 1529.
41. Yan M, Fan X, Feng H, Miller AJ, Shen Q, Xu G. Rice OsNAR2.1 interacts with OsNRT2.1, OsNRT2.2 and OsNRT2.3a nitrate transporters to provide uptake over high and low concentration ranges. *Plant Cell Environ.* 2011; 34: 1360-1372.
42. Ferreira LM, Souza V, Tavares O, Zonta E, Santa-catarina C, Souza SR, et al. OsAMT1.3 expression alters rice ammonium uptake kinetics and root morphology. *Plant Biotechnol Rep.* 2015; 9: 221-229.
43. Sonoda Y, Ikeda A, Saiki S, Von WN, Yamaya T, Yamaguchi J. Distinct expression and function of three ammonium transporter genes (OsAMT1;1-1;3) in rice. *Plant Cell Physiol.* 2003; 44: 726-734.
44. Xia X, Fan X, Wei J, Feng H, Qu H, Xie D, et al. Rice nitrate transporter OsNPF2.4 functions in low-affinity acquisition and long-distance transport. *J Exp Bot.* 2015; 66: 317-331.
45. Hu B, Jiang Z, Wang W, Qiu Y, Zhang Z, Liu Y, et al. Nitrate-NRT1.1B-SPX4 cascade integrates nitrogen and phosphorus signalling networks in plants. *Nat Plants.* 2019; 5: 401-413.
46. Chen P, Jing X, Liao B, Zhu Y, Xu J, Liu R, et al. BioNano Genome Map Resource for *Oryza sativa* ssp. japonica and indica and Its Application in Rice Genome Sequence Correction and Gap Filling. *Mol Plant.* 2017; 10: 895-898.
47. Kawahara Y, De LM, Hamilton JP, Kanamori H, McCombie WR, Ouyang S, et al. Improvement of the *Oryza sativa* Nipponbare reference genome using next generation sequence and optical map data. *Rice (N Y).* 2013; 6: 4.
48. Cai C, Wang JY, Zhu YG, Shen QR, Li B, Tong YP, et al. Gene structure and expression of the high-affinity nitrate transport system in rice roots. *J Integr Plant Biol.* 2008; 50: 443-451.
49. Ramamoorthy R, Jiang SY, Kumar N, Venkatesh PN, Ramachandran S. A comprehensive transcriptional profiling of the WRKY gene family in rice under various abiotic and phytohormone treatments. *Plant Cell Physiol.* 2008; 49: 865-879.
50. Guo H, Jiao Y, Tan X, Wang X, Huang X, Jin H, et al. Gene duplication and genetic innovation in cereal genomes. *Genome Res.* 2019; 29: 261-269.
51. Zhang Z, Li J, Zhao XQ, Wang J, Wong GK, Yu J. KaKs_Calculator: calculating Ka and Ks through model selection and model averaging. *Genom Proteom Bioinf.* 2006; 4: 259-263.
52. Yan N. Structural Biology of the Major Facilitator Superfamily Transporters. *Annu Rev Biophys.* 2015; 44: 257-283.

53. Hu HC, Wang YY, Yf T. AtCIPK8, a CBL-interacting protein kinase, regulates the low-affinity phase of the primary nitrate response. *Plant J.* 2009; 57: 264-278.
54. Marchive C, Roudier F, Castaings L, Bréhaut V, Blondet E, Colot V, et al. Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. *Nat Commun.* 2013; 4: 1713.
55. Chen G, Wang L, Chen Q, Qi K, Yin H, Cao P, et al. PbrSLAH3 is a nitrate-selective anion channel which is modulated by calcium-dependent protein kinase 32 in pear. *BMC Plant Biol.* 2019; 19: 190.
56. Sun SJ, Qi GN, Gao QF, Wang HQ, Yao FY, Hussain J, et al. Protein kinase OsSAPK8 functions as an essential activator of S-type anion channel OsSLAC1, which is nitrate-selective in rice. *Planta.* 2016; 243: 489-500.
57. Liu KH, Niu Y, Konishi M, Wu Y, Du H, Sun CH, et al. Discovery of nitrate-CPK-NLP signalling in central nutrient-growth networks. *Nature.* 2017; 545: 311-316.
58. Jia Z, Giehl R, Meyer RC, Altmann T, Von WN. Natural variation of BSK3 tunes brassinosteroid signaling to regulate root foraging under low nitrogen. *Nat Commun.* 2019; 10: 2378.
59. Asano T, Wakayama M, Aoki N, Komatsu S, Ichikawa H, Hirochika H, et al. Overexpression of a calcium-dependent protein kinase gene enhances growth of rice under low-nitrogen conditions. *Plant Biotechnol.* 2010; 27: 369-373.
60. Hsieh PH, Kan CC, Wu HY, Yang HC, Hsieh MH. Early molecular events associated with nitrogen deficiency in rice seedling roots. *Sci Rep.* 2018; 8: 12207.
61. Beier MP, Obara M, Taniai A, Sawa Y, Ishizawa J, Yoshida H, et al. Lack of ACTPK1, an STY kinase, enhances ammonium uptake and use, and promotes growth of rice seedlings under sufficient external ammonium. *Plant J.* 2018; 93: 992-1006.
62. Xing Y, Guo S, Chen X, Du D, Liu M, Xiao Y, et al. Nitrogen Metabolism is Affected in the Nitrogen-Deficient Rice Mutant *esl4* with a Calcium-Dependent Protein Kinase Gene Mutation. *Plant Cell Physiol.* 2018; 59: 2512-2525.
63. Khan MU, Li P, Amjad H, Khan AQ, Arafat Y, Waqas M, et al. Exploring the Potential of Overexpressed OsCIPK2 Rice as a Nitrogen Utilization Efficient Crop and Analysis of Its Associated Rhizo-Compartmental Microbial Communities. *Int J Mol Sci.* 2019; 20: 3636.
64. Du H, Yu Y, Ma Y, Gao Q, Cao Y, Chen Z, et al. Sequencing and de novo assembly of a near complete indica rice genome. *Nat Commun.* 2017; 8: 15324.
65. Wang M, Yu Y, Haberer G, Marri PR, Fan C, Goicoechea JL, et al. The genome sequence of African rice (*Oryza glaberrima*) and evidence for independent domestication. *Nat Genet.* 2014; 46: 982-988.
66. Yoshida S, Forno D, Cock JH. Laboratory manual for physiological studies of rice. IRRI, Philippines. 1976; pp: 62.
67. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* 2013; 14: R36.

68. Sun H, Qian Q, Wu K, Luo J, Wang S, Zhang C, et al. Heterotrimeric G proteins regulate nitrogen-use efficiency in rice. *Nat Genet.* 2014; 46: 652-656.
69. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014; 15: 550.

Table

Due to technical limitations, Table 1 is only available for download from the Supplementary Files section.

Supplementary Files Legend

Additional file 1: Table S1. Primer sequences used for qRT-PCR.

Additional file 2: Table S2. Identification of 96 PTR genes in Nipponbare.

Additional file 3: Table S3. Identification of 85 PTR genes in R498.

Additional file 4: Table S4. Identification of 78 PTR genes in *Oryza glaberrima*.

Additional file 5: Table S5. The distribution of PTR gene in Nipponbare, R498 and *Oryza glaberrima*.

Additional file 6: Table S6. The distribution of PTR genes in six species.

Additional file 7: Table S7. Ka/Ks ratio of PTR gene pairs in rice.

Additional file 8: Table S8. Sequencing data and mapping to Nipponbare genome.

Additional file 9: Table S9. The differentially expressed PTR genes at three stages.

Additional file 10: Table S10. The differentially expressed genes in the brown module.

Additional file 11: Table S11. The differentially expressed genes in the turquoise module.

Additional file 12: Figure S1. The correlation coefficient of 18 samples based on the gene expression levels. The colored box shows correlation coefficient.

Additional file 13: Figure S2. The qRT-PCR was used to validate the 18 expression genes identified from RNA-Seq. X-axis represents the stage of 3 treatments, the purple column represents qRT-PCR results in NIL19, the green column represents qRT-PCR results in NIL15, the blue column represents RNA-Seq results in NIL19, and the yellow column represents RNA-Seq results in NIL15. Y-axis represents the relative level of gene expression, qRT-PCR uses $2^{-\Delta\Delta Ct}$ value, and RNA-Seq uses FPKM value. Error bars indicate standard deviations of three biological repetitions.

Additional file 14: Figure S3. The DEGs were screened by DESeq2. A1 vs A2 represents 928 DEGs between NIL15 and NIL19 at 0 d. B1 vs B2 represents 1,467 DEGs between NIL15 and NIL19 at 3 d. C1 vs C2

represents 1,586 DEGs between NIL15 and NIL19 at 6 d.

Additional file 15: Figure S4. Sample cluster dendrogram and soft-thresholding (β) values. **a** Sample cluster dendrogram and clinical trait heatmap of 18 samples based on their expression profile. **b** Analysis of scale-free fit index of each β value from 1 to 20. **c** Analysis of mean connectivity of each β value from 1 to 20. $\beta = 10$ was chosen for subsequent analyses as it has the biggest mean connectivity when the scale-free fit index is up to 0.895.

Additional file 16: Figure S5. The heat-map of the gene network of 400 randomly selected genes. The gene dendrogram and module assignment are also shown along the top. Color scale: yellow indicates low correlation, and red indicates high correlation.

Additional file 17: Figure S6. Identification of modules associated with NUE in two near-isogenic lines. Color scale: yellow indicates low correlation, and red indicates correlation.

Figures

Figure 2

Distribution of PTR genes in different species. a Neighbor-joining tree of 259 PTR genes in Nipponbare, R498 and *Oryza glaberrima*, these genes classified into 8 groups. The different-colored arcs indicate different groups. Group I, II, III, IV, V, VI, VII and VIII contain 29, 73, 17, 46, 15, 23, 11 and 45 genes. b Neighbor-joining tree of 543 PTR genes in rice, maize, sorghum, peanut, soybean and *Arabidopsis*, these genes classified into 4 groups. Group I, II, III and IV contain 140, 87, 135 and 181 genes. Group I contains the most PTR genes of monocots, and group II had the most PTR genes in dicots. The scale bar indicates the simple matching distance.

Figure 4

Whole genome duplications of rice using PTR genes. X-axis indicates the synonymous mutation rate K_s . Y-axis indicates PTR paralogous gene pairs. Every peak represents a genome duplication event. The farther away from the origin represents the earlier the genome duplication event occurs.

Figure 5

PTR genes underwent positive selection. X-axis indicates Ka/Ks ratio. Y-axis indicates PTR paralogous gene pairs. $Ka/Ks < 1$ represents positive selection; $Ka/Ks = 1$ represent neutral selection; $Ka/Ks > 1$, represents purifying selection.

Figure 6

Gene structure and conserved motifs in PTR genes in rice. a Exon-intron structure of pineapple PTR genes. Green boxes indicate untranslated regions; yellow boxes indicate exons; black lines indicate introns. b The motif composition of rice PTR proteins. The eleven type motifs are displayed in different colored boxes. The information for each motif is shown on the right. The length of gene/protein can be estimated using the scale at the bottom.

Figure 7

GO functional annotation for DEGs at three stages.

Figure 8

KEGG functional annotation for DEGs at three stages.

Figure 9

The expression profiles of 96 PTR genes at three stages in two near-isogenic lines. The cluster of PTR gene expression profiles are showed on the left; the gene name is showed on the right.

Figure 10

Gene co-expression networks. a 59 DEGs with the highest weight are in the brown module. b 129 differentially expressed PTR genes with the highest weight are in the turquoise module. The top most highly connected genes were signed. The red and blue circles represent DEGs.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Fig.S1.jpg](#)
- [TableS11.xlsx](#)
- [TableS6.xlsx](#)
- [TableS5.xlsx](#)
- [TableS8.xlsx](#)
- [TableS7.xlsx](#)
- [TableS9.xlsx](#)
- [TableS10.xlsx](#)
- [TableS3.xlsx](#)
- [Fig.S2.jpg](#)
- [TableS4.xlsx](#)
- [TableS1.xlsx](#)
- [TableS2.xlsx](#)

- [Fig.S6.jpg](#)
- [Fig.S3.jpg](#)
- [Fig.S5.jpg](#)
- [Fig.S4.jpg](#)
- [Table1.xlsx](#)