

Effects of Exposure to Incense Smoke on Airway Function: An in Vitro and in Vivo Study

Norio Yamamoto

Kyushu University

Keiko Kan-o (✉ hanamura@kokyu.med.kyushu-u.ac.jp)

Department of Endoscopic Diagnostics and Therapeutics, Kyushu University Hospital, Fukuoka, Japan

Miyoko Tatsuta

Kyushu University

Yumiko Ishii

Kyushu University

Ogawa Tomohiro

Kyushu University

Seiji Shinozaki

Kyushu University

Satoru Fukuyama

Kyushu University

Yoichi Nakanishi

Kyushu University

Koichiro Matsumoto

Kyushu University

Research Article

Keywords: Incense smoke (IS), transepithelial electronic resistance (TEER), airway hyperresponsiveness (AHR), airway dysfunction

Posted Date: January 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-131170/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on March 31st, 2021. See the published version at <https://doi.org/10.1038/s41598-021-86745-7>.

Abstract

Recent clinical studies have suggested that inhalation of incense smoke (IS) may result in impaired lung function and asthma. However, there is little experimental evidence to link IS with airway dysfunction. Using mouse and cell culture models, we evaluated the effects of IS exposure on airway function, such as airway hyperresponsiveness (AHR), expression of multiple epithelial tight junction (TJ)- and adherens junction-associated mRNAs and proteins in the lungs, and the barrier function of bronchial epithelial cells assessed by transepithelial electronic resistance (TEER). Exposure of BALB/c mice to IS increased AHR and inflammatory macrophage recruitment to BALF; reduced claudin-1, -2, -3, -7, -10b, -12, -15, and -18, occludin, zonula occludens-1 [ZO-1], and E-cadherin mRNA expression; and caused discontinuity of claudin-2 and ZO-1 protein immunostaining in lung tissue. IS extract dose-dependently decreased TEER and increased reactive oxygen species production in bronchial epithelial cell cultures. Treatment with N-acetyl-L-cysteine, but not glucocorticosteroids or long-acting β_2 -agonists, prevented the detrimental effects of IS. IS exposure can be problematic for respiratory health, as evidenced by AHR, increased recruitment of inflammatory macrophages and disruption of TJ proteins in the lung, and damage to epithelial barrier integrity. However, antioxidants may be useful for the treatment of IS-induced airway dysfunction.

Introduction

Burning of incense in temples and homes is a common religious and cultural practice in many Asian and Middle Eastern countries. Incense smoke (IS) contains particulate matter of varying sizes; gases such as carbon monoxide, nitrogen dioxide and sulfur dioxide; and volatile organic compounds such as benzene, aldehydes and polycyclic aromatic hydrocarbons¹. Compared with cigarette smoking, incense burning has been reported to generate larger quantities of particulate matter of $\leq 2.5 \mu\text{m}$ diameter and particles that remain in the air for hours^{2,3}. Increasing evidence suggests that ambient air pollution from IS can cause health problems, especially airway dysfunction. Recent studies have reported that indoor exposure to IS increases the risk of wheezing and asthma and is associated with impaired lung function in adolescents⁴⁻⁷. Despite a number of clinical studies, however, there is little experimental evidence that inhalation of IS causes airway dysfunction.

Bronchial epithelial cells form a barrier against a wide range of inhaled substances and are at the front line of mucosal innate immunity. The epithelial barrier function is maintained by apical junctional protein complexes, composed of apical tight junctions (TJs) and underlying adherens junctions (AJs), that form between neighbouring cells⁸. TJ-associated proteins, which include members of the occludin, claudin, and junctional adhesion molecule families, and AJ-associated proteins such as E-cadherin are the major constituents of junctional complexes in the bronchial epithelium⁹. Members of the zonula occludens (ZO) protein family act as scaffolds that link the intracellular domains of TJ and AJ-associated proteins with the cytoskeleton. There is increasing evidence that junctional protein expression and airway barrier function are reduced in patients with asthma¹⁰. We recently reported that exposure of human bronchial epithelial cells to cigarette smoke *in vitro* disrupted epithelial barrier function and simultaneously

downregulated the expression of multiple TJ and AJ-associated proteins¹¹. However, the effects of inhalation of harmful substances on multiple junctional protein expression in animal models is unknown.

In this study, we evaluated the effect of a single exposure of mice to IS on airway responsiveness (AHR), inflammation and multiple TJ and AJ-associated protein expression in the lung, and we additionally analysed the effect of exposure to IS extract (ISE) on epithelial barrier function in human bronchial epithelial cells in air-liquid interface (ALI) cultures. We also investigated whether treatment with glucocorticosteroids (GCSs), long-acting β_2 -agonists (LABAs), or the antioxidant N-acetyl-L-cysteine (NAC) could protect against IS-induced airway dysfunction *in vitro* and *in vivo*.

Methods

Mice

Six-week-old female BALB/c mice were purchased from Japan SLC (Shizuoka, Japan) and housed under specific pathogen-free conditions. The study protocol was approved by The Kyushu University Animal Care and Use Committee (A19-200-0). All experiments were performed in accordance with our institutional guidelines and carried out in compliance with the ARRIVE guidelines.

Mouse experiments

The most commonly used incense sticks in Japan were used for the study (0.4 g/stick, Nippon Kodo brand). Mice were randomly assigned to two or three groups depending on the experimental protocol: (i) unexposed, (ii) exposed to a high dose of IS (IS^{high}), or (iii) exposed to a low dose of IS (IS^{low}). The mouse groups were housed separately to avoid cross-exposure to IS. Mice were placed in 44-L chambers and exposed to IS generated by burning of 3.2 g (IS^{high}) or 1.6 g (IS^{low}) of incense sticks for 60 min. Incense sticks were burnt in a separate 8-L chamber connected by a tube to the 44-L exposure chamber, and IS was drawn into the exposure chamber with fresh air at 4 L/min. Mice from the unexposed group were maintained for the same period in fresh air. The ratio of the exposure chamber volume to the mean body weight of mice was designed to recapitulate the ratio of a standard living room volume to the standard body weight of humans.

In some experiments, mice received intraperitoneal injections of NAC (320 mg/kg, Sigma-Aldrich, St. Louis, MO) at 6 h before and 6 h after the 1-h exposure to IS. NAC was dissolved in phosphate buffer saline (PBS) and neutralised to pH 7.4 with NaOH.

ISE preparation

ISE for *in vitro* experiments was prepared by bubbling IS from 1.6 g of incense sticks (Nippon Kodo brand) through 20 mL of culture medium. After adjustment of the pH to 7.4, the ISE was sterile-filtered (0.22- μ m pore size, 33-mm diameter Millex GV; Merck Millipore, Billerica, MA). This solution was considered to be 100% ISE and was diluted in medium containing 10% foetal bovine serum before use.

ISE preparations were standardized based on absorbance at 320 nm and were freshly prepared for each experiment.

Measurement of AHR

Airway responsiveness was measured according to our previously described protocol¹². Briefly, mice were anesthetized with a mixture of ketamine and sodium pentobarbital intraperitoneally, and then their tracheas were cannulated via tracheostomy. Mice were ventilated mechanically (tidal volume, 0.3 ml; frequency, 120 breaths/min) after a paralytic agent was administered. The airway opening pressure was measured with a differential pressure transducer and recorded continuously. Stepwise increase in acetylcholine dose (1.25 to 20 mg/ml) were given with an ultrasonic nebulizer (NE-U07; OMRON Co., Kyoto, Japan) for one minute. The data were expressed as the provocative concentration 200 (PC₂₀₀), the concentration at which airway pressure was 200% of its baseline value. PC₂₀₀ was calculated by log-linear interpolation for individual animals as described previously¹². Values of PC₂₀₀ were expressed as log (100 × PC₂₀₀).

Collection of bronchoalveolar lavage fluid (BALF), flow cytometry analysis and enzyme-linked immunosorbent assay (ELISA)

After measurement of AHR, mice were euthanised by administration of pentobarbital. For collection of BALF, the lungs were gently lavaged twice with 1 mL of 0.9% saline *via* a tracheal cannula. BALF was centrifuged at 250 × *g* for 10 min and the supernatants were stored at - 80 °C until analysed. Total and differential cell counts in BALF were performed as described previously¹³.

For flow cytometry analysis, lungs not subjected to bronchoalveolar lavage were minced and single-cell suspensions were prepared. Macrophages and inflammatory macrophages were identified and enumerated by flow cytometry (Becton Dickinson, Franklin Lakes, NJ), as described in detail in the Supplementary Methods.

Levels of interleukin (IL)-6 (detection limit 7.8–500 pg/mL), IL-1 β (detection limit 15.6–1000 pg/mL), tumour necrosis factor- α (TNF- α ; detection limit 31.3–2000 pg/mL) in BALF were measured using specific ELISA kits (R&D Systems, Minneapolis, MN).

Fluorometric TUNEL assay

Apoptotic cells in lung tissue sections were detected using a DeadEnd™ Fluorometric TUNEL System (Promega, Madison, WI) according to the manufacturer's instructions and observed by confocal laser microscopy (LSM700; Zeiss, Jena, Germany). Negative and positive control slides were prepared by omitting the TdT enzyme from the nucleotide mix and by treating tissue sections with DNase I, respectively.

Quantitative reverse-transcription PCR (qRT-PCR)

Total RNA was extracted from mouse lungs, reverse-transcribed, and subjected to qPCR as described in the Supplementary Methods. Primer sequences are provided in Supplementary Table S1.

Immunofluorescence staining

Immunofluorescence staining were performed according to modified protocol as described previously¹⁴. Briefly, freshly isolated lungs were washed in PBS and then lung tissue was embedded using optimal cutting temperature compound and frozen. Four μm tissue sections on microscope slides were fixed with cold methanol for 10 min at $-20\text{ }^{\circ}\text{C}$ and then blocked with PBS containing 1% BSA for 30 min at room temperature. The tissue sections were incubated with each primary antibody prepared in PBS containing 1% BSA at $4\text{ }^{\circ}\text{C}$ overnight, followed by incubation with Alexa Fluor 488-conjugated goat anti-rabbit IgG antibody (diluted 1:500; Abcam, Cambridge, UK), Alexa Fluor 568-conjugated goat anti-rat IgG antibody (diluted 1:500; Abcam) and nuclear staining with 4',6-diamidino-2-phenylindole (DAPI). Images of the stained tissue sections were obtained with a confocal laser microscope (LSM700; Zeiss). The primary antibodies were as follows: rabbit anti-claudin-1 polyclonal antibody (diluted 1:200; Thermo Fisher Scientific, Waltham, MA); rabbit anti-claudin-2 polyclonal antibody (diluted 1:200; Abcam); rabbit anti-claudin-10 polyclonal antibody (diluted 1:200, Thermo Fisher Scientific); rabbit anti-claudin-15 polyclonal antibody (diluted 1:200; Thermo Fisher Scientific); rabbit anti-occludin polyclonal antibody (diluted 1:200; Thermo Fisher Scientific); rat anti-ZO-1 monoclonal antibody (diluted 1:200; Santa Cruz Biotechnology, Inc., Santa Cruz, CA); and rabbit anti-E-cadherin polyclonal antibody (diluted 1:200; Thermo Fisher Scientific).

Cell culture and treatment

Calu-3 cells, a sub-bronchial human epithelial cell line (HTB-55; American Type Culture Collection, Manassas, VA) were cultured in Dulbecco's modified Eagle's medium/F-12 (Thermo Fisher Scientific) supplemented with 10% foetal bovine serum and 1% penicillin-streptomycin and maintained at $37\text{ }^{\circ}\text{C}$ in a humidified atmosphere of 5% CO_2 in air. The methods for ALI cultures are described in the Supplementary Methods. Preliminary experiments demonstrated that transepithelial electronic resistance (TEER) of the cells cultured under ALI conditions reached a plateau at about day 8 post-seeding and then decreased to about half-maximal levels at day 21¹¹.

For the present study, cells were cultured for 8 days, and on day 9, 200 μL of control medium, ISE, and/or pharmacological reagents were added to the apical chamber and the cells were cultured for the indicated times. For experiments with GCSs, LABAs, or NAC, cells were pretreated with 10 nM fluticasone propionate (FP; Tocris, Minneapolis, MN), 10 nM salmeterol (SAL; Tocris), 10 nM budesonide (BUD; Sigma-Aldrich), 10 nM formoterol (FOR; Sigma-Aldrich), or 1 mM NAC (Sigma-Aldrich) by addition to the apical and basal chambers for 2 h before addition of ISE. Drugs were present throughout the remainder of the incubation. The concentration of GCSs used reflects the estimated therapeutic levels achieved in the human lung during inhalation¹⁵.

Measurement of TEER

Bronchial epithelial cell layer integrity was evaluated by TEER measurements using a Millicell-ERS 2 V-Ohmmeter (Millipore Co., Bedford, MA). Medium was added to the apical chamber 1 h prior to TEER measurement. The electrode was soaked in 70% ethanol and rinsed with culture medium prior to use. TEER was calculated by the following Eq. 1⁶: $TEER (\Omega cm^2) = (R_{sample} - R_{blank}) \times \text{effective membrane area } (cm^2)$.

Permeability assay

Calu-3 cell permeability was assessed using a fluorescein isothiocyanate (FITC)-dextran (4 kDa, Sigma-Aldrich) assay as described in the Supplementary Methods.

Viability assay

Cell viability was assessed according to our previously described protocol¹¹. After 24 h of ISE exposure, the medium in the apical chamber was removed and Calu-3 cells were washed with PBS, detached with trypsin-EDTA, and stained with 0.4% trypan blue solution. Non-viable cells were counted in a LUNA™ Automated Cell Counter (Logos Biosystems, Annandale, VA).

Reactive oxygen species (ROS) assay

Calu-3 cells were grown in monolayer culture to approximately 50% confluency, incubated with 1 mM NAC or vehicle for 2 h and then incubated with or without 50% ISE for 1 h. Positive control cells were exposed to 200 μ M of the ROS inducer pyocyanin (10 mM stock in anhydrous dimethylformamide; Enzo Life Sciences, Farmingdale, NY). Total ROS and superoxide production were evaluated using a ROS-ID® Total ROS/Superoxide detection kit (Enzo Life Sciences) according to the manufacturer's instructions. The cells were then visualised by fluorescence microscopy (BZ-X800; KEYENCE, Osaka, Japan).

Statistical analysis

Unless otherwise stated, data are expressed as the mean \pm standard error (SEM). The Mann–Whitney U-test was used to compare data between two groups and one- or two-way analysis of variance (ANOVA) followed by Tukey's multiple comparisons test were used to compare three or more datasets. Statistical analyses were conducted with Prism 8 software (GraphPad Software, San Diego, CA). Differences were considered statistically significant at $p < 0.05$.

Results

IS exposure induces airway hyperresponsiveness and inflammation in mouse lungs

To determine the effects of a single exposure period to IS on airway function, groups of mice were exposed to fresh air (unexposed) or high or low doses of IS for 1 h, and AHR, recruitment of inflammatory cells to BALF, and apoptosis of lung cells were assessed over the following 24 h. IS exposure increased AHR, as reflected by PC₂₀₀ values, in an acetylcholine dose-dependent manner (Fig. 1a). The number of

macrophages, but not neutrophils, lymphocytes or eosinophils, in BALF was also significantly increased in IS-exposed compared with unexposed mice (Fig. 1b), and flow cytometry analysis of lung-derived cells showed that IS exposure specifically increased the Ly-6G^{low}/Ly-6C^{high} population of inflammatory macrophages (Fig. 1c). However, IL-6, IL-1 β and TNF- α concentrations in BALF from both unexposed and IS^{high}-exposed mice were below the assay levels of detection (Supplementary Fig. S1). Analysis of apoptosis in lung sections using a fluorometric TUNEL assay revealed no increase in cell apoptosis in the lungs of IS-exposed compared with unexposed mice (Fig. 1d).

IS exposure decreases expression of TJ- and AJ-associated genes in mouse lungs

Next, we performed qRT-PCR analysis of lung tissues to determine the effects of IS inhalation on transcription of TJ-associated (claudins, occludin and ZO-1) and AJ-associated (E-cadherin) genes in mouse lungs. IS exposure significantly decreased the mRNA levels of claudin-1, claudin-2, claudin-10b and claudin-12 at 6 h post-exposure; claudin-3, claudin-7, claudin-18, E-cadherin and ZO-1 at 9 h post-exposure; and claudin-15 and occludin at 24 h post-exposure (Fig. 2). Claudin-2, claudin-3 and claudin-7 mRNA levels returned to unexposed control levels by 24 h (Fig. 2).

Expression of claudin-2 and ZO-1 protein in mouse lungs is disrupted by IS exposure

To determine whether disruption of TJ- and AJ-associated genes was also observed at the protein level, we performed immunofluorescence staining of lung sections from unexposed and IS^{high}-exposed mice. Confocal microscopy revealed predominant staining of claudin-1, claudin-10 and E-cadherin in the bronchial epithelium of unexposed mice, while claudin-2, claudin-15, occludin and ZO-1 proteins were expressed in both bronchial epithelial and alveolar cells (Fig. 3). Notably, lung sections from IS^{high} mice showed discontinuous or reduced staining of claudin-2 and ZO-1 at 24 h after IS exposure, indicating disruption of TJs (Fig. 3).

IS impairs bronchial epithelial barrier integrity

TEER measurements and permeability assays were performed to assess the effects of IS on epithelial barrier function. Calu-3 sub-bronchial epithelial cells were differentiated in ALI cultures for 9 days, and then incubated with 0%, 25%, 50%, or 75% ISE concentrations for an additional 24 h. Treatment with ISE caused a dose-dependent reduction in TEER that was maintained for 24 h post-exposure (Fig. 4a). In contrast to the significant reduction in TEER induced by 50% and 75% ISE (Fig. 4b), the permeability and viability of Calu-3 cells was significantly decreased at 24 h only by exposure to 75% ISE (Fig. 4c, d). Thus, treatment with 50% ISE impaired epithelial barrier integrity without influencing cell permeability or viability.

ISE-induced impairment of bronchial epithelial barrier integrity is not affected by GCS or LABA treatment

The benefits of inhaled GCSs and LABAs in asthma are widely recognized. We previously demonstrated that GCSs protect against epithelial barrier dysfunction induced by cigarette smoke extract¹¹. Therefore, we investigated the effects on ISE-induced TEER reduction in Calu-3 cells after pretreatment with vehicle, 10 nM GCSs (FP or BUD), and/or 10 nM LABAs (SAL or FOR) for 2 h before exposure to 50% ISE (plus GCS/LABA) for an additional 24 h. However, we found no effects of GCSs and LABAs, alone or in combination, on ISE-induced reduction in TEER (Fig. 5a–c).

The antioxidant NAC protects against ISE-induced impairment of bronchial epithelial barrier integrity

Immunofluorescence staining of Calu-3 cells confirmed that generation of total ROS, including superoxide, could be detected within 1 h of exposure to 50% ISE, and pretreatment of cells with 1 mM NAC for 2 h before ISE exposure diminished ROS generation (Fig. 6a). Similarly, NAC pretreatment strongly suppressed the decrease in TEER observed at 12 and 24 h after Calu-3 cells exposure to 50% ISE (Fig. 6b), suggesting that ROS production was directly involved in IS-induced disruption of epithelial barrier function. Treatment of ISE-exposed or unexposed cells with NAC alone did not affect cell viability (Supplementary Fig. S2).

NAC treatment prevents disruption of TJ-associated proteins in mouse lungs exposed to IS

To determine whether NAC can also protect lungs against IS exposure *in vivo*, we examined TJ-associated protein expression in lung tissues. Immunofluorescence staining of claudin-2 and ZO-1 was performed on lung sections collected at 24 h after exposure. As noted earlier, IS^{high} exposure caused discontinuous or attenuated staining of claudin-2 and ZO-1 in both bronchial epithelial and alveolar cells (Fig. 7). Although 24 h of NAC treatment alone did not affect the pattern or intensity of protein expression in unexposed mice, NAC treatment effectively prevented the IS^{high}-induced disruption of claudin-2 and ZO-1 immunostaining patterns (Fig. 7).

NAC treatment prevents IS-induced AHR and inflammation in mouse lungs

Finally, we examined the effects of NAC on IS-induced AHR and accumulation of inflammatory macrophages in BALF of mice at 24 h after IS exposure. NAC treatment of IS^{high} mice reversed the effect of IS on acetylcholine-induced AHR to the same levels detected in unexposed mice (Fig. 8a). Similarly,

NAC treatment completely abrogated both the increase in macrophage abundance and the proportion of inflammatory macrophages in BALF after exposure to IS^{high} (Fig. 8b, c).

Discussion

In the present study, we showed that a single exposure of mice to IS aggravated AHR, provoked an influx of inflammatory macrophages into the lungs, and disrupted the expression and location of TJ-associated proteins in bronchial epithelium. *In vitro* experiments confirmed that exposure of bronchial epithelial cells to ISE induced ROS production and dose-dependently reduced TEER through a mechanism resistant to clinically relevant concentrations of GCSs and/or LABAs. Finally, NAC treatment ameliorated IS-induced effects on AHR, macrophage recruitment and claudin-2 and ZO-1 expression in the murine airway and reversed the effect of ISE on TEER *in vitro*.

The results in the present study indicate that oxidative stress may be responsible for IS-induced respiratory complications by inducing AHR, disassembly of TJ proteins, and epithelial barrier integrity. Previous reports suggest that increased ROS may be associated with aggravated AHR and impaired epithelial barrier function. Using animal models, ROS has been shown to contribute directly to AHR *via* damage to oxidant-sensitive β -adrenergic receptors and increases in vagal tone, and this effect was enhanced when the epithelium was injured¹⁷⁻¹⁹. Studies of cigarette smoke exposure in bronchial epithelium suggest that ROS production after exposure may induce disassembly of TJ proteins and impair epithelial barrier function through epidermal growth factor receptor (EGFR)-extracellular signal-regulated kinase 1/2 signaling pathway^{20,21}. Furthermore, Heijink *et al.* reported that the protective effect of GCSs against cigarette smoke induced-epithelial barrier dysfunction was mediated by affecting EGFR-downstream target glycogen synthase kinase-3 β ²². We also showed that cigarette smoke-induced reduction in TEER (approximately 35% compared with untreated control) was significantly attenuated in Calu-3 cells treated with 10 nM BUD (15.6% TEER reduction compared with untreated control) or 10 nM FP (9.7% TEER reduction compared with untreated control)¹¹. However, in the present study, IS-induced reduction in TEER (approximately 35% compared with untreated control) was affected by neither BUD nor FP. Together with those observations, our present results suggest that IS-induced ROS may impair epithelial barrier integrity through a different signal transduction pathway activated by cigarette smoke exposure and IS-induced barrier dysfunction might be resistant to treatment with inhaled GCSs.

In this study, we showed that IS exposure disrupted claudin-2 and ZO-1 and simultaneously downregulated the gene expression of multiple TJ-associated proteins in the lung. Bronchial epithelium from patients with asthma has also been shown to exhibit an irregular ZO-1 staining pattern and reduced barrier function that is further compromised by exposure to the Th2 cytokines IL-4 and IL-13^{23,24}. Although claudin-2 is expressed in healthy human bronchiolar and alveolar cells, its precise function in lung physiology is unknown²⁵. A limitation of our study is that the consequences of claudin-2 and ZO-1 disruption by IS exposure remain unclear. Further studies are needed to clarify the impact of these changes on the maintenance of airway responsiveness and epithelial barrier function.

In conclusion, our results suggest that inhalation of IS might be harmful to respiratory health, as evidenced by AHR, increased recruitment of inflammatory macrophages and disruption of TJ-associated proteins in the lung, and damage to epithelial barrier integrity. However, treatment with NAC reversed many of the detrimental effects of IS exposure, suggesting that antioxidants may be beneficial for the treatment of IS-related airway dysfunction.

Declarations

Acknowledgements

The authors would like to acknowledge Mikiko Nakano, Kayoko Ono, and the staff of The Research Support Center, Research Center for Human Disease Modeling, Kyushu University Graduate School of Medical Sciences for technical assistance. We also thank Anne M. O'Rourke, PhD, from Edanz Group (<https://en-author-services.edanzgroup.com/ac>) for editing a draft of this manuscript.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author, KK, upon reasonable request.

Author Contributions

K.K., Y.I., S.F., Y.N. and K.M. conceived the study design and supervised the scientific work. N.Y., M.T., T.O. and S.S. performed the experiments. N.Y. and K.K. analysed the data. All authors contributed to and approved the final manuscript.

Funding

This work was partly funded by a Novartis Pharma research grant in 2016.

Competing interests

The authors declare no competing interests.

References

- 1 Lin, T. C., Krishnaswamy, G. & Chi, D. S. Incense smoke: clinical, structural and molecular effects on airway disease. *Clin Mol Allergy* **6**, 3, doi:10.1186/1476-7961-6-3 (2008).
- 2 Mannix, R. C., Nguyen, K. P., Tan, E. W., Ho, E. E. & Phalen, R. F. Physical characterization of incense aerosols. *Sci Total Environ* **193**, 149-158, doi:10.1016/s0048-9697(96)05343-0 (1996).
- 3 Zhou, R. *et al.* Higher cytotoxicity and genotoxicity of burning incense than cigarette. *Environmental Chemistry Letters* **13**, 465-471, doi:10.1007/s10311-015-0521-7 (2015).

- 4 Al-Rawas, O. A., Al-Maniri, A. A. & Al-Riyami, B. M. Home exposure to Arabian incense (bakhour) and asthma symptoms in children: a community survey in two regions in Oman. *BMC Pulm Med***9**, 23, doi:10.1186/1471-2466-9-23 (2009).
- 5 Wang, I. J., Tsai, C. H., Chen, C. H., Tung, K. Y. & Lee, Y. L. Glutathione S-transferase, incense burning and asthma in children. *Eur Respir J***37**, 1371-1377, doi:10.1183/09031936.00137210 (2011).
- 6 Norback, D. *et al.* Sources of indoor particulate matter (PM) and outdoor air pollution in China in relation to asthma, wheeze, rhinitis and eczema among pre-school children: Synergistic effects between antibiotics use and PM10 and second hand smoke. *Environ Int***125**, 252-260, doi:10.1016/j.envint.2019.01.036 (2019).
- 7 Chen, Y. C., Ho, W. C. & Yu, Y. H. Adolescent lung function associated with incense burning and other environmental exposures at home. *Indoor Air***27**, 746-752, doi:10.1111/ina.12355 (2017).
- 8 Tsukita, S., Tanaka, H. & Tamura, A. The Claudins: From Tight Junctions to Biological Systems. *Trends Biochem Sci***44**, 141-152, doi:10.1016/j.tibs.2018.09.008 (2019).
- 9 Rezaee, F. & Georas, S. N. Breaking barriers. New insights into airway epithelial barrier function in health and disease. *Am J Respir Cell Mol Biol***50**, 857-869, doi:10.1165/rcmb.2013-0541RT (2014).
- 10 Georas, S. N. & Rezaee, F. Epithelial barrier function: at the front line of asthma immunology and allergic airway inflammation. *J Allergy Clin Immunol***134**, 509-520, doi:10.1016/j.jaci.2014.05.049 (2014).
- 11 Tatsuta, M. *et al.* Effects of cigarette smoke on barrier function and tight junction proteins in the bronchial epithelium: protective role of cathelicidin LL-37. *Respir Res***20**, 251, doi:10.1186/s12931-019-1226-4 (2019).
- 12 Kibe, A. *et al.* Differential regulation by glucocorticoid of interleukin-13-induced eosinophilia, hyperresponsiveness, and goblet cell hyperplasia in mouse airways. *Am J Respir Crit Care Med***167**, 50-56, doi:10.1164/rccm.2110084 (2003).
- 13 Fukuyama, S. *et al.* A zinc chelator TPEN attenuates airway hyperresponsiveness and airway inflammation in mice in vivo. *Allergol Int***60**, 259-266, doi:10.2332/allergolint.09-OA-0167 (2011).
- 14 Barth, K. *et al.* P2X7R-dependent regulation of glycogen synthase kinase 3beta and claudin-18 in alveolar epithelial type I cells of mice lung. *Histochem Cell Biol***146**, 757-768, doi:10.1007/s00418-016-1499-3 (2016).
- 15 Kelly, M. M. *et al.* Corticosteroid-induced gene expression in allergen-challenged asthmatic subjects taking inhaled budesonide. *Br J Pharmacol***165**, 1737-1747, doi:10.1111/j.1476-5381.2011.01620.x (2012).

- 16 Sekiyama, A. *et al.* Glucocorticoids enhance airway epithelial barrier integrity. *Int Immunopharmacol***12**, 350-357, doi:10.1016/j.intimp.2011.12.006 (2012).
- 17 Adam, L., Bouvier, M. & Jones, T. L. Nitric oxide modulates beta(2)-adrenergic receptor palmitoylation and signaling. *J Biol Chem***274**, 26337-26343, doi:10.1074/jbc.274.37.26337 (1999).
- 18 Owen, S., Pearson, D., Suarez-Mendez, V., O'Driscoll, R. & Woodcock, A. Evidence of free-radical activity in asthma. *N Engl J Med***325**, 586-587, doi:10.1056/nejm199108223250816 (1991).
- 19 Hulsmann, A. R. *et al.* Oxidative epithelial damage produces hyperresponsiveness of human peripheral airways. *Am J Respir Crit Care Med***149**, 519-525, doi:10.1164/ajrccm.149.2.8306055 (1994).
- 20 Petecchia, L. *et al.* Bronchial airway epithelial cell damage following exposure to cigarette smoke includes disassembly of tight junction components mediated by the extracellular signal-regulated kinase 1/2 pathway. *Chest***135**, 1502-1512, doi:10.1378/chest.08-1780 (2009).
- 21 Khan, E. M., Lanir, R., Danielson, A. R. & Goldkorn, T. Epidermal growth factor receptor exposed to cigarette smoke is aberrantly activated and undergoes perinuclear trafficking. *FASEB J***22**, 910-917, doi:10.1096/fj.06-7729com (2008).
- 22 Heijink, I. H. *et al.* Budesonide and fluticasone propionate differentially affect the airway epithelial barrier. *Respir Res***17**, 2, doi:10.1186/s12931-015-0318-z (2016).
- 23 Xiao, C. *et al.* Defective epithelial barrier function in asthma. *J Allergy Clin Immunol***128**, 549-556 e541-512, doi:10.1016/j.jaci.2011.05.038 (2011).
- 24 Wawrzyniak, P. *et al.* Regulation of bronchial epithelial barrier integrity by type 2 cytokines and histone deacetylases in asthmatic patients. *J Allergy Clin Immunol***139**, 93-103, doi:10.1016/j.jaci.2016.03.050 (2017).
- 25 Zou, J. *et al.* Idiopathic pulmonary fibrosis is associated with tight junction protein alterations. *Biochim Biophys Acta Biomembr***1862**, 183205, doi:10.1016/j.bbamem.2020.183205 (2020).

Figures




Figure 1

Effects of IS exposure on airway responsiveness, inflammation, and cell apoptosis in mouse lungs. Mice were unexposed or exposed for 1 h to high or low doses of IS and evaluated 24 h later. a. Acetylcholine-induced airway hyperresponsiveness. b. Total and differential cell counts in BALF. c. Flow cytometry analysis of macrophage subsets in lung-tissue-derived cells. d. Fluorometric TUNEL assay of apoptotic (green) cells in lung sections. DNA was stained with propidium iodide (red). Scale bar, 20 μ m. Data are

presented as the mean \pm SEM (n=7–14 per group) and are representative of at least two independent experiments. *p<0.05, **p<0.01, ***p<0.001 by the Mann–Whitney U-test or one- or two-way ANOVA as appropriate. ACh, acetylcholine; IS, incense smoke; PC200, provocative concentration causing 200% response.




Figure 2

Effects of IS exposure on TJ- and AJ-associated gene expression in mouse lungs. qRT-PCR analyses of the indicated mRNAs in lung tissues was performed at up to 24 h after a 1-h exposure to fresh air (unexposed) or IS^{high}. mRNA levels were normalised to GAPDH mRNA levels. Data represent the mean \pm SEM (n=6-9 per group) and are representative of at least two independent experiments. *p<0.05, **p<0.01, by two-way ANOVA. IS, incense smoke.




Figure 3

IS exposure-induced changes in TJ and AJ-associated protein expression in mouse lungs. Confocal immunofluorescence microscopy of ZO-1 (red) and additional AJ and TJ-associated proteins (green) in lung sections was performed at 24 h after a 1-h exposure to fresh air (unexposed) or IShigh. DAPI staining of nuclei is shown in blue. Scale bar, 100 μm . Results are representative of at least two independent experiments. DAPI, 4',6-diamidino-2-phenylindole; IS, incense smoke.




Figure 4

Effects of ISE on airway epithelial barrier function. a. Time course of TEER in Calu-3 cells cultured under ALI conditions for the indicated times after exposure to vehicle (0%) or ISE at the indicated concentrations. b. TEER measured at 24 h after exposure of Calu-3 cells to 0–75% ISE for 24 h. c. FITC-dextran permeability assay of Calu-3 cells after exposure to 0–75% ISE for 24 h. d. Trypan blue viability assay of Calu-3 cells after exposure to 0–75% ISE for 24 h. Data represent the mean \pm SEM (n=6–10 per group) and are representative of at least two independent experiments. *p<0.05, **p<0.01, ***p<0.001 by one- or two-way ANOVA as appropriate. ISE, incense smoke extract; TEER, transepithelial electrical resistance.




Figure 5

Effects of GCS and/or LABA treatment on ISE-induced reduction in TEER in Calu-3 cells. Cells were cultured under ALI conditions and pretreated with vehicle, 10 nM GCS and/or 10 nM LABA for 2 h prior to addition of vehicle or 50% ISE for 24 h. a. Cells incubated with or without 10 nM of the GCSs FP and BUD. b. Cells incubated with or without 10 nM of the LABAs SAL and FOR. c. Cells incubated with the indicated combinations of 10 nM GCSs and 10 nM LABAs. Data represent the mean \pm SEM (n=5–10 per group)

and are representative of at least two independent experiments. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ by one-way ANOVA. BUD, budesonide; FP, fluticasone propionate; FOR, formoterol; ISE, incense smoke extract; SAL, salmeterol; TEER, transepithelial electrical resistance.




Figure 6

Effects of NAC on ISE-induced ROS generation and TEER reduction in Calu-3 cells. Cells were pretreated with 1 mM NAC or vehicle for 2 h and then incubated with vehicle or 50% ISE. a. Detection of total ROS

(green) or superoxide (red) in Calu-3 cells at 1 h after exposure to ISE. Scale bar, 50 μm . b. Analysis of TEER at the indicated times before and after exposure to ISE. Data represent the mean \pm SEM (n=8–15 per group) and are representative of at least two independent experiments. *p<0.05, **p<0.01, ***p<0.001 by one- or two-way ANOVA as appropriate. ISE, incense smoke extract; NAC, N-acetyl-L-cysteine; ROS, reactive oxygen species; TEER, transepithelial electrical resistance.




Figure 7

Effects of NAC on IS-induced changes in TJ-associated protein expression in mouse lungs. Confocal immunofluorescence microscopy of ZO-1 (red) and claudin-2 (green) proteins in lung sections prepared 24 h after a 1-h exposure to fresh air (unexposed) or IS^{high}. Mice were administered NAC (320 mg/kg) or PBS (vehicle) by intraperitoneal injection 6 h prior to and 6 h after the 1-h IS exposure. DAPI staining of nuclei is shown in blue. Scale bar, 100 μm. Results are representative of at least two independent experiments. DAPI, 4',6-diamidino-2-phenylindole; IS, incense smoke; NAC, N-acetyl-L-cysteine; PBS, phosphate-buffered saline.




Figure 8

Effects of NAC on IS-induced AHR and inflammation in mouse lungs. Mice were administered NAC (320 mg/kg) or vehicle intraperitoneally 6 h prior to and 6 h after a 1-h exposure to fresh air (unexposed) or IShigh and evaluated 24 h later. a. Acetylcholine-induced AHR. b. Total and differential cell counts in BALF. c. Flow cytometry analysis of macrophage subsets in the lungs. Data represent the mean \pm SEM (n=5–7 per group) and are representative of at least two independent experiments. *p<0.05, **p<0.01, ***p<0.001 by one- or two-way ANOVA as appropriate. ACh, acetylcholine, AHR, airway hyperresponsiveness; IS, incense smoke; NAC, N-acetyl-L-cysteine; PBS, phosphate-buffered saline; PC200, provocative concentration causing 200% response.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [YamamotoSupplementaryInformation.pdf](#)