

Alpha Thalassemia Genotypes in Kuwait

Adekunle Adekile (✉ adekile@hsc.edu.kw)

Kuwait University <https://orcid.org/0000-0002-3388-6659>

Jalaja Sukumaran

Kuwait University

Diana Thomas

Kuwait University

Thomas De Souza

Kuwait University

Mohammad Haider

Kuwait University

Research article

Keywords: Alpha thalassemia, Hemoglobin H disease, Kuwait

Posted Date: June 22nd, 2020

DOI: <https://doi.org/10.21203/rs.2.22547/v3>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on August 24th, 2020. See the published version at <https://doi.org/10.1186/s12881-020-01105-y>.

Abstract

Background: The frequency of alpha thalassemia trait is about 40% in the Kuwaiti population, but there has been no comprehensive study of the prevalent alleles. This is a report of the patients who were referred for molecular diagnosis over a 20-year period.

Methods: This is a retrospective study of the α -globin genotypes obtained in the Hemoglobin Research Laboratory of the Department of Pediatrics, Kuwait University from 1994 to 2015. Genotyping was by a combination of PCR, allele-specific oligonucleotide hybridization and reverse dot-blot hybridization (Vienna Lab Strip Assay).

Results: Four hundred samples were characterized and analyzed from individuals aged <1 month to 80 years, with a median of 6 years from 283 unrelated families.. Most (90.8%) were Kuwaiti nationals. The common genotype was homozygosity for the polyadenylation-1 mutation ($\alpha^{\text{PA-1}}\alpha/\alpha^{\text{PA-1}}\alpha$) in 33.3%, followed by heterozygosity ($\alpha\alpha/\alpha^{\text{PA-1}}\alpha$) for the same mutation in 32.3%. The PA-1 was therefore the most frequent allele (0.59). The frequency of the α^0 , $-\text{MED}$ was 0.017. Rare alleles that were found in very low frequencies included the α^0 ($-\text{FIL}$) in a Filipino child, Hb Constant Spring, Hb Adana, and Hb Icaria.

Conclusion: There is a wide variety of alpha thalassemia alleles among Kuwaitis, but the nondeletional PA-1 is by far the most common cause of moderate to severe HbH (β_4 tetramer), disease phenotype. The α^0 ($-\text{MED}$) allele is also encountered, which has implications for pre-marital counselling especially the possibility of having babies with alpha thalassemia major (Barts hydrops fetalis).

Background

Alpha thalassemia is one of the most widespread genetic diseases worldwide, with frequencies of the carrier state reaching up to 80 – 90% in some areas [1]. It is about 25 – 30% in tropical Africa [2], while in the Arabian Peninsula, the frequency varies from a low of close to zero in the desert areas to as high as 60% in the agricultural zones of Eastern Saudi Arabia [3, 4].

The normal individual has a complement of four α -globin genes ($\alpha\alpha/\alpha\alpha$), any one of which can be absent, producing varying degrees of alpha thalassemia [1]. In α^0 alleles, both genes on a chromosome are deleted ($-$), while in α^+ , a variable portion of the α_2 and/or α_1 gene is deleted, resulting in reduction in chain synthesis. The size of the deletion in α^0 alleles is variable and each is named for the part of the world where it is prevalent. Thus $-\text{MED}$, $-\text{FIL}$, $-\text{SEA}$ are found in the Mediterranean, Philippines and South East Asia, respectively [5]. The α^+ deletions are designated by their sizes and the most common is the $-\alpha^{-3.7\text{kb}}$ allele. One- or 2-gene deletions ($-\alpha/\alpha\alpha$, $-\alpha/-\alpha$, $-/\alpha\alpha$), produce alpha thal trait, which is usually associated with mild microcytic, hypochromic anemia. The loss of 3 genes ($-/-\alpha$) produces the classical haemoglobin H (HbH, β_4 tetramer) disease, which is characterized by moderate to severe anemia. The loss of all 4 genes ($-/-$) causes Hb Bart's (γ_4) hydrops fetalis, which is incompatible with life in the absence of early onset of chronic transfusion or stem cell transplantation [6].

Apart from deletions, there are a few point mutations that affect the transcriptional efficiency of the α globin genes and are designated $\alpha^T\alpha$ or $\alpha\alpha^T$ depending on whether the α_2 or α_1 gene is affected [7]. These nondeletional α -thalassemia alleles commonly affect the promoter, the initiation codon (ATG), splicing signals (GT/AG), the termination codon (TAA), and the polyadenylation signal (AATAAA) of the affected gene. The most common of these are the $\alpha^{IVSI(-5\text{ nt})}\alpha$ (in the Mediterranean), polyadenylation site mutations α_2^{AATAAG} , α_2^{AATGAA} and $\alpha_2^{\text{AATA}^-}$ (in the Mediterranean and Middle East) [7-11], termination codon mutations leading to elongated Hb variants, such as Hb Constant Spring (HbCS), Hb Icaria, Hb Koya Dora, Hb Seal Rock and Hb Pakse [12-14]. Heterozygotes for these alleles have a thalassemia minor phenotype, but homozygotes and compound heterozygotes often have HbH disease of varying severity and blood transfusion requirements.

Alpha thalassemia frequency is about 30 – 40% in Kuwait and HbH disease is mostly linked to the polyadenylation (polyA, $\alpha^{\text{PA}^{-1}}$) mutation [15, 16]. However, there has been no comprehensive study of the spectrum of alpha thalassemia alleles in the country. The present study was carried out to document the frequencies of moderate to severe alpha thalassemia phenotypes among patients referred for molecular diagnosis. In particular, it aimed to investigate the presence of the α^0 allele in the population, to determine the possibility of having a newborn with alpha thalassemia major (Hb Bart's hydrops fetalis) [1, 17]. This will enable more purposeful counselling of patients and empower physicians in the identification of patients who require further investigations. To this end, we have analyzed all cases of suspected α -thalassemia referred to the hemoglobin research laboratory in Kuwait University over a 20-year period.

Methods

The hemoglobin research laboratory in the Department of Pediatrics, Kuwait University received samples from the hematology clinics in Mubarak Al-Kabeer Hospital and some of the other government hospitals in Kuwait. The study was approved by the Human Research Ethics Committee of the Faculty of Medicine, Kuwait University and the Ministry of Health, Kuwait. Patients were suspected to have some form of α -thalassemia if they have persistent microcytic, hypochromic anemia in the absence of iron deficiency and with normal hemoglobin electrophoresis. Some patients had H inclusions on cresyl blue staining of the peripheral erythrocytes or H band on high performance liquid chromatography (HPLC). Quite often, but not always, there was a record of the clinical presentation and the complete blood count (CBC). This study is a retrospective study of all such samples analyzed from 1994 to 2005.

Blood was obtained by venipuncture into EDTA, complete blood count was done using ABX Pentra 120 cell counter (ABX France, Montpellier). The hemoglobin quantitation was determined using cation-exchange high performance liquid chromatography (HPLC) - Shimadzu LC-20AT, Shimadzu Corporation, Kyoto, Japan. DNA was isolated from peripheral leucocytes using phenol extraction. All the samples were screened for the α -thal-2 (-3.7 kb) deletion [18] and the α_2 -globin gene polyadenylation site (AATAAA@AATAAG) mutation using a PCR method and allele-specific oligonucleotide hybridization respectively as previously reported [18, 19]. All other samples that were not solved with these methods,

were subjected to reverse dot-blot hybridization, using the Vienna Lab StripAssay technique (Vienna Lab Diagnostics, Vienna, Austria).

The ethnic origins of the patients were documented by their nationalities. The different α -globin genotypes were identified and the allele frequencies were calculated. The mean Hb level and RBC indices were compared among patients with different genotypes. Clinical data were collected from the HbH disease patients, especially whether they were being followed in a hematology clinic and if they were on any medications or how often they were transfused with blood.

The disease severity was classified as silent carrier, α -thalassemia trait or HbH disease depending on the number, type and location of mutations detected, in addition to the clinical phenotype and presence of HbH inclusions or H band on HPLC [1, 17, 20]

Data are presented as means \pm SD or as percentages, as appropriate. Statistical differences between mean values among the major groups were tested using Student t test. All the descriptive analyses were performed with International Business Machines Corporation Statistical Package for Social Sciences (IBM SPSS Statistics, New York, USA), version 25 software. The criterion for statistical significance was $p < 0.05$.

Results

Four hundred blood samples were received with the diagnosis of suspected moderate to severe alpha thalassemia or HbH disease over the period of the study from 1994 to 2015. There were 183 females and 217 males, with ages ranging from 1 to 80 years and median of 6 years. Table 1 shows the nationalities of the patients in whom these were documented; most (90.8%) were Kuwaitis. The patients were from 283 unrelated families.

Table 2 shows the distribution of α -globin genotypes in the study. The most common (33.3%) was homozygosity for the α^{PA-1} (c.*94A>G) nondeletional α -thal allele, which gives a HbH disease phenotype, followed by heterozygosity for the same allele (32.3%). The compound heterozygosity of $-\alpha^{3.7}/\alpha^{PA-1}$ was the next common at 20.5%. None of the other frequencies reached 10%.

Three (0.8%) patients were classified as silent carriers, 154 (38.5%) were α -thalassemia trait, while 243 (60.8%) had HbH disease. Table 3 shows the Hb and RBC indices in the 3 groups of patients. When values were compared between the patients with α -thal trait and those with HbH disease, all the indices were significantly lower in the latter ($P < 0.05$), although the mean Hb was not lower than 10 g/dl in all groups, showing that α -thalassemia phenotypes are relatively mild in this population. None of the patients was on a chronic transfusion therapy, but many had received sporadic transfusions. The patients with the most severe phenotypes in the study were 2 siblings with $-\text{MED}/\alpha^{CD19}\alpha$ genotype. They both had poor growth parameters and baseline Hb of < 6 g/dl and are now transfused every 6 to 8 weeks. Interestingly, they have 2 other affected siblings (included in Tables 2 and 4), who do not require transfusions.

Table 4 shows the frequencies of the common individual alleles in the whole population and among the Kuwaiti individuals in the study. The PA-1 allele was the most frequent, at ~0.60 in both groups, followed by the $\alpha^{-3.7\text{ kb}}$ deletion at 0.153. Interestingly, the $\alpha^0\text{-MED}$ (0.019), $\alpha^{-4.2}$, $\alpha 2^{\text{cd}142\text{ (T}\delta\text{C)}}$ (Hb Constant Spring, c.427T>Cp.X143Gln), $\alpha 2^{\text{cd}59\text{ (G}\delta\text{A)}}$ (Hb Adana, c.179G>Ap.Gly60Asp), $\alpha^{\text{PA-2}}$ (c.*92A>G), $\alpha 2^{\text{cd}142\text{ (T}\delta\text{A)}}$ (Hb Icaria, c.427T>Ap.X143Lys) alleles were all found only among Kuwaitis. When the frequencies were computed based on the total number of families in the study, again, PA-1 allele was most frequent at 0.0552, followed by $\alpha^{-3.7\text{ kb}}$ deletion in 0.18. This was approximately the same when only Kuwaiti families were considered.

Discussion

The molecular diagnosis of α -thalassemia is not straight forward and requires resources that are not readily available in many centers. Moreover, it is economically unviable to provide this service to all suspected cases in many countries with a high prevalence. It is therefore necessary to have a good understanding of the prevalent genotypes and phenotypes, which will facilitate genetic counselling and the identification of patients who really deserve molecular diagnosis. This report is limited to patients who were suspected to have moderate to severe alpha thalassemia based on persistent microcytic, hypochromic anemia, no laboratory evidence of iron deficiency and normal hemoglobin electrophoresis or HPLC. It also included patients with H inclusion bodies and/or H band on HPLC.

Given the selection criteria for screening patients in this study, it is not surprising that most patients had the HbH disease phenotype and few had thalassemia trait or silent carrier status. A previous study from Kuwait had shown that most alpha thalassemia silent carriers had Hb levels >10 g/dl. Therefore, individuals with mild microcytosis and hypochromia, but with Hb >10 g/dl and no HbH inclusions or band, should not be considered for molecular diagnosis, but counselling should be offered after screening the family with CBC and HPLC, to rule out β -thalassemia. Resources can then be channeled to individuals with a more severe picture to identify those with HbH disease that need to be followed in the clinic.

The study has revealed the extent of the diversity among individuals with alpha thalassemia in Kuwait. As expected, the vast majority of the patients had the $\alpha^{\text{PA-1}}$ nondeletional allele, which has been reported as the commonest cause of HbH disease in Kuwait and in the region. The next most common allele was the $\alpha^{-3.7\text{ kb}}$ deletion, although it was usually present as a compound heterozygote with other alleles. Interestingly the third most frequent allele in this study was the $\alpha^0, \text{-MED}$, which was found in 13 patients, in whom it mostly presented a HbH disease phenotype. Of the latter, it was in compound heterozygosity with the $\alpha 2^{\text{cd}19\text{-G}}$ in 6 patients from 2 related families, with the with the $\alpha^{-3.7\text{ kb}}$ in 2 patients, the $\alpha^{\text{IVS1-5nt}}$ and the $\alpha^{\text{PA-1}}$ in 1 patient each.

The $\alpha^{\text{PA-1}}$ allele was first described from Eastern Saudi Arabia[10] and previous studies from Kuwait[2, 16] had highlighted its role in the etiology of HbH disease among our patients. It has also been reported as

the commonest cause of HbH disease in Saudi Arabia[21], Jordan, [22], Bahrain[23] and UAE[24]. The allele is therefore widespread in the Arabian Gulf, especially in the countries adjoining Saudi Arabia. It is uniformly associated with a moderate thalassemia intermedia phenotype with only occasional blood transfusion requirement, usually with intercurrent infection [16]. HbH disease is mainly a disease of childhood in Kuwait. Most patients are referred to the clinic for investigation of persistent microcytic, hypochromic anemia not responding to iron therapy. Indeed, the anemia tends to improve as the child stops rapid growth and enters puberty.

This study is the first to report the presence of the α^0 , $-\text{MED}$ allele among Kuwaiti patients. It presented as compound heterozygotes with other α -thal alleles; however the phenotype was mild with moderate anemia. The exceptions were 2 patients with the $-\text{MED}/\alpha^{\text{cd}19}\alpha$ genotype, with stunted growth and on regular transfusion because of severe anemia. The other rare alleles found in the study included the α^0 ($-\text{FIL}$) in a Filipino child, HbCS ($\alpha^2\text{Cd}142\text{TAA} \rightarrow \text{CAA}$), Hb Adana ($\alpha^{\text{cd}59\text{G/A}}$), $\alpha^{\text{PA-2}}$ (AATAAA®AATGAA), Hb Icaria ($\alpha^{\text{Cd}142\text{TAA} \rightarrow \text{AAA}}$).

Conclusions

This study has demonstrated the diversity of the alpha thalassemia alleles among Kuwaitis. While the deletional allele is quite widespread, the nondeletional $\alpha^{\text{PA-1}}$ is responsible for most of the moderate to severe HbH disease. Interestingly, the $\alpha^{0-\text{MED}}$ allele is also encountered, with implications for pre-marital genetic counselling to prevent the birth of babies with alpha thalassemia major (Bart's hydrops fetalis).

Declarations

Ethics Approval and Consent to Participate

The study was approved by the Human Research Ethics Committees of the Faculty of Medicine, Kuwait University and the Ministry of Health, Kuwait.

Consent for Publication

No identifying images or other personal or clinical details of participants are presented. Hence, consent for publication is not applicable to our manuscript. Moreover, there is no objection from Kuwait University Research Administration that funded the study.

Competing Interests

There are no other competing financial or non-financial interests.

Funding

The study was supported by grant No. MK 034 from Kuwait University Research Administration. The funder had no other role in the study.

Authors' Contributions

1. AA was the lead physician who planned the study and carried out the data analysis and wrote the first draft
2. MH was the head of the laboratory and carried out the molecular characterization. He contributed to the writing
3. JS, DT and MS all contributed to sample and data input/analysis

All authors have read and approved the manuscript

Availability of Data and Materials

The dataset used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

The authors thank the doctors who referred their patients for molecular diagnosis.

List Of Abbreviations

PCR	Polymerase chain reaction
PA	Polyadenylation
MED	Mediterranean
FIL	Filipino
SEA	Southeast Asia
Kb	Kilobases
Hb	Hemoglobin
CS	Constant Spring
CBC	Complete blood count
EDTA	Ethylenediaminetetraacetic acid
SD	Standard deviation
ANOVA	Analysis of variance
UAE	United Arab Emirates

References

1. Piel FB, Weatherall DJ: **The alpha-thalassemiias.** *The New England journal of medicine* 2014, **371**(20):1908-1916.
2. Adekile AD, Liu JC, Sulzer AJ, Huisman TH: **Frequency of the alpha-thalassemia-2 gene among Nigerian SS patients and its influence on malaria antibody titers.** *Hemoglobin* 1993, **17**(1):73-79.
3. White JM, Byrne M, Richards R, Buchanan T, Katsoulis E, Weerasingh K: **Red cell genetic abnormalities in Peninsular Arabs: sickle haemoglobin, G6PD deficiency, and alpha and beta thalassaemia.** *Journal of medical genetics* 1986, **23**(3):245-251.
4. White JM, Byrne M, Richards R, Buchanan T, Sheik-Yousouf I, White YS, Frost B: **Thalassaemia genes in Peninsular Arabs.** *British journal of haematology* 1985, **60**(2):269-278.
5. Fucharoen S, Viprakasit V: **Hb H disease: clinical course and disease modifiers.** *Hematology American Society of Hematology Education Program* 2009:26-34.
6. Vichinsky EP: **Clinical manifestations of alpha-thalassemia.** *Cold Spring Harbor perspectives in medicine* 2013, **3**(5):a011742.
7. Hartevelde CL, Losekoot M, Haak H, Heister GA, Giordano PC, Bernini LF: **A novel polyadenylation signal mutation in the alpha 2-globin gene causing alpha thalassaemia.** *British journal of haematology* 1994, **87**(1):139-143.
8. Hartevelde CL, Higgs DR: **Alpha-thalassaemia.** *Orphanet journal of rare diseases* 2010, **5**:13.
9. Hall GW, Higgs DR, Murphy P, Villegas A, de Miguel A: **A mutation in the polyadenylation signal of the alpha 2 globin gene (AATAAA→AATA-) as a cause of alpha thalassaemia in Asian indians.** *British journal of haematology* 1994, **88**(1):225-227.
10. Higgs DR, Goodbourn SE, Lamb J, Clegg JB, Weatherall DJ, Proudfoot NJ: **Alpha-thalassaemia caused by a polyadenylation signal mutation.** *Nature* 1983, **306**(5941):398-400.
11. Yuregir GT, Aksoy K, Curuk MA, Dikmen N, Fei YJ, Baysal E, Huisman TH: **Hb H disease in a Turkish family resulting from the interaction of a deletional alpha-thalassaemia-1 and a newly discovered poly A mutation.** *British journal of haematology* 1992, **80**(4):527-532.
12. Clegg JB, Weatherall DJ, Contopolou-Griva I, Caroutsos K, Pountouras P, Tsevrenis H: **Haemoglobin Icaria, a new chain-termination mutant with causes alpha thalassaemia.** *Nature* 1974, **251**(5472):245-247.
13. Clegg JB, Weatherall DJ, Milner PF: **Haemoglobin Constant Spring—a chain termination mutant?** *Nature* 1971, **234**(5328):337-340.
14. De Jong WW, Meera Khan P, Bernini LF: **Hemoglobin Koya Dora: high frequency of a chain termination mutant.** *American journal of human genetics* 1975, **27**(1):81-90.
15. Haider M, Adekile A: **Alpha-2-globin gene polyadenylation (AATAAA→AATAAG) mutation in hemoglobin H disease among Kuwaitis.** *Med Princ Pract* 2005, **14** Suppl 1:73-76.
16. Adekile AD, Azab AF, Al-Sharida SI, Al-Nafisi BA, Akbulut N, Marouf RA, Mustafa NY: **Clinical and Molecular Characteristics of Non-Transfusion-Dependent Thalassemia in Kuwait.** *Hemoglobin* 2015,

39(5):320-326.

17. Viprakasit V, Ekwattanakit S: **Clinical Classification, Screening and Diagnosis for Thalassemia.** *Hematology/oncology clinics of North America* 2018, **32(2)**:193-211.
18. Baysal E, Huisman T: **Detection of common deletional alpha-thalassemia-2 determinants by PCR.** *Am J Hematol* 1994, **46**:208-213.
19. Adekile AD, Haider MZ: **Morbidity, beta S haplotype and alpha-globin gene patterns among sickle cell anemia patients in Kuwait.** *Acta Haematol* 1996, **96(3)**:150-154.
20. Gilad O, Shemer OS, Dgany O, Krasnov T, Nevo M, Noy-Lotan S, Rabinowicz R, Amitai N, Ben-Dor S, Yaniv I *et al*: **Molecular diagnosis of alpha-thalassemia in a multiethnic population.** *European journal of haematology* 2017, **98(6)**:553-562.
21. Al-Awamy BH: **Thalassemia syndromes in Saudi Arabia. Meta-analysis of local studies.** *Saudi medical journal* 2000, **21(1)**:8-17.
22. Abu-Ghoush M: **Subtypes of alpha thalassemia diagnosed at a medical center in Jordan.** *TAF Prev Med Bull* 2008, **7(5)**:373- 376.
23. Jassim N, Al-Arrayed S, Gerard N, Al-Mukharraq H, Al-Ajami A, Ducrocoq R, Nagel R, R K: **Molecular Basis of α -Thalassemia in Bahrain** *Bahrain Medical Bulletin*, 2001, **23(1)**:3 - 7.
24. El-Kalla S, Baysal E: **alpha-thalassemia in the United Arab Emirates.** *Acta haematologica* 1998, **100(1)**:49-53.

Tables

Table 1

Nationalities of the Patients in the study

Nationality	n	%
Kuwaiti	317	90.8
Iraqi	10	2.9
Egyptian	6	1.7
Lebanese	4	1.1
Non Kuwaiti Bedouin	3	0.9
Syrian	3	0.9
Others	6	1.7
Total	349*	100.0

- These are all the patients in whom nationality was documented

Table 2

Distribution of Alpha Globin Genotypes of the Patients in the Study

Genotype	Number	%
$\alpha^{PA-1}\alpha/\alpha^{PA-1}\alpha$	133	33.3
$\alpha^{PA-1}\alpha/\alpha\alpha$	129	32.3
$-\alpha^{3.7}/\alpha^{PA-1}\alpha$	81	20.5
$-\alpha^{3.7}/-\alpha^{3.7}$	20	5.0
$--MED/\alpha^{CD19}\alpha$	6	1.5
$\alpha^{CD59}\alpha/\alpha\alpha$	4	1.0
$-\alpha^{3.7}/\alpha\alpha$	3	0.8
$--MED/\alpha\alpha$	3	0.8
$\alpha^{5nt}\alpha/\alpha^{5nt}\alpha$	3	0.8
$-\alpha^{3.7}/\alpha^{5nt}\alpha$	2	0.5
$-\alpha^{3.7}/--MED$	2	0.5
$\alpha\alpha/\alpha^{CD19}\alpha$	2	0.5
$\alpha^{5nt}\alpha/\alpha^{PA-1}\alpha$	1	0.3
$--MED/\alpha^{5nt}\alpha$	1	0.3
$--MED/\alpha^{PA-1}\alpha$	1	0.3
$\alpha^{CS}\alpha/\alpha^{CD142}\alpha$	1	0.3

$\alpha^{PA-2}\alpha/\alpha\alpha$	1	0.3
$\alpha^{5nt}\alpha/\alpha\alpha$	1	0.3
$-\alpha^{4.2}/-\alpha^{4.2}$	1	0.3
$\alpha^{CD19}\alpha/\alpha^{CD19}\alpha$	1	0.3
$\alpha^{PA1}\alpha/\alpha^{PA2}\alpha$	1	0.3
$\alpha^{CD59}\alpha/\alpha\alpha$	1	0.3
$\alpha^{CD59}\alpha/\alpha^{PA-1}\alpha$	1	0.3
$\alpha^{CD59}\alpha/\alpha^{5nt}\alpha$	1	0.3
$--SEA/-\alpha^{3.7}$	1	0.3
Total	400	100

Table 3
Red Blood Cell Indices in Patients

Group	Hb g/dl	MCV fl	MCH pg
Silent Carrier	10.4 ± 0.8	60.7 ± 9.5	19.8 ± 2.5
a-Thalassemia Trait	11.3 ± 2.0	64.4 ± 11.0	21.1 ± 3.5
HbH	10.0 ± 1.6	59.2 ± 8.7	18.3 ± 3.1
Total	10.5 ± 1.9	61.2 ± 10.0	19.4 ± 3.5

Table 4
Frequency of the Common Alpha Thalassemia Alleles in the Study

Allele	All population		Kuwaiti	
	N = 698	Frequency	N = 634	Frequency
AATAAA□AATAAG (PA-1)	412	0.590	386	0.608
-α-3.7 single gene deletion	107	0.153	91	0.143
--MED double gene deletion	12	0.017	12	0.019
α2 cd19 (-G)	10	0.0143	9	0.0141
α2 IVS-1 (-5nt)	8	0.011	5	0.008
α2 cd59 (GàA)	4	0.006	4	0.006
-α-4.2 single gene deletion	2	0.003	2	0.003
AATAAAàAATGAA (PA-2)	2	0.003	2	0.003
α2 cd 142 (TàC) Hb Constant Spring	2	0.003	2	0.003