

# Confirmation of the absence of local transmission and geographic assignment of imported *falciparum* malaria cases to China using microsatellite panel

**Yaobao Liu**

Jiangsu Institute of Parasitic Diseases

**Sofonias K Tessema**

University of California San Francisco

**Maxwell Murphy**

University of California San Francisco

**Sui Xu**

Jiangsu Institute of Parasitic Diseases

**Alanna Schwartz**

University of California San Francisco

**Weiming Wang**

Jiangsu Institute of Parasitic Diseases

**Yuanyuan Cao**

Jiangsu Institute of Parasitic Diseases

**Feng Lu**

Jiangsu Institute of Parasitic Diseases

**Jianxia Tang**

Jiangsu Institute of Parasitic Diseases

**Yaping Gu**

Jiangsu Institute of Parasitic Diseases

**Guoding Zhu**

Jiangsu Institute of Parasitic Diseases

**Huayun Zhou**

Jiangsu Institute of Parasitic Diseases

**Qi Gao**

Jiangsu Institute of Parasitic Diseases

**Rui Huang**

Soochow University

**Jun Cao** (✉ [caojuncn@hotmail.com](mailto:caojuncn@hotmail.com))

**Bryan Greenhouse**

University of California San Francisco

## Research

**Keywords:** Malaria, imported malaria, local transmission, Jiangsu, China, microsatellite genotyping, geographic assignment

**Posted Date:** February 4th, 2020

**DOI:** <https://doi.org/10.21203/rs.2.22548/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.  
[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Malaria Journal on July 13th, 2020. See the published version at <https://doi.org/10.1186/s12936-020-03316-3>.

# Abstract

## Background

Current methods to classify local and imported malaria infections depends primarily on patients travel history, which can have limited accuracy. Genotyping has been investigated as a complementary approach to track the spread of malaria and identify the origin of imported infections.

## Methods

An extended panel of 26 microsatellites (16 new microsatellites) for *Plasmodium falciparum* was evaluated in 602 imported infections from 26 sub-Saharan African countries to the Jiangsu province of People's Republic of China. The potential of the 26 microsatellite markers to assign imported parasites to their geographic origin was assessed using a Bayesian method with MCMC (Markov Chain Monte Carlo) as implemented in the program Smoothed and Continuous Assignments (SCAT) with a modification to incorporate haploid genotype data.

## Results

The newly designed microsatellites were polymorphic and are not in linkage disequilibrium with the existing microsatellites, supporting previous findings of high rate of recombination in sub-Saharan Africa. Consistent with epidemiology inferred from patients travel history, we found no evidence for local transmission; nearly all genetically related infections were identified in people who traveled to the same country near the same time. The smoothing assignment method assigned imported cases to their likely geographic origin with an accuracy (Angola: 59%; Nigeria: 51%; Equatorial Guinea: 40%) higher than would be achieved at random, reaching statistical significance for Angola and Equatorial Guinea.

## Conclusions

Routine genotyping is valuable for malaria case classification and program evaluation in an elimination setting. Method for assigning geographic origin of mammals based on genetic data were adapted for malaria and showed potential for identification of the origin of imported infections.

## Background

China has made enormous progress in controlling malaria and has an ambitious goal of malaria elimination by 2020[1]. There has been a decline in indigenous malaria cases, and thus most areas of the country are now in the prevention of reintroduction phase[1]. Meanwhile, with increasing global human movement, imported malaria has become a major challenge for elimination in many regions[2, 3]. In countries approaching elimination, imported cases account for the majority of reported cases and threaten resurgence of transmission in receptive areas due to the persistence of malaria vectors[4, 5].

Imported malaria has significantly increased in China over the past few years, causing 19,574 cases and 98 deaths (11.3 per 1000 cases) over a five-year period of 2011–2015[6]. The proportion of cases which were imported reached 98.2% (3022/3078) and 98.8% (3253/3294) of all cases in 2014 and 2015, respectively[7]. Most imported cases are caused by *P. falciparum* and originated in sub-Saharan Africa from labor related Chinese travelers[8, 9]. Recent studies showed a significant association between the number of imported cases and the amount of China's financial investment and malaria transmission intensity in sub-Saharan African countries[6]. These investments are predicted to expand in scope and volume and thus increase the risk of malaria importation[10].

Local transmission of *P. falciparum* has been interrupted in the Jiangsu Province of China since 1989[11]. However, this success is vulnerable to a possible resurgence of local transmission given the potential receptivity and increasing volume of importation. This vulnerability has been demonstrated by studies which showed agreement between the location of imported cases and the environmental suitability for malaria transmission[12]. Genetic characterization of imported cases could potentially give insight on the origin of parasites, when this is not clear based on travel history, as well as detect any evidence of local transmission.

Despite recent advances in genome sequencing of malaria parasites[13], microsatellite markers remain an important source of data to understand the genetic diversity and relatedness of parasite populations and understanding of transmission dynamics[14, 15]. Microsatellite markers are abundant in the *P. falciparum* genome[16], and tend to be more reliably variable in multiple populations than other markers. However only ten microsatellite loci are frequently used in most studies on the diversity and genetic relatedness of parasite populations[17, 18]. In this study, we designed an additional 16 microsatellite markers flanking the most commonly used 10 microsatellite markers[17]. The expanded panel was used to characterize 602 *P. falciparum* isolates collected from Chinese returnees from 26 sub-Saharan African countries to the Jiangsu province of China. We assessed the genetic, spatial and temporal relatedness of imported infections, and population assignment methods was used to identify the geographic origin of the imported cases.

## Methods

### Sample collection and DNA extraction

Venous blood samples were collected from symptomatic patients presented between September 2011 and March 2015 to a local hospital or center for disease control (CDC) in 13 prefectures of the Jiangsu Province of China. All participants gave written informed consent and microscopy positive samples were submitted to the Jiangsu Provincial Reference Laboratory of Malaria Diagnosis for confirmation by microscopy and species-specific nested polymerase chain reaction (PCR). Genomic DNA was extracted from 200  $\mu$ L of blood with the QIAamp blood DNA extraction kit (Qiagen, Crawley, U.K.) according to the manufacturer's instructions. Species-specific PCR was done as described previously[19, 20].

### Microsatellite genotyping

Ten extensively studied microsatellite markers (Polya, TA81, TA87, TA1, TA109, TA40, ARA2, PfG377, PfPk2, and TA60), referred as the original panel, were used to guide and select additional tandem repeats in the up and down stream flanking regions[17, 18]. After excluding microsatellite markers that showed inconsistent amplification and low heterozygosity, 16 additional microsatellite markers were designed and included in this analysis (see Additional file 1). The 16 newly identified markers flank the 10 microsatellite markers allowing evaluation of short-range haplotypes here termed “meta-loci”.

A two-round PCR protocol was used to amplify the 26 microsatellite loci, referred to as the extended microsatellite panel. The multiplex primary PCR was performed in 4 groups using 2 different PCR conditions. 1  $\mu$ L of the amplified product was then used as a template for each individual secondary PCR. Primer sequences, PCR conditions are provided in Additional file 2. The labelled PCR products were then diluted and sized by denaturing capillary electrophoresis on an ABI 3730XL analyzer with GeneScan™ 400HD ROX™ Size Standard (Thermo Fisher Scientific). The resulting electropherograms were analyzed using microSPAT[21], which identifies true peaks from artifacts using a probability-based filter. All results were verified manually. Only peaks with  $\geq 300$  relative fluorescence units were filtered and processed for allele calling.

## Multiplicity of Infection (MOI)

Infections containing more than one allele for at least two markers were classified as multi-clone infections. The Multiplicity of Infection (MOI) for a given sample was defined as the maximum number of alleles observed at any of the 26 loci investigated after removing the locus with the highest number, to decrease susceptibility to outliers due to PCR artifact. Analyses were carried out using the original panel, the extended panel, and the extended panel grouped as meta-loci. For geographic analysis, samples were grouped into five regions including East Africa (Kenya, South Sudan, Sudan, Tanzania and Uganda), Central Africa (Central African Republic, Chad, Democratic Republic of the Congo, Equatorial Guinea, Gabon and Republic of the Congo), Central-West Africa (Cameroon and Nigeria), West Africa (Benin, Côte d'Ivoire, Ghana, Guinea, Liberia, Mali, Sierra Leone and Togo) and Southern Africa (Angola, Madagascar, Mozambique, South Africa and Zambia).

## Population genetic indices

Population-level genetic diversity was characterized by expected heterozygosity ( $H_E$ ) and allelic richness ( $R_S$ ) normalized by the smallest sample size. These two parameters were used to assess the level of genetic polymorphism at each locus and determine the overall genetic diversity.  $H_E$  was calculated using the formula,  $H_E = [n/(n - 1)][1 - \sum p^2]$  where  $n$  is the number of genotyped samples and  $p$  is the frequency of each allele at a locus. The pairwise Jost's  $D$  [22] and  $G_{ST}$  [23] metrics were used to estimate the genetic differentiation between pairs of populations. Multi-locus linkage disequilibrium (LD) was measured by the standardized index of association ( $I_A^S$ ) using the web-based LIAN 3.5 software[24]. Under the null hypothesis of linkage equilibrium, the significance of the  $I_A^S$  estimates was assessed using 10,000 random permutations of the data.

# Population structure and genetic relatedness

Spatial and temporal population structure of the imported cases were inferred using the admixture model as implemented in Maverick v1.0[25]. Genetic distance between isolates is often determined based on alleles detected in monoclonal infections or taking into account the dominant allele identified for each locus. We employed an approach to determine a modified identity by state (IBS) metric[26] between each pairs of isolates. IBS was calculated based on the number of alleles shared between isolates, rather than of their haplotypes, allowing for the inclusion of mono- and poly-clonal samples. Pairwise IBS was determined as describe elsewhere[27, 28]. Using this distance metric ( $1 - 1/BS$ ), a neighbor-joining tree and multidimensional scaling were calculated.

## Population assignment

The potential of the 26 microsatellite markers to assign imported parasites to their geographic origin was assessed using a Bayesian method with MCMC (Markov Chain Monte Carlo) as implemented in the program Smoothed and Continuous Assignments (SCAT)[29] with a modification to incorporate haploid genotype data. Following validation of the assignment tool, we investigated the assignment accuracy of imported cases to their likely origins in sub-Saharan Africa. We conducted a leave-one-out cross-validation procedure in 243 monoclonal *P. falciparum* isolates imported from Angola (n = 78), Equatorial Guinea (n = 124) and Nigeria (n = 41), the only three countries with at least 40 single-clone infections in this study. In these analyses, each sample was treated as the sample whose location was unknown, whereas the other samples were assumed to have a known location (i.e. the centroid location of the respective countries). Smoothing and continuous assignment methods were used to assign imported cases to their geographic origin as described previously[29]. We performed three independent assignment runs, each starting with a different seed. For each run the first 2,000 iterations were discarded as burn-in, and we stored every 10th of the following 3,000 iterations, resulting in 300 putative locations of origin for each isolate. The range of allowable locations for the continuous assignment of individuals was specified by creating a boundary around sub-Saharan Africa. The assignment tests were also conducted for randomly generated alleles to create a null distribution. For the continuous assignment method, the median distance of the assigned isolates was calculated from the centroid location of the country. The assignment accuracy of the smoothing method was evaluated by calculating the average accuracy for each run (i.e. the proportions of correctly assigned individuals within each country).

## Results

### Epidemiologic profile of imported malaria in Jiangsu province, China

A total of 946 *P. falciparum* malaria cases was identified in Chinese travelers returned from 26 African countries to the Jiangsu Province over a 5-year period: 2011 (57), 2012 (109), 2013 (332), 2014 (342) and 2015 (106). Of the 946 imported cases, 627 laboratory-confirmed *P. falciparum* malaria cases were

available for genotyping using the extended microsatellite panel. 96% (602) were successfully genotyped at 22 or more of the 26 loci. The majority of the genotyped cases were imported from Equatorial Guinea (232), Angola (137) and Nigeria (70). The median age of patients was 44 years (range: 15–61 years), and males accounted for 98.8% of the imported cases. The imported cases were grouped into five regions based on the geographic proximity of the 26 African countries (East, Central, Central-West, West and Southern Africa) (Fig. 1).

The travelers who acquired *P. falciparum* spent an average of 351 days (IQR: 196–655 days) in the respective African countries. 46.8% of the travelers reported at least one symptomatic episode during their stay in Africa with an average of 2.5 episodes. Chinese travelers to southern Africa had the lowest number of reported episodes (mean = 1.2), while travelers to central Africa had the highest number of reported episodes (mean = 3.9). The median parasite density for all imported cases was 19,820 parasites per  $\mu\text{L}$  (Table 1). As might be expected due to development of partial immunity, individuals with reported malaria episodes during their stay in Africa had significantly lower parasite density upon presentation in China (median = 12,920 [IQR: 3476–48966] parasites per  $\mu\text{L}$ ) than those who did not (median = 37,600 [IQR: 6500–118765] parasites per  $\mu\text{L}$ ;  $p = 0.0003$ , Wilcoxon rank sum test) and spent more time in sub-Saharan African countries (401 vs 280 days,  $p < 0.001$ ).

Table 1

Summary of imported malaria cases from sub-Saharan Africa to the Jiangsu province, China

Regions	Year reported	No. samples		Duration of stay (median [IQR])	No. reported malaria episodes mean [sd]	Parasite density (x1000) median [IQR]
		Total	Genotyped			
East Africa	2013–2014	25	22	286 [182, 390]	1.9 [1.8]	8.6 [3.8, 34.6]
Central Africa	2011–2015	512	287	394 [194, 688]	3.9 [4.6]	13.9 [3.6, 80]
Central-West Africa	2013–2014	113	84	254 [138, 371]	1.9 [2.2]	21 [4.5, 77]
West Africa	2013–2014	63	51	232 [96, 348]	1.8 [2.6]	27 [5.7, 75.7]
Southern Africa	2013–2014	233	158	471 [268, 748]	1.2 [1.3]	30 [6.6, 100]
Total	2011–2015	946	602	351 [196, 655]	2.5 [3.5]	19.8 [4.6, 81.1]

## Description and characteristics of the extended microsatellite panel

The extended microsatellite panel includes 16 newly identified loci flanking 10 previously described microsatellites panel[17]. The 26 microsatellites were evaluated for their utility as genotyping markers. For single clone infections, short range multi-locus haplotypes were constructed from flanking microsatellite markers to create a “meta-locus” for each of the 10 genomic regions (Table S1). The chromosomal location and frequency distribution of alleles are shown in Fig. 2. All the newly identified microsatellite markers were neutral and polymorphic in this study population. The number of unique alleles per locus ranged from 6 (AS1 and AS12) to 21 (AS25) with a mean expected heterozygosity of 0.65 (range: 0.38–0.85) and with 13/16 new markers exhibiting an expected heterozygosity greater than 0.5 (see Additional file 3 and Additional file 4).

Multi-locus linkage disequilibrium (LD) was calculated using the standardized index of association ( $I_A^S$ ), excluding samples with polyclonal infections. Overall, low linkage disequilibrium was detected in pairwise comparisons between the 26 loci in the extended panel ( $I_A^S = 0.006$ , range: 0–0.05). Associations between loci within the same chromosome or meta-locus were not significantly different from those with markers located on a different chromosome or a meta-locus, consistent with the high expected rate of recombination in parasites populations of sub-Saharan Africa (see Additional file 5). A very low yet significant multi-locus LD was observed in all five regional populations of sub-Saharan Africa for the 26 loci ( $I_A^S = 0.004$ ,  $p < 0.001$ ); 10 loci ( $I_A^S = 0.003$ ,  $p = 0.015$ ) and meta-loci ( $I_A^S = 0.004$ ,  $p < 0.001$ ) (see Additional file 5). Due to the low pairwise linkage disequilibrium between the extended microsatellite loci, including between flanking microsatellites, the rest of the analyses were performed by considering each locus as an independent unit.

## **Moderate complexity and high genetic diversity of imported cases**

The successfully genotyped isolates had a mean MOI of 1.6 (range: 1–5), and 59% were single-clone infections (see Additional file 6). Mean MOI and proportions of multi-clone infections were not significantly different among infections imported from the five regions of Africa. There was no association between MOI and parasite density ( $r = -0.007$ ,  $p = 0.87$ ). Furthermore, there was no difference in the mean MOI of those who reported malaria during their stay in Africa (mean MOI = 1.64) and those who did not (mean MOI = 1.60,  $p = 0.71$ ).

Allelic richness ( $R_S$ ) and expected heterozygosity ( $H_E$ ) of the monoclonal samples were compared among the five populations using the extended and original microsatellite panels[17, 18]. Across all samples, a total of 375 and 194 unique alleles were identified from the 26 and 10 loci, respectively. There was no significant difference in allelic richness and expected heterozygosity between infections imported from the five regions of sub-Saharan Africa (Table 2).

Table 2

Genetic diversity and allelic richness of 602 imported monoclonal *P. falciparum* isolates from five regional sub-populations of Africa

Population	No. samples	Extended panel (n = 26 loci)		Original panel (n = 10 loci)	
		$R_S \pm SE^*$	$H_E \pm SE^\dagger$	$R_S \pm SE$	$H_E \pm SE$
East Africa	12	4.2 ± 0.30	0.68 ± 0.038	3.12 ± 0.26	0.82 ± 0.033
Central Africa	156	4.1 ± 0.30	0.72 ± 0.035	3.31 ± 0.28	0.84 ± 0.022
Central West Africa	48	3.8 ± 0.28	0.70 ± 0.038	2.77 ± 0.22	0.83 ± 0.028
West Africa	29	4.0 ± 0.28	0.71 ± 0.034	3.03 ± 0.25	0.83 ± 0.025
Southern Africa	91	3.9 ± 0.29	0.72 ± 0.033	2.91 ± 0.27	0.85 ± 0.022
Total	336	4.0 ± 0.13	0.71 ± 0.016	3.02 ± 0.11	0.83 ± 0.011

\*  $R_S$  - Allelic richness based on the minimum sample size of 12 individuals (East Africa) and  $^\dagger H_E$  is the mean expected heterozygosity

## Low to moderate geographical differentiation of imported infections

Low to moderate genetic differentiations were observed between the five parasite subpopulations for the extended ( $G_{ST} = 0.004-0.017$ , Jost's  $D = 0.019-0.075$ ) and original microsatellite panels ( $G_{ST} = 0.003-0.017$ , Jost's  $D = 0.037-0.155$ ) (Table 3). The least differentiation was observed between parasite populations from Central and Central-West Africa ( $G_{ST} = 0.004$ ), Central and West Africa ( $G_{ST} = 0.008$ ) and Central and Southern Africa ( $G_{ST} = 0.006$ ). Jost's  $D$  values among the five regions indicated a wide range of private alleles ranging from 1.9–7.5% in the 26 loci and 3.7–15.5% in the 10 loci.

Table 3

Estimates of genetic differentiation among parasite isolates imported from five regional subpopulations of sub-Saharan Africa (upper triangle is Jost's D and bottom is  $G_{ST}$ )

A. Extended panel (n = 26 loci)					
Regions	East Africa	Central Africa	Central-West Africa	West Africa	Southern Africa
East Africa	-	0.046	0.05	0.054*	0.075*
Central Africa	0.011	-	0.019	0.038	0.031
Central-West Africa	0.012	0.004	-	0.055	0.046
West Africa	0.017*	0.008	0.013	-	0.058
Southern Africa	0.017*	0.006	0.01	0.013	-
B. Original panel (n = 10 loci)					
East Africa	-	0.105	0.104	0.129*	0.155*
Central Africa	0.012	-	0.038	0.037	0.035
Central-West Africa	0.012	0.004	-	0.08	0.051
West Africa	0.016*	0.004	0.009	-	0.071
Southern Africa	0.017*	0.003	0.005	0.007	-

\*significant genetic differentiations ( $p < 0.05$ )

## Imported *P. falciparum* malaria cases were diverse and not structured

Pairwise comparison of genetic relatedness between all pairs of infections revealed that most infections were genetically unrelated (median genetic relatedness = 0.29 [IQR: 0.24–0.34]) (see Additional file 7). Similarly, admixture model-based analyses did not find spatial and temporal structure in the imported infections (see Additional file 7). However, against this background, 39 highly related pairs of imported infections (genetic relatedness  $\geq 0.6$ ) were detected. All related pairs had travel history consistent with importation from the same or a nearby country (Fig. 3). In average, highly related infections were imported and reported in China within 59 days [IQR: 22–202 days], compared to 214 days [IQR: 98–371 days] for unrelated infections (Figure A and B). Similarly, highly related infections tended to be imported from similar region (645 km [IQR: 0–1793 km]) than unrelated infections (1827 km [IQR: 645–2437km]) ( $p = 0.0003$ ) (Fig. 3C). No relationship between parasite relatedness and patient's residence in Jiangsu, China was observed (Fig. 3D), supporting the absence of local transmission.

# Geographic assignment of imported infections

While unsupervised clustering methods were unable to detect any significant structure in the genetic data, we also evaluated methods for assignment of geographic origin using training data. The smoothing assignment method assigned imported cases to their likely geographic origin with an accuracy of 59%, 51% and 40% for Angola, Nigeria and Equatorial Guinea, respectively. Despite the relatively low accuracy, the assignment accuracy was higher than would be achieved at random, reaching statistical significance for Angola and Equatorial Guinea (Fig. 4A). Using the continuous assignment method, the Euclidian distances between the estimated origin of each sample from the centroid location of the respective countries was determined with observed and randomly generated allele frequencies. Overall, 50% of the samples were assigned within 390 km of the centroid location of the origin country. The median distance was 391 km for Angola; 297 km for Equatorial Guinea and 1197 km for Nigeria, all significantly closer than the expected assignment distance from randomly generated genotypes (Fig. 4B). While the genetic data available in this study did not allow precise assignment, these findings highlight the potential of genetic assignment methods to provide more information on the geographic origin of imported infections than unsupervised clustering methods.

## Discussion

Genetic surveillance has the potential to provide useful information for malaria transmission epidemiology, particularly in the elimination context where detecting ongoing local transmission and identifying the origin of imported infections may be relevant. Using an extended panel of microsatellite markers, our results reveal importation of diverse parasite populations from different regions of sub-Saharan Africa to Jiangsu province of China. Reassuringly, and consistent with epidemiology inferred from the travel history of cases, we found no evidence for local transmission – nearly all closely related infections were identified in people who traveled to the same or neighboring country near the same time. While the absence of local transmission has been well documented in Jiangsu, these results highlight the potential utility of related methods for surveillance in areas where the lack of transmission is less clear, e.g. in areas which have only recently achieved elimination and/or where surveillance systems are not as robust. Although the receptivity of *P. falciparum* malaria is limited in Jiangsu province[11, 30], persistent importation could result in a reemergence of local transmission[31], highlighting the need to sustain a stringent surveillance efforts in receptive areas of China[32, 33].

Chinese travelers importing malaria spent an average of a year in the respective African countries, and of those 47% reported symptomatic malaria episodes during their stay. Despite the duration of stay and the risk of infections, genetic characterization of imported cases revealed a limited number of genotypes per imported infection despite the high diversity of genotypes in the local parasite populations, suggesting limited superinfection of the travelers. In contrast to reports characterizing the diversity of locally detected cases, the average number of distinct genotypes per infection did not differ among cases imported from different regions of sub-Saharan Africa[17, 34, 35].

Due to the physical proximity of the newly designed markers with the existing loci, we expected strong pairwise linkage disequilibrium between flanking microsatellites, however the pairwise LD was very low even among loci within 1000 bp on the same chromosome, suggesting very high rate of recombination in *P. falciparum* parasites in sub-Saharan Africa as described previously[36, 37]. This observation is further evidenced by high genetic diversity in imported infection, similar to the level reported from local parasite populations[34, 38, 39] and the lack of detectable geographic clusters. Furthermore, imported infections exhibited significant differentiation only in those imported from East vs. West and East vs. Southern Africa countries, consistent with previous reports[17, 37].

Multi-locus genotyping data, combined with statistical methods of inference, have been employed to address key questions of genetic identity and population membership, even when the overall level of genetic differentiation among populations is low[29, 40–42]. Such methods have been commonly used to assess temporal and spatial clustering of *P. falciparum* as well as *P. vivax* populations at a continental as well as village levels[15, 43–46]. However, these methods have not been standardized for the classification of imported vs. local malaria cases. In this study, it was relatively trivial to document the lack of closely related infections, given the use of highly diverse genetic markers which provided robust data even in polyclonal infections, the setting of a robust surveillance system capturing all or nearly all cases, and importation of parasites from diverse, relatively high transmission areas. However, while one often has the agency to choose appropriate genetic markers, the other conditions are not always met highlighting the need to develop more formal methods for distinguishing local versus imported cases.

Current methods rely on reported travel history to identify the origin of imported infections, but such data are often incomplete in areas where surveillance is weak. The strong surveillance system in China allowed us to evaluate the value of genetic data to accurately assign imported cases to their geographic origins. We were able to identify the origins of infections better than expected by chance, highlighting the potential of genetic data to identify and assign the geographic origins of imported infections. However, the assignment in this case was not accurate enough to be useful for surveillance. The lower assignment accuracy observed in this study could be improved by 1) specifically identifying a panel of geographically informative loci; 2) developing sensitive and high throughput laboratory methods to genotype these loci and 3) development of formalized analytical tools that can incorporate polyclonal infections to perform accurate classification of local, imported and introduced cases as well as identify the origin of imported infections.

Expanding the genetic epidemiology toolkit is a timely task to obtain a better insight into the spatial and temporal dynamics of transmission as well as for accurate classification of local and imported infections. Using an expanded microsatellite panel, we confirmed the absence of local transmission in the Jiangsu province and demonstrated the potential for genetic data to identify the origin of imported malaria infections. More formalized methods would allow surveillance systems in eliminating countries to track infections and develop targeted policies to limit the risk of re-introduction of *P. falciparum* malaria from eliminating countries.

# Conclusions

There is no local transmission due to imported falciparum malaria cases in the Jiangsu province of China. Method for assigning geographic origin of mammals based on genetic data showed potential for identification of the origin of imported malaria infections. Routine genotyping is valuable for malaria case classification and program evaluation in an elimination setting.

# Abbreviations

MCMC

Markov Chain Monte Carlo

SCAT

Smoothed and Continuous Assignments

CDC

center for disease control

PCR

polymerase chain reaction (PCR)

MOI

Multiplicity of Infection

$H_E$

heterozygosity

$R_S$

allelic richness

LD

linkage disequilibrium

$I_A^S$

standardized index of association

IBS

identity by state

# Declarations

## Ethics approval and consent to participate

This study was reviewed and approved by the Institutional Ethics Committee of Jiangsu Institute of Parasitic Diseases (JIPD). The written informed consent was obtained from all participants.

## Consent for publication

Not applicable

## Availability of data and materials

Not applicable

## Competing interests

The authors declare that they have no competing interests.

## Funding

This work was supported by the National Research and Development Plan of China (2016YFC1200500), the Natural Science Foundation of Jiangsu Province (BK20150001), the Jiangsu Provincial Department of Science and Technology (BE2016631 and BM2018020), the Jiangsu Provincial Project of Invigorating Health Care through Science, Technology and Education (ZDXKA2016016). The funders had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

## Authors' contributions

YBL, ST, JC and BG conceived and designed the study; YBL, SX, WMW, YYC, FL, JXT, YPG, GDZ and HYZ conducted case investigation and sample collection; YBL and MM performed the experiment. SF, YBL, MM, AS, QG, RH, JC and BG analyzed and interpreted the data. YBL and ST drafted the manuscript, JC and BG revised the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

We thank the staffs from the County Centers for Disease Control and Prevention in Prefectures of Jiangsu Province for their assistance with this study.

## References

1. Hu T, Liu YB, Zhang SS, Xia ZG, Zhou SS, Yan J, Cao J, Feng ZC: **Shrinking the malaria map in China: measuring the progress of the National Malaria Elimination Programme.** *Infect Dis Poverty* 2016, **5**:52.
2. Moonen B, Cohen JM, Snow RW, Slutsker L, Drakeley C, Smith DL, Abeyasinghe RR, Rodriguez MH, Maharaj R, Tanner M, Targett G: **Operational strategies to achieve and maintain malaria elimination.** *Lancet* 2010, **376**:1592-1603.
3. Tatem AJ, Jia P, Ordanovich D, Falkner M, Huang Z, Howes R, Hay SI, Gething PW, Smith DL: **The geography of imported malaria to non-endemic countries: a meta-analysis of nationally reported statistics.** *Lancet Infect Dis* 2017, **17**:98-107.
4. Liu Y, Sturrock HJW, Yang H, Gosling RD, Cao J: **The challenge of imported malaria to eliminating countries.** *Lancet Infect Dis* 2017, **17**:141.
5. Golzari SE, Ghabili K, Khanli HM, Mahmoodpoor A, Sabermarouf B: **Imported malaria cases: a source of moving backward from elimination to the preelimination phase in malaria control.** *Clin Infect Dis* 2013, **57**:1061-1062.

6. Lai S, Wardrop NA, Huang Z, Bosco C, Sun J, Bird T, Wesolowski A, Zhou S, Zhang Q, Zheng C, et al: **Plasmodium falciparum malaria importation from Africa to China and its mortality: an analysis of driving factors.** *Sci Rep* 2016, **6**:39524.
7. Zhang L, Feng J, Zhang SS, Xia ZG, Zhou SS: **[Malaria Situation in the People's Republic of China in 2015].** *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* 2016, **34**:477-481.
8. Zhou S, Li Z, Cotter C, Zheng C, Zhang Q, Li H, Zhou S, Zhou X, Yu H, Yang W: **Trends of imported malaria in China 2010-2014: analysis of surveillance data.** *Malar J* 2016, **15**:39.
9. Liu Y, Hsiang MS, Zhou H, Wang W, Cao Y, Gosling RD, Cao J, Gao Q: **Malaria in overseas labourers returning to China: an analysis of imported malaria in Jiangsu Province, 2001-2011.** *Malar J* 2014, **13**:29.
10. S. S, Z. L, Y. L, Y. Z: **Attracting Chinese FDI in Africa: The role of natural resources, market size and institutional quality.** *Crit Perspect Int Bus* 2018, **14**:139-153.
11. Zhang X, Gao Q, Wu Z, Zhao Y, Ge J: **Investigation of falciparum malaria control in Jiangsu province (in Chinese).** *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* 1996, **9**:245-249.
12. Hundessa S, Li S, Liu L, Guo J, Guo Y, Zhang W, Williams G: **Projecting environmental suitable areas for malaria transmission in China under climate change scenarios.** *Environ Res* 2018, **162**:203-210.
13. Kirchner S, Power BJ, Waters AP: **Recent advances in malaria genomics and epigenomics.** *Genome Med* 2016, **8**:92.
14. Escalante AA, Ferreira MU, Vinetz JM, Volkman SK, Cui L, Gamboa D, Krogstad DJ, Barry AE, Carlton JM, van Eijk AM, et al: **Malaria Molecular Epidemiology: Lessons from the International Centers of Excellence for Malaria Research Network.** *Am J Trop Med Hyg* 2015, **93**:79-86.
15. Auburn S, Barry AE: **Dissecting malaria biology and epidemiology using population genetics and genomics.** *Int J Parasitol* 2017, **47**:77-85.
16. Su X, Ferdig MT, Huang Y, Huynh CQ, Liu A, You J, Wootton JC, Wellems TE: **A genetic map and recombination parameters of the human malaria parasite Plasmodium falciparum.** *Science* 1999, **286**:1351-1353.
17. Anderson TJ, Haubold B, Williams JT, Estrada-Franco JG, Richardson L, Mollinedo R, Bockarie M, Mokili J, Mharakurwa S, French N, et al: **Microsatellite markers reveal a spectrum of population structures in the malaria parasite Plasmodium falciparum.** *Mol Biol Evol* 2000, **17**:1467-1482.
18. Anderson TJ, Su XZ, Bockarie M, Lagog M, Day KP: **Twelve microsatellite markers for characterization of Plasmodium falciparum from finger-prick blood samples.** *Parasitology* 1999, **119 (Pt 2)**:113-125.
19. Snounou G, Viriyakosol S, Zhu XP, Jarra W, Pinheiro L, do Rosario VE, Thaithong S, Brown KN: **High sensitivity of detection of human malaria parasites by the use of nested polymerase chain reaction.** *Mol Biochem Parasitol* 1993, **61**:315-320.
20. Steenkeste N, Incardona S, Chy S, Duval L, Ekala MT, Lim P, Hewitt S, Sochantha T, Socheat D, Rogier C, et al: **Towards high-throughput molecular detection of Plasmodium: new approaches and molecular markers.** *Malar J* 2009, **8**:86.

21. Greenhouse-Lab/MicroSPAT: v2.0.3 [<https://doi.org/10.5281/zenodo.2564160>]
22. Jost L: **G(ST) and its relatives do not measure differentiation.** *Mol Ecol* 2008, **17**:4015-4026.
23. Nei M, Chesser RK: **Estimation of fixation indices and gene diversities.** *Ann Hum Genet* 1983, **47**:253-259.
24. Haubold B, Hudson RR: **LIAN 3.0: detecting linkage disequilibrium in multilocus data.** *Linkage Analysis. Bioinformatics* 2000, **16**:847-848.
25. Verity R, Nichols RA: **Estimating the Number of Subpopulations (K) in Structured Populations.** *Genetics* 2016, **203**:1827-1839.
26. A. J, B. C, D. C: *The Genetic Structure of Populations.*1974.
27. Tessema S, Wesolowski A, Chen A, Murphy M, Wilhelm J, Mupiri AR, Ruktanonchai NW, Alegana VA, Tatem AJ, Tambo M, et al: **Using parasite genetic and human mobility data to infer local and cross-border malaria connectivity in Southern Africa.** *Elife* 2019, **8**.
28. Pringle JC, Tessema S, Wesolowski A, Chen A, Murphy M, Carpi G, Shields TM, Hamapumbu H, Searle KM, Kobayashi T, et al: **Genetic Evidence of Focal Plasmodium falciparum Transmission in a Pre-elimination Setting in Southern Province, Zambia.** *J Infect Dis* 2019, **219**:1254-1263.
29. Wasser SK, Shedlock AM, Comstock K, Ostrander EA, Mutayoba B, Stephens M: **Assigning African elephant DNA to geographic region of origin: applications to the ivory trade.** *Proc Natl Acad Sci U S A* 2004, **101**:14847-14852.
30. Ju-Lin L, Guo-Ding Z, Hua-Yun Z, Jian-Xia T, Meng-Meng Y, Wei-Ming W, Jun C: **[Vector surveillance in Jiangsu Province during the stage of malaria elimination].** *Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi* 2018, **30**:390-395.
31. Cohen JM, Smith DL, Cotter C, Ward A, Yamey G, Sabot OJ, Moonen B: **Malaria resurgence: a systematic review and assessment of its causes.** *Malar J* 2012, **11**:122.
32. Cao J, Sturrock HJ, Cotter C, Zhou S, Zhou H, Liu Y, Tang L, Gosling RD, Feachem RG, Gao Q: **Communicating and monitoring surveillance and response activities for malaria elimination: China's "1-3-7" strategy.** *PLoS Med* 2014, **11**:e1001642.
33. Chen T, Zhang S, Zhou SS, Wang X, Luo C, Zeng X, Guo X, Lin Z, Tu H, Sun X, Zhou H: **Receptivity to malaria in the China-Myanmar border in Yingjiang County, Yunnan Province, China.** *Malar J* 2017, **16**:478.
34. Mobegi VA, Loua KM, Ahouidi AD, Satoguina J, Nwakanma DC, Amambua-Ngwa A, Conway DJ: **Population genetic structure of Plasmodium falciparum across a region of diverse endemicity in West Africa.** *Malar J* 2012, **11**:223.
35. Duffy CW, Ba H, Assefa S, Ahouidi AD, Deh YB, Tandia A, Kirsebom FCM, Kwiatkowski DP, Conway DJ: **Population genetic structure and adaptation of malaria parasites on the edge of endemic distribution.** *Mol Ecol* 2017, **26**:2880-2894.
36. Conway DJ, Roper C, Oduola AM, Arnot DE, Kremsner PG, Grobusch MP, Curtis CF, Greenwood BM: **High recombination rate in natural populations of Plasmodium falciparum.** *Proc Natl Acad Sci U S A*

- 1999, **96**:4506-4511.
37. Manske M, Miotto O, Campino S, Auburn S, Almagro-Garcia J, Maslen G, O'Brien J, Djimde A, Doumbo O, Zongo I, et al: **Analysis of Plasmodium falciparum diversity in natural infections by deep sequencing.** *Nature* 2012, **487**:375-379.
  38. Mara SE, Silue KD, Raso G, N'Guetta S P, N'Goran E K, Tanner M, Utzinger J, Ding XC: **Genetic diversity of Plasmodium falciparum among school-aged children from the Man region, western Cote d'Ivoire.** *Malar J* 2013, **12**:419.
  39. Mendes C, Salgueiro P, Gonzalez V, Berzosa P, Benito A, do Rosario VE, de Sousa B, Cano J, Arez AP: **Genetic diversity and signatures of selection of drug resistance in Plasmodium populations from both human and mosquito hosts in continental Equatorial Guinea.** *Malar J* 2013, **12**:114.
  40. Pritchard JK, Stephens M, Donnelly P: **Inference of population structure using multilocus genotype data.** *Genetics* 2000, **155**:945-959.
  41. Wasser SK, Brown L, Mailand C, Mondol S, Clark W, Laurie C, Weir BS: **Genetic assignment of large seizures of elephant ivory reveals Africa's major poaching hotspots.** *Science* 2015, **349**:84-87.
  42. Wasser SK, Mailand C, Booth R, Mutayoba B, Kisamo E, Clark B, Stephens M: **Using DNA to track the origin of the largest ivory seizure since the 1989 trade ban.** *Proc Natl Acad Sci U S A* 2007, **104**:4228-4233.
  43. Fola AA, Nate E, Abby Harrison GL, Barnadas C, Hetzel MW, Iga J, Siba P, Mueller I, Barry AE: **Nationwide genetic surveillance of Plasmodium vivax in Papua New Guinea reveals heterogeneous transmission dynamics and routes of migration amongst subdivided populations.** *Infect Genet Evol* 2018, **58**:83-95.
  44. Koepfli C, Rodrigues PT, Antao T, Orjuela-Sanchez P, Van den Eede P, Gamboa D, van Hong N, Bendezu J, Erhart A, Barnadas C, et al: **Plasmodium vivax Diversity and Population Structure across Four Continents.** *PLoS Negl Trop Dis* 2015, **9**:e0003872.
  45. Koepfli C, Timinao L, Antao T, Barry AE, Siba P, Mueller I, Felger I: **A Large Plasmodium vivax Reservoir and Little Population Structure in the South Pacific.** *PLoS One* 2013, **8**:e66041.
  46. Schultz L, Wapling J, Mueller I, Ntsuke PO, Senn N, Nale J, Kiniboro B, Buckee CO, Tavul L, Siba PM, et al: **Multilocus haplotypes reveal variable levels of diversity and population structure of Plasmodium falciparum in Papua New Guinea, a region of intense perennial transmission.** *Malar J* 2010, **9**:336.

## Additional Files

**Additional file 1.** Description of the 16 newly designed and 10 existing microsatellite markers used in this study.

**Additional file 2.** Primer sequences, PCR master mixes and cycling conditions for the primary and secondary PCRs. (A. List of primers B. Primary PCR C. Secondary PCR D. Dilution and CE)

**Additional file 3.** Genetic diversity and the number of unique alleles in 336 imported monoclonal *P. falciparum* isolates from 5 regional populations of sub-Saharan Africa. New microsatellite markers are highlighted in grey.


**Additional file 4.** Relationship between the number of unique alleles and expected heterozygosity in 602 *P. falciparum* cases imported to Jiangsu province from 26 African countries. Previously described markers are shown in red and new markers are shown in blue.

**Additional file 5. A.** Pairwise linkage disequilibrium between 26 microsatellite markers. Pairwise index of associations (IA) is indicated **B.** Estimates of multi-locus linkage disequilibrium in 336 *P. falciparum* isolates imported from five different regions of sub-Saharan Africa to Jiangsu province, China.  $I_A^S$  is the standardized index of association

**Additional file 6.** Mean multiplicity of infection (MOI) and percentage of polyclonal samples by the geographic origin of imported infections to the Jiangsu province, China.


**Additional file 7. A.** Neighbor-joining tree showing genetic relatedness of 602 imported *P. falciparum* isolates, branches are colored according to region of origin of the imported case. **B.** Plot of second principal component against the first, computed from a multidimensional scaling based on the same distance matrix used for the tree shown in panel A.

## Figures




## Figure 1

Origins of imported malaria cases from five regions of sub-Saharan Africa to the Jiangsu Province of China. A. Number of samples from each country on the map. B. The total number of genotyped samples per region and percentage of single-clone infections.




**Figure 2**

Allele size and frequency distribution of alleles for the extended microsatellites panel.



**Figure 3**

Characterization of highly related infections. A. Relationship between the proportion of highly related infections and distance between destination countries in Africa. B. Relationship between the proportion of highly related infections and date difference in the reporting of malaria cases in Jiangsu, China. C. Relationship between the proportion of highly related infections, distance and reporting time. Genetically related infections were usually imported from shared trips to a similar destination. D. Relationship between the proportion of highly related infections and distance between residence district in Jiangsu, China. For panels A, B and D, black dots and blue lines indicate the observed and fitted data respectively. The grey dots and red lines indicate a null distribution from a permutation test.



**Figure 4**

Population assignment of 243 imported *P. falciparum* cases from Angola, Equatorial Guinea and Nigeria. A. Assignment accuracy for imported cases using the smoothing-based assignment method. B. Estimated locations of origin of 243 imported *P. falciparum* cases using the continuous-based assignment method.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile2.xlsx](#)
- [Additionalfile1.docx](#)
- [Additionalfile6.pptx](#)
- [Additionalfile7.pptx](#)

- [Additionalfile3.docx](#)
- [Additionalfile5.pptx](#)
- [Additionalfile4.pptx](#)