Serosurvey of anti-*Toxoplasma gondii* antibodies in homeless persons of São Paulo city, southeastern Brazil

Laís Giuliani Felipetto

Federal University of Parana

Pedro Irineu Teider-Junior

Federal University of Parana

Felipe Fortino Verdan da Silva

Federal University of Parana

Ana Carolina Yamakawa

São Paulo State University

Louise Bach Kmetiuk

Federal University of Parana

Anahi Chechia do Couto

Federal University of Parana

Camila Marinelli Martins

Ponta Grossa State University

Eduarda Stankiwich Vaz

Federal University of Parana

Leila Sabrina Ullmann

São Paulo State University

Helio Langoni

São Paulo State University

Jorge Timenetsky

São Paulo University

Andrea Pires dos Santos

Purdue University

Alexander Weiker Biondo (✉ abiondo@ufpr.br)

Federal University of Paraná <https://orcid.org/0000-0002-4182-5821>

Research article

Keywords: homeless, *Toxoplasma gondii*, HIV, vulnerability

Posted Date: May 8th, 2020

DOI: <https://doi.org/10.21203/rs.2.22570/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Seroprevalence of *Toxoplasma gondii* has been extensively studied in a variety of vulnerable populations. However, no study has focused on homeless populations. Accordingly, the present study aimed to assess the seroprevalence of anti-*T. gondii* antibodies and the risk factors associated in homeless persons from homeless shelter of São Paulo city, southeastern Brazil. In addition, anti-HIV antibodies and associated risk of *T. gondii* and HIV coinfection have been evaluated.

Methods: Anti-*T. gondii* antibodies were detected by indirect fluorescent antibody test. In addition, anti-HIV levels were tested by chemiluminescence enzyme immunoassay, with positive samples confirmed by rapid immunoblot assay.

Results: Overall, IgG anti-*T. gondii* seropositivity was found in 43/120 (35.8%) homeless persons, with endpoint titers varying from 16 to 1,024. The only two pregnant women tested were negative for IgM by chemiluminescence enzyme immunoassay, with normal parturition and clinically healthy newborns in both cases. There were no statistical differences in the risk factors for anti-*T. gondii* serology ($p > 0.05$). Anti-HIV seropositivity was found in 2/120 (1.7%) homeless persons, confirmed as HIV-1. One HIV seropositive individual was also sero-reactive to IgG anti-*T. gondii*, and both were negative to IgM anti-*T. gondii*.

Conclusions: This is the first study that reports the serosurvey of *T. gondii* in homeless persons worldwide. Despite the limited sample size available in the present study, our findings have shown that the prevalence of anti-*T. gondii* antibodies in homeless persons herein was lower than the general population, probably due to homeless diet habit of eating mainly processed food intake. No statistical differences were found regarding risk factors for anti-*T. gondii* exposure in homeless persons.

Background

Homeless persons have been described as one of the three most vulnerable populations, along with refugees and incarcerated persons [1]. Morbidity and mortality of diseases have been reportedly higher in homeless than general population, probably due to social inequality associated with lack of settled home, job opportunity, and a series of family problems including drug addiction, mental health disorders, and social justice issues, mostly exacerbated by absence of health assistance [2]. A population of 1.6 billion people without adequate housing has been estimated worldwide, of which 100 million are homeless [3, 4]. In Brazil, the nationwide homeless population has been estimated in 101,854 individuals, with about 40.1% living in cities with more than 900,000 inhabitants and about 16,000 living on streets of São Paulo city [5, 6].

Toxoplasma gondii is a coccidian parasite relying on cats and other *Felidae* as definitive hosts, which may shed fecal oocysts infecting a variety of homeothermic intermediate hosts [7]. Human infection has been typically subclinical or asymptomatic, with the time of infection and transmission route not known in most cases [8, 9]. Despite that, in immunodeficient people, such as in HIV-toxoplasmosis combination, the protozoan can cause severe clinical manifestations, with invasion into the central nervous system and encephalitis [10, 11].

Although human *T. gondii* seroprevalence has been extensively reported, ranging from 0.8% to 77.5% worldwide [12], few reports are available for vulnerable populations including 123/597 (20.6%) aboriginal individuals of Thailand, 236/628 (37.6%) prisoners of Turkey, mostly (92.8%) males, and 63/199 (31.7%) pregnant refugee and borderline migrant women of Asia [13-15]. In Brazil, seropositivity of *T. gondii* has been ranged from 14/65 (21.5%) urban students of northeastern region to 113/116 (97.4%) farmers of dairy cattle farm in central-western region [16, 17]. In vulnerable populations of Brazil, the positivity reported was 131/231 (56.7%) inhabitants living in riverside communities of northern region and 119/148 (80.4 %) indigenous of central-western region [18, 19].

Although *T. gondii* seroprevalence has been reported in a variety of vulnerable populations, no study has focused on homeless populations. Accordingly, the present study aimed to assess the seroprevalence of anti-*T. gondii* and the associated risk factors for exposure in homeless persons from homeless shelter of São Paulo city, southeastern Brazil. In addition, anti-HIV antibodies and associated risk of *T. gondii* and HIV coinfection have been evaluated.

Material And Methods

Local of study

The present study represents a descriptive cross-sectional seroepidemiological approach of the homeless population from the western São Paulo city (23°33'1"S, 46°38'2"W) shelter (Social Center "Our Lady of Good Delivery"), responsible for daytime attendance of all the city region. The shelter is a Non-Governmental Organization (NGO), sponsored by a city partnership daily attending around 800 to 1,200 homeless persons, providing meals, medical assistance, job opportunities, and recreational activities.

São Paulo city, capital of São Paulo State, southeastern Brazil, has been ranked as the most populated city of Latin America with 11,253,500 people and the tenth-largest Gross Domestic Product (GDP) of the world, with a very high Human Development Index (HDI) (0.805). The city is located under a humid subtropical climate with average temperatures ranging from 19 °C (winter) to 25 °C (summer) [20].

The present study was conducted along with the city multi-task professionals team at the São Paulo City Secretary of Health, called "street outreach office", which includes physicians, nurses, dentists, social assistants, and psychologists, based on the strategy of the Brazilian Unified Health System [21]. This city official team offers permanent assistance and save clinical records of the homeless population, promoting health actions on a continuing care bond.

Epidemiological data collection

Epidemiological analyses were performed based on a questionnaire associated with homeless person exposure to *T. gondii* and HIV, which included: 1) Demographic profile: sex, marital status, racial self-declaration, age, educational background, income, and city of origin; 2) Social profile: travel to other cities, communication with family, causes for becoming homeless, homelessness time, resting place, have children, have own children, pregnant woman, live alone, pet owner, use of licit and illicit drugs, alcohol consumption, tobacco use, marijuana use, cocaine use, crack use, assistance by the Psychosocial Care Centers (CAPS) as part of the free national Unified Health System; 3) Hygiene profile: bath frequency, change of clothes frequency, wash clothes, body lice (*Pediculus humanus humanus*) bites, and body lice presence (S1 Dataset). Refusal to fully or partially answer any question or incomplete answers were accepted and registered.

Sample collection

A minimal sampling of 71 individuals was calculated using commercially available software (Epi Info 7.7.7.6) based on an estimative of 16,000 homeless persons in São Paulo City and homeless HIV infection prevalence of 4.9% [22], the sampling was simple with 95% confidence and 5% accuracy. Blood samples of homeless persons were conveniently collected in São Paulo city from June to August 2018. Homeless persons were recruited by government health officials and invited to participate voluntarily of research, and blood collection was performed by cephalic puncture. Samples were placed in tubes without anticoagulant and kept at 25 °C until visible clot retraction. Serum was separated by centrifugation at 3,000 revolutions per minute for 10 minutes and stored at -20 °C until processing.

In addition, the packed cell volume (PCV) by [capillary tube](#) centrifugation and total plasma protein (TPP) by refractometry were performed on the day of sampling and immediately given to the shelter administration. Due to the shelter demand, homeless persons were also examined for body lice (*Pediculus humanus humanus*) bites and presence, as previously described at the same shelter [23]. The University made a clothing donation drive during the study and researchers offered clean clothes to all lice-infested homeless persons.

Serological diagnosis

Detection of *T. gondii* antibodies was performed by indirect immunofluorescent antibody test (IFAT) [24], with serial serum dilutions of 1:16 to 1:4,096 performed in pH 7.2 phosphate-buffered saline solution (PBS) with the cut-off titer of ≥ 16 IU. Immunofluorescence slides were previously sensitized with 0.1% formaldehyde to inactivated tachyzoites of *T. gondii* (RH strain) obtained from an intraperitoneal lavage in Swiss mice after three days of inoculation. A commercial anti-human IgG antibody,

conjugated with fluorescein isothiocyanate (Bethyl Laboratories, Montgomery, TX, USA) was used as secondary antibody. For positive samples, the highest titer was considered with at least 50% of fluorescence on the border of tachyzoites.

In addition, the detection of HIV was performed by chemiluminescent microparticle immunoassay (CMIA) (Alinity® sHIV Ag/Ab Combo Reagent Kit, Abbott Laboratories, Chicago, IL, USA) used for the simultaneous qualitative detection of HIV p24 antigen and antibodies to HIV type 1 (HIV-1 group M and group O) and/or type 2 (HIV-2) in human serum. The resulting chemiluminescent reaction was measured as relative light units (RLU). Samples of the pregnant woman and the HIV-positive individuals were tested for *T. gondii* IgM presence by CMIA (Anality® s Toxo IgM Reagent Kit, Abbott Laboratories, Chicago, IL, USA).

Statistical analysis

Statistical analysis was performed using SPSS 20.0 [25]. Frequencies of *T.gondii* and HIV seropositivity (absolute and relative) were determined by the stratification of the observations according to demographic, social, and hygiene profiles. The *Chi-Square* test was used to determine the bivariate association between studied variables, and odds ratios (OR) were used for the association of *T. gondii* prevalence and potential risk factors.

Ethics approval and consent to participate

This study was approved by the Ethics Committee in Human Research at the Federal University of Paraná (CAAE: 80099017.3.0000.0102, protocol number: 2.512.196), by the Ethics in Human Health Committee at the São Paulo City Secretary of Health (CAAE: 80099017.3.3004.0086, protocol number: 3.366.684) and by Ethics Committee in Human Research of the Clinics Hospital at the Federal University of Paraná (CAAE: 80099017.3.3005.0096, protocol number: 3.623.845), all subordinate to the National Human Ethics Research Committee of the Brazilian Ministry of Health. The Informed Consent Form was applied to all homeless persons, according to the ethical guidelines and principles of Federal University of Paraná. All participants research provided written informed consent.

Results

Overall, anti-*T.gondii* antibodies were detected in 43/120 (35.8%, CI 95% 26,7- 43,0%) homeless persons, with endpoint titers varying from 16 to 1,024. No statistical differences were found regarding risk factors for anti-*T. gondii* exposure ($p>0.05$) in homeless persons (Table 1).

Associated risk factors for the presence of anti-*T. gondii* were not statistically significant regarding educational background ($p=0.438$), income ($p=0.805$), resting place (hostels, street, occupancy) ($p>0.05$), pregnancy ($p=0.567$), pet owner (p=0.399); cat owner ($p=0.916$), bath frequency ($p=0.652$), age ($p=0.223$), and homelessness time ($p=0.827$) (Table 2). The homeless persons sampled were mostly men counting 107/120 (89.2%) individuals, with 39/107 (36.8%) seropositive samples for *T. gondii*. On the other side, women accounted for 13/120 (10.8%) with 3/13 (23.1%) positive samples. Despite in lower number, eight women were within the reproductive age of 24 to 35 years old, and 7/8 (87.5%) presented negative serology for *T. gondii*, including the two pregnant homeless women exposed to infection. Fortunately, the two pregnant women tested negative for anti-*T. gondii* antibodies in both IgG by IFAT and IgM by CMIA, with normal parturition and clinically healthy newborns in both cases.

In addition, a total of 2/120 (1.7%, CI 95% 0.0- 4.2%) anti-HIV seropositive homeless persons were detected by CMIA and confirmed by rapid immunoblot assay tests. No evaluation of HIV risk factors was made due to low seropositive rate.

Discussion

To the authors' knowledge, this is the first study that reports the serosurvey of *T. gondii* in homeless persons and the associated risk factors.

The seroprevalence of anti- *T. gondii* antibodies herein (35.8%) was higher than other vulnerable populations, such as aborigines (20.6%) and pregnant refugee and borderline migrant women (31.7%) but similar to incarcerated populations (37.6%) [13-15]. In Brazil, the anti-*T. gondii* seroprevalence herein was higher than the general population of the northeastern region, with 14/65

(21.5%) seropositive urban students, but lower than the central-western region, with 113/116 (97.4%) farmers from a single dairy cattle farm with domestic cats and potentially contaminated environment [16, 17]. Interestingly, the seroprevalence of anti-*T. gondii* antibodies in the present study was lower than other Brazilian neglected populations, such as 131/231 (56.7%) persons of riverside communities in the northern and 119/148 (80.4%) indigenous persons in the central-western region [18, 19]. In São Paulo, similar results were found, with 110/339 (32.4%) seropositive children from a low-socioeconomic community [26]. Not surprisingly, a previous study has shown an association between high seropositivity for *T. gondii* and socioeconomic vulnerability in southern Brazil, with 526/715 (73.57%) seropositive individuals, particularly in low-income families [27].

Although low education and socioeconomic status have been associated with increased risk of *T. gondii* infection in different Brazilian studies [28-30], no statistical association with *T. gondii* infection was previously found regarding educational background and income, probably due to the broadly variable classification of and the low population homogeneity [31, 32]. Similarly, no association was found in either education or income, likely associated to the impact of the vulnerable living style, with mostly drug addicts with poor eating habits.

Since the low socioeconomic status may be associated to malnutrition and might impair the host defense against protozoan infection, the relatively low seroprevalence of anti-*T. gondii* antibodies in homeless herein may be consequence of mainly consumption of ready-to-eat foods, as already indicated by previous studies on homelessness and food preparation facilities, which have reported dependence on charity meals such as pre-prepared foods, processed foods or popular snacks [33–36]. Not surprisingly, pre-processed ready-to-eat and meat-based foods have been shown to inactivate *T. gondii* cysts [37].

In addition, healthier and more expensive items such as meat, fish, vegetables, and fruits have been less often consumed by homeless [33, 36, 38, 39], which may be a contributing factor to the low *T. gondii* seroprevalence found in this study. Hence, it is reasonable to speculate that the beneficial shelters, hostels, and meal services may have offered protection to the homeless population [36, 40] but not as nutritional good food habits when compared to the general population. Although no homeless person has been diagnosed with either anemia by packed cell volume (PCV) or hypoproteinemia by refractometry, such tests may not have enough sensitivity to detect chronic alimentary deficiencies, which should be further investigated.

A previous study with pregnant women has shown high seroprevalence of specific anti-*T. gondii* antibodies (68.4%; 333/487) and vertical transmission associated with social vulnerability in central Brazil [41]. In the present study, despite negative for both IgG and IgM anti-*T. gondii* antibodies, the two pregnant women sampled, fortunately, gave birth to clinically healthy babies. *T. gondii* infection during pregnancy has been a significant problem, especially during the first months, and may result in spontaneous abortion, fetal and/or neonatal death or several congenital disabilities such as hydrocephalus, central nervous system disorders, and chorioretinitis [42, 43]. In the second and third trimester, newborns have usually been asymptomatic, with symptoms appearing late in childhood or early in adulthood, and may sporadically cause visual impairment [42, 44, 45]. In addition, congenital toxoplasmosis may also be associated with reactivation of the chronic maternal infection, particularly in HIV-infected and immunosuppressed women [46]. As 7/8 (87.5%) women herein were within reproductive age and presented negative serology for *T. gondii*, the homeless may be highly unprotected to infection during pregnancy.

Although the present study has shown no association between *T. gondii* infection and pet ownership, including stray cat owners, corroborating with previous studies in rural and other vulnerable populations of southern Brazil [27, 31], only 6/120 (0.5%) homeless persons owned a total of 11 cats. Outdoor lifestyle of stray cats may include hunting of birds and rodents, leading to raw meat dietary habits and increased risk of *T. gondii* ingestion [47]. Nonetheless, human toxoplasmosis outbreaks may be attributed to exposure to infected cats, which may indicate an important role of cat oocyst excretion on infection spreading [48, 49], and homeless might be daily overexposed to environmental contamination. However, as mentioned above, homeless dietary habits of high intake of processed foods and low fresh meat, fish, vegetables and fruits may have led to lower *T. gondii* exposure.

Body lice (*Pediculus humanus humanus*) has been recently considered as a reemerging problem among homeless populations in France, Italy, USA, Colombia, and Brazil [23, 50–53]. Even though body lice presence suggests social vulnerability and 17/120 (14.2%) of the homeless herein were infested with lice, no statistically risk of *T. gondii* exposure was found.

In addition to the *T. gondii* serosurvey, the HIV seroprevalence has been assessed herein. In the present study, 2/120 (1.7%) homeless persons was positive to anti-HIV antibodies, above the estimated prevalence of 0.4% for the general Brazilian population and lower than a recent study with 69/1,402 (4.9%) seropositive homeless. Despite previous studies have shown a high prevalence of *T. gondii* and HIV co-infection, with 35.8% worldwide in a meta-analysis study and up to 88.4% of individuals coinfecting in Ethiopia [54, 55], the analysis of associated risk factors herein has been impaired due to low HIV positive frequency.

Another limitation of the present study has surveyed low number of individuals, which has generated insufficient data to provide robust statistical description and analyses. However, mostly due to difficulties in accessing homeless and their refusal to participate, no study to date has been performed in toxoplasmosis and associated risk factors. In addition, questionnaires to assess homeless information may be problematic, particularly regarding food intake and dietary habits, once such a population has often shown a chaotic lifestyle and a high prevalence of drug use and mental health disorders [36].

Finally, further studies should be conducted in higher number and different homeless populations to fully establish the exactly impact of *T. gondii* in homeless persons worldwide.

Conclusions

In conclusion, this is the first study that reports the serosurvey of *T. gondii* in homeless persons worldwide. Despite the limited sample size available in the present study, our findings have shown that the prevalence of anti-*T. gondii* antibodies in homeless persons herein was lower than the general population, probably due to homeless diet habit of eating mainly processed food intake. No statistical differences were found regarding risk factors for anti-*T. gondii* exposure in homeless persons.

List Of Abbreviations

CAPS: Psychosocial Care Centers; CMIA: chemiluminescent microparticle immunoassay; GDP: Gross Domestic Product; HDI: Human Development Index; HIV: Human immunodeficiency virus; IFAT: immunofluorescent antibody test; NGO: Non-Governmental Organization; PBS: phosphate-buffered saline solution; PCV: packed cell volume; RLU: relative light units; OR: odds ratios; SD: standard deviation; TPP: total plasma protein.

Declarations

Acknowledgments

This research has been partially supported by the Veterinary Science Graduate Program at Federal University of Paraná, Clinical Analysis Laboratory Unit, Clinics Hospital (UFPR), Institute of Biotechnology (UNESP), Department of Veterinary Hygiene and Public Health (UNESP), and Purdue University. The authors are thankful to Dr. Mara Lúcia Gravinatti and the Social Center “Our Lady of Good Delivery” for helping with collection, sampling, and follow-up information.

Authors' Contributions

Drafting and revision of the manuscript: LGF, PIT-Jr, ACY, LBK, CMM, LSU, AWB. Initiation, conception, design and coordination of the research project: LGF, LSU, HL, JT, APS, AWB. Development of the intervention and evaluation materials: LGF, FFVS, ACC, CMM, ESV. Implementation of the intervention: LGF, PIT-Jr, LBK, CMM, LSU, APS, AWB. All authors read and approved the final version of the manuscript.

Funding

MSc Felipetto has been supported by a PhD fellowship and research funding from Capes DS through the Graduate College of Veterinary Science, Federal University of Paraná (UFPR).

Availability of data and materials

All data generated or analysed during this study are included in this published article [and its supplementary information files].

Consent for publication

Not applicable.

Competing interests

The authors have declared that no competing interests exist.

Author details

¹Departament of Veterinary Science, Federal University of Paraná (UFPR), Curitiba, Paraná (PR), 80035-050, Brazil. ²Clinical Analysis Laboratory Unit, Clinics Hospital, Federal University of Paraná (UFPR), Curitiba, Paraná (PR), 80060-900, Brazil. ³Department of Veterinary Hygiene and Public Health, São Paulo State University (UNESP), Botucatu, São Paulo (SP), 18618-681, Brazil. ⁴Departament of Cellular and Molecular Biology, Federal University of Paraná (UFPR), Curitiba (PR), 81531-970, Brazil. ⁵Department of Nursing and Public Health, State University of Ponta Grossa, Ponta Grossa (UEPG), Paraná (PR), 84030-900, Brazil, and AAC&T Research Consulting LTDA, Curitiba, Paraná (PR), 82540-096, Brazil. ⁶Institute of Biotechnology, São Paulo State University (UNESP), Botucatu, São Paulo (SP), 18607-440, Brazil. ⁷Department of Medical Microbiology, University of São Paulo (USP), São Paulo, São Paulo (SP), 05508-900, Brazil. ⁸Department of Comparative Pathobiology, Purdue University (PU), West Lafayette, Indiana (IN), 47907-2073, USA.

References

1. Aldridge RW, Story A, Hwang SW, Nordentoft M, Luchenski SA, Hartwell G, et al. Morbidity and mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders in high-income countries: a systematic review and meta-analysis. *Lancet*. 2018;391(10117):241-50. doi:10.1016/s0140-6736(17)31869-x.
2. Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: descriptive epidemiology, health consequences, and clinical and policy recommendations. *Lancet*. 2014;384(9953):1529–40. doi:10.1016/S0140-6736(14)61132-6.
3. United Nations Organization. Commission on Human Rights. Economic, Social and Cultural Rights. Report of the Special Rapporteur on adequate housing as a component of the right to an adequate standard of living, Miloon Kothari. 2005. <https://documents-dds-ny.un.org/doc/UNDOC/GEN/G05/117/55/PDF/G0511755.pdf?OpenElement>. Accessed 20 Dec 2019.
4. United Nations Organization. Expert Group Meeting. Affordable Housing and Social Protection Systems for All to Address Homelessness. Department of Economic and Social Affairs. <https://www.un.org/development/desa/dspd/wp-content/uploads/sites/22/2019/10/summary-egm-final-9sep.pdf>. Accessed 20 Dec 2019.
5. Institute of Economic Research Foundation. Pesquisa Censitária da População em Situação de Rua, Caracterização Socioeconômica da População Adulta em Situação de Rua e Relatório Temático de Identificação das Necessidades desta População na Cidade de São Paulo. 2015. https://www.prefeitura.sp.gov.br/cidade/secretarias/upload/00-publicacao_de_editais/0005.pdf. Accessed 10 Dec 2007.
6. Institute for Applied Economic Research. Atlas de Vulnerabilidade Social dos Municípios Brasileiros. http://ivs.ipea.gov.br/images/publicacoes/lvs/publicacao_atlas_ivs.pdf. Accessed 14 Sept. 2007.
7. Hill DE, Dubey JP. *Toxoplasma gondii*. Biology of Foodborne Parasites. CRC Press. 2015;209-222 pp.
8. Hill D, Dubey JP. *Toxoplasma gondii*: transmission, diagnosis and prevention. *Clin Microbiol Infect*. 2002;8(10):634-40. doi: 10.1046/j.1469-0691.2002.00485.x.
9. Petersen E, Vesco G, Villari S, Buffolano W. What do we know about risk factors for infection in humans with *Toxoplasma gondii* and how can we prevent infections?. *Zoonoses Public Health*. 2010;57(1):8-17. doi: 10.1111/j.1863-2378.2009.01278.x.
10. Deeks SG, Overbaugh J, Phillips A, Buchbinder S. HIV infection. *Nat Rev Dis Primers*. 2015;1:15035. doi: 10.1038/nrdp.2015.35.

11. Cong W, Dong XY, Meng QF, Zhou N, Wang XY, Huang SY, et al. *Toxoplasma gondii* infection in pregnant women: A seroprevalence and case-control study in Eastern China. *Biomed Res Int*. 2015;2015:170278. doi: 10.1155/2015/170278.
12. Pappas G, Roussos N, Falagas ME. Toxoplasmosis snapshots: global status of *Toxoplasma gondii* seroprevalence and implications for pregnancy and congenital toxoplasmosis. *Int J Parasitol*. 2009;39(12):1385-94. doi: 10.1016/j.ijpara.2009.04.003.
13. Fan CK, Liao CW, Wu MS, Su KE, Han BC. Seroepidemiology of *Toxoplasma gondii* infection among Chinese aboriginal and Han people residing in mountainous areas of northern Thailand. *J Parasitol*. 2003;89(6):1239-42. doi: 10.1645/GE-3215RN.
14. Yaman O, Yazar S, Çetinkaya Ü, Özcan Temel H, Balci E, Pehlivan İ, et al. Kayseri Kapalı.Cezaevi. Mahkumlarında *Toxoplasma gondii* Seroprevalansı. *Türkiye Parazitolo Derg*. 2009;33(1):15-9.
15. van Enter BJD, Lau YL, Ling CL, Watthanaworawit W, Sukthana Y, Lee WC, et al. Seroprevalence of *Toxoplasma gondii* Infection in Refugee and Migrant Pregnant Women along the Thailand-Myanmar Border. *Am J Trop Med Hyg*. 2017;97(1):232-5. doi:10.4269/ajtmh.16-0999.
16. de Amorim Garcia CA, Oréfice F, de Oliveira Lyra C, Gomes AB, França M, de Amorim Garcia Filho CA. Socioeconomic conditions as determining factors in the prevalence of systemic and ocular toxoplasmosis in Northeastern Brazil. *Ophthalmic Epidemiol*. 2004;11(4):301-17.
17. Santos GM, Silva SA, Barbosa AP, Campos DM. Investigação soropidemiológica sobre a larva migrans visceral por *Toxocara canis* em usuários de serviços de saúde de Goiânia – GO. *Ver Patol Trop*. 2009;38(3):197-206.
18. Amendoeira MRR, Sobral CAQ, Teva A, Lima JN, Klein CH. Inquérito sorológico para a infecção por *Toxoplasma gondii* em ameríndios isolados, Mato Grosso. *Rev. Soc. Bras. Med. Trop*. 2003;36(6):671-6. doi: 10.1590/S0037-86822003000600005.
19. Vitaliano SN, Mendonça GM, Sandres FAM, Camargo JSAA, Tarso P, Basano AS, et al. Epidemiological aspects of *Toxoplasma gondii* infection in riverside communities in the Southern Brazilian Amazon. *Rev. Soc. Bras. Med. Trop*. 2015;48(3):301-6. doi: 10.1590/0037-8682-0040-2015.
20. Brazilian Institute of Geography and Statistics. Censo Demográfico 2010: Panorama geral, São Paulo, Brasil. 2010. <https://cidades.ibge.gov.br/brasil/sp/sao-paulo/panorama>. Accessed 14 Sept. 2019.
21. Paula HC, Daher DV, Koopmans FF, Faria MGA, Brandão PS, Scoralick GBF. Implementation of the Street Outreach Office in the perspective of health care. *Revista Brasileira de Enfermagem*. 2018;71(6):2843-7. doi: 1590/0034-7167-2017-0616.
22. Grangeiro A, Holcman MM, Onaga ET, Alencar HDR, Placco ALN, Teixeira PR. Prevalência e vulnerabilidade à infecção pelo HIV de moradores de rua em São Paulo, SP. *Rev. Saúde Públ*. 2012;46(4):674-84. doi: 10.1590/S0034-89102012005000037.
23. Gravinatti ML, Faccini-Martínez ÁA, Ruys SR, Timenetsky J, Biondo AW. Preliminary report of body lice infesting homeless people in Brazil. *Rev Inst Med Trop S. Paulo*. 2018;60:e9. doi: 10.1590/s1678-9946201860009.
24. Camargo ME. Introdução às técnicas de imunofluorescência. *Rev. Bras. Patol. Clín*. 1974;10(3):143–69.
25. IBM Corp. Released in 20 IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.
26. Francisco FM, Souza SLP, Gennari SM, Pinheiro SR, Muradian V, Soares RM. Seroprevalence of toxoplasmosis in a low-income community in the São Paulo municipality, SP, Brazil. *Rev. Inst. Med. trop. S. Paulo*. 2006;48(3):167-70. doi: 10.1590/S0036-46652006000300009.
27. Mareze M, Benitez ADN, Brandão APD, Pinto-Ferreira F, Miura AC, Martins FDC, et al. Socioeconomic vulnerability associated to *Toxoplasma gondii* exposure in southern Brazil. *PLoS One*. 2019;14(2):e0212375. doi: 10.1371/journal.pne.0212375.
28. Bahia-Oliveira LM, Jones JL, Azevedo-Silva J, Alves CC, Oréfice F, Addiss DG. Highly endemic, waterborne toxoplasmosis in north Rio de Janeiro state, Brazil. *Emerg Infect Dis*. 2003;9(1):55-62. doi: 10.3201/eid0901.020160.
29. Avelino MM, Campos D Jr, Parada JB, Castro AM. Risk factors for *Toxoplasma gondii* infection in women of childbearing age. *Braz J Infect Dis*. 2004;8(2):164-74. doi: 10.1590/s1413-86702004000200007.
30. Lopes-Mori FM, Mitsuka-Breganó R, Bittencourt LHFB, Dias RCF, Gonçalves DD, Capobiango JD, et al. Gestational toxoplasmosis in Paraná State, Brazil: prevalence of IgG antibodies and associated risk factors. *Braz J Infect Dis*. 2013;17(4):405-9. doi: 10.1016/j.bjid.2012.12.003.
31. Araújo AC, Villela MM, Sena-Lopes Â, Farias NAR, Faria LMJ, Avila LFC, et al. Seroprevalence of *Toxoplasma gondii* and *Toxocara canis* in a human rural population of Southern Rio Grande do Sul. *Rev. Inst. Med. Trop. S. Paulo*. 2018;60:e28. doi:

10.1590/S1678-9946201860028.

32. Passos ADC, Bollela VR, Furtado JMF, Lucena MM, Bellissimo-Rodrigues F, Paula JS, et al. Prevalence and risk factors of toxoplasmosis among adults in a small Brazilian city. *Rev. Soc. Bras. Med. Trop.* 2018;51(6):781-7. doi: 10.1590/0037-8682-0214-2017.
33. Hickey C, Downey D. *Hungry for Change: Social Exclusion, Food Poverty and Homelessness in Dublin*. Dublin: Focus Ireland. 2003; 133pp.
34. Food Standards Agency. *Research into food poverty and homelessness in Northern Ireland – final report*. Belfast: Deloitte MCS Limited. 2006.
<http://webarchive.nationalarchives.gov.uk/20111206074236/http://www.food.gov.uk/multimedia/pdfs/homelessnifood.pdf>. Accessed 20 Dec 2019.
35. Alvarado-Esquivel C, Torres-Castorena A, Liesenfeld O, Estrada-Martínez S, Urbina-Álvarez JD. High seroprevalence of *Toxoplasma gondii* infection in a subset of Mexican patients with work accidents and low socioeconomic status. *Parasit Vectors*. 2012;5:13. doi: 10.1186/1756-3305-5-13.
36. Sprake EF, Russell JM, Barker ME. Food choice and nutrient intake amongst homeless people. *J Hum Nutr Diet*. 2014;27(3):242-50. doi: 10.1111/jhn.12130.
37. Mie T, Pointon AM, Hamilton DR, Kiermeier A. A qualitative assessment of *Toxoplasma gondii* risk in ready-to-eat smallgoods processing. *J Food Prot*. 2008;71(7):1442-52. doi: 10.4315/0362-028x-71.7.1442.
38. Rushton CM, Wheeler IE. The dietary intake of homeless males sleeping rough in Central London. *J Hum Nutr Diet*. 1993;6(5):443-56. doi: 10.1111/j.1365-277X.1993.tb00389.x.
39. Fallaize R, Seale JV, Mortin C, Armstrong L, Lovegrove JA. Dietary intake, nutritional status and mental wellbeing of homeless adults in Reading, UK. *Br J Nutr*. 2017;118(9):707-14. doi: 10.1017/S0007114517002495.
40. Li A, Dachner N, Tarasuk V. Food intake patterns of homeless youth in Toronto. *Can J Public Health*. 2009;100(1):36-40.
41. Gontijo da Silva M, Clare Vinaud M, de Castro AM. Prevalence of toxoplasmosis in pregnant women and vertical transmission of *Toxoplasma gondii* in patients from basic units of health from Gurupi, Tocantins, Brazil, from 2012 to 2014. *PLoS One*. 2015;10(11):e0141700. doi: 10.1371/journal.pone.0141700.
42. Kieffer F, Wallon M. Congenital toxoplasmosis. *Handb Clin Neurol*. 2013;112:1099-101. doi: 10.1016/B978-0-444-52910-7.00028-3.
43. Abamecha F, Awel H. Seroprevalence and risk factors of *Toxoplasma gondii* infection in pregnant women following antenatal care at Mizan Aman General Hospital, Bench Maji Zone (BMZ), Ethiopia. *BMC Infect Dis*. 2016;16(1):460. doi: 10.1186/s12879-016-1806-6.
44. Montoya JG, Liesenfeld O. Toxoplasmosis. *Lancet*. 2004;363(9425):1965-76. doi: 10.1016/S0140-6736(04)16412-X.
45. Berrébi A, Assouline C, Bessières MH, Lathière M, Cassaing S, Minville V, et al. Long-term outcome of children with congenital toxoplasmosis. *Am J Obstet Gynecol*. 2010;203(6):552.e1–552.e6. doi: <https://doi.org/10.1016/j.ajog.2010.06.002>.
46. Azevedo KML, Setúbal S, Lopes VGS, Camacho LAB, Oliveira AR. Congenital toxoplasmosis transmitted by human immunodeficiency-virus infected women. *Braz J Infect Dis*. 2010;14(2):186-9. doi: <http://dx.doi.org/10.1590/S1413-86702010000200014>.
47. Ding H, Gao YM, Deng Y, Lamberton PHL, Lu DB. A systematic review and meta-analysis of the seroprevalence of *Toxoplasma gondii* in cats in mainland China. *Parasit Vectors*. 2017;10:27. doi: 10.1186/s13071-017-1970-6.
48. Teutsch SM, Juranek DD, Sulzer A, Dubey JP, Sikes RK. Epidemic toxoplasmosis associated with infected cats. *N Engl J Med*. 1979;300(13):695-9. doi: 10.1056/NEJM197903293001302.
49. Torrey EF, Yolken RH. Toxoplasma oocysts as a public health problem. *Trends Parasitol*. 2013;29(8):380-4. doi: 10.1016/j.pt.2013.06.001.
50. Bonilla DL, Cole-Porse C, Kjemtrup A, Osikowicz L, Kosoy M. Risk factors for human lice and bartonellosis among the homeless, San Francisco, California, USA. *Emerg Infect Dis*. 2014;20(10):1645-51. doi: 10.3201/eid2010.131655.
51. Ly TDA, Touré Y, Calloix C, Badiaga S, Raoult D, Tissot-Dupont H, et al. Changing Demographics and Prevalence of Body Lice among Homeless Persons, Marseille, France. *Emerg Infect Dis*. 2017;23(11):1894-7. doi: 10.3201/eid2311.170516.

52. Faccini-Martínez ÁA, Márquez AC, Bravo-Estupiñan DM, Calixto OJ, López-Castillo CA, Botero-García CA, et al. *Bartonella quintana* and Typhus Group Rickettsiae Exposure among Homeless Persons, Bogotá, Colombia. *Emerg Infect Dis*. 2017;23(11):1876-9. doi: 10.3201/eid2311.170341.

53. Liberato C, Magliano A, Romiti F, Menegon M, Mancini F, Ciervo A, et al. Report of the human body louse (*Pediculus humanus*) from clothes sold in a market in central Italy. *Parasit Vectors*. 2019;12(201). doi: 10.1186/s13071-019-3458-z.

54. Wang ZD, Wang SC, Liu HH, Ma HY, Li ZY, Wei F, et al. Prevalence and burden of *Toxoplasma gondii* infection in HIV-infected people: a systematic review and meta-analysis. *Lancet HIV*. 2017;4(4):e177-e188. doi: 10.1016/S2352-3018(17)30005-X.

55. Feleke DG, Gebreweld A, Zewde G. Toxoplasmosis in Pregnant Women and HIV/AIDS Patients in Ethiopia: A Systematic Review and Meta-Analysis. *J. Parasitol. Res*. 2019;2019(Article ID 4670397):8pages. doi: 10.1155/2019/4670397.

Tables

Table 1: Statistical results of univariate and multiple logistic regression models of associated risk factors for seropositivity of IgG anti-*T. gondii* antibodies in homeless persons.

<i>T. gondii</i>											
Risk Factor		Total		Positive		Negative		<i>p</i> -value	OR		
		N	% of total N	N	% of line	N	% of line				
1. Demographic profile											
Sex	Male	107	89.2	41	38.3	66	61.7	0.282	0.48	(0.12–1.85)	
	Female	13	10.8	3	23.1	10	76.9				
Marital Status	Unmarried	108	90.0	41	38.0	67	62.0	0.377	0.54	(0.13–2.12)	
	Accompanied	12	10.0	3	25.0	9	75.0				
Racial self-declaration	White	28	23.3	8	28.6	20	71.4	0.310	1.60	(0.64–4.03)	
	Non-white	92	76.7	36	39.1	56	60.9				
Educational background	None to 8th grade	91	75.8	34	37.4	57	62.6	0.438	0.72	(0.32–1.64)	
	High School and University	29	24.2	9	31.0	20	69.0				
Income	No income	99	84.6	37	37.0	63	63.0	0.879	1.08	(0.38–3.03)	
	With income	18	15.4	7	38.9	11	61.1				
City of origin	São Paulo city	38	32.2	12	31.6	26	68.4	0.404	1.41	(0.62–3.02)	
	Other cities	80	67.8	32	39.5	49	60.5				
Travel to other cities	Yes	23	20.2	33	36.3	58	63.7	0.799	1.13	(0.44–2.89)	
	No	91	79.8	9	39.1	14	60.9				
2. Social profile											
Contact with family	Yes	66	55.9	23	34.8	43	65.2	0.582	1.22	(0.58–2.59)	
	No	52	44.1	21	39.6	32	60.4				
Causes for becoming homeless	Alcohol and drugs	28	23.3	9	32.1	19	67.9	0.736	1.16	(0.47–2.89)	
	Family conflicts	48	40.0	19	39.6	29	60.4				
	Housing loss	13	10.8	3	23.1	10	76.9				
	Other	18	15.0	9	50.0	9	50.0				
	Unemployment	34	28.3	10	29.4	24	70.6				
Resting place	Hostel	68	45.3	26	38.2	42	61.8	0.683	0.85	(0.40–1.81)	
	Street	52	34.7	18	34.6	34	65.4				
	Occupancy	30	20	10	33.3	20	66.7				
Pregnant woman	Yes	2	1.7	0	0.0	2	100.0	0.278	*		
	No	118	98.3	44	37.3	74	62.7				
Have children	Yes	81	67.5	33	40.7	48	59.3	0.182	0.57	(0.25–1.30)	
	No	39	32.5	11	28.2	28	71.8				
Have own children	Live together	4	5.1	0	0.0	4	100.0	0.087	*		
	Other people	74	94.9	32	43.2	42	56.8				
Live alone	Yes	52	46.0	19	36.5	33	63.5	0.815	0.91	(0.42–1.97)	
	No	61	54.0	21	34.4	40	65.6				
Pet owner	Yes	26	22.4	12	46.2	14	53.8	0.231	0.58	(0.24–1.41)	
	No	90	77.6	30	33.3	60	66.7				
Dog owner	Yes	21	18.1	11	52.4	10	47.6	0.088	0.44	(0.16–1.14)	
	No	95	81.9	31	32.6	64	67.4				
Cat owner	Yes	6	5.2	2	33.3	4	66.7	0.880	1.14	(0.20–6.52)	
	No	110	94.8	40	36.4	70	63.6				
Use of licit and/or illicit drugs	Yes	91	75.8	35	38.5	56	61.5	0.470	0.72	(0.29–1.75)	
	No	29	24.2	9	31.0	20	69.0				
Alcohol consumption	Yes	52	43.3	18	34.6	34	65.4	0.471	0.76	(0.36–1.60)	
Tobacco use	Yes	32	26.7	16	50.0	16	50.0	0.068	0.46	(0.20–1.06)	
Marijuana use	Yes	31	25.8	12	38.7	19	61.3	0.784	0.88	(0.38–2.06)	
Cocaine use	Yes	34	28.3	12	35.3	22	64.7	0.844	1.08	(0.47–2.48)	
Crack use	Yes	16	13.3	7	43.8	9	56.2	0.528	0.71	(0.24–2.06)	
Other drugs	Yes	5	4.2	3	60.0	2	40.0	0.269	0.36	(0.05–2.30)	
Assistance by Psychosocial Care Centers (CAPS)	Yes	31	25.8	10	32.3	21	67.7	0.554	1.29	(0.54–3.08)	
	No	89	74.2	34	38.2	55	61.8				
3. Hygiene profile											
Bath frequency	Daily	99	82.5	34	34.3	65	65.7	0.252	1.73	(0.67–4.50)	
	Sporadic	21	17.5	10	47.6	11	52.4				
Wash clothes	Yes	82	69.5	30	36.6	52	63.4	0.978	1.01	(0.45–2.24)	
	No	36	30.5	14	36.8	24	63.2				

Change clothes frequency	Daily	50	42.7	17	34.0	33	66.0	0.838	1.08 (0.50–2.33)
	Sporadic	67	57.3	24	35.8	43	64.2		
Body lice bite	Yes	63	59.4	23	36.5	40	63.5	0.752	1.13 (0.51–2.52)
	No	43	40.6	17	39.5	26	60.5		
Presence of body lice	Yes	17	14.2	4	23.5	13	76.5	0.225	2.06 (0.62–6.77)
	No	103	85.8	40	38.8	63	61.2		

*The percentages can go higher than 100% because individuals could answer more than one option.

Table 2: Average, median and standard deviation (SD) of *T. gondii* positive and negative homeless persons according to age (years), homelessness time (months), number of children, number of dogs, number of cats, packed cell volume (PCV), total plasma protein (TTP).

<i>ndii</i>							
Factor	Negative			Positive			<i>p</i> -value
	Average	Median	SD	Average	Median	SD	
(years)	43.55	44.50	14.03	41.18	42.50	11.96	0.412
elessness time (months)	67.68	36.00	78.43	85.10	36.00	103.60	0.557
ber of children	2.19	1.00	4.02	2.07	2.00	2.08	0.385
ber of dogs	0.34	0.00	1.27	0.62	0.00	1.40	0.081
ber of cats	0.05	0.00	0.23	0.17	0.00	0.82	0.928
	42.46	42.00	4.00	42.84	44.00	4.40	0.629
	7.65	7.60	0.63	7.64	7.60	0.52	0.918

SD, standard deviation; PCV, packed cell volume; TTP, total plasma protein.

Supplementary Information

Additional file 1: S1 Dataset. Data source, risk factors of demographic, social and hygiene profile and serological results for *T. gondii* the HIV inhomeless persons.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [S1File.xlsx](#)