Fall Predictors in Hospitalized Patients Living with Cancer: A Case-Control Study

Anita Zeneli (✉ anita.zeneli@irst.emr.it)

Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori Srl <https://orcid.org/0000-0002-8556-378X>

Sandra Montalti

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Itria Masciangelo

Cancer Institute of Romagna: IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Gloria Manieri

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Monica Golinucci

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Oriana Nanni

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Maria Teresa Montella

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Giovanni Martinelli

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Elisabetta Petracci

IRCCS Istituto Romagnolo per lo Studio dei Tumori Dino Amadori

Research Article

Keywords: hospitalized cancer patients, fall predictors, fall risk evaluation, risk assessment tool, fall with injury

Posted Date: March 1st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1317687/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Purpose

to identify fall predictors and develop an assessment tool to be used for screening hospitalized cancer patients at risk for fall.

Methods

a retrospective case-control study was conducted in 2018 at a cancer center in Northern Italy. The study participants were 448 adult cancer patients admitted to the oncology ward from 2009 to 2013. The cases group consisted of 112 patients presenting at least one fall, while controls were randomly chosen by matching each case for age, sex and admission period with three patients who did not fall. Data for the fall predictors were extracted from the electronic medical records. Conditional logistic regression was used to evaluate the association between patient's characteristics and fall risk.

Results

the overall prevalence of patients having at least one candidate fall predictor was high (98%). Seven of the studied variables showed an independent association with fall risk at multivariate analysis. These were tumor site, the presence of neurologic diseases, gait imbalance disorders, fatigue and the assumption of certain medications such as diuretics, hypnotics and opioids (odds ratios and 95% confidence intervals in brackets were 3.78 (1.78-8.13), 2.26 (1.08-4.77), 4.22 (1.87-9.52), 2.76 (1.45-5.26), 2.66 (1.52-4.66), 2.41 (1.20-4.85), and 3.03 (1.68-5.45), respectively).

Conclusions

results of our study highlight the need for developing risk assessment tools based on the characteristics of cancer centres. The validation with a prospective independent cohort is necessary before implementing our screening tool in clinical practice.

Introduction

Falls are a constant safety issue for healthcare providers, determining a heavy burden in terms of social, medical and financial outcomes. Despite several studies have previously investigated fall predictors and evaluated fall prevention strategies, this event continues to be a frequently reported critical incident in acute care hospitals [1, 2]. Oncological diseases and their treatments may increase the risk of falls; indeed, patients with cancer present higher fall and injury rates compared to other patients. This population should be considered unique [3, 4] and in particular in older patients living with cancer the problem will continue to be of increasing importance [5].

Several studies on the topic have previously been conducted on hospitalized patients living with cancer [3–18]. However, studies evaluating fall risk tools in this clinical setting are few and the characteristics of study centers are not described in the published works. Given the great variability between the oncology wards in different countries, the lack of such information makes it difficult to apply the results of these studies in clinical practice [8].

The aim of the present study was to identify predictors of accidental falls and to develop an assessment/screening tool for evaluating the risk of falls in hospitalized patients living with cancer.

Methods

Study design: this was a retrospective case-control study conducted in 2018 at Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori (IRST) in Meldola, a research cancer center in the Northern Italy.

Inclusion criteria: The target population consisted of the adult cancer patients that had at least one admission to the Medical Oncology ward of IRST between January 1, 2009 and December 31, 2013.

Selection of cases

In the present study, an accidental fall event was defined as "an unexpected event in which patients come to rest on the ground, floor, or other lower level" [19]. All patients who had at least an accidental fall during the study period were included in the case group. Falls were identified by the institutional fall incident report forms (FIRFs), routinely completed by the nurse and the physician at the time of each event. Date, time, location, circumstances of the fall, vital signs measurement, patient physical examination and type of harm are recorded in the FIRFs. The Patient Safety Network definitions are used to categorize the severity of harms described in the FIRFs. According to these definitions harms are classified as: no injury - fall resulted in no signs or symptoms of injury as determined by post fall evaluation; minor injury - fall resulted in application of a dressing, ice, cleaning of a wound, limb elevation, topical medication, bruise, or abrasion; moderate injury - fall resulted in suturing, application of steri-strips/skin glue, and splinting of muscle/joint strain; major injury - fall resulted in surgery, casting, traction, required consultation for neurological (basilar skull fracture, small subdural hematoma) or internal injury (rib fracture, liver laceration) or patients with coagulopathy who receive blood products as a result of the fall; Death – death as a result of injuries from the fall and not the physiological events causing the fall. [19–21].

Selection of controls

Patients with at least one admission to the oncology ward during the study period that did not fall were randomly selected as controls and matched (3:1 ratio) with cases by age (± 1 year), sex and the admission date (± 1 months).

The electronic admissions database was used for this purpose.

Explored risk factors

The risk factors investigated in this study were selected through a literature review and consisted of tumor site, comorbidities, sensory-neurologic deficits, mood and cognitive disorders (i.e., depression and delirium), previous fall history, treatment-induced fatigue, motor disorders requiring mobility assistance, medications (opioids, hypnotics, hypoglycemic agents, antihypertensives, diuretics), blood product use in the previous 24 hours, medical devices limiting ambulation, ambulatory aid use (i.e., crutches, cane or walker) [3–18].

Oncology ward characteristics

The oncology ward involved in this study is located inside the newly built cancer center in Meldola city. The facility was designed and furnished with the intent to maximize the humanization of care. The ward has 36 beds, of which 26 in double rooms and 10 in single rooms. Six of the single rooms are located in a low microbial load environment to allow care and protective isolation of the immunocompromised patients. All rooms are equipped with toilets and height-adjustable patient beds. In the inpatient setting nursing activities are scheduled in three 8-hour shifts to cover the 24-hours. Depending on the distribution of clinical activities during each shift, the nursing staffing foresees the presence in the morning shift of 8 nurses. This includes the head nurse, one case manager and one admission nurse. There are five nurses during the afternoon shift and three of them for the night shift. Three carers (health workers) for each shift during the day support nursing activities. The organizational model is based on the subdivision of responsibilities by sectors, in which there is a reference nurse in charge of nursing care of patients in each sector during working shifts. An institutional policy aimed at preventing accidental falls of patients is in place in our hospital. The average number of yearly admissions to the oncology ward is nearly 1700 patients with different types of advanced cancers. The most frequent diseases treated at our center are breast, lung, gastro-enteric, uro-gynecologic, hematologic and rare tumors.

Data collection

Data on risk factors were extracted from the electronic medical records (EMRs) by two trained research nurses. The EMRs contain the documentation of all clinical activities and assessments made during the patient admission period. Such activities are timely registered in the dedicated sections of the EMRs: i.e., vital signs, nursing assessment scheme, nursing and doctor diaries and therapy cards. In our clinical practice, an accurate and holistic patient evaluation is made by the admission nurse, who daily interviews all new admitted patients and registers data related to the patient health status, vital signs, symptoms evaluation using the Edmonton Symptom Assessment System (ESAS), the presence and status of medical and walking aid devices, sensor or cognitive disorders, motor disorders, and the use of medications. Patient assessment data are updated daily by the staff nurses, and in case of changes in patient clinical conditions, additional focused assessments are made. Data related to symptoms as potential fall predictors in this study (e.g., fatigue, pain, depression and anxiety) were collected as categorical depending on whether patients reported controlled/uncontrolled of the specific symptoms evaluated (i.e., the symptom fatigue was considered present in all patients reporting fatigue ≥ 4 when

evaluated with ESAS scale). The medications in this study were collected as categorical, independently from the daily dose and frequency use i.e., the “prescription and administration” (traceable from the therapy schema) in the 24 hours preceding the event of the following medication categories: opioids, hypnotics, anxiolytics, neuroleptics, antidepressants, antidiabetics as well as antihypertensives and diuretics. Data related to patient comorbidities were reported as categorical for each secondary diagnosis that was reported by the oncologists in the anamnesis section of the EMRs.

Statistical analysis

Continuous variables were reported as median, minimum and maximum values and were compared between case patients and controls using the Student t-test or the Wilcoxon-Mann-Whitney test, as appropriate. Categorical variables were reported as natural frequencies and percentages and were compared using the Chi-squared or Fisher’s Exact test, as appropriate. Correlation among variables was mainly assessed by means of tetrachoric correlation. Conditional logistic regression was used to evaluate the association between patient’s characteristics and fall risk. Results were reported as odds ratios (ORs) and 95% confidence intervals (CIs). From an initial full model, containing all factors listed in Table 1, a final parsimonious model was obtained by means of backward selection procedure with significance level for variable removal of 0.05. From the final multivariable model, a score to easily screen oncologic patients at the moment of hospital admission was derived as follows: *i)* a partial score was computed dividing the value of each regression coefficient by the smallest regression coefficient and the results were rounded to the nearest integer, or to the nearest integer +0.5; *ii)* a total score for each patient was obtained by adding their appropriate partial scores. To help the clinical applicability of the score, a cut-point was chosen by using the Youden index computed on the predicted probabilities obtained by fitting a logistic model for the risk of fall on the total score plus the matching variables [22] and taking the total score value corresponding to the found cut-point.

Table 1
Main characteristics of cases and controls (column percentages)

	Total (n = 448)		Cases (n = 112)		Controls (n = 336)		P- value
	n	%	n	%	n	%	
Female sex	180	40.2	44	39.3	136	40.5	0.824
Median age , years [min - max]	66.2 [20.6 – 83.2]		66.17 [23.3 – 82.9]		66.2 [20.6 – 83.2]		0.998
Tumor site							0.001
Solid tumors	389	86.8	87	77.7	302	89.9	
Hematologic malignancies	59	13.2	25	22.3	34	10.1	
Previous fall*	13	2.9	6	5.4	7	2.1	0.077
Need help for walking *	89	20.0	39	34.8	50	15.0	<0.001
Comorbidities							
Cardiovascular diseases*	218	49.0	59	52.7	159	47.8	0.367
Endocrine diseases*	113	25.3	27	24.1	86	25.7	0.742
Respiratory diseases*	144	32.3	41	36.6	103	30.8	0.259
Sensor-cognitive disorders							
CNS diseases*	56	12.6	26	23.2	30	9.0	<0.001
Peripheral neuropathy (neotoxicity)	22	4.9	7	6.3	15	4.5	0.449
Impaired vision/hearing	19	4.2	10	8.9	9	2.7	0.004
Plegie	6	1.3	2	1.8	4	1.2	0.643
Delirium	31	6.9	16	14.3	15	4.5	<0.001
Anxiety	169	37.7	52	46.4	117	34.8	0.028
Motor disorders							
Gait/balance disorders*	55	12.3	30	26.8	25	7.5	<0.001
Fatigue	208	46.4	72	64.3	136	40.5	<0.001
Hyposthenia of lower limbs	46	10.3	22	19.6	24	7.1	<0.001
Medications							

* Frequency of missing values for each variable were < 0.9%.

	Total		Cases		Controls		P-value
	(n = 448)		(n = 112)		(n = 336)		
Diuretics*	183	40.9	64	57.1	119	35.5	<0.001
Antihypertensives*	120	26.9	27	24.1	93	27.8	0.450
Hypnotics*	67	15.0	31	27.7	36	10.8	<0.001
Hypoglycaemic agents *	61	13.6	16	14.3	45	13.4	0.811
Opioids	221	49.3	80	71.4	141	42.0	<0.001
Blood transfusion, previous 24 hours*	52	11.7	11	9.8	41	12.3	0.478
Enemas	37	8.3	11	9.8	26	7.8	0.493
Medical devices							
Has IV line or access*	288	64.4	61	54.5	227	67.8	0.011
Ambulatory devices use*	40	9.0	19	17.0	21	6.3	0.001
* Frequency of missing values for each variable were < 0.9%.							

Multinomial logistic regression with fixed effects was used to investigate predictors of falls differentiating between falls without injuries and falls with a minor or moderate injury. The reference outcome category was represented by the control group.

All the statistical analyses were conducted using STATA statistical software version 15.1 (StataCorp, College Station, Texas, USA).

Results

Four hundred and forty eight adult cancer patients with at least one admission between 2009 and 2013 were included in the study. Of these, 112 were cases, i.e., patients that presented at least one event of accidental fall during the hospitalization period and 336 controls, i.e., patients without falls. Among the cases, 63 patients (56%) did not report injuries whereas 49 patients (44%) reported minor or moderate injuries. No events with major injuries or deaths were observed. The median hospital stay for cases was 12 days [range: 1-59] whereas for controls it was 3 days [range: 1-45]. The total number of admissions to the oncology ward during the five-year period considered in this study was 9,404, while the total number of patient hospital days was 53,876. The 5-year observed fall rate was 2.07 per 1000 patient hospital days and 1.19 per 100 patient hospital admissions. Data on patient characteristics and fall risk factors are shown in Table 1. About 60% of the patients were males with a median age of 66 years. The percentage of patients having at least one of the candidate fall predictors was 98.2. The number of

previous falls was low for both groups (overall frequency less than 3%) even if case patients reported a higher frequency as compared to controls ($P = 0.077$). Cases also had a significantly longer median hospital length of stay than controls, 12 days vs 3 days ($P < 0.001$). While solid tumors were the most frequent diagnoses in the study population (among them gastrointestinal and rare tumors were the most frequent), cases had a higher frequency of hematologic malignancies compared to controls, 22% vs. 10% ($P = 0.001$). With regards to other potential risk factors considered in this study, their overall prevalence was higher in cases than in controls. This was true in particular, for sensor-cognitive and mood disorders such as delirium, anxiety and depression, neurological diseases (including CNS diseases such as brain tumors or brain metastases), motor disorders, the use of ambulation devices and medications such as diuretics, hypnotics and opioids. Table 1.

Results from univariable conditional logistic regression are shown in Figure 1. Among studied risk factors, tumor type (haematological vs solid malignancies), sensorial-cognitive and mood disorders such as impaired vision/hearing, delirium, anxiety and depression, motor disorders, the use of ambulation devices, and the use of some medications (diuretics, hypnotics and opioids) resulted as predictors of accidental falls. The only factor showing a protective effect was having an intravenous (IV) access with infusion pump.

At multivariate analysis, seven variables showed an independent association with fall risk (Table 2). These were tumor site, the presence of CNS diseases, gait imbalance disorders, fatigue and the assumption of certain medications such as diuretics, hypnotics and opioids. Table 2 also shows the regression coefficients as well as the partial scores for each factor included in the multivariable model. The total score for a given patient was obtained by summing their appropriate partial scores. For example, for a patient with a solid tumor, a gait imbalance disorder and taking opioids, the total score was: $0+0+2+0+0+0+1.5=3.5$. The median total score computed on all patients was 2.5 [range: 0 - 7]. The total score cut-point was equal to 2. This was associated with a sensitivity of 92% and a specificity of 54%.

Table 2
Results from multivariable conditional logistic regression

	Coef.	OR (95% CI)	P-value	Partial score
Tumor site				
Solid	0	1 (ref)		0
Hematologic	1.33	3.78 (1.76-8.13)	0.001	1.5
neurologic diseases				
No	0	1 (ref)		0
Yes	0.82	2.26 (1.07 – 4.77)	0.032	1.0
Gait/balance disorders				
No	0	1 (ref)		0
Yes	1.44	4.22 (1.87 – 9.52)	0.001	2.0
Fatigue				
No	0	1 (ref)		0
Yes	1.02	2.76 (1.45 – 5.26)	0.002	1.0
Diuretics				
No	0	1 (ref)		0
Yes	0.98	2.66 (1.52 – 4.66)	0.001	1.0
Hypnotics				
No	0	1 (ref)		0
Yes	0.88	2.41 (1.20 – 4.85)	0.014	1.0
Opioids				
No	0	1 (ref)		0
Yes	1.11	3.03 (1.68 – 5.45)	<0.001	1.5

Additional exploratory analyses were performed distinguishing the cases between those who reported a minor or moderate injury and those who did not report an injury. The Supplementary Table S1 shows the distribution of patient characteristics among the three groups. Hospitalization length was longer for cases reporting an injury compared to the other patients (median of 10 days [range: 1-59] vs 3 [range: 1-45], respectively). In addition, such patients reported a slightly higher prevalence of hematologic tumors and need for a walking aid. The frequency of sensor-cognitive and mood disorders was higher in patients

not reporting a fall injury compared to other patients with the exception the presence of anxiety or depression. A similar pattern was observed with regards to motor disorders with the exception of fatigue, more prevalent among patients reporting an injury. Hypnotics were more prevalent among cases without injuries whereas hypoglycaemic medications among those reporting a minor or moderate injury.

To investigate predictors of fall without injury and fall with minor or moderate injury, a multinomial logistic regression was applied. Controls were chosen as reference category. Results from univariate analyses showed in Supplementary Table S2 were substantially consistent to those obtained by considering the binary outcome. However, few differences were observed. Previous falls, the use of infusion pumps in patients with IV lines and of medical devices as well as the presence of a neurologic disorder resulted as predictors only for the risk of fall without injury. Conversely, anxiety/depression resulted associated only to the risk of fall with injury (minor or moderate). At multivariate analysis, tumor site, the presence of gait/balance disorder, and the use of opioids were factors associated with the risk of fall with and without injury whereas fatigue was a predictor of risk of fall with injury and the use of hypnotics and diuretics of the risk of fall without harm (Table S3).

Discussion

Our study was aimed at identifying fall predictors in hospitalized patients with cancer and developing an assessment tool for screening patients at risk for fall at admission to the hospital. Even though several studies conducted in different countries have investigated the issue before, ours is the first study conducted in an Italian population in the cancer setting [3–18]. Risk for fall varies according to type and specificity of oncology wards [8]. The need to implement an assessment tool based on our patients and organizational characteristics justified the study. Hospitalized patients with different types of advanced cancer diseases were included in this study. The median age of the study population was in line with Stone et al, while the prevalence of falls with minor or moderate injuries (44%) was in line with two previous studies [9, 23], but lower if compared with Tsai (84.8%) [12]. Forty per cent of study participants were women, 86.8% had solid tumors and 13.2% were affected from hematologic malignancies.

Because the fall rates and injuries and the environment and organizational characteristics of included centres are not constantly described in the previous studies focusing on fall predictors, a full comparison of our findings with them is not possible. The fall rate per 1000 patient days in this study resulted lower than in Weed-Pfaff, Fischer and Pautex et al: respectively 2.07 vs. 3.41, 3.83 and 6.89. This also applies to the fall rate per 100 patients compared to Lorca, Weed-Pfaff and Capone's studies (i.e., respectively 1.19% vs. 1.98%, 2.4% and 3%) [3, 8, 11, 15, 23]. Clinical characteristics of patients and organizational variables may partially explain such differences in fall rates between studies.

In line with previous studies, we found for the most of the candidate fall predictors higher prevalences in the cases group as compared to controls. This was applies to the need for walking aid, having a neurologic disease, impaired vision/hearing, delirium, anxiety/depression, gait imbalance disorders, fatigue, hyposthenia of lower limbs, use of ambulatory devices, diuretics, hypnotics and opioids [3, 8- 11,

13]. Conversely, in contrast with previous studies, we found no statistically significant differences in prevalences of having comorbidities such as cardiovascular, endocrine and respiratory diseases, treatment induced neuropathy, the presence of plegies, and the use of antihypertensive, hypoglycemic agents and having received blood products transfusions in the last 24 hours. History of previous fall prevalence in the study population resulted low (2,6%) with no statistically significant difference between fallers and not fallers.

Having a hematologic malignancy (tumor site), neurologic disease (including brain metastases, or other neurologic diseases), gait imbalance disorders, fatigue and the use of diuretics, hypnotics and opioids resulted as independent predictors for fall.

Patients with hematologic malignancies at our centre are hospitalized mainly for receiving high intensity treatments (e.g., high dose chemotherapies, bone-marrow transplantation) or for the management of serious toxicities that can't be managed in outpatient settings. Treatment regimens of such patients could explain having hematologic malignancies as a fall predictor. This result is in line with the studies of Capone et al, Weed-Pfaff et al and Jun et al, who reported that hematologic patients receiving bone marrow transplantation present higher risk for falls [3, 11, 13].

Having a neurologic disease resulted as an independent fall predictor. Indirectly, this finding confirms the study conducted by Stone et al in 2012, in which having primary brain cancer or brain metastases resulted as a hospital fall predictor [9].

Gait imbalance disorders, asthenia of lower limbs and fatigue resulted as independent fall predictors as well. These factors are mutually influenced and are related with disease itself as well as cancer treatment side effects. Besides, patients experiencing fatigue likely have asthenia of lower limbs which may result in gait imbalance. These fall predictors were frequently found in most if not all of previous studies [3, 6, 7, 11–13, 25–28].

The use of diuretics resulted as an independent predictor of fall in the study population. It is known that the need for elimination as result of the effect of this medication category and the hypotensive effect of diuretics, which can explain such predictor. In fact, having elimination issues have resulted as fall predictor in other studies before [7, 24]. Among studied medications, the use of hypnotics and opioids resulted as independent fall predictors in line with findings reported from previous authors [3, 8–10, 28–30]. Both these two medication categories have effects on patient's cognitive status by influencing their space and time orientation and motor capabilities, which may explain this finding [7–9, 11].

The presence of medical devices in the study population was analysed singularly for each medical device limiting ambulation and by considering having at least one of them (i.e., nose-gastric tube, having infusion lines, drainage tubes, urinary catheter). In our study population having an IV line resulted protective for fall since the prevalence of patients with IV lines was higher in the controls group (OR 0.54; 95% CI 0.34-0.85; P-value = 0.008). This finding may be explained by the characteristics of the drip stands

used in our hospital for infusions and with the fact that patients with an IV line probably use them as an ambulatory aid while walking.

Among fall predictors, tumor site, gait imbalance disorders, fatigue and sensor-cognitive disorders such as anxiety/depression resulted related with falls with injury, while previous falls, use of diuretics, hypnotics and opioids were associated with falls without harms.

It is worth noting that despite the retrospective study design, the quality of the information contained in the EMRs helped minimize the known risk of recall bias.

In conclusion, the definition of a screening tool able to identify patients at higher risk for fall based on patients, organizational and facility characteristics represents the first step for implementing tailored evidence-based strategies for fall prevention. We believe that having described the characteristics of the study center, including facility, staffing and nursing organization could help other researchers to evaluate whether the results of our study apply or not to their own context. Thus, future study is needed to externally validate our results and investigate the clinical utility of our screening tool.

Declarations

Funding: No funding, grants or other support was received for this work.

Conflicts of interest/Competing interests: The authors have no competing interests to declare that are relevant to the content of this article

Data availability statement: All data relevant to the study are included in the article or uploaded as supplemental information. The authors confirm that the data supporting the findings of this study are available within the article (and/or) its supplemental materials. The authors have full control of primary data and agree to allow the journal to review their data if requested.

Code availability: Not applicable

Statement of authorship: AZ, SM, GM and MTM designed the study and provided the training of research nurses involved in the study. MG, GM, and IM were involved in data collection and registration. EP was responsible for data analysis. AZ, ON and EP contributed to the drafting of the manuscript and all authors critically reviewed the draft and approved the final version for submission.

Ethics approval: The study protocol was approved by the relevant medical, scientific and ethics committees of Area Vasta of Romagna (Prot. N. 2741/2013/1.5/113) and has been conducted in accordance with good clinical practice and the principles of the Helsinki Declaration.

Informed consent: Given the retrospective nature of the study, it was not possible to obtain the informed consensus from patients because most of them were not reachable (dead or in critical conditions). In

research centers and universities in Italy, the use of patient data for research purposes can be authorized by the General patients Authorization 2018 [31].

Consent to publish: Not applicable. Specific precautions were put in place to warranty the confidentiality of participants. In particular, the study data extracted from the EMRs were initially registered in a dedicated encrypted database with limited access to study nurses only. The completed database was permanently anonymized by deleting patients' identity data and by assigning a unique numerical code to each study participant.

Acknowledgements: The authors are grateful to Ing. Nicola Gentili for providing the electronic selection of controls.

References

1. National Patient Safety Agency NPSA (2007) The third report from the Patient Safety Observatory: *Slips trips and falls in hospital* available from: <https://cupdf.com/document/falls.html>
2. Mikos M, Banas T, Czerw A, Banas B, Strzpek Ł, Curyło M (2021) Hospital inpatient falls across clinical departments. *Int J Environ Res Public Health* 18(15):8167. doi: 10.3390/ijerph18158167
3. Capone LJ, Albert NM, Bena JF, Tang AS (2012) Predictors of a fall event in hospitalized patients with cancer. *Oncol Nurs Forum* 39(5):E407–E415. doi: 10.1188/12.ONF.E407-E415
4. Allan-Gibbs R (2010) Falls and hospitalized patients with cancer: a review of the literature. *Clin J Oncol* 14(6):784–792. doi: 10.1188/10.CJON.784-792
5. Spoelstra SL, Given BA, Schutte DL, Sikorskii A, You M, Given CW (2013) Do older adults with cancer fall more often? A comparative analysis of falls in those with and without cancer. *Oncol Nurs Forum* 40(2):E69–78. doi: 10.1188/13.ONF.E69-E78
6. Holley S (2002) A look at the problem of falls among people with cancer. *Clin J Oncol Nurs* 6(4):193–197. doi: 10.1188/02.CJON.193-197
7. O'Connell B, Cockayne M, Wellman D, Baker L (2005) Fall risk factors and the nature of falls in inpatient oncology and palliative care settings. *Contemp Nurse* 18(3):247–257. doi: 10.5172/conu.18.3.247
8. Pautex S, Herrmann FR, Zulian GB (2008) Factors associated with falls in patients with cancer hospitalized for palliative care. *J Palliat Med* 11(6):878–884. doi: 10.1089/jpm.2007.0241
9. Stone CA, Lawlor PG, Kenny RA (2011) How to identify patients with cancer at risk of falling: a review of the evidence. *J Palliat Med* 14(2):221–230. doi: 10.1089/jpm.2010.0326
10. Wildes TM, Dua P, Fowler SA, Miller JP, Carpenter CR, Avidan MS, Stark S (2015) Systematic review of falls in older adults with cancer. *J Geriatr Oncol* 6(1):70–83. doi: 10.1016/j.jgo.2014.10.003
11. Weed-Pfaff SH, Nutter B, Bena JF, Forney J, Field R, Szoka L, Karius D, Akins P, Colvin CM, Albert NM Validation of Predictors of Fall Events in Hospitalized Patients With Cancer. *Clin J Oncol Nurs* 20(5):E126-31. doi: 10.1188/16.CJON.E126-E131

12. Tsai LY, Campbell M, Chen CJ, Hsieh RK, Chien HH, Tsai JM (2017) Falls and related injuries in hospitalized patients with cancer in Taiwan. *J Nurs Res* 4:310–318. doi: 10.1097/JNR.0000000000000174
13. Jun MD, Lee KM, Park SA (2018) Risk factors of falls among inpatients with cancer. *Int Nurs Rev* 65(2):254–261. doi: 10.1111/inr.12381
14. Zhang X, Sun M, Liu S, Leung CH, Pang L, Popat UR, Champlin R, Holmes HM, Valero V, Dinney CP, Tripathy D, Edwards BJ (2018) Risk factors for falls in older patients with cancer. *BMJ Support Palliat Care* 8(1):34–37. doi: 10.1136/bmjspcare-2017-001388
15. Lorca LA, Sacomori C, Balagué-Ávila VP, Pino-Márquez LP, Quiroz-Vidal FA, Ortega L (2019) Incidence and risk of falls in patients treated for hematologic malignancies in the Intensive Hematology Unit. *Rev Lat Am Enfermagem* 29:27:e3145. doi: 10.1590/1518-8345.2953-3145
16. Sattar S, Alibhai SM, Spoelstra SL, Fazelzad R, Puts MT (2016) Falls in older adults with cancer: a systematic review of prevalence, injurious falls, and impact on cancer treatment. *Support Care Cancer* 24(10):4459–4469. doi: 10.1007/s00520-016-3342-8
17. Sattar S, Kenis C, Haase K, Burhenn P, Stolz-Baskett P, Milisen K, Ayala AP, Puts MTE (2020) Falls in older patients with cancer: Nursing and Allied Health Group of International Society of Geriatric Oncology review paper. *J Geriatr Oncol* 11(1):1–7. doi: 10.1016/j.jgo.2019.03.020
18. Sattar S, Haase K, Kuster S, Puts M, Spoelstra S, Bradley C, Wildes TM, Alibhai S (2021) Falls in older adults with cancer: an updated systematic review of prevalence, injurious falls, and impact on cancer treatment. *Support Care Cancer* 29(1):21–33. doi: 10.1007/s00520-020-05619-2
19. Lamb SE, Jørstad-Stein EC, Hauer K, Becker C, Prevention of Falls Network Europe and Outcomes Consensus Group (2005) Development of a common outcome data set for fall injury prevention trials: the Prevention of Falls Network Europe consensus. *J Am Geriatr Soc* 53(9):1618–1622. doi: 10.1111/j.1532-5415.2005.53455.x
20. Burns Z, Khasnabish S, Hurley AC et al (2020) Classification of Injurious Fall Severity in Hospitalized Adults. *J Gerontol A Biol Sci Med Sci* 75(10):e138–e144. doi: 10.1093/gerona/glaa004. PMID: 31907532
21. Health Research & Educational Trust (2016), October Preventing patient falls: A systematic approach from the Joint Commission Center for Transforming Healthcare project. Chicago, IL: Health Research & Educational Trust. <https://www.aha.org/ahahret-guides/2016-10-04-preventing-patient-falls> Accessed 26 June 2021
22. Pepe MS, Fan J, Seymour CW (2013) Estimating the receiver operating characteristic curve in studies that match controls to cases on covariates. *Acad Radiol* 20(7):863–873. doi: 10.1016/j.acra.2013.03.004
23. Fischer ID, Krauss MJ, Dunagan WC, Birge S, Hitcho E, Johnson S, Costantinou E, Fraser VJ (2005) Patterns and predictors of inpatient falls and fall-related injuries in a large academic hospital. *Infect Control Hosp Epidemiol* 126(10):822–827. doi: 10.1086/502500

24. Dharmarajan TS, Avula S, Norkus EP (2007) Anemia increases risk for falls in hospitalized older adults: an evaluation of falls in 362 hospitalized, ambulatory, long-term care, and community patients. *J Am Med Dir Assoc* 8(3 Suppl 2):e9–e15. doi: 10.1016/j.jamda.2006.12.015
25. Winters-Stone KM, Horak F, Jacobs PG, Trubowitz P, Dieckmann NF, Stoyles S, Faithfull S (2017) Falls, functioning, and disability among women with persistent symptoms of chemotherapy-induced peripheral neuropathy. *J Clin Oncol* 10(23):2604–2612. doi: 10.1200/JCO.2016.71.3552
26. Gewandter JS, Fan L, Magnuson A, Mustian K, Peppone L, Heckler C, Hopkins J, Tejani M, Morrow GR, Mohile SG (2013) Falls and functional impairments in cancer survivors with chemotherapy-induced peripheral neuropathy (CIPN): a University of Rochester CCOP study. *Support Care Cancer* 21(7):2059–2066. doi: 10.1007/s00520-013-1766-y
27. Tofthagen C, Overcash J, Kip K (2012) Falls in persons with chemotherapy-induced peripheral neuropathy. *Support Care Cancer* 20(3):583–589. doi: 10.1007/s00520-011-1127-7
28. Vela CM, Grate LM, McBride A, Devine S, Andritsos LA (2018) A retrospective review of fall risk factors in the bone marrow transplant inpatient service. *J Oncol Pharm Pract* 24(4):272–280. doi: 10.1177/1078155217697485
29. Li Y, Zhang Q, Yang X, Zheng L, Yang J, Zhao H, Yang D (2018) Research of falls risk of taking central nervous system drugs in oncology inpatients. *Curr Probl Cancer* 42(2):261–267. doi: 10.1016/j.crrproblcancer.2018.01.008
30. Herzig SJ, Rothberg MB, Moss CR, Maddaleni G, Bertisch SM, Wong J, Zhou W, Ngo L, Anderson TS, Gurwitz JH, Marcantonio ER (2021) Risk of in-hospital falls among medications commonly used for insomnia in hospitalized patients. *Sleep* 13(9):zsab064. doi: 10.1093/sleep/zsab064
31. DECRETO LEGISLATIVO 10 agosto (2018) n. 101 “Disposizioni per l'adeguamento della normativa nazionale alle disposizioni del regolamento (UE) 2016/679 del Parlamento europeo e del Consiglio, del 27 aprile 2016, relativo alla protezione delle persone fisiche con riguardo al trattamento dei dati personali, nonché alla libera circolazione di tali dati e che abroga la direttiva 95/46/CE (regolamento generale sulla protezione dei dati). Retrieved from <https://www.gazzettaufficiale.it/eli/id/2018/09/04/18G00129/sg> Accessed 26 June 2021

Figures

Figure 1

Results from univariate conditional logistic regression analysis

‡ Referred to the previous 24 hours

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS1.docx](#)
- [TableS2.docx](#)
- [TableS3.docx](#)