Interactive Effects of Melatonin and Salicylic Acid on Brassica napus Under Drought Condition

Noshin Ilyas (✉ noshinilyas@yahoo.com)

<https://orcid.org/0000-0002-5942-5439>

Noman Rafique

Muhammad Aqeel

Naveed Iqbal Raja

Ghulam Shabbir

Muhammad Ajaib

Research Article

Keywords: Phyto-Melatonin, Abiotic stress, Antioxidant response, Salicylic acid, Canola yield, Brassica napus

Posted Date: August 3rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1321383/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

AIMS:

It is imperative to understand the synergetic actions of melatonin (MET) and salicylic acid (SA) in counteracting the negative effects of drought. A pot experiment was performed to investigate the interactive effects of exogenously applied MET and SA on morphology, physiology, and biochemical characteristics of drought stressed canola (*Brassica napus*).

METHODS:

Two cultivars (Super canola and Faisal canola) were used. MET (0.1 μ M) and SA (0.50 mM) were applied as seed priming, and as foliar spray. Plants were exposed to severe drought stress (45% field capacity) at reproductive stage.

RESULTS:

Subjecting canola cultivars to drought caused significant reduction in shoot length (19%), plant fresh (15%) and dry (17%) biomass, yield (30%) and photosynthetic attributes (43%). Increased activity of free amino acids (20%), total soluble sugars (5%), and different antioxidant enzymes like catalase (27%), peroxidase (23%), superoxide dismutase (20%) and ascorbate peroxidase (35%) were recorded under drought. Seed priming, and foliar application of MET and SA increased growth and yield of canola cultivars. Priming of seeds with MET as well as synergetic application of MET priming and SA foliar spray were the most effective treatments among all, showing the greatest growth criteria (22% increase in plant height, 46% in fresh and 40% in dry biomass), yield components (49%) and endogenous MET (17%) and SA (16%) activity. Super canola cultivar showed better growth than Faisal canola.

CONCLUSION:

Study showed that synergetic application of MET and SA has the potential to induce stress resistance and enhance canola plants growth under drought.

Introduction

As the world population which is currently about 7 billion is expected to reach 9 billion by 2050, food security to support this much population is threatened to a larger extent by different types of environmental stresses (Khan et al. 2020a). Among these, drought is a severe problem for the agriculture of Pakistan, affecting crop growth and yield. More than 1.2 billion hectares of agricultural land of the world are affected by drought stress (Cui et al. 2017; Urban et al. 2017). Limited supply of irrigation water is the main factor responsible for the low yield and production of crops in Pakistan (Liang et al. 2019). Water stress shows its effect by reducing root growth, seed number, and size of leaves as well as stomatal opening (Kamran et al. 2018). It also delays flowering and fruiting in plants thus reducing their productivity (Cui et al. 2017; Naeem et al. 2018; Osakabe et al. 2014; Xu et al. 2016). Canola (*Brassica*

napus) is normally grown in Pakistan in areas that depend upon rainwater for good crop growth, so the availability of water remains a major limiting factor for plant growth and development in these rainfed areas (Tariq et al. 2019).

Brassica crops commonly grown as oilseed crops in Pakistan include Rapeseed (*B. rapa* and *B. napus*) and mustard (*B. juncea*). Among these, mustard is traditionally used in Pakistan, however, its oil is not preferred in the preparation of vegetable oil or ghee due to the presence of very high amounts (40–70%) of erucic acid in oil, which is not suitable for human consumption (Onemli 2014). The annual edible oil requirement of Pakistan is about 1.95 million tonnes, of which only 18% is obtained from the local resources and the rest 88% is imported (Mustafa et al. 2017). *B. napus*, being indigenous species, has all the potentials to overcome the problems of low local oil production. It contains 44–46% of good quality oil, with its meal having 38–40% of protein contents rich in lysine, methionine, and cystine amino acids (Onemli 2014).

The yield of crops in most areas of Pakistan is drastically affected by drought. Canola is cultivated in high rainfall regions and shows low production in areas with low rainfall (Ahmadi and Bahrani 2009; Tariq et al. 2019). To overcome drought, plant cells show different mechanisms which help them to survive under mild water stress (Osakabe et al. 2013). Among these, the production of plant growth regulators plays a vital role in plants' resistance against drought. Plant resistance to drought is found to be significantly boosted by exogenous application of these plant growth regulating substances (Peleg and Blumwald 2011).

Melatonin and Salicylic acid are two important plant growth regulators. Melatonin (N-acetyl-5-methoxytryptamine), being a relatively 'new' compound discovered in plants, has become a focus of plant researchers due to its tremendous potential as a regulator of many physiological processes. Melatonin application not only promotes plant growth and development, but it also induces stress resistance in plant cells (Kaya et al. 2022; Eaisalou et al. 2021; Amjadi et al. 2021; Liang et al. 2019; Martinez et al. 2018). Salicylic acid is also involved in the regulation of developmental processes in plants. It is a phenolic compound that regulates plants' responses to biotic and abiotic stresses (Kaya et al. 2022; Amjadi et al. 2021; Zaid et al. 2019). Many physiological processes in plants like photosynthesis, nitrogen metabolism, synthesis of glycine betaine, and production of antioxidants are regulated by salicylic acid, providing plants resistance against abiotic stresses (Eaisalou et al. 2021; Amjadi et al. 2021; Hasanuzzaman et al. 2019; Zaid et al. 2019).

Being biostimulants of photosynthetic processes as well as acting as enhancers of crop yield by modulating anti stress responses, both melatonin and salicylic acid have tremendous potential to enhance plant resistance against biotic and abiotic stresses (Hernandez-Ruiz and Arnao 2018). They share a number of similarities in their mode of action and have common precursor in their biosynthetic pathways. Combined potential of these growth regulators in reducing the severity of abiotic stresses (Kaya et al. 2022; Eaisalou et al. 2021; Amjadi et al. 2021) as well as biotic stress (Lee et al. 2015) is well documented. Individual or synergetic application of melatonin and salicylic acid cause significant

increase in the activity of enzymes involved in ascorbate-glutathione cycle, inducing abiotic stress resistance in plants (Amjadi et al. 2021). Synergetic application of melatonin and salicylic acid, through seed priming or any other method might enhance plant resistance to drought, maintaining proper metabolism during this stress (Hernandez-Ruiz and Arnao, 2018).

Individual effects of melatonin and salicylic acid in counteracting drought stress in plants is well documented; however, their combined effect is not studied yet. Present research work aimed to evaluate the response of two *B. napus* cultivars to drought stress concerning their growth, yield, and biochemical responses. In particular, our goal was to investigate the potential of the combined application of melatonin and salicylic acid, via pre-sowing seed treatment and foliar spray, in inducing drought resistance and mitigating the drought stress effects on two canola cultivars.

Materials And Methods

Experimental setup and procurement of seeds:

Present experiment was designed to assess the interactive actions of melatonin (MET) and salicylic acid (SA), potential plant growth regulators, in reducing the adversity of drought stress on the growth of canola (*Brassica napus* L.). For this purpose, exogenous application of these compatible solutes as seed priming, and foliar spray methods were employed in soil culture pot experiment. An experimental area of the department of Botany, PMAS Arid Agriculture University, Rawalpindi was used for the experiment. Two canola cultivars, namely Super canola (V1) and Faisal canola (V2) were used. Seeds were attained from National Agriculture Research Center Islamabad, Pakistan.

Treatment's schedule:

Before sowing, seeds were surface sterilized with 10% Clorox. Potassium, Phosphorous, and nitrogen fertilizers were applied to the soil prior to seed sowing. Soil analysis was also carried out before starting the experiment. Two plant growth regulators, Melatonin (Sigma-Aldrich, Lot# SLBZ6359) and Salicylic acid (Sigma-Aldrich, Lot# STBH3314) were used in present study. MET (0.1 μ M) and SA (0.5 mM) were applied in two different ways, firstly, seeds were primed with these growth regulators for 24 hours, and later at adult growth stages, foliar application of MET (0.1 μ M) and SA (0.5 mM) was done in both control and water stressed plants. 4 hours seed soaking in distilled water (for control) was also done, before sowing. Plastic pots were used for experimentation which were filled with sandy loam soil (sand: 44.2%, silt: 4.6%, and clay: 51.2%). Height and diameter of each pot was 24cm. Every pot contained 5 plants initially, which were later reduced to 3 plants per pot. Each treatment was replicated three times. Drought stress was applied by withholding water at reproductive stage, pots were weighed on daily basis and soil field capacity was maintained at 45% (Eq. 1). Plants were grown in the 2020–2021 growing season.

Relevant water content (RWC) % = $[(FM - DM)/(TM - DM)] \times 100$ (Eq. 1)

The treatments were applied as described below:

S. No.	Treatment Title	Treatment Description
1	T ₀	Controlled Irrigation
2	T ₁	MET (0.1 µM) Priming
3	T ₂	MET (0.1 µM) Foliar Spray
4	T ₃	SA (0.5 mM) Priming
5	T ₄	SA (0.5 mM) Foliar Spray
6	T ₅	MET (0.1 µM) plus SA (0.5 mM) Combined Priming
7	T ₆	MET (0.1 µM) plus SA (0.5 mM) Combined Foliar Spray
8	T ₇	MET (0.1 µM) Priming combined with SA (0.5 mM) Foliar Spray
9	T ₈	SA (0.5 mM) Priming Combined with MET (0.1 µM) Foliar Spray

10 days after the imposition of drought, data for different photosynthetic attributes of each plant was recorded by using portable infrared gas analyzer (IRGA). Leaf samples (5 healthy leaves each pot) were then collected for further physiological and biochemical analysis.

Morphological And Yield Attributes:

Data for different morphological parameters, i.e., plant height (cm), root length (cm), plant fresh and dry biomass (g), and yield attributes i.e., no. of siliqua per plant, no. of seeds per siliqua were recorded.

Gas Exchange/ Water Relation Parameters:

Infrared gas analyzer (LCA-4 ADC) was used for the estimation of gas exchange characteristics. These measurements were carried out from 10:00 to 14.00 hours. Measurement of photosynthetic rate, transpiration rate, water use efficiency and stomatal conductance were made using the young leaf. Methodology given by Capell and Doerffling (1993) was used for the estimation of osmotic potential. An osmometer (Model: Wescor Vapro 5520, WESCOR, USA) was used for this purpose. Water potential was estimated by using Scholander Pressure chamber (SAPS 11, Model 3115, Soil Moisture Equipment Corp. USA) according to the methodology given by Scholander et al. (1965).

Biochemical Attributes:

Bates et al. (1973) method was followed for estimation of protein contents. 0.2g of fresh leaf material were mixed with 10ml of phosphate buffer. 0.5ml of this extract was then added with 3ml of bio-red dye and 0.5ml of distilled water. Sample absorbance was checked at 595nm. Total amino acids were estimated according to the methodology given by Sohail et al. (2018). 0.2g of fresh leaf material was mixed with 10ml of phosphate buffer. 1ml of this extract was then added with 1ml of ninhydrin and 1ml

of pyridine solution. Obtained mixture was then boiled at 100 °C for 30 mins and sample absorbance was checked at 570nm by using Cecil 2021 Photo spectrometry. Dubois et al. (1956) methodology was used for this purpose. 0.2g of fresh leaf material was mixed with 10ml of 80% ethanol. After filtration, 0.5ml of filtrate was mixed with 0.5ml of distilled water and 1ml of 18% phenol solution. After keeping the mixture as such for an hour, 2.5ml of sulphuric acid was added. Absorbance was noted at 490nm.

Antioxidant Enzymes:

Fresh leaf material was grounded in liquid nitrogen, crushed with pestle mortar, and powdered material was stored at -80°C for further use. Methodology given by Aebi (1984) was used to check catalase activity. Reaction mixture contained 20µl of enzymes extract mixed with 50mM of potassium phosphate buffer (pH 6.8), 1mM EDTA and 15mM H₂O₂. Catalase activity was measured by decrease in H₂O₂ in 1 min at the absorbance range of 240 nm.

Methodology given by Nakano and Asada (1981) was used to check peroxidase activity. Reaction mixture was prepared by dissolving 5 mM H₂O₂, 15 mM guaiacol and 40mM phosphate buffer having pH 6.8. H₂O₂ was added and absorbance at 470nm was recorded for 1 minute. SOD activity was estimated according to the methodology given by Sohail et al. (2018). 0.5g of plant material was grinded in sodium phosphate buffer. Centrifugation for 15 minutes at 15000rpm was then performed. 0.1ml of extract was mixed with 0.1ml riboflavin and 3ml SOD buffer. Another test tube set was taken having 0.1ml riboflavin and 3ml SOD buffer, without plant extract. Plant extract containing test tubes was placed under fluorescent lamp to initiate the reaction while the 2nd set of test tubes was kept in dark. Absorbance of both sets at 560 nm wavelength was recorded.

APX content was measured using the methodology of Nakano and Asada (1981). For this purpose, 150 µL enzyme extract was taken along with phosphate buffer (50 mM), ascorbic acid (0.5 mM), and EDTA (0.1 mM). Light absorption was read at 290 nm using spectrophotometer. Enzyme activity was calculated by measuring changes in absorption rate per minute, per gram of fresh weight of sample.

Extraction and quantification of Melatonin (MET) and salicylic acid (SA)

Sturtz et al. (2011) method will be used for measuring plant melatonin contents. 2ml acetone will be added in 1g of grounded plant tissue. Mixture will be shaken at 25C for half an hour followed by its centrifugation at 4C for 15 minutes. High performance liquid chromatography (HPLC) system will be used for quantitative analysis of melatonin in plant samples.

Methodology given by Khan et al (2020b) was used for the estimation of endogenous salicylic acid activity. 0.3g of freeze-dried plant tissue was first mixed with 100% methanol, followed by its centrifugation at 10,000 x g. The extract was then dried and treated with 2.5mL of 5% trichloroacetic acid (TCA). Resulting solution was again centrifuged at 10,000 x g. cyclopentane and isopropanol were then

used for separation of supernatant, after which quantification was done by using HPLC at 305 and 365 nm.

Statistical Analysis:

The experiment was executed in a Completely Randomized Design (CRD) with two factors (variety and treatment) with 3 replications for each treatment. Pearson's correlation co-efficient and principal component analysis was plotted by using statistical program "R (v 4.0.4)" (R Development Core Team 2020), and Costat-v6.303 (Cohort software, Monterey, CA, USA) was used to execute two-way analysis of variance (ANOVA) and Duncan's multiple range test (DMRT) at $p \leq 0.001$ significance level for comparison of means (Kim 2014; Steel et al. 1997).

Results

Morphological Attributes:

Drought stress affected different morphological attributes of plants, and 19% reduction was recorded in shoot lengths of both canola cultivars as compared to control (Fig. 1-a). The application of MET and SA caused a significant increase in plant height in both normal and stressed conditions. MET priming showed better results among all treatments, as an increase of 22% in plant height was recorded in MET primed (T_1), super canola plants compared with non-treated drought-stressed plants (Fig. 1-a). Plant height was also improved by 17% in plants treated with MET priming synergetic with foliar spray of SA (T_7). Shoot fresh and dry biomass also showed a declining trend when plants were exposed to drought and a reduction of about 14% in fresh biomass and 17% in dry weight was recorded in both canola cultivars (Fig. 1-c, d). Application of plant growth regulators, reduced to some extent the negativity caused by drought on shoot fresh and dry biomass. MET priming showed better results under drought, where about 46% better fresh biomass and 40% better dry biomass was recorded in both canola cultivars (Fig. 1-c, d). These results were statistically non-significant from the plants treated with combined application of melatonin priming and salicylic acid foliar spray. Overall, Super Canola (V1) showed better growth under drought than Faisal Canola (V2).

Photosynthetic Attributes:

Photosynthetic rate showed a significant reduction (about 43% in both cultivars) after canola cultivars were exposed to drought. Seed priming treatments with plant growth regulators improved the photosynthetic rate of both cultivars under normal as well as stressed conditions (Fig. 2). Seed priming treatment with MET (T_1) caused an increase of 43% in photosynthetic rate of both canola cultivars, followed by those plants which were treated with combined MET and SA (T_5) priming treatment (Fig. 2-a). Both cultivars of canola showed a declining trend in transpiration rate (73%) and stomatal conductance (55%) after exposure to drought (Fig. 2-b, c). Seed priming treatment with MET (T_1) and SA (T_3) showed better results and improved transpiration rates and stomatal conductance in stressed

conditions (Fig. 2-b, c). Foliar spray of SA (T4) and combined foliar treatment of SA and MET (T6) slightly improved all photosynthetic attributes in stressed environment, but their impact was not as significant as of other treatments involved. Among the cultivars, Super Canola (V1) showed better results than Faisal Canola (V2).

Water Relations:

Declining trend was recorded in water and osmotic potential values of canola plants after exposure to drought (Fig. 3-a, b). Seed priming and foliar spray application of both plant growth regulators improved water potential and osmotic potential under both controlled and stressed conditions. Maximum water and osmotic potential values were recorded in MET primed plants (T1), followed by synergetic treatment of MET priming and SA foliar spray (T7). Combined foliar spray treatment (T6) of both growth regulators didn't show as significant effect on these parameters as shown by other treatments involved (Fig. 3-a, b).

Bio-Chemical Attributes:

An increase in compatible solutes concentration i.e., soluble proteins, free amino acids, and total soluble sugars were recorded after drought treatment in canola (Fig. 4). An increase of 15% was recorded in protein contents under drought, compared to water-treated plants. These Proteins contents were further increased by MET and SA application, and the maximum increase was recorded in MET primed (T1) plants (Fig. 4-a), where an increase of 25% in protein contents of both canola cultivars was recorded under drought, compared to non-treated drought-stressed plants (T0). Similarly, about 20% more free amino-acid contents were recorded in drought stress, as plants accumulate more amino acids to cope with this stress. Synergetic treatment of MET priming and SA foliar spray (T7) further enhanced these amino-acid contents where about 34% more amino-acid activity was found compared with non-treated (T0) water-stressed plants (Fig. 4-b). A slight increase (5%) was recorded in total soluble sugar contents in drought-stressed cultivars, compared to well-watered plants. Priming treatments with plant growth regulators further enhanced these contents, and MET priming treatment combined with SA foliar spray (T7) showed better results under drought. Foliar spray of SA (T4) and combined foliar treatment of SA and MET (T6) slightly improved all biochemical attributes under drought stressed conditions, but their impact was not as significant as of all other treatments involved (Fig. 4-c). Among the cultivars, Super Canola (V1) showed better biochemical activity than Faisal Canola (V2).

Antioxidant activity:

Antioxidant levels i.e., Peroxidase (POD), Superoxide Dismutase (SOD), Catalase (CAT) and Ascorbate peroxidase (APX) activity increased in drought exposed canola plants, compared to well-watered plants (Fig. 5). An increase of 23% in POD levels was observed in drought-stressed canola cultivars. Application of plant growth regulators further enhanced POD contents under both normal and stressed conditions. Maximum results were shown by synergetic treatment of MET as seed priming and SA as foliar spray (T7) under both experimental conditions (Fig. 5-a). These results were statistically non-significant from MET priming treated plants (T1). Similarly, an increase of 20% in SOD contents was recorded in drought-stressed plants. MET priming combined with SA foliar spray (T7) further enhanced these contents and

about 17% increase under normal and 28% increase under stressed conditions was recorded (Fig. 5-b). CAT and APX contents also showed a significant increase (27% & 42% respectively) after drought exposure (Fig. 5-c, d). MET priming (T1) further increased catalase contents by 35% and APX contents by 42% in stressed environment and by 14% and 36% respectively, under normal conditions, in both canola cultivars under study. Foliar spray of SA (T4) and synergetic foliar treatment of SA and MET (T6) slightly improved SOD, POD, CAT and APX levels in stressed environment, but their impact was not as significant as of all other treatments involved (Fig. 5-a, b, c, d). Among the cultivars, Super Canola (V1) showed better antioxidant activity than Faisal Canola (V2).

Yield Attributes:

Yield attributes of canola showed significant reduction after exposure to drought stress, where about 30% less silique per plant and 18% and 30% lower number of seeds per silique were recorded in Super and Faisal canola respectively, as compared to control (Fig. 6-a, b). However, a considerable increase (49% and 47%) in silique per plant and an increase of 40% and 56% in the number of seeds per silique was recorded in Super Canola (V1) and Faisal canola (V2) respectively, after treating plants with MET and SA under drought. Maximum no. of silique per plant under drought conditions were shown by MET priming treated plants (T1), followed by plants treated with SA (T3) priming (Fig. 6-a). Similarly, the maximum number of seeds per silique was recorded in those plants which were treated with MET priming combined with foliar spray of SA (T7) in a stressed environment (Fig. 6-b). Foliar spray of SA (T4) and combined foliar treatment of SA and MET (T6) slightly improved plant yield attributes in both controlled and stressed environment, but their impact was not as significant as of other treatments involved. Among the cultivars, Super Canola (V1) showed slightly better yield than Faisal Canola (V2).

Hormonal activity:

Exogenous application of MET didn't cause any significant increase in endogenous levels of MET and SA under normal, well irrigated conditions (Fig. 7a). Much increase in MET levels was observed in the plants exposed to drought, and maximum MET contents were recorded in MET priming treated plants (T1), followed by plants treated with synergetic priming application of MET and SA (Fig. 7a). Endogenous SA contents were also positively influenced by exogenous MET application (Fig. 7b, T1), which showed that MET application does have some sort of effect on endogenous SA activity in plant tissues. Similarly, non-significant effect of exogenously applied SA on endogenous activity of MET and SA was observed under controlled well-watered conditions (Fig. 7b). Exposure to drought stress caused marked increase in endogenous SA levels, and maximum SA content was recorded in SA priming treated plants (T3). Obtained results suggested that both MET, and SA are involved in responses of canola cultivars to drought stress, and exogenously applied MET and SA could result in changing accumulation patterns of endogenous MET and SA under stress.

Relationship and multivariate analysis:

In order to verify the relationship between different studied attributes, we have plotted Pearson correlation and principal component analysis (PCA-biplot) (Fig. 8–9). This analysis disclosed correlation (positive

and negative) among various parameters of canola cultivars under drought and normal conditions. It has been noticed that drought stress imposed significant impact on all the parameters. Figure 8 (a-b) showed significantly positive correlations among biochemical attributes, enzymatic antioxidant, phyto-hormones and plant water relations both under drought and controlled conditions. On the other side, biochemical parameters showed a negative correlation with root length, and root biomass. Plant height and fresh biomass were positively impacted by water relation attributes, but transpiration rate and stomatal conductance showed negative relationship with them. Overall transpiration rate (E) had negatively correlation with all other attributes under control conditions but showed almost zero relationship with number of seeds per silique and number of siliqua per plant, while photosynthetic rate had shown positive correlation both under drought as well as controlled conditions due to application of melatonin and salicylic acid. Similarly, Melatonin and salicylic acid had shown positive impact on plant growth and photosynthetic characteristics under normal and drought stressed conditions (Fig. 8a, b).

Our Pearson's correlation among various attributes of both canola varieties have been further validated by PCA-Biplot. Figure 9 represents PCA-Biplot of different attributes in canola subjected to drought stress with foliar and priming seed treatments of growth regulators. Principal component analysis executed the degree of association between variables under drought and controlled environment. Both dimensions of PCA i.e., Dim-1 and Dim-2 jointly elaborated 83.6% variability in the dataset. They showed obvious and marked separation of attributes under stressed and non-stressed conditions. Dim-1 showed 50.8%, while Dim-2 observed 32.8% variation to the total variance. There was a clear separation in studied parameters under normal and drought stress. Biochemical and antioxidant traits showed more relationship and impact under drought while growth attributes showed better relation in controlled conditions. Moreover, melatonin and salicylic acid have shown positive impact in mitigation of drought stress (Fig. 9).

Discussion

Keeping in view the drastic effect of drought stress on plant growth, the present study was conducted to check out the efficacy of plant growth regulators (MET and SA) in enhancing morphological, biochemical, and physiological attributes of canola cultivars grown under drought stress conditions. The role of hormones in enhancing plant growth under stressed environments is a well-documented phenomenon.

Water stress is the major limiting factor responsible for the reduced growth and yield of crops (Sattar et al. 2021). Results obtained from our study showed all the growth attributes i.e., shoot and root length, plant fresh and dry biomass, as well as yield parameters, showed significant reduction after exposure to water stress in both canola cultivars. These results are in agreement with those obtained by Abd Elhamid et al. (2016); Dawood et al. (2019); Elewa et al. (2017) and Sadiq et al. (2018). Dawood and Sadak (2014) observed that canola growth and yield showed significant reduction after exposure to drought which might be due to the production of ROS in plant cells during stress. Drought negatively affects cell elongation, cell turgor, and cell volume, which together cause a decrease in plant height (Sadak et al. 2020). The negative effects of drought on fresh and dry biomass of shoot and roots of canola might be due to reduced rates of photosynthesis in drought exposed plants (Haq et al. 2014). Reduced yield in

canola cultivars might be due to the reduced photosynthetic rate under stress conditions. Reduced photosynthesis in drought-stressed canola leaves leads to lower accumulation of carbohydrates in mature leaves, resulting in their lower rates of transport to developing organs, so the reduction in yield results (Sadak et al. 2020). However, the reduction in growth and yield of canola caused by drought is significantly reduced by seed priming treatment with melatonin, as well as priming and foliar spray of salicylic acid. Melatonin caused enhanced growth and yield is reported by various researchers (Kaya et al. 2022; Eaisalou et al. 2021; Amjadi et al. 2021; Huang et al. 2019; Kabiri et al. 2018).

Melatonin acts both as a growth promotor (Arnao and Hernandez-Ruiz 2019) as well as a protector against abiotic stresses (Amjadi et al. 2021; Ahmad et al. 2019). Yield enhancing role of melatonin can be attributed to its role in enhancing photosynthesis, thus increasing translocation to sink from source organs. Melatonin application further causes ion homeostasis and stimulates vegetative growth, leading to increased seed yield in plants (Çolak 2018; El-Awadi et al. 2017; Li et al. 2017). SA application also resulted in much increase in morphological and physiological attributes under drought, which may be attributed to its role in increased uptake of nutrients and more photosynthetic rate in drought stressed plants (Abdelaal et al. 2020). Significant increase in yield is recorded in SA treated plants, as SA has major role in flower formation, leading to more production of grains. Similar sort of results was recorded by Abdelaal (2015), who observed significant increase in seed weight and pod number of faba beans under drought after SA application. This increase in yield in SA treated plants in stressed environments could be due to water reserving actions of SA in plant cells, which results in increased enzymatic actions in stressed conditions resulting in better metabolism and yield (Pirasteh-Anosheh et al. 2017).

Among the different adaptive mechanisms shown by plants to cope with water stress, the closure of stomata to reduce water loss is the primary one. This resulted in a reduced photosynthetic rate under drought stress due to decreased stomatal conductance (Ahmad et al. 2021). Recent research work on melatonin on different crop species shows that melatonin increases plant resistance to drought by improving stomatal functions, where reopening of stomata is recorded under drought after application of exogenous melatonin at optimum levels (Ahmad et al. 2021; Fereiduni et al. 2019; Sharma et al. 2020). Present research work showed a significant reduction in net photosynthetic rate and stomatal conductance in both canola cultivars after exposure to drought stress. Seed priming with melatonin and salicylic acid increased photosynthetic rate and stomatal conductance, both under normal and stressed conditions. Foliar spray of salicylic acid in combination with melatonin priming also showed statistically significant results, signifying the synergetic effect of these growth regulators on photosynthetic attributes under stressed conditions. Our results are in line with those obtained by Cui et al. (2017), who observed similar sort of melatonin effects on photosynthetic rate and stomatal conductance while working on stress-exposed wheat plants.

Exposure to abiotic stress results in excessive production of ROS inside plant tissues, which causes breakdown and peroxidation of membrane lipids (Kar 2011). For maintaining cell homeostasis, plants show different resistive mechanisms against this ROS-caused oxidative damage under abiotic stresses. Activation of many enzymes like SOD, CAT, APX and POD occurs which inhibit further production of ROS

in plant cells (Imran et al. 2021; Khan et al.

2020a <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8255075/> - CIT0065). SOD first converts O_2^- to H_2O_2 , which is then broken down to water molecules by the action of POD and CAT (Hu et al. 2016; Imran et al. 2021). Findings of present research work show declined enzymatic antioxidants levels in drought-stressed canola cultivars. However, pre-sowing seed treatments with melatonin enhanced enzymatic antioxidants activity under controlled conditions, and combined application of Melatonin as seed priming and SA as foliar spray showed a marked increase in CAT, POD, APX and SOD concentrations under water stress conditions. This might be attributed to the fact that melatonin application reduces H_2O_2 levels and electrolyte leakage, caused by drought stress (Li et al. 2015). Melatonin application, on one hand, stimulates antioxidant enzymes activity, and at the same time, it maintains intracellular H_2O_2 at a constant level (Cui et al. 2017; Imran et al. 2021). The present study also signifies the importance of foliar application of salicylic acid in enhancing ROS scavenging via stimulated antioxidant action, in combination with pre-sowing melatonin application. SA treated plants showed increased enzymatic antioxidant activities under stress, as SA boosts plant resistance to stress by reducing oxidative stress, resulting in better plant growth under drought (Mutlu et al. 2016). Our study, thus, confirms that these two plant growth regulators act in a synergetic way to boost plant resistance against drought stress by enhancing antioxidant enzyme activity.

Defensive responses in plants to drought stress include accumulation of low molecular weight, osmotically active compounds called compatible solutes which include soluble proteins, free amino acids, and total soluble sugars (Ahmad et al. 2021; Ibrahim et al. 2020; Liu et al. 2018; Zhao et al. 2021). The present study showed an increase in levels of these compatible solutes under drought. Seed priming application of melatonin and salicylic acid caused a further significant increase in levels of these compatible solutes, highlighting the significance of MET and SA in coping with drought. Enhanced synthesis of amino acids and proteins, under drought, stabilize cellular structures and maintain osmotic pressure (Meng et al. 2014; Zhong et al. 2018; Zhao et al. 2021). An increase in proteins and amino acids by melatonin application shows that melatonin stimulates the synthesis of additional compatible compounds which ensure membrane stability at stress times (Georgiadou et al. 2018; Zhao et al. 2021). Although much reduction in photosynthetic rate is observed in drought-stressed plants, an important resistive mechanism in plants to drought is the accumulation of soluble sugars. Different researchers working on different crop plants exposed to drought recorded an increase in total soluble sugar contents (Elewa et al. 2017; Ezzo et al. 2018; Sadak and Bakry 2020). Seed priming with MET as well as synergetic application of MET seed priming and SA foliar spray further enhanced soluble sugar contents in drought-stressed canola. Elevated levels of soluble carbohydrates maintain turgor and stabilize cellular membranes by removing ROS (Hosseini et al. 2015; Sadak and Bakry 2020).

Conclusion

Present research work has proved that exogenous application of melatonin and salicylic acid has the potential to induce drought tolerance in canola plants, though melatonin appeared to be more effective

than salicylic acid. The synergetic application of melatonin seed priming and salicylic acid foliar spray showed significant effect on plant growth and can be effective in inducing drought resistance in canola. Among the cultivars, Super canola showed better growth and yield than Faisal canola, so Super canola can be a better choice in drought affected areas and can be used for future studies.

Declarations

Conflict of interest:

The authors declare no conflict of interest.

Data availability:

Not applicable

Declaration of competing interest:

Not applicable

Author contribution:

NI: Conceived the Idea and designed the experiment; **NR: Set and performed the experiment, took data,** wrote first draft; **NR, NI,** and **MA:** Writing, Review, editing; **NR,** and **MA:** analyzed the data, and helps in interpretation of results; **NI:** Critically revised the MS and provide technical support. All authors approved the final version of the manuscript.

Acknowledgment:

This paper is a part of PhD research work of the first author. Authors are highly thankful to the Higher Education of Pakistan (HEC) for providing scholarship and funding for this project. Authors would like to pay their gratitude to PMAS-Arid Agriculture University for providing experimental station and lab facilities.

References

1. Abd Elhamid EM, Sadak MS, Tawfik M (2016) Physiological response of Fenugreek plant to the application of proline under different water regimes. *Res J Pharm Biol Chem Sci* 7: 580-594.
2. Abdelaal KAA, Attia KA, Alamery SF, El-Afry MM, Ghazy AI, Tantawy DS, Al-Doss AA, El-Shawy E-SE, M. Abu-Elsaoud A, Hafez YM (2020) Exogenous Application of Proline and Salicylic Acid can Mitigate the Injurious Impacts of Drought Stress on Barley Plants Associated with Physiological and Histological Characters. *Sustainability* 12: 1736. <https://doi.org/10.3390/su12051736>
3. Abdelaal KAA (2015) Effect of Salicylic acid and Abscissic acid on morpho-physiological and anatomical characters of faba bean plants (*Vicia faba* L.) under drought stress, *J Plant Prod* 6: 1771-

1788.

4. Aebi H (1984) Catalase in vitro. *Methods in enzymology* 105: 121-126.
5. Ahmad S, Kamran M, Ding R, Meng X, Wang H, Ahmad I, Fahad S, Han Q (2019) Exogenous melatonin confers drought stress by promoting plant growth, photosynthetic capacity and antioxidant defense system of maize seedlings. *Peer J* 7: e7793.
6. Ahmad S, Muhammad I, Wang GY, Zeeshan M, Yang L, Ali I, Zhou XB (2021) Ameliorative effect of melatonin improves drought tolerance by regulating growth, photosynthetic traits and leaf ultrastructure of maize seedlings. *BMC Plant Biol* 21: 1-14.
7. Ahmadi M, Bahrani M (2009) Yield and yield components of rapeseed as influenced by water stress at different growth stages and nitrogen levels. *Americ-Eur J Agric Environ Sci* 5: 755-761.
8. Amjadi Z, Namdjoyan S, Abolhasani Soorki A (2021) Exogenous melatonin and salicylic acid alleviates cadmium toxicity in safflower (*Carthamus tinctorius* L.) seedlings. *Ecotoxicol* 30: 387-401. doi: 10.1007/s10646-021-02364-y.
9. Arnao MB, Hernandez-Ruiz J (2019) Melatonin and reactive oxygen and nitrogen species: A model for the plant redox network. *Melatonin Res* 2: 152-168.
10. Bates LS, Waldren RP, Teare I (1973) Rapid determination of free proline for water-stress studies. *Plant Soil* 39: 205-207.
11. Capell B, Doerffling K (1993) Genotype-specific differences in chilling tolerance of maize in relation to chilling-induced changes in water status and abscisic acid accumulation. *Physiol Plant* 88: 638-46. doi:10.1111/j.1399-3054.1993.tb01383.x.
12. Colak AM (2018) Effect of melatonin and gibberellic acid foliar application on the yield and quality of Jumbo blackberry species. *Saudi J Biol Sci* 25: 1242-1246.
13. Cui G, Zhao X, Liu S, Sun F, Zhang C, Xi Y (2017) Beneficial effects of melatonin in overcoming drought stress in wheat seedlings. *Plant Physiol Biochem* 118: 138-149.
14. Dawood M, El-Awadi M, Sadak M, El-Lethy S (2019) Comparison between the physiological role of carrot root extract and β -carotene in inducing *Helianthus annuus* L. drought tolerance. *Asian J Biol Sci* 12: 231-241.
15. Dawood MG, Sadak MS (2014) Physiological role of glycinebetaine in alleviating the deleterious effects of drought stress on canola plants (*Brassica napus* L.). *Middle East J Agric Res* 3: 943-954.
16. Dubois M, Gilles KA, Hamilton JK, Rebers PT, Smith F (1956) Colorimetric method for determination of sugars and related substances. *Analytic Chem* 28: 350-356.
17. Eaisalou AV, Namdjoyan S, Soorki AA (2021) The combined effect of melatonin and salicylic acid improved the tolerance of safflower seedlings to zinc toxicity. *Acta Physiol Plant* 43, 138. <https://doi.org/10.1007/s11738-021-03311-4>
18. El-awadi M, Dawood M, Abdel-Baky Y, Hassan EA (2017) Physiological effect of melatonin, IAA and their precursor on quality and quantity of chickpea plants grown under sandy soil conditions. *Agric Eng Int: CIGR J*: 35-44.

19. Elewa TA, Sadak MS, Saad AM (2017) Proline treatment improves physiological responses in quinoa plants under drought stress. *Biosci Res* 14: 21-33.
20. Ezzo M, Ebtihal M, Elhamid A, Sadak MS, Abdalla AM (2018) Improving drought tolerance of Moringa plants by using trehalose foliar treatments. *Biosci Res* 15: 4203-4214.
21. Fereiduni E, Ghasemi A, Elbestawi M (2019) Characterization of composite powder feedstock from powder bed fusion additive manufacturing perspective. *Materials* 12: 3673.
22. Georgiadou EC, Kowalska E, Patla K, Kulbat K, Smolinska B, Leszczynska J, Fotopoulos V (2018) Influence of heavy metals (Ni, Cu, and Zn) on nitro-oxidative stress responses, proteome regulation and allergen production in basil (*Ocimum basilicum* L.) plants. *Front Plant Sci* 9: 862.
23. Haq T, Ali A, Nadeem SM, Maqbool MM, Ibrahim M (2014) Performance of canola cultivars under drought stress induced by withholding irrigation at different growth stages. *Soil Environ* 33: 43-50.
24. Hasanuzzaman M, Matin MA, Fardus J, Hasanuzzaman MD, Hossain, MS, Parvin K (2019) Foliar application of salicylic acid improves growth and yield attributes by upregulating the antioxidant defense system in *Brassica campestris* plants grown in lead-amended soils. *Acta Agrobot* 72:1765. <https://doi.org/10.5586/aa.1765>
25. Hernandez-Ruiz J, Arnao MB (2018) Relationship of Melatonin and Salicylic Acid in Biotic/Abiotic Plant Stress Responses. *Agron* 8:33.
26. Hosseini SM, Hasanloo T, Mohammadi S (2015) Physiological characteristics, antioxidant enzyme activities, and gene expression in 2 spring canola (*Brassica napus* L.) cultivars under drought stress conditions. *Turk J Agric Forest* 39: 413-420.
27. Hu Z, Fan J, Xie Y, Amombo E, Liu A, Gitau MM, Khaldun A, Chen L, Fu J (2016) Comparative photosynthetic and metabolic analyses reveal mechanism of improved cold stress tolerance in bermudagrass by exogenous melatonin. *Plant Physiol Biochem* 100: 94-104.
28. Huang B, Chen Y-E, Zhao Y-Q, Ding C-B, Liao J-Q, Hu C, Zhou L-J, Zhang Z-W, Yuan S, Yuan M (2019) Exogenous melatonin alleviates oxidative damages and protects photosystem II in maize seedlings under drought stress. *Front Plant Sci* 10: 677.
29. Ibrahim MF, Elbar OHA, Farag R, Hikal M, El-Kelish A, El-Yazied AA, Alkahtani J, El-Gawad HGA (2020) Melatonin counteracts drought induced oxidative damage and stimulates growth, productivity and fruit quality properties of tomato plants. *Plants* 9: 1276.
30. Imran M, Latif Khan A, Shahzad R, Aaqil Khan M, Bilal S, Khan A, Kang S-M, Lee I-J (2021) Exogenous melatonin induces drought stress tolerance by promoting plant growth and antioxidant defence system of soybean plants. *AoB Plants* 13: plab026.
31. Kabiri R, Hatami A, Oloumi H, Naghizadeh M, Nasibi F, Tahmasebi Z (2018) Foliar application of melatonin induces tolerance to drought stress in Moldavian balm plants (*Dracocephalum moldavica*) through regulating the antioxidant system. *Folia Horti* 30: 155.
32. Kamran M, Wennan S, Ahmad I, Xiangping M, Wenwen C, Xudong Z, Siwei M, Khan A, Qingfang H, Tiening L (2018) Application of paclobutrazol affect maize grain yield by regulating root morphological and physiological characteristics under a semi-arid region. *Scient Rep* 8: 1-15.

33. Kar RK (2011) Plant responses to water stress: role of reactive oxygen species. *Plant Sig Behav* 6: 1741-1745.
34. Kaya C, Sarioglu A, Ashraf M, Nasser M, Ahmad AP (2022) The combined supplementation of melatonin and salicylic acid effectively detoxifies arsenic toxicity by modulating phytochelatins and nitrogen metabolism in pepper plants. *Environ Polut* 297: 118727.
<https://doi.org/10.1016/j.envpol.2021.118727>.
35. Khan A, Numan M, Khan AL, Lee I-J, Imran M, Asaf S, Al-Harrasi A (2020a) Melatonin: Awakening the defense mechanisms during plant oxidative stress. *Plants* 9: 407.
36. Khan A, Khan AL, Imran M, Asaf S, Kim YH, Bilal S, Numan M, Al-Harrasi A, Al-Rawahi A, Lee IJ (2020b) Silicon-induced thermotolerance in *Solanum lycopersicum* L. via activation of antioxidant system, heat shock proteins, and endogenous phytohormones. *BMC Plant Biol* 20: 1-18.
37. Kim HY (2014) Analysis of variance (ANOVA) comparing means of more than two groups. *Rest Dent End* 39: 74.
38. Lee HY, Byeon Y, Tan DX, Reiter RJ, Back K (2015) *Arabidopsis* serotonin N-acetyltransferase knockout mutant plants exhibit decreased melatonin and salicylic acid levels resulting in susceptibility to an avirulent pathogen. *J Pin Res* 58: 291-299.
39. Li C, Tan D-X, Liang D, Chang C, Jia D, Ma F (2015) Melatonin mediates the regulation of ABA metabolism, free-radical scavenging, and stomatal behaviour in two *Malus* species under drought stress. *J Exp Bot* 66: 669-680.
40. Li X, Yu B, Cui Y, Yin Y (2017) Melatonin application confers enhanced salt tolerance by regulating Na⁺ and Cl⁻ accumulation in rice. *Plant Growth Reg* 83: 441-454.
41. Liang D, Ni Z, Xia H, Xie Y, Lv X, Wang J, Lin L, Deng Q, Luo X (2019) Exogenous melatonin promotes biomass accumulation and photosynthesis of kiwifruit seedlings under drought stress. *Sci Hortic* 246: 34-43.
42. Liu Z, Cai J-s, Li J-j, Lu G-y, Li C-s, Fu G-p, Zhang X-k, Liu Q-y, Zou X-l, Cheng Y (2018) Exogenous application of a low concentration of melatonin enhances salt tolerance in rapeseed (*Brassica napus* L.) seedlings. *J Integ Agric* 17: 328-335.
43. Martinez V, Nieves-Cordones M, Lopez-Delacalle M, Rodenas R, Mestre TC, Garcia-Sanchez F, Rubio F, Nortes PA, Mittler R, Rivero RM (2018) Tolerance to stress combination in tomato plants: New insights in the protective role of melatonin. *Molec* 23: 535.
44. Meng JF, Xu TF, Wang ZZ, Fang YL, Xi ZM, Zhang ZW (2014) The ameliorative effects of exogenous melatonin on grape cuttings under water-deficient stress: antioxidant metabolites, leaf anatomy, and chloroplast morphology. *J Pin Res* 57: 200-212.
45. Mustafa H, Hasan E, Mahmood T, Aftab M (2017) Quantitative and Qualitative Evaluation of Rapeseed (*Brassica napus* L.) genotypes for the development of high yielding canola quality cultivars. *Discovery* 53: 380-387.
46. Mutlu S, Okkes N, Barbaros ME (2016) Exogenous salicylic acid alleviates cold damage by regulating antioxidative system in two barley (*Hordeum vulgare* L.) cultivars. *Front Life Sci*: 1-10.

10.1080/21553769.2015.1115430.

47. Naeem M, Naeem MS, Ahmad R, Ahmad R, Ashraf MY, Ihsan MZ, Nawaz F, Athar H-u-R, Ashraf M, Abbas HT (2018) Improving drought tolerance in maize by foliar application of boron: water status, antioxidative defense and photosynthetic capacity. *Arch Agron Soil Sci* 64: 626-639.
48. Nakano Y, Asada K (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* 22: 867-880.
49. Onemli F (2014) Fatty acid content of seed at different development stages in canola on different soil types with low organic matter. *Plant Prod Sci* 17: 253-259.
50. Osakabe Y, Arinaga N, Umezawa T, Katsura S, Nagamachi K, Tanaka H, Ohiraki H, Yamada K, Seo S-U, Abo M (2013) Osmotic stress responses and plant growth controlled by potassium transporters in *Arabidopsis*. *The Plant Cell* 25: 609-624.
51. Osakabe Y, Osakabe K, Shinozaki K, Tran L-SP (2014) Response of plants to water stress. *Front Plant Sci* 5: 86.
52. Pirasteh-Anosheh H, Emam Y, Rousta M, Ashraf M (2017) Salicylic Acid Induced Salinity Tolerance Through Manipulation of Ion Distribution Rather than Ion Accumulation. *J Plant Growth Reg* 36: 10.1007/s00344-016-9633-y.
53. Peleg Z, Blumwald E (2011) Hormone balance and abiotic stress tolerance in crop plants. *Current Opinion Plant Biol* 14: 290-295.
54. R Development Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <http://www.R-project.org>. Accessed 15 December, 2021.
55. Sadak MS, Abdalla AM, Abd Elhamid EM, Ezzo M (2020) Role of melatonin in improving growth, yield quantity and quality of *Moringa oleifera* L. plant under drought stress. *Bulletin of the National Research Centre* 44: 1-13.
56. Sadak MS, Bakry BA (2020) Alleviation of drought stress by melatonin foliar treatment on two flax varieties under sandy soil. *Physiol Mol Biol Plants* 26: 907-919. doi: 10.1007/s12298-020-00789-z.
57. Sadiq M, Akram NA, Ashraf M (2018) Impact of exogenously applied tocopherol on some key physio-biochemical and yield attributes in mungbean [*Vigna radiata* (L.) Wilczek] under limited irrigation regimes. *Acta Physiol Plant* 40: 1-14.
58. Sattar A, Wang X, Abbas T, Sher A, Ijaz M, Ul-Allah S, Irfan M, Butt M, Wahid MA, Cheema M (2021) Combined application of zinc and silicon alleviates terminal drought stress in wheat by triggering morpho-physiological and antioxidants defense mechanisms. *Plos one* 16: e0256984.
59. Scholander P, Hammel H, Bradstreet E, Hemmingson E (1965) Sap pressure in vascular plants. *Plant Sci* 148:339–45.
60. Sharma A, Wang J, Xu D, Tao S, Chong S, Yan D, Li Z, Yuan H, Zheng B (2020) Melatonin regulates the functional components of photosynthesis, antioxidant system, gene expression, and metabolic pathways to induce drought resistance in grafted *Carya cathayensis* plants. *Science of the Total Environment* 713: 136675.

61. Sohail AU, Shad S, Ilyas N, Manaf A, Raja NI (2018) In vitro germination and biochemical profiling of *Brassica napus* in response to biosynthesised zinc nanoparticles. IET Nanobiotech 13: 46-51.
62. Steel RG, Torrie JH, Dickey DA (1997) Principles and procedures of statistics. McGraw-Hill, New York. USA.
63. Sturtz M, Cerezo AB, Cantos-Villar E, Garcia-Parrilla MC (2011) Determination of the melatonin content of different varieties of tomatoes (*Lycopersicon esculentum*) and strawberries (*Fragaria ananassa*). Food Chem 127: 1329–1334. doi: 10.1016/j.foodchem.2011.01.093
64. Tariq M, Saad H, Mustafa HS, Aftab M, Ali Q, Malik D (2019). Super Canola: Newly Developed High Yielding, Lodging and Drought Tolerant Double Zero Cultivar of Rapeseed (*Brassica napus* L.). 19. 16039951. 10.4238/gmr16039951.
65. Urban MO, Vasek J, Klima M, Krtkova J, Kosova K, Prasil IT, Vitamvas P (2017) Proteomic and physiological approach reveals drought-induced changes in rapeseeds: Water-saver and water-spender strategy. J Proteom 152: 188-205.
66. Xu Z, Jiang Y, Jia B, Zhou G (2016) Elevated-CO₂ response of stomata and its dependence on environmental factors. Front Plant Sci 7: 657.
67. Zaid A, Mohammad F, Wani SH, Siddique KMH (2019) Salicylic acid enhances nickel stress tolerance by up-regulating antioxidant defense and glyoxalase systems in mustard plants. Ecotoxicol Environ Saf 180:575–587
68. Zhao C, Guo H, Wang J, Wang Y, Zhang R (2021) Melatonin Enhances Drought Tolerance by Regulating Leaf Stomatal Behavior, Carbon and Nitrogen Metabolism, and Related Gene Expression in Maize Plants. Front Plant Sci 12: 779382-779382.
69. Zhong C, Cao X, Bai Z, Zhang J, Zhu L, Huang J, Jin Q (2018) Nitrogen metabolism correlates with the acclimation of photosynthesis to short-term water stress in rice (*Oryza sativa* L.). Plant Physiol Biochem 125: 52-62.

Figures

Figure 1

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) plant height, (b) Root Length, (c) Plant Fresh Biomass, and (d) Plant Dry Biomass of two canola cultivars under normal and drought stress. Small letter above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T_0), MET priming (T_1), MET foliar spray (T_2), SA priming (T_3), SA

foliar spray (T_4), MET plus SA combined priming (T_5), MET plus SA combined foliar spray (T_6), MET priming combined with SA foliar spray (T_7), SA priming combined with MET foliar spray (T_8).

Figure 2

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) Photosynthetic Rate, (b) Transpiration Rate, (c) Stomatal Conductance and (d) Water Use Efficiency of two canola cultivars under normal and drought stress. Small letter above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T_0), MET priming (T_1), MET foliar spray (T_2), SA priming (T_3), SA foliar spray (T_4), MET plus SA combined priming (T_5), MET plus SA combined foliar spray (T_6), MET priming combined with SA foliar spray (T_7), SA priming combined with MET foliar spray (T_8).

Figure 3

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) Water Potential and (b) Osmotic Potential of two canola cultivars under normal and drought stress. Small letter

above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T_0), MET priming (T_1), MET foliar spray (T_2), SA priming (T_3), SA foliar spray (T_4), MET plus SA combined priming (T_5), MET plus SA combined foliar spray (T_6), MET priming combined with SA foliar spray (T_7), SA priming combined with MET foliar spray (T_8).

Figure 4

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) Total Proteins, (b) Total Amino acids, and (c) Total Soluble Sugars of two canola cultivars under normal and drought stress. Small letter above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T_0), MET priming (T_1), MET foliar spray (T_2), SA priming (T_3), SA foliar spray (T_4), MET plus SA combined priming (T_5), MET plus SA combined foliar spray (T_6), MET priming combined with SA foliar spray (T_7), SA priming combined with MET foliar spray (T_8).

Figure 5

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) POD activity, (b) SOD activity, (c) Catalase activity and (d) APX activity of two canola cultivars under normal and drought stress. Small letter above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T_0), MET priming (T_1), MET foliar spray (T_2), SA priming (T_3), SA foliar spray

(T₄), MET plus SA combined priming (T₅), MET plus SA combined foliar spray (T₆), MET priming combined with SA foliar spray (T₇), SA priming combined with MET foliar spray (T₈).

Figure 6

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) Number of Siliqua per plant and (b) Number of seeds per Siliqua of two canola cultivars under normal and drought stress. Small letter above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T₀), MET priming (T₁), MET foliar spray (T₂), SA priming (T₃), SA foliar spray (T₄), MET plus SA combined priming (T₅), MET plus SA combined foliar spray (T₆), MET priming combined with SA foliar spray (T₇), SA priming combined with MET foliar spray (T₈).

Figure 7

Effect of different treatments of priming and foliar spray of Melatonin and Salicylic acid on (a) Melatonin content and (b) SA content of two canola cultivars under normal and drought stress. Small letter above each bar shows statistical differences at $p < 0.05$. Treatments include: controlled irrigation (T₀), MET priming (T₁), MET foliar spray (T₂), SA priming (T₃), SA foliar spray (T₄), MET plus SA combined priming (T₅), MET plus SA combined foliar spray (T₆), MET priming combined with SA foliar spray (T₇), SA priming combined with MET foliar spray (T₈).

Figure 8

Pearson correlation under (a) control, and (b) drought condition showing relationship and impact of different measured attributes on each other. PH: Plant Height; RL: Root Length; PFB: Plant Fresh Biomass; PDB: Plant Dry Biomass; WP: Water Potential; OS: Osmotic Potential; NOS-P: No. of Siliqua Per Plant; NOG-S: No. of seeds per Siliqua; A: Photosynthetic Rate; E: Transpiration Rate; gs: Stomatal Conductance; WUE: Water Use Efficiency; TP: Total Proteins; TAA: Total Amino acids; TSS: Total Soluble Sugars; POD: Peroxidase; SOD: Superoxide Dismutase; CAT: Catalase; APX: Ascorbate Peroxidase, MET: Melatonin, SA: Salicylic acid

Figure 9

Principal component analysis (PCA) showing the relationship between different attributes of canola cultivars under normal and drought stress condition. PH: Plant Height; RL: Root Length; PFB: Plant Fresh Biomass; PDB: Plant Dry Biomass; WP: Water Potential; OS: Osmotic Potential; NOS-P: No. of Siliqua Per Plant; NOG-S: No. of seeds per Siliqua; A: Photosynthetic Rate; E: Transpiration Rate; gs: Stomatal Conductance; WUE: Water Use Efficiency; TP: Total Proteins; TAA: Total Amino acids; TSS: Total Soluble Sugars; POD: Peroxidase; SOD: Superoxide Dismutase; CAT: Catalase, APX: Ascorbate Peroxidase, MET: Melatonin, SA: Salicylic acid

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1.docx](#)