Knowledge user survey and Delphi process to inform development of a new risk of bias tool to assess systematic reviews with network meta-analysis (RoB NMA tool)

Carole Lunny (✉ carole.lunny@ubc.ca)

St Michael's Hospital, Unity Health Toronto <https://orcid.org/0000-0002-7825-6765>

Areti Angeliki Veroniki

Institute of Health Policy Management and Evaluation, University of Toronto

Brian Hutton

Ottawa Hospital Research Institute <https://orcid.org/0000-0001-5662-8647>

Ian R White

MRC Clinical Trials Unit at UCL <https://orcid.org/0000-0002-6718-7661>

Julian PT Higgins

Population Health Sciences, Bristol Medical School, University of Bristol

James M Wright

Cochrane Hypertension Review Group and the Therapeutics Initiative, University of British Columbia

Ji Yoon Kim

McGill University

Sai Surabi Thirugnanasampanthar

University of Toronto

Shazia Siddiqui

Lorenzo Moja

World Health Organisation

Nichole Taske

NICE, UK

Robert Lorenz

Federal Joint Committee (G-BA)

Savannah Gerrish

Simon Fraser University

Sharon Straus

Dalla Lana School of Public Health & Institute of Health Policy, Management, and Evaluation, University of Toronto

Virginia Minogue

Health Service Executive <https://orcid.org/0000-0002-0171-8642>

Franklin Hu

University of British Columbia

Ayah Kapani

University of British Columbia

Samin Nagi

University of British Columbia

Lillian Chen

University of British Columbia

Mona Akbar-nejad

University of British Columbia

Andrea C. Tricco

Knowledge Translation Program, Li Ka Shing Knowledge Institute, St. Michael's Hospital, Unity Health Toronto

Research Article

Keywords: collaboration, network-meta-analysis, knowledge user engagement, co-production, quality tool, Health Systems Research

Posted Date: February 13th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-1324758/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Network meta-analysis (NMA) is increasingly used in guideline development and other aspects of evidence-based decision-making. We aimed to develop a risk of bias (RoB) tool to assess NMAs (RoB NMA tool). An international steering committee intended the RoB NMA tool to be used in combination with ROBIS (recommend as it was designed to assess biases only) or other similar quality appraisal tools (e.g., AMSTAR 2) to assess quality of systematic reviews. The RoB NMA tool will assess NMA biases and limitations regarding how the analysis was planned, data were analysed, and results were presented, including the way in which the evidence was assembled and interpreted.

Objectives: Conduct (a) a Delphi process to determine expert opinion on an item's inclusion, and (b) a knowledge user survey to widen its impact.

Design: Cross sectional survey and Delphi process

Methods: Delphi panellists were asked to rate whether items should be included. All agreed-upon item were included in a second round of the survey (defined as 70% agreement). We surveyed knowledge users' views and preferences about the importance, utility, and willingness to use the RoB NMA tool to evaluate evidence in practice and in policymaking. We included 12 closed and 10 open-ended questions, and we followed a knowledge translation plan to disseminate the survey through social media and professional networks.

Results: 22 items were entered into a Delphi survey of which 28 respondents completed round 1, and 22 completed round 2. Seven items did not reach consensus in round 2. A total of 298 knowledge users participated in the survey (14% respondent rate). 75% indicated that their organization produced NMAs, and 78% showed high interest in the tool, especially if they had received adequate training (84%). Most knowledge users and Delphi panellists preferred a tool to assess *both* bias in individual NMA results *and* authors' conclusions. Response bias in our sample is a major limitation as knowledge users working in higher income countries were more represented. One of the limitations of the Delphi process is that it depends on the purposive selection of experts and their availability, thus limiting the variability in perspectives and scientific disciplines.

Conclusions: This Delphi process and knowledge user survey informs the development of the RoB NMA tool.

Registration: We uploaded our study protocol on the Open Science Framework at <https://osf.io/da4uy/>

Summary

What is already known about this subject?

The development of new tools to inform evidence-based medicine requires the feedback of knowledge users and experts. The purpose of the knowledge user survey and Delphi process was to ask respondents about the structure of a proposed tool for assessing biases in an network meta-analysis (NMA), and about the concepts related to bias in NMAs to potentially include in the tool.

What are the new findings?

The majority of knowledge users and Delphi respondents preferred a tool to assess the bias in NMA results and in authors' conclusions. Delphi respondents agreed to potentially include 15 out of 22 concepts about bias in NMAs.

How might it impact clinical practice in the foreseeable future?

This Delphi and knowledge user surveys inform the development of a new RoB NMA tool.

Introduction

Guidance about how to systematically develop quality and bias assessment tools is well established (Moher [1] and Whiting [2]), multi-staged, and includes the involvement of knowledge users and experts. The benefits of engaging knowledge users in tool development is a key factor associated with knowledge translation and the reduction of research waste [3-8]. Specifically, the benefits include: greater public acceptance [9]; identifying and prioritising topics for research [10]; providing feedback on the tool's usability [10]; wider dissemination, uptake and communication of findings [10]; and increased likelihood of impact [10, 11]. Identifying an external group of experts to obtain a multitude of perspectives will produce a more valid tool than a judgment given by an individual expert, or a group of experts heavily involved in the development process. Engaging with knowledge users and experts during development ensures that new tools will be relevant and applicable.

The risk of bias in network meta-analysis (NMA) tool project aims to develop the first tool to assess risk of bias (RoB) in a review with NMA. We intended the RoB NMA tool to be used in combination with ROBIS [12] (which we recommend as it was designed to assess biases specifically) or other similar tools (e.g., AMSTAR 2 [13]) to assess quality of systematic reviews. The RoB NMA tool will assess NMA biases and limitations regarding how the analysis was planned, data were analysed, and results were presented, including the way in which the evidence was assembled and interpreted. Our proposed RoB NMA tool has several uses. It can help knowledge users: (i) decide whether to believe the results from a single NMA; and (ii) help choose between NMAs on the same question with discordant results or conclusions.

Development of the tool follows five stages. In the first stage, we conducted and published a methodological review to identify items related to bias in NMAs [14]; second, the steering group made conceptual decisions about the type of tool that will be developed, and refined the items into concepts from the methodological review; third, a knowledge user survey was developed to solicit feedback on the structure of the tool from potential users; and fourth, expert opinion was obtained through a Delphi survey to select and define the concepts. The final and future stage will involve compile the items into a tool; and conducting pilot testing to refine the items in the tool.

In this paper, we report on the knowledge user survey and Delphi process. We define “knowledge user” as an individual who is likely to be able to use research results to make informed decisions about health policies, programs and/or practices [15]. A knowledge user can be, but is not limited to, a practitioner, a policy maker, an educator, a decision maker, a health care administrator, a community leader or an individual in a health charity, patient group, private sector organization or media outlet [15]. Our definition of experts is based on an individual's scientific/professional expertise, in our case in methods for NMAs, bias in systematic reviews with or without NMAs, and risk of bias tool development.

The purpose of the knowledge user survey was to ask knowledge users about the structure of our proposed tool and about their potential use of the tool in evidence-informed practice, policymaking, guideline development, or research. We also aimed to conduct a multi-round Delphi process to solicit expert opinion about concepts to potentially include in the tool.

Methods

Management of the project

At the start of our project to develop a risk of bias tool for NMAs, we first convened a steering group of nine experts in NMA, bias and tool development (**Appendix A**) [12]. The steering group is responsible for the management of the project and has executive power over all decisions related to the proposed tool, which is still under development.

Protocol

We uploaded our study protocol on the Open Science Framework at <https://osf.io/da4uy/>. The knowledge user survey complied with the Checklist for Reporting Results of Internet E-Surveys (**Appendix B**) [16]. Important definitions are found in **Box 1**.

A cross-sectional survey design was used for the knowledge user survey using Qualtrics [17, 18]. Approval from the University British Columbia Ethics Board was obtained for both the survey and the Delphi process, and consent was implied when respondents completed the online survey. Unique site visitors were identified via IP address and personal information was collected on a voluntary basis from respondents (no incentives were offered). Knowledge users and experts were identified using a purposive sampling strategy.

Box 1. Important definitions

Network meta-analysis (NMA)

We adopted a broad definition of an NMA as a method that aims to, or intends to, synthesise simultaneously the evidence from multiple studies investigating more than two health care interventions of interest. We used the Cochrane Handbook definition of an NMA: “*Any set of studies that links three or more interventions via direct comparisons forms a network of interventions. In a network of interventions there can be multiple ways to make indirect comparisons between the interventions. These are comparisons that have not been made directly within studies, and they can be estimated using mathematical combinations of the direct intervention effect estimates available [19].*” A network is composed by at least three nodes (interventions or comparators) and these are connected (graphically depicted as lines/edges) when at least one study compares the underlying two interventions - that is the direct comparisons. Reviews that intend to compare multiple treatments with an NMA but then find that the expectations or assumptions are violated (e.g. underlying assumptions of the method are not met), and hence an NMA is not possible or optimal, are also considered in our definition.

NMA risk of bias assessment

A risk of bias assessment would evaluate limitations in the way in which the NMA analysis was planned, analysed, and presented. If these methods are inappropriate, the validity of the findings can be compromised. Our tool aims either/or to assess the biases in the individual results of the NMA, and the authors’ conclusions.

Bias in results of an NMA

Network meta-analysis of effect estimates from primary studies can result in over-estimation or under-estimation of the effects of specific intervention comparisons [20, 21]. For example, Chaimani et al. conducted a network meta-epidemiological study and found that, in the majority of the 32 networks they analysed, small studies tended to exaggerate the true effect estimate of the intervention, possibly due to small-study effects and publication bias [22]. Our tool will focus on the results of an NMA [e.g. network characteristics (including geometry, effect modifiers) [23]. This is the approach taken in tools such as the Cochrane RoB 2 tool for assessing risk of bias in randomized trials [24].

Bias in the conclusions of an NMA

Bias may be introduced when interpreting the NMA results to draw conclusions. Conclusions may include ‘spin’ (e.g. biased mis-representation of the evidence, perhaps to facilitate publication) or (erroneous) mis-interpretation of the evidence [25]. Ideally, potential biases identified in the results of the NMA might be addressed appropriately when drawing conclusions. Similarly, a well conducted systematic review draws conclusions that are appropriate to the included evidence and can therefore be free of bias even when the primary studies included in the review have high risk of bias.

KNOWLEDGE USER SURVEY

Design

An English-language survey with 15 (12 closed and 10 open-ended) questions was developed by the investigative team (**Appendix C**). Five authors piloted the survey and modified it iteratively to improve content validity. Respondents were allowed to skip questions they did not wish to answer. The knowledge user survey ran from June 28 to August 1, 2021.

There were two parts to the survey: (1) demographic information and information about whether the knowledge users' organization used or produced NMAs; (2) purpose of the RoB-NMA tool, namely whether knowledge users preferred to assess the bias in the results, the authors' conclusions of an NMA, or both. Further sections asked about interest and engagement in development, piloting, dissemination, and training.

Email list development

We created an email list of journal editors publishing NMAs, using one bibliometric study of NMAs [26]. From this list, we extracted the journal names, and names of authors of NMAs. We also developed a list of organizations and institutions producing NMAs (**Appendix D**). We also included in the email list respondents from a UBC Methods Speaker Series on evidence synthesis methods (<https://www.ti.ubc.ca/2022/01/01/methods-speaker-series-2022/>).

Dissemination

All potential survey respondents were sent an email describing the purpose of the study, requesting their participation and providing a link to the survey (**Appendix E**). A knowledge translation plan was followed to disseminate and advertise the survey (**Appendix F**). We used twitter cards (i.e. advertisements with pictures) and targeted hashtags to increase awareness of the survey (see the Twitter Campaign in **Appendix F**). In addition, we advertised through the e-newsletters of Knowledge Translation Canada, SPOR Evidence Alliance and Therapeutics Initiative.

Data analysis

Questionnaires that were terminated early, where respondents did not go through all questionnaire pages, were included in analyses, but those that were entirely blank were excluded. We measured the time respondents took to fill in a questionnaire regardless of whether it was complete.

Descriptive statistics were calculated for each closed response question including count, frequency, *with* denominators taken as the number who provided a response to the question. One researcher coded the open-ended questions independently by identifying themes. Respondents' comments on questions 9 and 10 were merged as they were similar in nature.

DELPHI PROCESS

Design

Our published a methodological review to identify items related to bias in NMAs [14] identified 22 items related to bias in NMAs. These items were reworded into concepts by the steering committee because past Delphi groups focused on the wording of the item, and not its main idea. The concepts and questions about the structure of the tool (domains, signalling questions, rating scales) were entered into a Qualtrics survey platform.

Delphi panellists were asked to rated concepts based on a 5-point Likert scale of importance from 1 (not important - should be dropped as a concept to consider) to 5 (very important – must be included) or unable to score [27, 28]. If respondents did not provide a rating, the concept was recorded as missing. Respondents were asked to comment on whether they preferred to modify or reword the concepts [29]. Free text comment boxes allowed experts to provide additional comments. Non-responders or those failing to complete each round were sent three email reminders [28].

The respondents completed two survey rounds to reach a high level of agreement, defined as at least 70% scored 4 or above on the 5-point Likert scale [28] [30] (**Table 1**). After round 1, we generated reports of group verses individual responses. Respondents were also provided with anonymized free text comments from the last round [31].

Table 1. Decision criteria for inclusion, exclusion, and further consideration of potential concepts*

Scenario (rounds 1 and 2)	Handling of information
Concept scored 4-5 by ≥70% of respondents	Consensus achieved for potential inclusion in the RoB NMA tool. Further consideration in a subsequent Delphi round not needed.
Concept scored 3 by ≥70% of respondents	Include in Delphi round 2
Concept scored 1 or 2 (not important) by ≥70% of respondents	Consider excluding the concept
New concepts nominated by respondents	Include in round 2. Follow decision criteria scenarios above.

Data analysis

We reported the number of respondents completing each round. An overall response rate was calculated as well as summary statistics for each concept. The qualitative data from the free text questions was analysed through thematic analysis by one author (CL) and read by one or more of the other co-authors. The steering committee used their executive power to decide whether excluded concepts that reached consensus by the Delphi panel should be excluded or not.

Results

KNOWLEDGE USER SURVEY

Recruitment results

A total of 2,821 emails were sent out to advertise the survey, 87 failed to reach the recipients due to incorrect addresses, resulting in 2,734 emails that reached the intended individual (**Appendix H**). Most respondents completed the survey through our Qualtrics email survey link (n = 390, response rate = 14%) and 27 completed the survey through an anonymous link distributed over social media and e-newsletters (n = 27).

After consolidating duplicates (using IP addresses, n = 33) and blank responses (n = 86), a total of 298 responses were included in the analysis. Of the 298 respondents, 252 (85%) answered all the survey questions, and 46 (15%) completed half of the questions. The mean time to complete the survey was 2.27 minutes (SD 1.33).

Characteristics of respondents

Of the 298 respondents, 136 (45.6%) self-identified as a systematic review expert, 122 (40.9%) as a guideline developer, 98 (32.9%) as a health care professional (**Table 2**). Half of the respondents had primary affiliations at a university (50%). Most respondents resided in North America (41%) and/or Europe (34%) (**Table 2**). Three quarters (75%) of respondents indicated that their organization produced systematic reviews with NMAs, but only 54% of knowledge users said they used an NMA in their work (**Table 2**).

Table 2: Characteristics of knowledge user respondents and familiarity with NMAs

Characteristics	Overall (N=298)
Primary and current roles*	
Systematic reviewer	136 (45.6%)
Academic	122 (40.9%)
Clinician or health care professionals	98 (32.9%)
Graduate student/postdoctoral researcher	60 (25.1%)
Epidemiologist	54 (18.1%)
Guideline developer	44 (14.8%)
Independent researcher	42 (14.1%)
Health Technology Assessment (HTA) producer or specialist	39 (13.1%)
Statistician	38 (12.8%)
Journal editor	31 (10.4%)
Research support	19 (6.4%)
Decision/policymaker	9 (3.0%)
Information scientist/medical librarian	6 (2.0%)
Funding agency representative and clinician	3 (1.0%)
Patient partner	3 (1.0%)
Other (methodologist, non-profit organisation worker, knowledge translation specialist, scientific officer, health economist, etc.)	11 (3.7%)
Primary affiliation	
University	149 (50.0%)
Hospital and university hospital	61 (20.5%)
Research institute	25 (8.4%)
Government	19 (6.4%)
Non-profit organization (e.g., NGO, charity)	23 (7.7%)
For-profit private organization (e.g. industry)	10 (3.4%)
Other (e.g. clinic, HTA organization, blood service, independent researcher)	11 (3.7%)
Geographic region	
North America/Central America	121 (40.6%)
Europe	101 (33.9%)
Asia	50 (16.8%)
South America	17 (5.7%)
Africa	2 (0.67%)
Pacific Islands	1 (0.34%)
Australia	4 (1.3%)
Other (i.e. Middle East, Oceania)	2 (0.67%)
Does your organization or institution (or work colleagues) produce systematic reviews with NMA?	
Yes	223 (75.1%)
Unsure	32 (10.8%)
No	42 (14.1%)
Missing	1 (0.3%)
Have you used systematic reviews with NMA as a source of evidence in decision making?	
Yes	193 (65.4%)
No	73

	(24.7%)
Unsure	29 (9.83%)
Missing	3 (1.0%)
Have you used a systematic review with NMA in your work?	
Yes	160 (54.2%)
No	160 (54.2%)
Unsure	75 (25.4%)
Missing	3 (1.0%)
If you have used one or more systematic reviews with NMA in your work, did you use:	
Both individual results of NMAs and conclusions	127 (57.7%)
Individual analysis results from the NMA to draw your own conclusions (e.g. pooled effect estimate)	72 (32.7%)
NMA authors' conclusions	21 (9.55%)
Missing	78 (26.2%)

*Percentages add to more than 100% because respondents could provide more than one response.

HTA: Health Technology Assessment; NGO: Non-governmental organization; NMA: Network meta-analysis

Interest and type of tool preferred

Most knowledge users (84%) reported they would use the RoB NMA tool if they received adequate training on how to use it (**Figure 2**). When asked about their level of interest in our tool, 182/298 (61.1%) had high interest, 53/298 (17.8%) had low interest, and only one person had no interest. Many respondents said they would use the RoB NMA tool's bias assessment when conducting an overview of reviews, Health Technology Assessment (HTA), or guideline; and to distinguish between NMAs at higher or lower risk of bias.

When we asked knowledge users about the type of tool that might be useful to them or their organisation, half of the respondents (145/298) reported they preferred a tool to assess *both* the bias in individual NMA results and authors' conclusions (**Figure 1**).

Open-ended questions are summarised in **Appendix I Tables 1 to 4**. We also report in **Appendix I Table 4** respondents' interest in dissemination and engagement activities. The majority of respondents (153/231; 66%) said they would want to read the final study reports, receive updates (147/231; 64%), and receiving training in using the new tool (140/231; 61%).

DELPHI SURVEY

Recruitment results

The steering committee invited 53 experts to participate in the Delphi surveys, and 19 emails failed for various reasons, resulting in 37 emails that reached the intended individual. Of these, 28 completed round 1, and 22 completed round 2 (Flowchart in **Appendix J**). The response rate of panelists participating to our study was 28/37 (75.7%) in round 1, and 22/28 (78.6%) in round 2.

Characteristics of round 1 respondents

Of the 28 round 1 respondents, 15 (53.7%) self-identified as statisticians, 10 (35.7%) as academics, 4 (14.3%) as systematic review specialists or scientists, epidemiologists, or graduate students/postdoctoral researchers (**Table 3**). More than half of the respondents had a primary affiliation at a university (68%). Most respondents resided in Europe (53.6%) or North America (39.2%) (**Table 3**). Most (96.4%) respondents indicated that their organization produced systematic reviews with NMAs.

Table 3: Characteristics of Delphi round 1 respondents and expertise in NMAs

Characteristics	Overall (N=28)
Primary and current roles*	
Statistician	15 (53.7%)
Academic	10 (35.7%)
Systematic review specialist or scientist	4 (14.3%)
Epidemiologist	4 (14.3%)
Graduate student/postdoctoral researcher	4 (14.3%)
Clinician or allied health professional	2 (%)
HTA producer or specialist	1 (%)
Health economist	1 (%)
Independent researcher	1 (%)
Journal editor	1 (%)
Geographic region	1 (%)
North America/Central America	11 (39.2%)
Europe	15 (53.6%)
Australasia	3 (%)
Primary affiliation*	
University	19 (67.9%)
Research institute	5 (%)
Hospital	4 (%)
Government	1 (%)
For-profit private organization (e.g. industry)	4 (%)
Non-profit organization (e.g. NGO, charity)	1 (%)
Primary affiliation produces systematic reviews with NMA	27 (%)
Type of experience with NMAs	24 (%)
Subject-matter expert or experienced researcher with knowledge of a variety of evidence synthesis methods and practical experience with systematic reviews with NMA	23 (%)
Author with publications relevant to systematic reviews with NMA	19 (%)
Reader and user of systematic reviews with NMA	17 (%)
Author of a tool for systematic reviews with NMA	12 (%)
Perceived expertise in methods used in the conduct of systematic reviews with NMAs on a scale from 0 to 100, where 0 represents 'beginner' and 100 represents 'expert'	Mean 80.3 (SD 13.9)
Perceived expertise in identification of potential biases in NMAs on a scale from 0 to 100, where 0 represents 'beginner' and 100 represents 'expert'. -	Mean 71 (SD 22.7)

*Percentages add to more than 100% because respondents could provide more than one response.

HTA: Health Technology Assessment; NGO: Non-governmental organization; NMA: Network meta-analysis

Rating of concepts

Of the 22 concepts, seven did not reach consensus in round 2 (indicated in red in **Table 4**). **Table 4** lists all concepts in the left column that respondents rated from strongly disagree to strongly agree. The second column indicates whether the concept was included based on 70% of agreement (agree and strongly agree combined). The next columns indicate the number of responses over the denominator (number of people who answered) for each rating, percentage responses, and the group median. The list of concepts in Table 4 is not intended to be used to assess biases in NMAs, but to inform the development of items to be included in our tool.

Table 4: Rating of concepts by Delphi respondents in round 1 and 2

Domains	Concepts related to risk of bias in network meta-analysis	Round	Included (70% consensus)	Responses						Group Median (Q1, Q3)
Strongly Disagree	Disagree	Neither Agree nor Disagree	Agree	Strongly Agree	Unable to Score					
Network characteristics/geometry	1: Whether all interventions in the network (including comparators) were potentially suitable for all eligible study respondents	1	Y	0/27	0/27	2/27 (7%)	10/27 (37%)	15/27 (56%)	1	5 (4,5)
	2: Whether any interventions were inappropriately excluded from the network (e.g. through eligibility criteria or after seeing the results)	1	Y	0/27	1/27 (4%)	3/27 (11%)	11/27 (41%)	12/27 (44%)	1	4 (4,5)
	3: Whether importantly different intervention strategies were kept as distinct nodes in the network (i.e. whether appropriate groupings were made of interventions -- lumping vs splitting)	1	Y	0/28	2/28 (7%)	5/28 (18%)	11/28 (39%)	9/28 (32%)	0	4 (3,5)
Effect modifiers	4: Whether effect-modifying participant characteristics are sufficiently similar across the whole network	1	Y	0/28	0/28	2/28 (7%)	11/28 (39%)	15/28 (53%)	0	4.5 (4,5)
	5: Whether outcomes and time points are sufficiently similar across the whole network	1	Y	0/28	0/28	4/28 (14%)	14/28 (50%)	10/28 (36%)	0	4 (4,5)
	6: Whether study-level risks of bias are sufficiently similar across the whole network	1	N	0/28	9/28 (32%)	5/28 (18%)	8/28 (29%)	5/28 (18%)	0	3 (2,4)
		2	N	2/22 (9%)	8/22 (36%)	5/22 (23%)	5/22 (23%)	2/22 (9%)	0	3 (2,4)
	7: Whether other trial characteristics are sufficiently similar across the whole network	1	N	0/24	4/24 (17%)	9/24 (38%)	7/24 (29%)	4/24 (17%)	4	3 (3,4)
		2	N	0/21	4/21 (19%)	8/21 (38%)	8/21 (38%)	1/21 (5%)	1	3 (3,4)
Statistical synthesis	8: Whether an appropriate pre-specified approach was used in node making	1	Y	1/25 (4%)	1/25 (4%)	5/25 (20%)	13/25 (52%)	5/25 (20%)	3	4 (3,4)
	9: Whether a process was used to define nodes in the network (e.g. undertaken independently by 2 reviewers, following a pre-	1	N	0/25	2/25 (8%)	11/25 (44%)	9/25 (36%)	3/25 (12%)	3	3 (3,4)

Domains	Concepts related to risk of bias in network meta-analysis	Round	Included (70% consensus)	Responses						Group Median (Q1, Q3)
	planned node-making process)	2	N	1/20 (5%)	1/20 (5%)	5/20 (25%)	11/20 (55%)	2/20 (10%)	2	4 (3,4)
	10: Whether effect metric(s) for each outcome (e.g. odds ratios, risk difference) in the network were presented with confidence/credible intervals	1	N	3/28 (11%)	4/28 (14%)	2/28 (7%)	6/28 (21%)	13/28 (46%)	0	4 (2.25,5)
		2	N	3/22 (14%)	7/22 (32%)	2/22 (9%)	2/22 (9%)	8/22 (36%)	0	3 (2,5)
	11: If disconnected networks were connected to perform the analysis, whether methods to do this were appropriate	1	Y	0/25	2/25 (8%)	4/25 (16%)	13/25 (52%)	6/25 (24%)	3	4 (3.5,4)
	12: Whether methods used to represent multi-arm studies in the dataset and in the analysis are appropriate	1	Y	0/28	4/28 (14%)	3/28 (11%)	10/28 (36%)	11/28 (39%)	0	4 (3.25,5)
	13: Whether assumptions across the network about homogeneity/heterogeneity of effects within comparisons are appropriate	1	Y	0/27	1/27 (4%)	6/27 (22%)	13/27 (48%)	7/27 (26%)	1	4 (3,4)
	14: Whether a valid approach was used to determine whether there was conflict between direct and indirect sources of evidence on the same comparisons (often called inconsistency or incoherence)	1	Y	0/28	2/28 (7%)	0/28	8/28 (29%)	18/28 (64%)	0	5 (4,5)
	15: If inconsistency detected, then whether methods such as re-evaluation of the choice of scale, effect modification, and similarity of the contributing randomized controlled trials were investigated	1	N	0/28	4/28 (14%)	5/28 (18%)	14/28 (50%)	5/28 (18%)	0	4 (3,4)
		2	N	0/22	4/22 (18%)	4/22 (18%)	9/22 (41%)	5/22 (23%)	0	4 (3,4)
	16: If a Bayesian analysis was conducted, whether the selection of prior distributions was justified	1	Y	0/28	3/28 (11%)	2/28 (7%)	13/28 (46%)	10/28 (36%)	0	4 (4,5)

Domains	Concepts related to risk of bias in network meta-analysis	Round	Included (70% consensus)	Responses						Group Median (Q1, Q3)
	17: Whether the analysis appropriately addressed any differences in effect modifiers across different parts of the network	1	Y	0/28	2/28 (7%)	3/28 (11%)	18/28 (64%)	5/28 (18%)	0	4 (4,4)
	18: Whether there was evidence of conflicting results between direct and indirect evidence (often called inconsistency and incoherence in results)	1	Y	0/28	3/28 (11%)	0/28	10/28 (36%)	14/28 (50%)	0	4 (4,5)
	19: If there were conflicting results between direct and indirect evidence was this addressed appropriately (e.g. meta-regression, data extraction errors, redefining the network)	1	Y	0/27	2/27 (7.4%)	1/27 (4%)	13/27 (48%)	11/27 (41%)	1	4 (4,5)
	20: Evidence that the statistical model, as it was used to get the key results, was not suitable for the data (e.g. from analysis of residuals or information criteria such as DIC)	1	N	0/28	7/28 (25%)	6/28 (21%)	9/28 (32%)	6/28 (21%)	0	3.5 (2.25,4)
		2	N	0/14	1/21 (5%)	9/21 (43%)	7/21 (33%)	4/21 (19%)	1	4 (3,4)
	21: Whether sensitivity analyses demonstrate that findings were robust to the statistical model and estimation methods (including prior distributions if Bayesian methods were used)	1	N	0/28	2/28 (7%)	9/28 (32%)	8/28 (29%)	9/28 (32%)	0	4 (3,4.75)
		2	N	0/22	1/22 (5%)	8/22 (36%)	7/22 (32%)	6/22 (27%)	0	4 (3,4.75)
	22: Whether limitations at study and outcome level (e.g. risk of bias), and at review level (e.g., incomplete retrieval of identified research, reporting bias) were discussed	1	Y	0/27	1/27 (4%)	4/27 (15%)	8/27 (30%)	13/27 (48%)	1	4 (4,5)

* Blank responses and “Unable to score” responses were not counted in the denominator. DIC: Deviance Information Criterion)

Structure of the RoB NMA Tool

When asked about the structure of the RoB NMA tool, the majority of respondents agreed that a domain-based structure (25/28; 89.3%) with signalling questions (20/28; 71.4%) was preferred. The domain-based structure would be similar to that used in Cochrane Risk of Bias tool and the ROBIS tool. Signalling questions flag aspects of study design related to the potential for bias and aim to help reviewers judge risk of bias. They also agreed (19/28; 67.9%) that the steering committee should provide guidance on how to produce a risk of bias assessment for NMAs, outcomes within a network, or authors' conclusions of NMAs.

When asked about their preference for a tool to assess the risk of bias in NMA results and/or the author's conclusions, the majority of respondents (15/28; 53.6%) preferred a tool to assess bias in both results and conclusions, one third (10/28; 35.7%) preferred a tool to assess bias in the results only, and a minority (3/28; 10.7%) preferred to assess only the NMA authors conclusions.

Discussion

A majority of knowledge users responded that they had high interest in the RoB NMA tool if they received adequate training on how to use it and said they would use the tool to distinguish between NMAs at higher or lower risk of bias, and to assess an NMA in an overview of reviews, HTA, or guideline. Delphi respondents articulated a clear preference for a tool that is domain-based with signalling questions which would be used to assess biases in the results and the authors' conclusions. Seven out of 22 concepts did not reach consensus by the Delphi group, and these concepts and accompanying comments will be reviewed and considered by the steering committee for eligibility in the tool. The tool is still under development and the list of concepts is not intended to be used to assess biases in NMAs.

Respondents also indicated the need for guidance on how to use the tool to assess biases in the NMA. These results highlight the necessity for clear and easy to understand elaboration and explanation materials plus training, and perhaps the development of more structured guidelines for reaching domain-based RoB judgements (for example algorithms) [32]. Many knowledge users erroneously thought the RoB NMA tool's final assessment would be used in an evaluation of the certainty of the evidence (e.g. CINeMA [33] or GRADE [34]), even though we clearly stated that our proposed tool is intended for the assessment of the potential biases in an NMA. Only the quantitative results of an NMA (i.e. the analysis) are used in a certainty of the evidence evaluation.

We aimed to engage knowledge users and NMA experts early in the tool development for multiple reasons. Engagement would ensure that our tool will be relevant, useable and accepted [9]. The Delphi expert responses provided us with feedback on which concepts would be most relevant, and the knowledge user responses emphasised the desire for future training [10] and desire to help us with dissemination and communication of findings [10, 11].

Implications of this study

Knowledge user [5, 7, 35-38] and Delphi [39-43] surveys have been successfully used to inform the development of other types of tools, systematic reviews and guidelines. Online surveys have been used to evaluate the reliability and face validity of tools, and the use of a risk of bias tool in practice [32]. However, we are not aware of similar published surveys conducted prior to the development of a risk of bias or quality appraisal tool, targeted specifically at knowledge users.

Strengths and limitations

A strength of our research was that we developed a protocol which we followed in the conduct of the research (<https://osf.io/da4uy/>). We aimed to engage knowledge users and NMA experts early in the tool development for multiple reasons. Engagement would ensure that our tool will be relevant, useable and accepted [9]. The responses provided us with feedback on which concepts are most relevant, training needs [10] and future dissemination a findings. We combined newsletter, email distribution lists, and social media to reach a wide range of knowledge users from across the globe. We attempted to maximise the response rate by sending email reminders and repeating messages through social media. Response bias in our sample is a major limitation as knowledge users working in higher income countries were more represented, and respondents (i.e. systematic reviewers (45.6%), academics (40.9%) or clinician/ health care professionals) may have been more likely to have responded to a survey about a new tool to assess the bias in NMAs.

A limitation is we did not ask knowledge users to define what their role was and whether they considered themselves: (i) decision makers; (ii) purchasers of services/pharma products; (iii) professional service providers; (iv) evidence generators; or (v) advocates of health promotion. Another limitation is that our targeted emails and social media advertisement may have missed important knowledge users that use NMAs (e.g. members of the Canadian Institutes of Health Research [CIHR] Drug Safety and Effectiveness Network [DSEN] Methods and Applications Group for Indirect Comparisons [MAGIC] Group).

A strength of the Delphi process was that by performing the surveys online, experts from around the world were able to participate. The response rate of Delphi panellists participating to our study was high in both rounds. All comments were thoroughly read by one of the authors (CL), and part of them were read by one or more of the other co-authors. In addition to a full feedback report, a summary of the comments in round 1 was also given in the next round to panellists. Delphi processes have many limitations, one of which is that they depend too much on the purposive selection of "experts" and their availability, thus raising the question of whether all relevant perspectives and scientific disciplines have been taken into consideration. Our Delphi panel consisted of a small double digit number, which may risk collecting certain thought collectives and may be an issue of reliability [44]. 11

Future research

The results of the survey will inform a new tool to assess biases in NMAs. Our proposed RoB NMA tool will allow knowledge users to assess the biases in an NMA. Our tool is not targeted at authors of NMAs, as it does not outline methods that should be followed to conduct an NMA. It is targeted at knowledge users such as healthcare providers, policymakers, physiotherapists, who want to determine if the results of an NMA can be trusted to be at low risk of bias.

The steering committee will use the results of the knowledge user survey to determine preferences around the tools structure, and the Delphi process to choose and refine the concepts in the tool. Concepts will be reworded into signalling questions and

categorised into domains. The new tool will then be pilot tested with different knowledge user groups: patients, healthcare providers, and researchers. Further research will involve reliability and validity testing.

Conclusions

The knowledge user and Delphi surveys provided feedback from knowledge users and experts on their preferences for the structure, focus and concepts included in a proposed RoB NMA tool, which is under development. Both knowledge users and Delphi panelists preferred a tool to assess both the bias in individual NMA results and authors' conclusions. The majority of knowledge users had high interest in the tool and reported they would use it if they received adequate training.

Declarations

Availability of data and materials

The datasets used and/or analysed during the current study are available freely at <https://osf.io/da4uy/>.

Competing interest

AAV was an Associate Editor for the journal, but was not involved with the decision or peer-review process. All other authors declared that they had no competing interests. Brian Hutton has previously received honoraria from Eversana Incorporated for the provision of methodologic advice related to the conduct of systematic reviews and meta-analysis.

Patient and Public Involvement

No patients were involved in our study.

Ethics approval

This study involved human participants but the University of British Columbia Behavioural Research Ethics Board exempted this study (ID H20-02013).

Funding

Funding was received from a 2020 CIHR Project Grant (2021-2024; ID 451190). Andrea Tricco currently holds a Tier 2 Canada Research Chair in Knowledge Synthesis. Ian White was supported by the Medical Research Council [programme MC_UU_00004/06]. Brian Hutton has previously received honoraria from Eversana Incorporated for the provision of methodologic advice related to the conduct of systematic reviews and meta-analysis.

JPTH is a National Institute for Health Research (NIHR) Senior Investigator (NF-SI-0617-10145), is supported by NIHR Bristol Biomedical Research Centre at University Hospitals Bristol and Weston NHS Foundation Trust and the University of Bristol, is supported by the NIHR Applied Research Collaboration West (ARC West) at University Hospitals Bristol and Weston NHS Foundation Trust, and is a member of the MRC Integrative Epidemiology Unit at the University of Bristol. The views expressed

in this article are those of the authors and do not necessarily represent those of the NHS, the NIHR, MRC, or the Department of Health and Social Care.

Authors' contributions

CL conceived of the study; ACT, AAV, BH, CL, IW, JPTH, JMW, PW, SD contributed to the design of the study; CL drafted the survey; CL, LC, and SST inputted the questions into Qualtrics; CL, SS, SST analysed the data; CL wrote the draft manuscript; ACT, AAV, BH, CL, IW, JPTH, JMW, PW, SD revised the manuscript; all authors edited the manuscript; and all authors read and approved the final manuscript.

Acknowledgement

We acknowledge the contribution of our collaborators: Sofia Dias, Penny Whiting, Adrienne Stevens, Bob Nakagawa, Gayatri Jayaraman, Georgia Salanti, Karla Soares-Weiser, Larry Mróz, Lucy Turner, Nicole Mittmann, Nicolette McGuire, and Matthew Tunis. We also acknowledge our Delphi panellists: Simon Turner, Dimitris Mavridis, Virginia Chiocchia, Ian Shrier, Adriani Nikolakopoulou, Dan Jackson, Richard Riley, Becky Turner, Bruno Roza da Costa, Petros Pechlivanoglou, Steve Kanter, Ferran Catala-Lopez, Tianjing Li, Matt Page, Chris Cameron, Anna Chaimani, Kerry Dwan, Audrey Beliveau, Kristian Thorlund, Jonathan Sterne, Cinzia del Giovane, Guido Schwarzer, Joanne McKenzie, Lehana Thabane, Theodoros Papakonstantinou, Isabelle Bourtron, Sharon Strauss, Jenn Watt.

References

1. Moher, D., et al., *Guidance for developers of health research reporting guidelines*. PLoS medicine, 2010. **7**(2): p. e1000217.
2. Whiting, P., et al., *A proposed framework for developing quality assessment tools*. Syst Rev, 2017. **6**(1): p. 204.
3. Bullock, A., Z.S. Morris, and C. Atwell, *Collaboration between health services managers and researchers: making a difference?* Journal of health services research & policy, 2012. **17**(2_suppl): p. 2-10.
4. Hanney, S.R., et al., *The utilisation of health research in policy-making: concepts, examples and methods of assessment*. Health research policy and systems, 2003. **1**(1): p. 1-28.
5. Innvaer, S., et al., *Health policy-makers' perceptions of their use of evidence: a systematic review*. Journal of health services research & policy, 2002. **7**(4): p. 239-244.
6. Lavis, J., et al., *Towards systematic reviews that inform health care management and policy-making*. Journal of health services research & policy, 2005. **10**(1_suppl): p. 35-48.
7. Vokó, Z., et al., *Similarities and differences between stakeholders' opinions on using HTA information across five European countries*. Improving the Use of an Economic Decision Support Tool, 2018. **14**: p. 149.
8. Vokó, Z., et al., *Similarities and differences between stakeholders' opinions on using Health Technology Assessment (HTA) information across five European countries: results from the EQUIPT survey*. Health research policy and systems, 2016. **14**(1): p. 1-17.
9. Richards, C., Carter, C., Sherlock, K., *Practical approaches to participation*. 2004: Macaulay Institute.

10. Cottrell, E., et al., *Defining the benefits of stakeholder engagement in systematic reviews*. 2014, Agency for Healthcare Research and Quality (US): Rockville (MD).
11. Deverka, P.A., et al., *Stakeholder participation in comparative effectiveness research: defining a framework for effective engagement*. Journal of comparative effectiveness research, 2012. **1**(2): p. 181-194.
12. Whiting, P., et al., *ROBIS: A new tool to assess risk of bias in systematic reviews was developed*. J Clin Epidemiol, 2016. **69**: p. 225-34.
13. Shea, B.J., et al., *AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both*. bmj, 2017. **358**: p. j4008.
14. Lunny, C., et al., *Methodological review to develop a list of bias items used to assess reviews incorporating network meta-analysis: protocol and rationale*. BMJ Open, 2021. **11**(6): p. e045987.
15. CIHR, *Knowledge User Engagement*. 2016, Canadian Institute for Health Research.
16. Eysenbach, G., *Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES)*. J Med Internet Res, 2004. **6**(3): p. e34.
17. Gupta, K., *A practical guide to needs assessment*. 2011: John Wiley & Sons.
18. Dillman, D.A., *Mail and internet surveys*. 2nd edition ed. 2007, Hoboken, New Jersey: John Wiley & Sons Inc. 18.
19. Higgins, J.P., et al., *Cochrane handbook for systematic reviews of interventions*. 2019: John Wiley & Sons.
20. Mavridis, D., et al., *A selection model for accounting for publication bias in a full network meta-analysis*. Statistics in medicine, 2014. **33**(30): p. 5399-5412.
21. Mavridis, D., et al., *Publication bias and small-study effects magnified effectiveness of antipsychotics but their relative ranking remained invariant*. Journal of clinical epidemiology, 2016. **69**: p. 161-169.
22. Savović, J., et al., *Association Between Risk-of-Bias Assessments and Results of Randomized Trials in Cochrane Reviews: The ROBES Meta-Epidemiologic Study*. American Journal of Epidemiology, 2017. **187**(5): p. 1113-1122.
23. Cote, M.P., et al., *Understanding Network Meta-analysis (NMA) Conclusions Requires Scrutiny of Methods and Results: Introduction to NMA and the Geometry of Evidence*. Arthroscopy, 2021. **37**(7): p. 2013-2016.
24. Sterne, J.A., et al., *RoB 2: a revised tool for assessing risk of bias in randomised trials*. bmj, 2019. **366**.
25. Boutron, I. and P. Ravaud, *Misrepresentation and distortion of research in biomedical literature*. Proceedings of the National Academy of Sciences, 2018. **115**(11): p. 2613.
26. Ban, J.K., et al., *History and publication trends in the diffusion and early uptake of indirect comparison meta-analytic methods to study drugs: animated coauthorship networks over time*. BMJ open, 2018. **8**(6): p. e019110.
27. Tricco, A.C., et al., *An international survey and modified Delphi approach revealed numerous rapid review methods*. Journal of clinical epidemiology, 2016. **70**: p. 61-67.
28. Pandor, A., et al., *Delphi consensus reached to produce a decision tool for Selecting Approaches for Rapid Reviews (STARR)*. Journal of Clinical Epidemiology, 2019. **114**: p. 22-29.
29. Page, M.J., et al., *Mapping of reporting guidance for systematic reviews and meta-analyses generated a comprehensive item bank for future reporting guidelines*. J Clin Epidemiol, 2020. **118**: p. 60-68.

30. Stevens, A., et al., *Developing PRISMA-RR, a reporting guideline for rapid reviews of primary studies (Protocol)*. 2018, EQUATOR Network: Oxford, UK: <https://www.equator-network.org/wp-content/uploads/2018/02/PRISMA-RR-protocol.pdf>.
31. Boukredid, R., et al., *Using and Reporting the Delphi Method for Selecting Healthcare Quality Indicators: A Systematic Review*. PLOS ONE, 2011. **6**(6): p. e20476.
32. Savović, J., et al., *Evaluation of the Cochrane Collaboration's tool for assessing the risk of bias in randomized trials: focus groups, online survey, proposed recommendations and their implementation*. Systematic Reviews, 2014. **3**(1): p. 37.
33. Nikolakopoulou, A., et al., *CINeMA: An approach for assessing confidence in the results of a network meta-analysis*. PLoS Medicine / Public Library of Science, 2020. **17**(4): p. e1003082.
34. Puhan, M.A., et al., *A GRADE Working Group approach for rating the quality of treatment effect estimates from network meta-analysis*. BMJ, 2014. **349**: p. g5630.
35. Haddaway, N.R., et al., *A framework for stakeholder engagement during systematic reviews and maps in environmental management*. Environmental Evidence, 2017. **6**(1): p. 11.
36. Helbig, N., et al., *Stakeholder engagement in policy development: observations and lessons from international experience*, in *Policy practice and digital science*. 2015, Springer. p. 177-204.
37. Sultan, S., et al., *An International Needs Assessment Survey of Guideline Developers Demonstrates Variability in Resources and Challenges to Collaboration between Organizations*. Journal of general internal medicine, 2021: p. 1-9.
38. Sibbald, S.L., H. Kang, and I.D. Graham, *Collaborative health research partnerships: a survey of researcher and knowledge-user attitudes and perceptions*. Health research policy and systems, 2019. **17**(1): p. 92-92.
39. Booth, A., et al., *Establishing a minimum dataset for prospective registration of systematic reviews: an international consultation*. PloS one, 2011. **6**(11): p. e27319.
40. Harrison, R., et al., *Quality assessment with diverse studies (QuADS): an appraisal tool for methodological and reporting quality in systematic reviews of mixed-or multi-method studies*. BMC health services research, 2021. **21**(1): p. 1-20.
41. Mokkink, L.B., et al., *COSMIN Risk of Bias tool to assess the quality of studies on reliability or measurement error of outcome measurement instruments: a Delphi study*. BMC medical research methodology, 2020. **20**(1): p. 1-13.
42. Whiting, P., et al., *The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews*. BMC medical research methodology, 2003. **3**(1): p. 1-13.
43. Whiting, P., et al., *ROBIS: a new tool to assess risk of bias in systematic reviews was developed*. Journal of clinical epidemiology, 2016. **69**: p. 225-234.
44. Niederberger, M. and J. Spranger, *Delphi Technique in Health Sciences: A Map*. Front Public Health, 2020. **8**: p. 457.

Figures

Figure 1

Flowchart of emails sent and responses

Legend: Flowchart of emails sent and responses

Figure 2

(A) Use of a RoB NMA tool in knowledge users work, and (B) interest in a RoB NMAs tool

Legend: The left figure (A) depicts the proportion of responses to the question of whether knowledge users would use our proposed RoB NMA tool to assess the NMA analysis results, the authors conclusions, or both results and conclusions. Figure 1 (B) shows the proportion of responses to the question about interest in a tool for appraising risk of bias in NMAs.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AppendicesAtoJ20220226.pdf](#)