Relations between family cohesion and adolescent-parent's neural synchrony in response to emotional stimulations

Xinmei Deng (✉ xmdeng@szu.edu.cn)

School of Psychology, Shenzhen University

Mingping Lin

School of Psychology, Shenzhen University

Lin Zhang

School of Psychology, Shenzhen University

Xiaoqing Li

School of Psychology, Shenzhen University

Qiufeng Gao

Department of society, School of Law, Shenzhen University

Research Article

Keywords: EEG hyperscanning, family cohesion, adolescents, interbrain synchrony, emotional processing

Posted Date: February 8th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1324944/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The interaction between parent and adolescent is more challenging than other age periods. Family cohesion seriously impacts parent-adolescent emotional interactions. However, the underlying neural mechanism has not been fully examined. This study examined the differences in the neural synchrony in response to emotional film clips between high and low family cohesion adolescent-parent dyads by using the electroencephalograph (EEG) hyperscanning.

Results: Simultaneously electroencephalograph (EEG) was recorded while 15 low family cohesion parent-adolescent dyads (LFCs) and 14 high family cohesion parent-adolescent dyads (HFCs) received different emotional induction when viewing film clips. Interbrain phase-locking-value (PLV) in gamma band was used to calculate parent-adolescent dyads' interbrain synchrony. Results showed that higher gamma interbrain synchrony was observed in the HFCs than the LFCs in the positive conditions. However, there was no significant difference between the HFCs and LFCs in other conditions. Also, the HFCs had significantly higher gamma interbrain synchrony in the positive conditions than in the negative conditions.

Conclusion: Interbrain synchrony may represent an underlying neural mechanism of the parent-adolescent emotional bonding, which is the core of family cohesion.

1 Introduction

Family cohesion refers to the emotion, support, helpfulness and caring among family members (Olson, 2011). Adolescence can be considered a sensitive period for social development, with psychological and social cognitive changes (Blakemore & Mills, 2014). During adolescence, the social world and the interpersonal interactions become increasingly important. Previous research indicated that family cohesion may help adolescents to overcome the challenges of social interactions by receiving social and emotional supports from family members (Johnson et al., 2001). However, the underlying neural mechanism for the impacts of family cohesion on adolescent-parent emotional interactions has not been fully examined. As a biomarker of social and emotional interaction, the electroencephalogram (EEG) interbrain synchrony may represent the behavioral and emotional synchrony, which reflects the emotional bond within adolescent-parent dyads (Konvalinka & Roepstorff, 2012). Making further exploration of the impacts of family cohesion on the neural synchrony within adolescent-parent dyads during an emotional related task may shed light on the understanding of the biological base of social interaction. Therefore, this study examined the differences in the neural synchrony in response to different emotional stimulus simultaneously between the adolescent-parent dyads who had different level of family cohesion.

Family cohesion is defined as the emotional bonding that family members have toward each other (Olson & Craddock, 1980). It is the core component of family functioning. Previous studies indicated that balanced level of family cohesion (e.g., from moderately low to moderately high level) is beneficial and

viable for healthy family functioning (Olson, 2011). In a functional and cohesive family, family members can be emotionally connected to, provide support and caring for their families (Starke & Svensson, 2001).

Family cohesion has significant impacts on adolescents' behavior, socio-emotional development, well-being in their later life, and caregivers' parenting behaviors (Rahgozar et al., 2012; Behnke et al., 2008). It is highly related to individuals' positive well-being. For example, adolescents in families with a high level of family cohesion would report more positive emotional experience, more life satisfied, and more meaning and purpose in life (Fosco & Lydon-Staley, 2020). Adverse family processes may operate to increase adolescents' vulnerability to depression (Sheeber et al., 2001). Previous research also indicated that family cohesion is associated with a lower level of negative behavioral outcomes during development (e.g., externalizing behaviors) (Hong et al., 2021). The influence of family cohesion on children and adolescents' behavioral problems may be moderated by the caregivers' health condition and parenting behaviors. Family cohesion can provide important emotional support and bonding to reduce the negative impacts of the maladaptive parenting behavior on children's behavioral problems (Leidy et al., 2010). It is considered to be an important protective buffer as it meets the individual's psychological need for affiliation, security and emotional connection with others (Roth & Brooks-Gunn, 2000).

Not only the positive impacts of adolescents' behavior and mental health, family cohesion also has the protective impacts on social and emotional interaction for adolescents. Family cohesion is positively correlated with close emotional bonding and depend attachments. A cohesive family reflects a symbiotic and interdependent relationship between family members (Alavi et al., 2020). For example, a child's mental and emotional status would depend on and be impacted by the other members. Family cohesion is also related to the family member's feelings (i.e., loneliness), which reflect their family environment and the quality of the interactions with other family members (Johnson et al., 2001). Previous research indicated that family cohesion is negatively correlated with the parent-adolescent conflict (Sui et al., 2015). Higher family cohesion which is characterized as a higher family engagement is negatively correlated with the adolescents' depressive symptoms, higher levels of family support, and less psychological control from mothers (Starke & Svensson, 2001). Adolescents who grow in a family with moderate cohesion levels would have more positive communication and are more comfortable to be close with other during social interactions. Adolescents with high family cohesion would show better communication skills and feel less anxious and negative experiences during socio-emotional interactions (Olson, 2000). Previous research suggested that family cohesion is positively associated with parental support, responsiveness, and greater positive parent-child engagement. Family members have significantly more contact and communication with one another. It is possible that the family rules in a cohesive family make family members stay more in contact with one another (Reeves et al., 2020). On the contrary, a lack of family cohesion is associated with the negative parent-child interaction which may increase the social stress and reduce emotional support from the parents during social interaction (Hong et al., 2015). Family members with low cohesion would have less involvement in one another's lives and show less emotional bonding with one another (Olson, 2000).

Adolescence is characterized as the increased independence and the development of self. With the pursuit of independence out of the family environment, adolescents spend more time alone or with their peers, and the conflict affective intensity interactions between adolescents and their parents increases (Steinberg, 1991). As stated above, family cohesion may serve as a protective factor for parent-adolescent conflicts during socio-emotional interactions and seriously influence the effectiveness and quality of parent-adolescent emotional interactions (Johnson et al., 2001). The most important impacts of family cohesion could increase the psychological and emotional bonding between parent-adolescent dyads. During socio-emotional interactions, the behavioral, emotional, physiological and neural synchronization has been considered as an evolutionary adaptive outcome of interpersonal bonding (Hove & Risen, 2009). In this case, when engaging in social situations, family members in a high cohesive family might present a higher level of similarities in behavioral, emotional, physiological and neural levels because of their inherently biological and emotional bonding. However, there is little neural evidences to suggest that the level of family cohesion is associated with neural interbrain synchrony between adolescents and parents during socio-emotional interactions.

As the most important social relationship of each individual, the interaction between infant and caregiver is thought to be the first experience of social interaction. The interbrain synchrony between infant and caregiver in biological rhythms and social signals reflects the inherently biological and emotional bonding, which is also an important feature of the early brain matures (Djalovski et al., 2021). Previous research demonstrated that the moment-to-moment interbrain synchrony is a sensitive maker that can predict dynamic socio-emotional interactions (Dikker et al., 2017). It is likely driven by the shared attention and emotional processing mechanism within the dyads during moments of social contact and communication.

Generally, there are four types of interbrain synchrony, induced synchrony is associated with the phenomenon that two brains tend to be synchronized with the influence of common external stimuli (Burgess, 2013). For instance, two participants will show synchrony without information transmission and interaction, such as viewing a movie (Hasson et al., 2008). According to the phase resetting theory, each individual is sending out conspicuously social signals in the process of social interaction, such as gaze, voice, body posture. As the trigger of neural synchrony, these signals trigger the phase reset of the ongoing neural oscillations of the interacting parties, thus causing the phase difference between the two signal strings to remain constant (Wang et al., 2021). Previous research suggested that the interbrain synchrony, which can be measured by the EEG-based functional connectivity provides a useful tool for studying the simultaneous brain activity between dyads during interactions in different emotional states (Costa et al., 2006). For example, in a study, participants viewed film clips that evoked different emotional states (e.g., neutral, positive, or negative). There was an overall increase of interbrain synchrony during emotional stimulation (e.g., during viewing positive or negative film clips). However, the interbrain synchrony indices were significantly different among emotional states (Lee & Hsieh, 2014). Another study also found that there was a significant difference of hyper-connectivity existed in the gamma frequency band between positive and negative stimulus conditions (Zhu et al., 2018). During social interactions,

compared with the negative emotional states, the inter-brain network showed significantly higher strength for positive emotional states for the parent-infant dyads (Santamaria et al., 2020).

This biobehavioral synchrony could be a consequence of the coordination of physiological and behavioral processes when individual and their caregiver engagement in each other's social life (Djalovski et al., 2021). Increased interbrain synchrony facilitates emotional sharing, social understanding, psychological support, empathy between individual and their caregiver because of the accessibility to each other's internal state (Wheatley et al., 2012). In this case, such interbrain synchrony could be an important neural indicator of the cohesion level for the family members.

The quality and connection between parent and adolescents during social interaction could reflect the level of family cohesion and family atmosphere. However, little is known about the neural underpinnings about how the level of family cohesion is associated with the interpersonal neural connectivity between adolescents and parents. As mentioned above, induced synchrony could be reflected from the phenomenon that two brains tend to be synchronized with the influence of common external stimuli (Burgess, 2013). Thus, in the current study, the hyperscanning method was used to examine the neural synchrony in response to different emotional film clips between adolescent-parent dyads in different level of family cohesion. Because of the positive impacts of family cohesion on socio-emotional interactions between family members (Konrad, 2016), we hypothesized that high family cohesion parent-adolescent dyads would have high levels of interbrain synchrony in response to emotional film clips. Moreover, previous study demonstrated that the hyper-connectivity (e.g., in the gamma frequency band) in the positive stimulation condition was greater than in the negative stimulation condition (Zhu et al., 2018). We hypothesized that how the level of family cohesion associated with the interbrain synchrony between adolescents and parents would be different in the positive and negative emotional conditions.

2 Methods

2.1 Participants

Adolescent participants were recruited via flyers that invited healthy volunteers to participate in a study of parenting and emotion. Interested families were invited to visit the university laboratory to take part in the study. Final samples of the study were recruited on the basis of their scores on the Family Adaptability and Cohesion Scale, Chinese Version, (FACES-CV; Fei et al., 1991). The 16-item Family Cohesion Subscale of FACES-CV was used to assess parent-adolescent dyads' level of family cohesion and emotional bonding among family members on a 5-point Likert scale (1= never or very rarely true, 5= very often or always true; e.g., "All the family members get together for activities"). ($\alpha = .71$).

Given the poor EEG data quality and technical error, 2 parent-adolescent dyads were excluded from the study. Thus, the final sample was composed of 15 low family cohesion parent-adolescent dyads (LFCs; 11 male and 4 female, aged from 11 to 14 years old, $M_{\text{age}}=12.00$, $SD=1.25$, 6 mothers, $M_{\text{age}}=43.56$, $SD=5.43$; 9 fathers, $M_{\text{age}}=44.22$, $SD=4.63$) and 14 high family cohesion parent-adolescent dyads (HFCs;

11 male and 3 female, aged from 11 to 14 years old, $M_{\text{age}}=12.36$, $SD=1.08$, 7 mothers, $M_{\text{age}}=42.57$, $SD=1.90$; 7 fathers, $M_{\text{age}}=42.76$, $SD=3.21$). HFCs ($M_{\text{HFCs}}=76.93$, $SD=3.71$) and LFCs ($M_{\text{LFCs}}=66.33$, $SD=3.64$) differed significantly on the overall family cohesion score ($p<.001$). The sample size in this study was also in line with typical hyperscanning EEG studies (Yan et al., 2016).

All of the adolescents came from urban communities in Shenzhen city in China. Approximately 93.10% of fathers and 89.66% of mothers had received a college education, whereas other parents had received an education of high school or lower. All of the participants were right-handed and had a normal or corrected-to-normal vision, and were in good neurological and psychiatric condition. No participant had a history of neurological or psychiatric disorder, as determined by self- and/or parent report. The research protocol was approved by the local Institutional Review Board. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from the participants and their parents before the study, and the parent-adolescent dyads were fully debriefed after the experiment.

2.2 Stimuli

Six film clips were selected from the Chinese Affective Film System (CAFS; Xu et al, 2010). The film clips were age-appropriate for adolescents. Negative film clips consisted of 2 100-s segments from “My Beloved” and “My Sisters and Brothers”. The negative film clips were previously found to elicit a strong level of sadness (between 4 and 5 on the 5-point rating scale, $M=4.5$) with low variation across subjects (SD between 2 and 3, mean SD 2.5). The negative film clips included unpleasing social situations, such as being separated from loved ones. For example, in film clip of “My Sisters and Brothers”, the older brother had to give his youngest sister to others for adoption because of poverty. The youngest sister cried out and begged her brother not to abandon her. Positive film clips consisted of 2 100-s segments from “A big potato” and “Eat Hot Tofu Slowly”. The positive film clips were previously found to elicit a strong level of happiness (between 4 and 5 on the 5-point rating scale, $M=4.5$) with low variation across subjects (SD between 1 and 2, mean SD 1.5). For example, in the film clip of “Eat Hot Tofu Slowly”, a student sleeping in class was asked to sing a song by the teacher. Because his singing was out of tune, the whole class laughed. Neutral film clips consisted of 2 100-s segments from “Computer Maintenance” and “IP Encapsulation and IDE Interface Fixes”. The neutral film clips were previously found to elicit a medium level of emotional intensity (between 1 and 4 on the 5-point rating scale, $M=1.5$) with low variation across subjects (SD between 1 and 2, mean SD 1.5). The neutral film clips included one person completes the operation of repairing the computer and repairing the IDF interface. For example, in the film clip of “IP Encapsulation and IDE Interface Fixes”, an engineer turns the screws, removes the keyboard and trackpad, checks the wires and so on.

2.3 Procedures

We employed a 2 (Group: HFCs vs. LFCs) × 3 (Valence: positive vs. negative vs. neutral) repeated measures design. After receiving the demographic information from the parent-adolescent dyads, the parent-adolescent dyads were asked to see the film clips together. Electroencephalograph (EEG) sensors were attached on both of the parent-adolescent dyads and they were introduced to the procedures of the task before the experiment. After research assistant confirmed the parent-adolescent dyads fully understood the procedure, the experiment began. At the beginning of each trial, a fixation point was presented for one minute in the middle of the screen. Following the fixation, a film clip was presented for 100s. Then, participants were asked to rate each film clip on the valence and arousal scales of the self-assessment manikin (SAM, Lang et al., 2005), ranging from low pleasure or arousal (1) to high pleasure or arousal (5). The parent-adolescent dyads were told to rate the valence of each film clip based on how pleasant the film clip made them feel (e.g., How pleasant did you feel after viewing the film clip?). Participants were told to rate arousal based on the strength of their feelings in response to the film clip (e.g., How strongly did you feel after viewing the film clip?). The parents and adolescents didn't see each others' rating.

There were six film clips in the experiment. Each of the six film clips were displayed once in a random order. E-Prime software was used to present all stimuli against a black background on a 21-inch monitor, with a viewing distance of approximately 80 cm. An experimental session took 30-35 min for each participant (see Figure 1). The parent-adolescent dyads were fully debriefed the study after the whole study.

2.4 Dual-EEG Recording and Data Analysis

The parent-adolescent dyads were comfortably seated side by side in an electrically shielded and dimly lit room. Simultaneously and continuously EEG signals of each dyad during viewing six film clips were recorded using two 32-channel portable EEG systems (BrainAmp, Brainproducts GmbH, Germany) at a sampling rate of 500 Hz. Electrode impedance was kept under 30 kohms for all recordings. Online reference electrodes were placed at TP9 on parents' head. EEG signals were referenced offline to the averaged mastoid references and band-pass filtered in the range of 1 to 40 Hz. Epochs with eye movement or other movement artifacts were manually removed by inspection. Independent component analysis (ICA) was performed for ocular artifact reduction was performed by using EEGLAB. The artifact scored epochs were eliminated from all subsequent analyses. Onsets were set as the points where the film clips started. The artifact-free EEG signal from each trial was segmented from 0s before to 100 s after the onset of the film clips. The mean number of valid epochs for LFCs was 173.40 ($SD=24.7$) for positive stimuli, 184.07 ($SD=26.13$) for negative stimuli and 174.27 ($SD=29.03$) for neutral stimuli. The mean number of valid epochs for HFCs was 172.93 ($SD=34.52$) for positive stimuli, 180.29 ($SD=34.78$) for negative stimuli and 177.57 ($SD=25.72$) for neutral stimuli

In previous studies, inter-brain synchrony has been measured by the phase synchronization of EEG signals (Santamaria et al., 2020). We used an interbrain phase-locking-value (PLV) index to estimate the interbrain phase synchrony between parent-adolescent dyads in the positive, negative and neutral

conditions. This interbrain phase synchrony index has been developed to measure whether the signals from the two interacting individuals are phase locked across time (Lachaux et al., 1999). In line with previous hyperscanning research (Barraza et al., 2020), the band-pass filtering is applied to the EEG data of each segment of the parent's electrode (ϕ) and the adolescent's electrode (ψ) to obtain the EEG data of the frequency band of interest. The Hilbert transform is used to analyze the signals of ϕ and ψ poles in a specific frequency band, extract phase and subtract, and get PLV of each epoch. Average the PLV of electrodes ϕ and ψ in the frequency band of interest is obtained by average for epoch under each condition. Therefore, PLV can be expressed by formula, where N is the number of trials, $\phi_{(t,n)}$ is the phase on trial, n at time t , in channel ϕ and $\psi_{(t,n)}$ in channel ψ . The averaged interbrain PLVs in gamma frequency bands (31-40Hz) in the frontal (Fz), central (Cz) and parietal areas (Pz) were used for the inter-brain synchrony analyses. The averaged interbrain PLVs in the following four frequency bands were calculated for further statistical analysis: delta (1-3 Hz), theta (4-7 Hz), alpha (8-13 Hz), beta (14-30 Hz). Results of the averaged interbrain PLV in other frequency bands were shown in Appendix A. The frequency band has been considered to be highly related to emotional interaction (Kinreich et al., 2017; Mu et al., 2017). PLVs were averaged across the positive, negative and neutral trials.

$$PLV_t = T^{-1} \left| \sum_{n=1}^T e^{i[\phi(t,n) - \psi(t,n)]} \right|$$

The interbrain PLV were assessed using a 2 (group: HSAs vs. LSAs) $\times$ 3 (valence: positive vs. negative vs. neutral) $\times$ 3 (electrode: Cz vs. Fz vs. Pz) ANOVAs, with valence and electrode as within-subject variable and group as between-subject variable. The interbrain PLV were statistically evaluated using SPSS 20.0. The significance level was set at $p < .05$, and Greenhouse-Geisser correction was applied to p values associated with multiple-df comparisons. Partial eta squared was reported as a measure of effect size.

To rule out the possible impact of relevant variables of the parent-adolescents emotional interactions on our findings, demographic variables, adolescents' level of depression, adolescents' level of anxiety, adolescents' level of social support, parents' level of depression, parents' level of anxiety, and the level of parent involvement between the high and low social anxiety parent-adolescents dyads were examined. Repeated measures ANOVAs were used and these relevant variables were set as the covariates. The results of the repeated measures ANOVA indicated that there were no significant main effects of demographic variables and other examined variables on the interbrain PLV between the high and low social anxiety groups (see Appendix B).

3 Results

To examine the interbrain synchrony between the parent-adolescent dyads when watching positive, negative and neutral emotional films, we calculated the interbrain phase-locking-value (PLV) which has been developed to measure whether the signals from the two interacting individuals are perfectly phase locked across time.

Results of the repeated measures ANOVA on the interbrain phase synchrony in the gamma band showed that the main effect of Group was not significant, $F(1, 27) = .04, p = .852, \eta_p^2 = .001$. The main effect of Valence was not significant, $F(2, 54) = 1.43, p = .249, \eta_p^2 = .05$. The main effect of Electrode was also not significant, $F(2, 54) = .79, p = .430, \eta_p^2 = .03$. The interaction of Group and Valence was not significant, $F(2, 54) = .07, p = .933, \eta_p^2 = .00$. The interaction of Group and Electrode was not significant, $F(2, 54) = 1.40, p = .254, \eta_p^2 = .05$. The interaction of Valence and Electrode was also not significant, $F(4, 108) = .81, p = .520, \eta_p^2 = .03$.

The interaction of Group $\times$ Valence $\times$ Electrode was significant, $F(4, 108) = 3.00, p = .022, \eta_p^2 = .10$. Post hoc analysis reflected that at Pz (see Figure 2), greater gamma interbrain synchrony was observed in the HFCs than the LFCs in the positive conditions ($p=.002$). At Pz, higher gamma interbrain synchrony was observed in the positive conditions than in the negative conditions in the HFCs. ($p=.005$). In the HFCs, higher gamma interbrain synchrony was observed in the Pz than in the Cz ($p=.044$) and Fz ($p=.037$) in positive conditions. There were no significant differences between different conditions at other electrode in the LSAs and HSAs ($ps>.05$).

Table 1 *Gamma Interbrain Phase-locking-value (PLV) between Low and High Family Cohesion Parent-adolescent Dyads in Different Conditions at Fz, Cz and Pz*

Electrode	Conditions	LSAs	HSAs	<i>t</i>	<i>d</i>	<i>p</i>	95%CI	
		(<i>M</i> $\pm$ <i>SD</i>)	(<i>M</i> $\pm$ <i>SD</i>)					
Fz	Positive	.22 $\pm$.01	.22 $\pm$.01	1.30	.00	.205	-.003	.012
	Negative	.22 $\pm$.01	.22 $\pm$.01	-.42	.00	.676	-.010	.007
	Neutral	.22 $\pm$.01	.22 $\pm$.01	-.90	.00	.373	-.010	.004
Cz	Positive	.23 $\pm$.01	.22 $\pm$.01	.99	1.00	.333	-.004	.011
	Negative	.22 $\pm$.01	.22 $\pm$.01	-.48	.00	.635	-.010	.006
	Neutral	.22 $\pm$.01	.22 $\pm$.01	.62	.00	.544	-.006	.012
Pz	Positive	.22 $\pm$.01	.23 $\pm$.01	-3.52	-1.00	.002	-.004	.013
	Negative	.22 $\pm$.01	.22 $\pm$.01	1.10	.00	.282	-.004	.013
	Neutral	.22 $\pm$.01	.22 $\pm$.01	-.54	.00	.593	-.013	.008

Note. LFCs = low family cohesion parent-adolescent dyads, HFCs = high family cohesion parent-adolescent dyads.

4 Discussion

Family cohesion has been regarded as an important component of family functioning. It reflects the affective and supportive relationship among family members (Olson, 2011). Previous research has demonstrated the protective impact of family cohesion on the negative socio-emotional interactions (e.g., emotional conflicts) between parents and adolescents during “the period of storm and stress” (Konrad, 2016). Nevertheless, there is less evidence has described how family cohesion is related to the synchronous activations and the interbrain connectivity during the emotional interaction between adolescent-parent dyads by using the neuroscientific method. The present study explored the differences in the interbrain synchrony between adolescent-parent dyads during synchronous positive and negative emotional stimulation with different family cohesion levels. By identifying the psychophysiological evidence for this relationship between family cohesion and the interbrain synchrony during emotional interactions between parents and adolescents, we extended the understanding of the neural underpinnings of family cohesion that influence the socio-emotional interactions within family members.

Interbrain synchrony is demonstrated as a neural marker for social interaction. The moment-to-moment interactions between individual’s brains can be understood as a bidirectional behavioral stimulus-to-brain coupling (Konvalinka & Roepstorff, 2012). Even without the real information transmission and interaction, induced synchrony could be observed with the influence of common external stimuli according to the phase resetting theory (Burgess, 201; Hasson et al., 2008). In this case, the interbrain synchrony within the dyads in response to emotional film clips in our present study might reflect the degree of the emotional connection and the quality of shared attention between parents and adolescents during the socio-emotional processing at the neural level (Reindl et al., 2018). The simultaneous emotional stimulation might tap into the similar tendencies for emotional perception or responses across parents and adolescents. The results of the present study were consistent with the hypothesis that family cohesion is related to interbrain synchrony between the adolescent-parent dyads during emotional interactions. Specifically, we found that greater gamma interbrain synchrony in response to emotional film clips was observed in the HFCs than the LFCs in the positive conditions. However, there was not significant group difference between the HFCs and LFCs in the negative and neutral conditions. The emotional bonding that family members experience toward each other is the core of family cohesion. Low level of family cohesion suggests emotional disengagement, less emotional closeness, and high independence. High level of family cohesion suggests emotional engagement, high emotional support and dependence (Olson, 1986). Family cohesion increases the communication and the emotional bonding with each other, which may promote their interbrain synchrony and coupling (Babiloni, 2014). Moreover, previous research has found that children and adolescents in the positive emotional atmosphere families would tend to cooperate more and more efficiently with parents in positive emotional context compared with the individuals in negative atmosphere families (Konrad, 2016). This is maybe a possible explanation of the significant higher interbrain synchrony in HFCs compared with the LFCs in positive simulation conditions.

Also, we found a significant higher interbrain synchrony in HFCs in positive simulation conditions than in negative conditions. Previous research indicated emotional sharing and expressiveness is more apparent and easier for positive emotions than negative emotions, because of few social constraints regarding positive emotional expression (Halberstadt & Eaton, 2002). In this case, during socio-emotional

interactions, positive emotional expression would be easier between children and parents, in which may result in a higher level of emotional sharing and interbrain synchrony within dyads. In a higher cohesive family, when the mother has a more emotional engagement, the child would be more willing to express positive emotions than negative emotions. The parents' active participation will increase the child's safety attachment, which may result in the positive emotional sharing and expression to their parents (Bretherton, 1990; Diener & Mangelsdorf, 1999). In this case, significant higher interbrain synchrony between high family cohesion adolescent-parent dyads could be observed in the positive emotional contexts rather than the negative emotional contexts.

In the current study, we only found high and low family cohesion adolescent-parent dyads differences in the gamma band interbrain synchrony indexed by the PLV in the parietal areas (Pz) when the dyads responded to emotional film clips. Parietal Gamma activities associated with emotion processing, including emotion recognition, emotion regulation and empathic sharing of others' emotions. Gamma frequency bands in the frontal and parietal areas have been identified to provide discriminative information associated with emotion processing (Lin et al., 2010). Parietal Gamma activity may be involved in emotional regulation, as this process integrates the top-down cognitive reappraisal process and the bottom-up sensory-motional process (Kang et al., 2012). Right inferior parietal lobe is an important hub for both sensory-emotion integration and empathic sharing of others' emotions (Maffei et al., 2019). In line with the previous study, findings from the present study have also demonstrated that functional communication between brains during socio-emotional interaction (e.g., simultaneous emotional processing) relies on the oscillatory synchronization of gamma band responses (Liu et al., 2018; Martini et al., 2012). The gamma oscillations are functional and prominent in brain regions that underline emotional processing and functions (e.g., amygdala) (Stujenske et al., 2014). For example, research indicated that increase in gamma band activity and its synchronization in the amygdala, prefrontal, and posterior cingulate cortices to emotional relative to neutral stimuli (Luo et al., 2009; Matsumoto et al., 2006; Matsumoto et al., 2010). When individual experience negative social emotions (e.g., social threat), interbrain gamma synchrony associated with social coordination would be increased. Instead of triggering the alpha synchrony, the interbrain gamma synchrony may foster a shared emotional representation among individual and promote the social coordination when facing negative emotional stimulus. (Mu et al., 2017). Together with the results from previous studies, gamma interbrain synchrony as an important neural indicator of socio-emotional interaction, the present study warrants further investigation of socio-emotional interaction in this special age.

The present study examined the relationship between family cohesion and interbrain synchrony during emotional interactions between adolescent-parent dyads. As the main limitation, the sample size of the present study is relatively small. It is necessary to include more adolescent-parent dyads in the study to increase the generalizability of our findings. To rule out the possible impact of relevant variables of the parent-adolescent emotional interactions on our findings, demographic variables, adolescents' level of depression, adolescents' level of anxiety, adolescents' level of social support, parents' level of depression, parents' level of anxiety, and the level of parent involvement between the high and low family cohesion parent-adolescent dyads were examined. However, we didn't include more variables about the parent-

adolescent relationship (e.g., parenting style and family atmosphere) which might influence their emotional interactions. It is important and necessary for future research to explore the impacts of other variables about the parent-adolescent relationship. Moreover, to increase the generalizability of our findings, longitudinal design as a potential approach is necessary to employ in future research to examine how the relationship between family cohesion and interbrain synchrony during emotional interactions accompanied by the brain development and maturity of adolescents.

Limitations notwithstanding, the findings of the current study contribute to the literature on deepening our understanding of the relationship between family cohesion and socio-emotional interaction for a special blood relationship in a special period – adolescence. The current findings indicate that the impacts of family cohesion during socio-emotional interaction might be differently related to the gamma interbrain synchrony within dyads in the positive and negative emotional situations. Since high interbrain synchrony of high family cohesion adolescents in the positive emotional situations reflected a higher quality of socio-emotional interactions, cultivating a positive family atmosphere and increase emotional sharing among family members may have important value in promoting social skills for adolescents. Last but not the least, our findings in gamma interbrain synchrony again demonstrated that examining gamma band activity and its synchronization could be a feasible way in the study of socio-emotional interactions in different age groups (e.g., adolescents).

Abbreviations

EEG: electroencephalograph

LFCs: low family cohesion parent-adolescent dyads

HFCs: high family cohesion parent-adolescent dyads

PLV: phase-locking-value

FACES- α -CV: the Family Adaptability and Cohesion Scale, Chinese Version

Declarations

Ethics approval and consent to participate

The research protocol was reviewed and approved by the Institutional Review Board of Shenzhen University. All procedures performed in studies involving human participants were conducted in accordance with the ethical standards of the Institutional Review Board of Shenzhen University and with the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. Informed consent was obtained from the participants and their parents before the study.

Consent for publication

Not applicable.

Availability of data and materials

The data are currently not publicly available due to participant privacy, but they are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Educational Science Foundation in Shenzhen [grant number ybfz20010] and the Natural Science Foundation of SZU [grant number 000311].

Authors' c contributions

XD contributions to the conception and design of the work; ML and LZ performed the experiment and analyzed the data; XD and ML interpreted the data and drafted the manuscript; XL and QG critically revised the manuscript for intellectually important content, supervised the study. All authors read and approved the final submitted manuscript.

Acknowledgements

Not applicable.

References

1. Alavi, M., Latif, A. A., Ninggal, M. T., Mustaffa, M. S., & Amini, M. (2020). Family Functioning and Attachment Among Young Adults in Western and Non-Western Societies. *Journal of Psychology: Interdisciplinary and Applied*, 154(5), 346–366. <https://doi.org/10.1080/00223980.2020.1754153>
2. Babiloni, F., & Astolfi, L. (2014). Social neuroscience and hyperscanning techniques: Past, present and future. *Neuroscience and Biobehavioral Reviews*, 44, 76–93. <https://doi.org/10.1016/j.neubiorev.2012.07.006>
3. Barraza, P., Perez, A., & Rodriguez, E. (2020). Brain-to-brain coupling in the gamma-band as a marker of shared intentionality. *Frontiers in Human Neuroscience*, 14, 295. <https://doi.org/10.3389/fnhum.2020.00295>
4. Behnke, A. O., MacDermid, S. M., Coltrane, S. L., Parke, R. D., Duffy, S., & Widaman, K. F. (2008). Family cohesion in the lives of Mexican American and European American parents. *Journal of Marriage and Family*, 70(4), 1045–1059. <https://doi.org/10.1017/S0954579418000901>
5. Blakemore, S. J., & Mills, K. L. (2014). Is adolescence a sensitive period for sociocultural processing? *Annual Review of Psychology*, 65, 187–207. <https://doi.org/10.1146/annurev-psych-010213-115202>

6. Bretherton, I. (1990). Open communication and internal working models: Their role in the development of attachment relationships. In R. A. Thompson (Ed.), *Socioemotional development*. Lincoln: University of Nebraska Press.
7. Burgess, A. P. (2013). On the interpretation of synchronization in EEG hyperscanning studies: A cautionary note. *Frontiers in Human Neuroscience*, 7, 881.
<https://doi.org/10.3389/fnhum.2013.00881>
8. Costa, T., Rognoni, E., & Galati, D. (2006). EEG phase synchronization during emotional response to positive and negative film stimuli. *Neuroscience Letters*, 406(3), 159–164.
<https://doi.org/10.1016/j.neulet.2006.06.039>
9. Djalovski, A., Dumas, G., Kinreich, S., & Feldman, R. (2021). Human attachments shape interbrain synchrony toward efficient performance of social goals. *NeuroImage*, 226(7).
<https://doi.org/10.1016/j.neuroimage.2020.117600>
10. Diener, M. L., & Mangelsdorf, S. C. (1999). Behavioral strategies for emotion regulation in toddlers: Associations with maternal involvement and emotional expressions. *Infant Behavior and Development*, 22(4), 569–583. [https://doi.org/10.1016/S0163-6383\(00\)00012-6](https://doi.org/10.1016/S0163-6383(00)00012-6)
11. Dikker, S., Wan, L., Davidesco, I., Kaggen, L., Oostrik, M., McClintock, J., et al. (2017). Brain-to-Brain synchrony tracks real-world dynamic group interactions in the classroom. *Current Biology* 27(9), 1375-1380. <https://doi.org/10.1016/j.cub.2017.04.002>
12. Fei, L. P., Shen, Q. J., Zhen, Y. P., Zhao, J. P., Jiang S. A., Wang, L. W. & Wang, X. D. (1991). Preliminary Evaluation of Family Adaptability and Cohesion Scale and Family Environment Scale: A Comparative Study of Family Members with Schizophrenia and Family Members with Normal Families. *Chinese Journal of Mental Health* 5(5), 198-202.
13. Fosco, G. M., & Lydon-Staley, D. M. (2020). Implications of Family Cohesion and Conflict for Adolescent Mood and Well-Being: Examining Within- and Between-Family Processes on a Daily Timescale. *Family Process*, 59(4), 1672–1689. <https://doi.org/10.1111/famp.12515>
14. Halberstadt, A. G., & Eaton, K. L. (2002). A meta-analysis of family expressiveness and children's emotion expressiveness and understanding. *Marriage and Family Review*, 34(1–2), 35–62.
https://doi.org/10.1300/J002v34n01_03
15. Hasson, U., Furman, O., Clark, D., Dudai, Y., & Davachi, L. (2008) Enhanced inter subject correlations during movie viewing correlate with successful episodic encoding. *Neuron*, 57(3), 452-462.
<https://doi.org/10.1016/j.neuron.2007.12.009>
16. Hong, R. Y., Tan, C. S., Lee, S. S. M., Tan, S. H., Tsai, F. F., Poh, X. T., Zhou, Y., Sum, E. L., & Zhou, Y. (2015). Interactive Effects of Parental Personality and Child Temperament with Parenting and Family Cohesion. *Parenting*, 15(2), 92–118. <https://doi.org/10.1080/15295192.2015.1020143>
17. Hong, J. S., Hsieh, Y. P., Burlaka, V., Ortega, R. M., Espelage, D. L., & Pineros-Leano, M. (2021). Do Caregivers' Health Conditions Impact Children's Externalizing Behavior? The Roles of Caregivers' Parenting Frustration and Family Cohesion. *Child and Adolescent Social Work Journal*, 18(1).
<https://doi.org/10.1007/s10560-020-00736-7>

18. Hove, M. J., & Risen, J. L. (2009). It's all in the timing: Interpersonal synchrony increases affiliation. *Social Cognition, 27*(6), 949–960. <https://doi.org/10.1521/soco.2009.27.6.949>
19. Johnson, H. D., LaVoie, J. C., & Mahoney, M. (2001). Interparental conflict and family cohesion: Predictors of loneliness, social anxiety, and social avoidance in late adolescence. *Journal of Adolescent Research, 16*(3), 304–318. <https://doi.org/10.1177/0743558401163004>
20. Kajal, D. S., Fioravanti, C., Elshahabi, A., Ruiz, S., Sitaram, R., & Braun, C. (2020). Involvement of top-down networks in the perception of facial emotions: A magnetoencephalographic investigation. *NeuroImage, 222*, 117075. <https://doi.org/10.1016/j.neuroimage.2020.117075>
21. Kinreich, S., Djalovski, A., Kraus, L., Louzoun, Y., and Feldman, R. (2017). Brain-to-brain synchrony during naturalistic social interactions. *Scientific Reports, 7*, 1–12. <https://doi.org/10.1038/s41598-017-17339-5>
22. Konrad, S. Č. (2016). Family emotional expressiveness and family structure. *Psihologija, 49*(4), 319–333. <https://doi.org/10.2298/PSI1604319C>
23. Konvalinka, I., & Roepstorff, A. (2012). The two-brain approach: How can mutually interacting brains teach us something about social interaction? *Frontiers in Human Neuroscience, 6*(7), 1–10. <https://doi.org/10.3389/fnhum.2012.00215>
24. Lachaux, J. P., Rodriguez, E., Martinerie, J., & Varela, F. J. (1999). Measuring phase synchrony in brain signals. *Human Brain Mapping, 8*(4), 194–208. [https://doi.org/10.1002/\(SICI\)1097-0193\(1999\)8:4<194::AID-HBM4>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1097-0193(1999)8:4<194::AID-HBM4>3.0.CO;2-C)
25. Lang, P. J., Bradley, M. M., Cuthbert, B. N. (2005). *International Affective Picture System (IAPS): Instruction manual and affective ratings*. Gainesville: University of Florida.
26. Lee, Y. Y., & Hsieh, S. (2014). Classifying different emotional states by means of eeg-based functional connectivity patterns. *PLoS ONE 9*(4): e95415. <https://doi.org/10.1371/journal.pone.0095415>
27. Leidy, M. S., Guerra, N. G., & Toro, R. I. (2010). Positive parenting, family cohesion, and child social competence among immigrant Latino families. *Journal of Family Psychology, 24*(3), 252–260. <https://doi.org/10.1037/a0019407>
28. Lin, Y. P., Wang, C. H., Jung, T. P., Wu, T. L., Jeng, S. K., Duann, J. R., & Chen, J. H. (2010). EEG-based emotion recognition in music listening. *IEEE Transactions on Biomedical Engineering, 57*(7), 1798–1806. <https://doi.org/10.1109/TBME.2010.2048568>
29. Liu, D., Liu, S., Liu, X., Zhang, C., Li, A., Jin, C., Chen, Y., Wang, H., & Zhang, X. (2018). Interactive brain activity: Review and progress on EEG-based hyperscanning in social interactions. *Frontiers in Psychology, 9*(11), 1–11. <https://doi.org/10.3389/fpsyg.2018.01862>
30. Luo, Q., Mitchell, D., Cheng, X., Mondillo, K., McCaffrey, D., Holroyd, T., Carver, F., Coppola, R., & Blair, J. (2009). Visual awareness, emotion, and gamma band synchronization. *Cerebral Cortex, 19*(8), 1896–1904. <https://doi.org/10.1093/cercor/bhn216>
31. Maffei, A., Spironelli, C., & Angrilli, A. (2019). Affective and cortical EEG gamma responses to emotional movies in women with high vs low traits of empathy. *Neuropsychologia, 133*, 107175. <https://doi.org/10.1016/j.neuropsychologia.2019.107175>

32. Martini, N., Menicucci, D., Sebastiani, L., Bedini, R., Pingitore, A., Vanello, N., Milanese, M., Landini, L., & Gemignani, A. (2012). The dynamics of EEG gamma responses to unpleasant visual stimuli: From local activity to functional connectivity. *NeuroImage*, 60(2), 922–932. <https://doi.org/10.1016/j.neuroimage.2012.01.060>
33. Matsumoto, A., Ichikawa, Y., Kanayama, N., Ohira, H., & Iidaka, T. (2006). Gamma band activity and its synchronization reflect the dysfunctional emotional processing in alexithymic persons. *Psychophysiology*, 43, 533-540. <https://doi.org/10.1111/j.1469-8986.2006.00461.x>
34. Mu, Y., Han, S., and Gelfand, M. J. (2017). The role of gamma interbrain synchrony in social coordination when humans face territorial threats. *Social Cognitive and Affective Neuroscience*. 12(10), 1614–1623. <https://doi.org/10.1093/scan/nsx093>
35. Olson, D. (2000). Circumplex model of marital and family systems. *Journal of Family Therapy*, 22(2), 144–167. <https://doi.org/10.1111/1467-6427.00144>
36. Olson, D. (2011). FACES IV and the Circumplex Model: Validation Study. *Journal of Marital and Family Therapy*, 37(1), 64–80. <https://doi.org/10.1111/j.1752-0606.2009.00175.x>
37. Olson, D. H., & Craddock, A. E. (1980). Circumplex model of marital and family systems. *Australian Journal of Sex, Marriage and Family*, 1(2), 53–69. <https://doi.org/10.1080/01591487.1980.11004154>
38. Phillips-Salimi, C. R., Robb, S. L., Monahan, P. O., Dossey, A., & Haase, J. E. (2014). Perceptions of communication, family adaptability and cohesion: A comparison of adolescents newly diagnosed with cancer and their parents. *International Journal of Adolescent Medicine and Health*, 26(1), 19–26. <https://doi.org/10.1515/ijamh-2012-0105>
39. Rahgozar, H., Yousefi, S., Mohammadi, A., & Piran, P. (2012). The impact of family cohesion and flexibility on university students' identity: The case of Shiraz Branch, Islamic Azad University. *Asian Social Science*, 8(2), 95–103. <https://doi.org/10.5539/ass.v8n2p95>
40. Reeves, T., Horne, S. G., Rostosky, S. S., Riggle, E. D. B., Baggett, L. R., & Aycok, R. A. (2010). Family members' support for GLBT issues: The role of family adaptability and cohesion. *Journal of GLBT Family Studies*, 6(1), 80–97. <https://doi.org/10.1080/15504280903472857>
41. Reindl, V., Gerloff, C., Scharke, W., & Konrad, K. (2018). Brain-to-brain synchrony in parent-child dyads and the relationship with emotion regulation revealed by fNIRS-based hyperscanning. *NeuroImage*, 178(11), 493–502. <https://doi.org/10.1016/j.neuroimage.2018.05.060>
42. Roth, J., & Brooks-Gunn, J. (2000). What do adolescents need for healthy development? Implications for youth policy. *Social Policy Report*, 14(1), 3–19. <https://doi.org/10.1002/j.2379-3988.2000.tb00012.x>
43. Santamaria, L., Noreika, V., Georgieva, S., Clackson, K., Wass, S., & Leong, V. (2020). Emotional valence modulates the topology of the parent-infant inter-brain network. *NeuroImage*, 207(11), 1-16. <https://doi.org/10.1016/j.neuroimage.2019.116341>
44. Sheeber, L., Hops, H., & Davis, B. (2001). Family processes in adolescent depression. *Clinical Child and Family Psychology Review*, 4(1), 19–35. <https://doi.org/10.1023/A:1009524626436>

45. Starke, M., & Svensson, E. (2001). Construction of a global assessment scale of family function, using a questionnaire. *Social Work in Health Care*, 34(1–2), 131–142.
<https://doi.org/10.1080/00981380109517022>
46. Steinberg, L. (1990). Autonomy, conflict, and harmony in the family relationship. In S. S. Feldman & G.R. Elliott (Eds.), *At the threshold: The developing adolescent*, 255 – 276, Cambridge, MA: Harvard University Press.
47. Stujenske, J. M., Likhtik, E., Topiwala, M. A., & Gordon, J. A. (2014). Fear and Safety Engage Competing Patterns of Theta-Gamma Coupling in the Basolateral Amygdala. *Neuron*, 83(4), 919–933. <https://doi.org/10.1016/j.neuron.2014.07.026>
48. Sui, G. Y., Wang, J. N., Liu, G. C., & Wang, L. (2015). The effects of being an only child, family cohesion, and family conflict on behavioral problems among adolescents with physically ill parents. *International Journal of Environmental Research and Public Health*, 12(9), 10910–10922.
<https://doi.org/10.3390/ijerph120910910>
49. Wang, Y. W., Yang, X., Yao, X. Q., & Fu, C. (2021). Computational methods of brain-to-brain coupling during human interaction (in Chinese). *Chinese Science Bulletin-Chinese*, 66(4-5), 501-514.
50. Wheatley, T., Kang, O., Parkinson, C., and Looser, C. E. (2012). From mind perception to mental connection: synchrony as a mechanism for social understanding. *Social and Personality Psychology Compass* 6, 589–606. <https://doi.org/10.1111/j.1751-9004.2012.00450.x>
51. Yan, M., GUO, C. Y., & Han, S. H. (2016). Oxytocin enhances inter-brain synchrony during social coordination in male adults. *Social Cognitive and Affective Neuroscience*, 11(12), 1882-1893.
<https://doi.org/10.1093/scan/nsw106>
52. Zhu, L., Lotte, F., Cui, G., Li, J., Zhou, C., Cichocki, A., (2018). Neural mechanisms of social emotion perception: an EEG hyper-scanning study. *Proceedings - 2018 International Conference on Cyberworlds, CW 2018*, 199–206. <https://doi.org/10.1109/CW.2018.00045>

Figures

Figure 1

Sample of stimulus and procedure

Figure 2

Gamma Interbrain Phase-locking-value (PLV) in the Parietal areas (Pz) between Low and High Family Cohesion Parent-adolescent Dyads (LFCs and HFCs) in Different Conditions. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Appendix.docx](#)