Prevalence of Sexually Transmitted Infections among cervical cancer suspected women at University of Gondar Comprehensive Specialized Hospital, North-west Ethiopia

Milkias Abebe (✉ mickyelfe21@gmail.com)

Wollega University

Setegn Eshetie

University of Gondar

Belay Tessema

University of Gondar

Research Article

Keywords: HIV, HBV, HCV, syphilis, cervical cancer, Ethiopia

Posted Date: January 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-132667/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on April 23rd, 2021. See the published version at <https://doi.org/10.1186/s12879-021-06074-y>.

1 **Prevalence of Sexually Transmitted Infections among cervical cancer suspected women**
2 **at University of Gondar Comprehensive Specialized Hospital, North-west Ethiopia**

3 Milkias Abebe^{1*}, Setegn Eshetie², Belay Tessema²

4

5 ¹Department of Medical Laboratory Sciences, Institute of Health Sciences, Wollega
6 University, P.O. Box: 395, Nekemte, Ethiopia.

7 ²Department of Medical Microbiology, School of Biomedical and Laboratory Sciences,
8 College of Medicine and Health Sciences, University of Gondar, P.O. Box: 196, Gondar,
9 Ethiopia.

10 *Corresponding author

11 Email addresses:

12 MA: mickyelfe21@gmail.com

13 SE: wolet03.2004@gmail.com

14 BT: bt1488@yahoo.com

15

16

Abstract

Background: Sexually transmitted infections (STIs) such as Human Immunodeficiency Virus (HIV), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), and Syphilis have continued a significant public health problem, especially in areas with substandard infection prevention and control programs. It is known that STIs are largely associated with the increased occurrence of cervical cancer. To date, little is known about the burden of STIs among cervical cancer suspected women in Ethiopia.

Objectives: To assess the seroprevalence of STIs and associated risk factors among cervical cancer suspected women with special emphasis on HIV, HBV, HCV, and Syphilis.

Methods: A hospital-based cross-sectional study was conducted among cervical cancer suspected women at the University of Gondar Comprehensive Specialized Hospital from February to April 2017. A pre-tested and structured questionnaire was used to collect data on patients' characteristics. The patient's serum or plasma samples were tested for HIV, HBV, HCV, and syphilis using enzyme-linked immunosorbent assay. In all aspects, the standard operational procedure was strictly followed. Data were analyzed using SPSS version 20 software and presented using tables. Statistical associations were measured using bivariate and multivariable logistic regression. A p-value of below 0.05 was considered statistically significant.

Result: A total of 403 cervical cancer suspected women with the mean age of 42.54 (SD±11.24) years were enrolled in the study. The overall prevalence of STIs was 16.6% (67/403) and the prevalence of HIV, HBV, HCV, and syphilis was 36/403 (8.9 %), 10/403 (2.5 %), 4/403 (1 %), and 29/403 (7.2 %) respectively. History of multiple sexual partners (Adjusted

OR=3.02, 95%CI 1.57-5.79, P=0.001), alcohol addiction (Adjusted OR=2.2, 95%CI 1.07-4.5, P=0.031), history of STI (Adjusted OR= 3.38; 95% CI: 1.76-6.47, P= 0.00) and not use of condom (Adjusted OR= 4.99; 95% CI: 1.5-16.16, P= 0.007) were significantly associated with STIs.

Conclusion: The prevalence of STIs was generally higher among cervical cancer suspected patients. Health education is encouraged to promote awareness about the prevention of STIs.

Key words: HIV, HBV, HCV, syphilis, cervical cancer, Ethiopia

Introduction

Sexually transmitted infections (STIs) are infections caused by pathogens that are mainly transmitted through unsafe sexual activity. They can also spread by non-sexual contact with contaminated blood, tissues, through breastfeeding, and during childbirth. Different types of bacteria, viruses, and parasites can able to cause STIs (1). Viral STIs like human immunodeficiency virus (HIV), human papillomavirus (HPV), hepatitis B virus, and herpes simplex virus are preventable, but not treatable (2). At present, such STIs (HIV, HBV, HCV, and syphilis) has continued to be a community health problem in high, middle and low countries that causes acute illness, infertility, long-term disability, and death, with severe medical, and psychological consequences for millions of women and infants (1). Different reports showed that the prevalence of STIs related with early age of initiation of sexual activity, drug use before sex, and multiple sexual partners (3).

Though, HPV is a necessary principal cause of cervical cancer, but it is not a sufficient cause (4). Tobacco smoking, high parity, long-term contraceptive use, and co-infection with HIV and other STIs (syphilis, HBV, and HCV), have been identified as established cofactors

necessary for progression from cervical HPV infection to cancer (5). Women infected with HIV have been shown to have a 2 to 12 fold risk of developing cervical cancer (6). This relationship between HIV infection and HPV infection is also due to the fact that both are sexually transmitted diseases and behaviors that put women at risk for one also put them at risk for the other (6). Co-infection with other sexually transmitted diseases (syphilis, hepatitis B virus, and hepatitis C virus) and vaginal infections are associated with increased susceptibility to HPV infection (7).

Sexually transmitted diseases, especially HIV, HBV, HCV, and Syphilis infections are the main contributing factors for cervical cancer, which is the second leading cause of cancer death in women (5–7). Hence, the result of the study is helpful to design a strategy for the prevention of sexually transmitted infection in cervical cancer suspected women, draw the attention of stakeholders to focus on such life-threatening but preventable infections, it also serves as a baseline data for the researchers who are interested in related topics. Therefore, the aim of the study is to determine the prevalence of sexually transmitted infection and its associated risk factors among women suspected of cervical cancer attending at University of Gondar Comprehensive Specialized Hospital, North-west Ethiopia.

Methods

Study design, area, and population

A hospital-based cross-sectional study was conducted among women suspected of cervical cancer from February to April 2017 at the University of Gondar Comprehensive Specialized Hospital. University of Gondar Comprehensive Specialized Hospital is one of the largest comprehensive, specialized hospitals served as a teaching as well as patient care in the

Amhara region. It is located in Gondar town, 750 km far from Addis Ababa in Northwest Ethiopia. The hospital has an accredited laboratory, more than 1200 beds, and provides health care referral services for more than 5 million people from the surrounding zones and nearby regions (8).

Inclusion and exclusion criteria

Cervical cancer suspected women (women screened for cervical cancer who had abnormal vaginal bleeding such as inter-menstrual bleeding, post-coital bleeding) who are attending at University of Gondar Comprehensive Specialized Hospital during the study period were included in the study while patients with known STIs were excluded from the present study.

Sample size and sampling procedure

The sample size was calculated based on the assumption of 5% expected margins of error and a 95% confidence interval, taking the prevalence of 50% by using a single population proportion formula as follows.

$$N = \frac{(Z_{\alpha/2})^2 P(1-P)}{d^2}$$

Where N is the calculated sample size of cervical cancer suspected women; Z is the standard normal deviate at 95%, confidence interval = 1.96; P is the prevalence of STI to get maximum sample size = 50%; d is the precision level = 0.05. Then 5% contingency= 19 were added. Finally, the total sample size = 384+19= 403 study participants.

The study participants were enrolled consecutively using a convenience sampling technique until a sample size of 403 study participants was achieved. The detailed information of socio-

demographic characteristics, behavioral characteristics, clinical and environmental characteristics were collected prior to laboratory diagnosis from each of the study participants using structured and pre-tested questionnaires.

Laboratory diagnosis

Study participants sera was tested for the presence of sexually transmitted infection (HBV, HIV, HCV and syphilis) by using the Enzyme-Linked Immunosorbent Assay (ELISA). For HIV1/2: Vironostika ELISA (Bio-Merieux, Boxtel, Netherlands); Furthermore, for HBsAg: using ELISA, Hepanostika HBsAg (Bio-Merieux, Boxtel, Netherlands); Moreover, for HCV: Human anti-HCV ELISA (Human Gesellschaft for Bio-chemical and diagnostic MbH, Germany). Besides, anti-syphilis Ab of syphilis was checked by using DIALAB ELISA (Nora Kampitsch, MSc, India).

Data analysis and interpretation

Data was collected, summarized, tabulated, and analyzed using Statistical Package Epi-Info Version7 and Statical package for social science version 20 Software. The results were presented through graphs and tables. The statistical significance, the association was measured by using bivariate and multivariate analysis, odds ratio at 95% confidence intervals. A p-value < 0.05 was considered as statistically significant.

Results

Socio-Demographic Data

A total of 403 cervical cancer suspected women were included in the study. The majority of the respondents were in the age groups of 36-49 years, 179 (44.4%), and the mean age of

study subjects was 42.54 ± 11.24 years. Nearly 70% of the participants were married, and most of the 230 (57.1%) the participants were unable to read and write and 199 (49.4%) of were housewives, 252 (67.5%) of participants were from the urban (Table1).

Prevalence of Sexually Transmitted Infection

Among 403 cervical cancer suspected women 36 (8.9%, 95% CI= 6.2-11.9) were HIV positive, 10 (2.5%, 95% CI= 1-4.2) were HBV positive, 4 (1%, 95% CI= 0.2-2.2) were HCV positive, and 29 (7.2%, 95% CI= 4.7-9.7) of were syphilis positive.

Of the study participants, 67 (16.6%) (95% CI, 12.9-20.6) were found to be positive for STIs. Positive STIs defined as a positive status either for single or combination of infections specific to HIV, HBV, HCV, and syphilis. Of these STIs positive women, 12/67 (17.9%) had serological evidence of multiple infections. Among those with multiple infections, 9/67 (13.4%) were HIV- syphilis positive, 1/67 (1.5%) were HBV-syphilis positive, 1 (1.5%) were HCV-HIV positive and 1 (1.5%) were HCV-syphilis positive.

Risk Factors for the acquisition of Sexually Transmitted Infections

Firstly, bivariate logistic regression analysis was done and variables with P-value ≤ 0.2 were further tested by multivariable logistic regression analysis. History the multiple sexual partners, history of an STI, not using condoms, alcohol addiction were an independent risk factors for STI. Patients who had multiple sexual partners were three times more likely to develop STI compared to those who had no multiple sexual partner (Adjusted OR=3.02, 95% CI 1.57-5.79, P= 0.001). Similarly, the odds of developing STI were 3.4 times higher among patients who had a history of STI compared to the counterparts (Adjusted OR= 3.38; 95% CI: 1.76-6.47, P= 0.00). Besides patients, who had a habit of drinking alcohol were 2.2 times more likely to have STI compared to those who were not (Adjusted OR=2.2, 95% CI 1.07-4.5, P=0.031) (Table 2).

Discussion

The burden of sexually transmitted infections is high among cervical cancer suspected women this is due to the fact that HPV is a sexually transmitted infection and they share a common route of transmission and risk factors. This study has sought the seroprevalence of STI, HIV, HBV, HCV, and syphilis infections and contributing factors for STI among cervical cancer suspected women attending at the University of Gondar Comprehensive Specialized Hospital.

The overall prevalence of STIs in the current study was 16.6% (95% CI= 12.9 to 20.6%). The finding of this study was higher than studies conducted in Gondar which was 10.5% (9) and 11.7% in India (10). The high prevalence of STIs among cervical cancer suspected women is due to the fact that cervical cancer-causing virus and sexually transmitted infections share a common route of transmission and risk factors.

The present study shows that the prevalence of HIV was 8.9% (95% CI= 6.2 to 11.9%). It was higher compared to the study conducted in Cameroon which was 4.2% (11) and 4.1% in Nigeria (12). The variation could be due to the fact that cervical cancer suspected women are more risk group for STIs namely HIV. An individual infected with one STIs become a high risk to be infected by HIV. Human Papilloma Virus causes inflammation around the genital area which increases the concentration of “activated” immune cells. Although the inflammatory response is meant to help fight the HPV, HIV likes to infect some of these recruited immune cells, also known as CD4 cells. Therefore, if someone has an STI (HPV) in the genitals and that area is exposed to HIV, the higher concentration of “activated” CD4 cells facilitates HIV infection, replication, and spread throughout the body (13). In contrast to this; higher prevalence observed in South Africa, 14% (14). This may be due to differences in socio-cultural behavior.

The seroprevalence of HBV was also noted in the current study and it was 2.5% (95% CI=1.0 to 4.2%). Comparable results were also reported in Ethiopia ranged from 3% - 3.8% (15–17). However, the higher prevalence was reported in Gondar (4.7 %), Dessie (4.9%), and Congo (5.9%) (18–20). This difference might be attributable to differences in risk for HBV acquisition and the impact of education level. In contrast to this, the lower prevalence was reported in India (0.9%) (21) and Iran (0.7%) (22). The discrepancies might be a result of differences in sampling population, methodological difference, and geographical variation.

The prevalence of HCV in this study was also determined and it was 1% (95% CI= 0.2 to 2.2%). The finding of this study was in line with studies conducted in the Gondar Health Center 1.3% (9), 0.26% North-west Ethiopia (23). However; a higher prevalence of HCV was documented

among the attendants of voluntary counseling and testing for HIV in Ethiopia 9.1%, 6.0%, and 4.3% in Hawassa, Mekelle, and Adwa, respectively (24–26). The variation is due to differences in risk behavior, sampling size, and study population. Besides, HCV infection is majorly acquired via blood transfusion other than other STIs.

Moreover, the prevalence of syphilis among cervical cancer suspected women in this study was 7.2% (95% CI, 4.7-9.7). The results from this study were in agreement with the report in South Africa 5% (14), 6.4% in North Tanzania (27), 5% in Nigeria (12). Studies with lower prevalence compared to our findings include; 2.3% in Gondar Health Center (9) and 2.9% in Addis Ababa (28). The high prevalence of syphilis among cervical cancer suspected women is due to the fact that cervical cancer-causing virus and *T. palladium* have a common route of transmission and risk factors. On the other hand; an extremely higher prevalence (17.4%) was reported in Cameroon (29) and 10% in South Western Nigeria (30). The variations could be largely due to geo-cultural differences and the decrease time trend of syphilis is also claimed in Ethiopia.

Lastly, the present study was also aimed to identify independent risk factors of STIs. The distribution of STIs was significantly higher among alcohol user women than their counterparts in the current study. In brief, patients, who had habit of drinking alcohol, were 2.2 times more likely to have STI compared to those who were not. This might be due to the fact that alcohol use is one of the common factors that lead people to exhibit risky sexual behavior and lowers a person's inhibitions, which may influence their sexual behavior, and thus a compromise safe sex practices (sex without condom) (31–33).

Additionally, the prevalence of STI was significantly higher among women who had a history of STIs ($P < 0.001$) which conforms to previous studies conducted in Jimma (15) and Peru (34).

Moreover, the prevalence of STI was also significant and three times higher among the study subjects who had a history of multiple sexual partners compared to their counterparts. This is due to women who have multiple sexual partners who had a chance of getting sex with an infected person.

Furthermore, STI prevalence was five times higher among study participants who have not used condoms compared to their counterparts. Without any doubt, the condom is one of the preventive strategies against STIs. Evidently, consistent and correct use of condoms is 90% effective in reducing HIV, up to 66% effective in reducing syphilis (35). Similarly, studies have done in New York and Southwestern China support the advantage of condoms in preventing STI (36,37). In this study, however, STI did not show any correlation with the history of blood transfusion, tattooing, surgical operation, smoking, and genital mutilation in the present study. This is similar to previous studies in Ethiopia (16,38).

Conclusion

This study shows the prevalence of sexually transmitted infections was generally higher among cervical cancer suspected women. Having multiple sexual partners, not using condoms, alcohol addiction, and history of STIs found to be independent risk factors for sexually transmitted infections.

Since STIs are major health problems in cervical cancer suspected women, all cervical cancer suspected women should be screened for STIs (syphilis, HBV, HCV), increase awareness of people on modes of transmission of sexually transmitted infections and educate people how to prevent sexually transmitted infections could help in reducing the burden of STIs.

227 **Declarations**

228 **Acknowledgments**

229 We would like to thank all study participants.

230 **Authors' contributions**

231 MA was conceived the study, designed data collection, conducted data analysis and
232 interpretation. BT and SE interpreted the results and reviewed the initial and final drafts of the
233 manuscript. All authors read and approved the final manuscript.

234 **Funding**

235 Not applicable.

236 **Availability of data and materials**

237 All data generated or analyzed during this study were included in this article.

238 **Ethics approval and consent to participate**

239 This research was conducted after obtaining ethical clearance from School of Biomedical and
240 Laboratory Sciences of the University of Gondar ethical review committee with reference
241 number SBMLS/625/08. Written informed consent was obtained from Study participants after
242 explaining the purpose and objective of the study. Any patient who was not willing to participate
243 in the study wasn't forced to participate. All data and samples obtained from them were kept
244 confidential by using codes instead of any personal identifiers and were meant only for the
245 purpose of the study. The laboratory results of the study participant were communicated to their
246 physicians for medical care. All methods were performed in accordance with the relevant guidelines
247 and regulations in the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Medical Laboratory Sciences, Institute of Health Sciences, Wollega University, P.O. Box: 395, Nekemte, Ethiopia.

²Department of Medical Microbiology, School of Biomedical and Laboratory Sciences, College of Medicine and Health Sciences, University of Gondar, P.O. Box: 196, Gondar, Ethiopia.

References

1. Geremew RA, Agizie BM, Bashaw AA, Seid ME, Yeshanew AG. Prevalence of Selected Sexually Transmitted Infection (STI) and Associated Factors among Symptomatic Patients Attending Gondar Town Hospitals and Health Centers. *Ethiop J Health Sci.* 2017;27(6):589–600.
2. De Schryver A, Meheus A. Epidemiology of sexually transmitted diseases: The global picture. *Bull World Health Organ.* 1990;68(5):639–54.
3. Diclemente R, Wingood G, Sionean C, Crosby R, Harrington K, Davies S, et al. Association of Adolescents' History of Sexually Transmitted Disease (STD) and Their Current High-Risk Behavior and STD Status. *Sex Transm Dis.* 2002;29(9):503–9.
4. Carlson LM, Gonzalez S. Knowledge of cervical cancer pathology of high school students in San Carlos, Costa Rica. *Rev Biol Trop.* 2014;62(3):877–86.
5. Smith JS, Herrero R, Bosetti C, Muñoz N, Bosch FX, Eluf-Neto J, et al. Herpes simplex virus-2 as a human papillomavirus cofactor in the etiology of invasive cervical cancer. *J Natl Cancer Inst.*

270 2002;94(21):1604–13.

271 6. Mapanga W, Girdler-Brown B, Feresu SA, Chipato T, Singh E. Prevention of cervical cancer in
 272 HIV-seropositive women from developing countries through cervical cancer screening: A
 273 systematic review. *Syst Rev*. 2018;7(1):198.

274 7. King CC, Jamieson DJ, Wiener J, Cu-Uvin S, Klein RS, Rompalo AM, et al. Bacterial vaginosis
 275 and the natural history of human papillomavirus. *Infect Dis Obstet Gynecol*. 2011;2011:1–8.

276 8. Molla R, Tiruneh M, Abebe W, Moges F. Bacterial profile and antimicrobial susceptibility
 277 patterns in chronic suppurative otitis media at the University of Gondar Comprehensive
 278 Specialized Hospital, Northwest Ethiopia. *BMC Res Notes* [Internet]. 2019;12(1):1–6. Available
 279 from: <https://doi.org/10.1186/s13104-019-4452-4>

280 9. Tiruneh M. Seroprevalence of multiple sexually transmitted infections among antenatal clinic
 281 attendees in Gondar Health Center , northwest Ethiopia. *Ethiop Med J*. 2008;46(4):359–66.

282 10. Hussain T, Kulshreshtha KK, Sinha S, Yadav VS, Katoch VM. HIV , HBV , HCV , and syphilis
 283 co-infections among patients attending the STD clinics of district hospitals in Northern India. *Int J*
 284 *Infect Dis*. 2006;10:358–63.

285 11. Goh BT. Syphilis in adults. *sex Transm Infect*. 2005;448–52.

286 12. Buseri F, Jeremiah Z, Seiyaboh E. Surveying infections among pregnant women in the Niger
 287 Delta, Nigeria. *J Glob Infect Dis*. 2010;2(3):203–11.

288 13. Quinn TC, Wawer MJ, Sewankambo N, Serwadda D, Li C, Wabwire-Mangen F, et al. Viral load
 289 and heterosexual transmission of human immunodeficiency virus type 1. *N Engl J Med*.
 290 2000;342(13):921–9.

291 14. Dinh T, Centers K, Control D, Diseases C. Integration of Preventing Mother-To-Child

292 Transmission of HIV and Syphilis Testing and Treatment in Antenatal Care Services in the
293 Northern Cape and Gauteng Provinces, South Africa. *Sex Transm Dis.* 2013;40(11):846–51.

294 15. Awole M, Gebre-Selassie S. Seroprevalence of HBsAg and its risk factors among pregnant
295 women in Jimma, Southwest Ethiopia. *Ethiop J Heal Dev.* 2005;19(1):45–50.

296 16. Zenebe Y, Mulu W, Yimer M, Abera B. Sero-prevalence and risk factors of hepatitis B virus and
297 human immunodeficiency virus infection among pregnant women in Bahir Dar city , Northwest
298 Ethiopia : a cross sectional study. *BMC Infect Dis.* 2014;14(1):1–7.

299 17. Tegegne D, Desta K, Tegbaru B, Tilahun T. Seroprevalence and transmission of Hepatitis B virus
300 among delivering women and their new born in selected health facilities , Addis Ababa , Ethiopia :
301 a cross sectional study. *BMC Res Notes.* 2014;7(1):1–7.

302 18. Tessema B, Yismaw G, Kassu A, Amsalu A, Mulu A, Emmrich F, et al. Seroprevalence of HIV ,
303 HBV , HCV and syphilis infections among blood donors at Gondar University Teaching Hospital ,
304 Northwest Ethiopia : declining trends over a period of five years. *BMC Infect Dis.* 2010;10:111.

305 19. Seid M, Gelaw B, Assefa A. Sero-prevalence of HBV and HCV Infections Among Pregnant
306 Women Attending Antenatal Care Clinic at Dessie Referral Hospital , Ethiopia. *Adv Life Sci Heal.*
307 2014;1(2):109–20.

308 20. Kabinda JM, Akilimali TS, Miyanga AS, Donnen P, Michèle D. Hepatitis B , Hepatitis C and HIV
309 in Pregnant Women in the Community in the Democratic Republic of Congo. *World J AIDS.*
310 2015;5:124–30.

311 21. Dwivedi M, Misra SP, Misra V, Pandey A, Pant S, Singh R, et al. Seroprevalence of hepatitis B
312 infection during pregnancy and risk of perinatal transmission. *Indian J Gastroenterol.*
313 2011;30(2):66–71.

- 314 22. Mohebbi SR, Sanati A, Cheraghipour K, Nejad MR, Shalmani HM, Zali MR. Hepatitis C and
315 hepatitis B virus infection: Epidemiology and risk factors in a large cohort of pregnant women in
316 Lorestan, West of Iran. *Hepat Mon.* 2011;11(9):736–9.
- 317 23. Molla S, Munshea A, Nibret E. Seroprevalence of hepatitis B surface antigen and anti HCV
318 antibody and its associated risk factors among pregnant women attending maternity ward of
319 Felege Hiwot Referral Hospital , northwest Ethiopia : a cross-sectional study. *Virol J.*
320 2015;12:204.
- 321 24. Alemayehu A, Tassachew Y, Sisay Z, Shimelis T. Prevalence and risk factors of Hepatitis C
322 among individuals presenting to HIV testing centers , Hawassa city, Southern Ethiopia. *BMC Res*
323 *Notes.* 2011;4:193.
- 324 25. Hadush H, Gebre-Selassie S, Mihiret A. Hepatitis C virus and Human Immunodeficiency Virus
325 coinfection among attendants of Voluntary Counseling and Testing Centre and HIV follow up
326 clinics in Mekelle Hospital. *Pan Afr Med J.* 2013;14:107.
- 327 26. Atsbaha AH, Dejen TA, Belodu R, Getachew K, Saravanan M, Wasihun AG. Sero - prevalence
328 and associated risk factors for hepatitis C virus infection among voluntary counseling testing and
329 anti retroviral treatment clinic attendants in Adwa hospital , Northern Ethiopia. *BMC Res Notes.*
330 2016;9:121.
- 331 27. Kumogola Y, Slaymaker E, Zaba B, Mngara J, Isingo R, Chagalucha J, et al. Trends in HIV &
332 syphilis prevalence and correlates of HIV infection : results from cross-sectional surveys among
333 women attending ante-natal clinics in Northern Tanzania. *BMC Res Notes.* 2010;10:553.
- 334 28. Kebede E. and Chamiso. Prevalence of syphilis in pregnancy in Addis Ababa. *East Afr Med J.*
335 2000;77(4):212–6.
- 336 29. Mbopi Keou FX, Mbu R, Andela A, Tetanye E, Leke R, Chaouat G, Barre-Sinoussi F, Martin P

337 BL. Antenatal HIV prevalence in Yaounde, Cameroon. *Int J STD AIDS*. 1998;9(7):400–2.

338 30. Ojo DA, Oyetunji IAO. Sero-prevalence of syphilis among pregnant women in Osogbo in South-
339 western Nigeria. *ASSET Ser B*. 2007;6(1):61–5.

340 31. Regassa N, Kedir S. Attitudes and practices on HIV preventions among students of higher
341 education institutions in Ethiopia : the case of Addis Ababa University. *East Afr J Public Heal*.
342 2011;8(2):141–54.

343 32. Malaju MT, Asale GA. Association of Khat and alcohol use with HIV infection and age at first
344 sexual initiation among youths visiting HIV testing and counseling centers in Gamo-Gofa Zone,
345 South West Ethiopia. *BMC Int Health Hum Rights*. 2013;13:10.

346 33. Gelibo T, Belachew T, Tilahun T. Predictors of sexual abstinence among Wolaita Sodo University
347 Students, South Ethiopia. *Reprod Health*. 2013;10:18.

348 34. Alarcon JO, Johnson KM, Courtois B, Rodriguez C, Sanchez J, Watts DM, et al. Determinants and
349 prevalence of HIV infection in pregnant Peruvian women. *AIDS*. 2003;17(4):613–8.

350 35. Hocking J FC. Associations between condom use and rectal or urethral chlamydia infection in
351 men. *Sex Transm Dis*. 2006;33(4):256–8.

352 36. Caico C. Sexually Risky Behavior in College-Aged Students. *Open J Prev Med*. 2014;4:354–64.

353 37. Hu Y, Liang S, Zhu J, Qin G, Liu Q, Song B, et al. Factors Associated with Recent Risky Drug
354 Use and Sexual Behaviors among Drug Users in Southwestern China *AIDS & Clinical Research*. *J*
355 *AIDS Clin Res*. 2011;2(3):1–6.

356 38. Abera B, Zenebe Y, Mulu W, Kibret M, Kahsu G. Seroprevalence of hepatitis B and C viruses and
357 risk factors in HIV infected children at the felgehiwot referral hospital , Ethiopia. *BMC Res Notes*.
358 2014;7:838.

Table 1: - Socio-demographic characteristics of cervical cancer suspected women at University of Gondar Comprehensive Specialized Hospital.

Characteristics	Frequency	Percent
Age		
<=35 years	123	30.5
36- 49years	179	44.4
<=50years	101	25.1
Total	403	100
Marital status		
Married	280	69.5
Divorce	63	15.6
Widowed	60	14.9
Total	403	100
Educational level		
Unable to read and write	230	57.1
Primary	71	17.6
Secondary	52	12.9
Diploma	39	9.7
Degree and above	11	2.7
Total	403	100
Occupational status		

Self employed	15	3.7
House Wife	199	49.4
Merchant	47	11.7
Government employed	46	11.4
Farmer	96	23.8
Total	403	100

Residence

Rural	151	37.5
Urban	252	62.5
Total	403	100

362 **Table 2:** - Risk factors of STIs among cervical cancer suspected at University of Gondar
363 Comprehensive Specialized Hospital.

Variables	STIs status		COR (95%CI)	P-vale	AOR (95% CI)	P-value
	Neg (N)	Pos (N)				
Multiple sexual partner						
No	197	20	1			
Yes	139	47	3.33(1.89-5.87)	0.00	3.02(1.57-5.79)	0.001*
Tattooing						
No	126	18	1			
Yes	210	49	1.63(0.91-2.93)	0.099	1.05(0.54-2.03)	0.9
History of Abortion						
No	244	39	1			
Yes	92	28	1.9(1.11-3.27)	0.020	1.5(0.79-2.85)	0.217

Tooth extraction

No	223	29	1			
Yes	113	38	2.59(1.52-4.41)	0.00	1.4(0.72-2.71)	0.33

History of STI

No	264	32	1			
Yes	72	35	4(2.32-6.92)	0.00	3.38(1.76-6.47)	0.00*

Using Condom

Yes	61	4	1			
No	275	63	3.5(1.23-9.96)	0.019	4.99(1.5-16.16)	0.007*

Alcohol addiction

No	148	14	1			
Yes	188	53	2.98(1.59-5.58)	0.001	2.2(1.07-4.5)	0.031*

Smoke Cigarette

No	307	54	1			
Yes	29	13	2.55(1.25-5.21)	0.01	1.34(0.51-3.56)	0.55

364 **Key-***statically significant ($P \leq 0.05$), 1= reference value, pos= positive, Neg= negative N=
365 number, COR= crude odd ratio, AOR= adjusted odd ratio, CI= confidence interval,