Diagnostic utility of absolute neutrophil count as a new marker of spontaneous bacterial peritonitis; multicenter study

Nasser Mousa (✉ mousa_medic@yahoo.com)

Mansoura University

Ahmed Abdel-Razik

Mansoura University

Tarek Sheta

Mansoura University

Ahmed Deiab

Mansoura University

Khaled Taha

Mansoura University

Rasha Mahmoud

Mansoura University

Ahmed Yassen

Mansoura University

Alaa Habib

Mansoura University

Amany Amer

Menoufia University

Eman Mousa

Mansoura University

Niveen El-Wakeel

Mansoura University

Amr El-Sabbagh

Mansoura University

Mohmoud Awad

Mansoura University

Muhammad Diasty

Mansoura University

Aya Fathy

Mansoura University

Ola El-Emam

Mansoura University

Mostafa Abdelsalam

Mansoura University

Ayman Elgamal

Mansoura University

Sherif Elbaz

Aswan University

Essam Elmahdi

Mansoura University

Dina Elhammady

Mansoura University

Research Article

Keywords: Ascites, spontaneous bacterial peritonitis, absolute neutrophil count, C reactive protein, bacterial peritonitis, and liver cirrhosis.

Posted Date: February 15th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1331927/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Diagnostic utility of absolute neutrophil count as a new marker of spontaneous bacterial peritonitis; multicenter study

Nasser Mousa^{1,2}, Ahmed Abdel-Razik¹, Tarek Sheta³, Ahmed Deiab³, Khaled Taha³, Rasha Mahmoud³, Ahmed Yassen¹, Alaa Habib³, Amany Amer⁴, Eman Mousa⁵, Niveen El-Wakeel⁶, Amr El-Sabbagh⁶, Mohmoud Awad³, Muhammad Diasty¹, Aya Fathy⁷, Ola El-Emam⁸, Mostafa Abdelsalam⁹, Ayman Elgamal⁴, Sherif Elbaz¹⁰, Essam Elmahdi³, Dina Elhammady¹.

¹ Tropical Medicine Department, Mansoura University, Mansoura, Egypt.

² Hepatology and Gastroenterology Unit, Damietta Cardiology and Gastroenterology Center, Damietta, Egypt.

³ Internal Medicine Department, Mansoura University, Mansoura Egypt.

⁴ Department of Tropical Medicine, Menoufia University, Menoufia, Egypt.

⁵ Faculty of Dentistry, Mansoura University, Mansoura, Egypt.

⁶ Medical Microbiology and Immunology, Mansoura University, Mansoura, Egypt.

⁷ Public Health and Community Medicine, Mansoura University, Mansoura, Egypt.

⁸ Clinical Pathology Department, Mansoura University, Mansoura, Egypt.

⁹ Nephrology and Dialysis Unit, Internal Medicine Department, Faculty of Medicine, Mansoura University, Mansoura, Egypt.

¹⁰ Endemic Diseases and Gastroenterology Department, Aswan University, Aswan, Egypt

Corresponding author: Prof: Nasser Mousa, MD Tropical Medicine, Faculty of medicine. Mansoura University, Egypt. Mobile: 00201227029213-00201024501181.

E-mail.mousa_med@yaho.com

Abstract.

background. Spontaneous bacterial peritonitis (SBP) is a fatal complication of liver cirrhosis with high mortality rates.

Objective: The aim of this study is to investigate the diagnostic utility of absolute neutrophil count (ANC) as a non-invasive marker for SBP diagnosis.

Methods. Six hundred patients with cirrhotic ascites were included in the study. All patients underwent abdominal paracentesis and the ascitic fluid was processed for cell count and culture.

Results. Absolute neutrophil count was significantly higher in SBP versus non-SBP and in culture positive SBP versus culture negative SBP. ANC at cutoff value > 2.804 has 84% sensitivity and 78% specificity for diagnosis of SBP with positive and negative predictive values (79.4% and 83.6 respectively). At a cut-off point > 5.6 , ANC is capable of differentiating culture positive SBP from culture negative SBP cases with 62.07% sensitivity and 60.87 % specificity. Increased ANC, white blood cell (WBC), C reactive protein (CRP), creatinine and decreased platelet emerged as independent risk factors for SBP development, while increased ANC, WBC and decreased platelets were independent predictors of culture positive SBP.

Conclusion. This study demonstrates that, ANC count is simple, non-invasive diagnostic marker for SBP. Increased ANC, WBC, CRP, creatinine and decreased platelet emerged as independent risk factors for SBP development.

Keywords. Ascites, spontaneous bacterial peritonitis, absolute neutrophil count, C reactive protein, bacterial peritonitis, and liver cirrhosis.

Introduction

Ascites is a serious complication of decompensated liver cirrhosis. Although many pathogenic processes have been implicated in the development of ascites, about 75% likely occur as a result of portal hypertension in the setting of liver cirrhosis with the remainder due to infectious, inflammatory and infiltrative conditions.¹⁻⁶ In cirrhotic patients, bacterial infection increases the morbidity and mortality.⁷⁻⁹ Spontaneous bacterial peritonitis (SBP) is a risky complication that occurs amongst cirrhotic patients with ascites. SBP develops in approximately 10 to 30% of cirrhotic patients.¹⁰ The mortality rate in patients with spontaneous bacterial peritonitis ranges from 40-70% with hospital-mortality rate about 20%. However, nowadays the mortality from spontaneous bacterial peritonitis may be decreasing because of advances in its diagnosis and treatment.^{11,12} SBP results from qualitative and quantitative alterations in microbiota of the gut, bacterial translocation and increased permeability of the intestine.¹² Additionally, the impairment of the immune system observed in advanced cirrhotic patients plays an important role.¹³ The significance of SBP is mainly due to the late onset of symptoms, high rate of recurrence and the multiple causative organisms.¹⁴⁻¹⁶ The blood neutrophil count have diagnostic utilities in evaluation of severity and intensive care unit admission of many types of infection including SBP.^{17, 18} Serum CRP is synthesized mainly in liver cell under the control of many cytokines produced by macrophages at the time of bacterial infection.¹⁹

Although many non-invasive and invasive markers were studied as diagnostic tools for SBP.²⁰⁻²⁴, the diagnosis of SBP remains based on diagnostic paracentesis. However, diagnostic paracentesis is an invasive maneuver with many complications such as wound infection, abdominal wall hematoma, spontaneous hemoperitoneum that necessitating other non-invasive diagnostic tools.

Therefore, the aim of this study is to evaluate the clinical utility of absolute neutrophil count as a novel, simple, cheap, and noninvasive biochemical test for diagnosis of SBP.

Patients and methods

In this prospective study, we recruited 600 patients with liver cirrhosis and ascites referred to Tropical and Internal Medicine Departments, Mansoura University,

Department of Tropical Medicine, Menoufia University, and Damietta Cardiology and Gastroenterology Center from October 2018 to Jun 2020. All patients were subjected to complete history taking and physical examination include, abdominal ultrasonography and triphasic CT when indicated. The exclusion criteria were non-cirrhotic ascites (e.g. malignant ascites, tuberculus ascites), immunocompromised patients, and patients who had taken antibiotics prior to hospital admission or on prophylactic antibiotics for SBP or on anticoagulant medications. Furthermore, patients with heart or renal failure, neoplastic disorders, hematological disorders, clinically evident autoimmune disorders, secondary bacterial peritonitis due to any surgical causes and patients with unrelated infection that may influence the levels of blood WBC, e.g. skin and lung infection, were also excluded.

Sampling

1- The ascitic fluid samples (15 ml) were aspirated under the complete aseptic condition at the bedside, using the standard paracentesis technique. Ten ml was immediately inoculated in bedside aerobic and anaerobic blood culture bottles. The remaining amount of ascitic fluid was sent for biochemical and cytological examination in tubes containing EDTA and analyzed within 3 h of aspiration.²⁵ Ziehl-Neelsen staining of the ascitic fluid was done when needed.

Regarding microbiological examination, inoculated blood culture bottles were incubated for 3 successive days at 37°C with daily subculture on blood, MacConkey and chocolate agars. Antimicrobial susceptibility testing and bacterial identification were carried out using standard procedures.

According to international guidelines, SBP was diagnosed if the polymorph nuclear neutrophil (PMN) cell count in the ascitic fluid is $\geq 250/\text{mm}^3$ with positive culture of ascitic fluid (culture positive SBP) or culture negative neutrocytic ascites (CNNA) with $\text{PMN} > 250/\text{mm}^3$ and a negative ascitic fluid culture in the absence of other causes of peritonitis and hemorrhagic ascites.^{26,27} Patients with ascitic fluid PMN cell count < 250 cells/ mm^3 and a culture-negative ascitic fluid were assigned as the control group.

2. At the time of paracentesis, five ml of venous blood was withdrawn; one ml in polystyrene EDTA tube for CBC and four ml delivered in another tube and allowed to

clot. Non-hemolyzed sera were separated by centrifugation and used for determination of liver functions (ALT, AST, and albumin bilirubin and prothrombin time) blood sugar, creatinine, urea, alpha-fetoprotein and carcinoembryonic antigen. CRP was measured by particle-enhanced immunoturbidimetric assay (C-reactive protein (Roche Diagnostics). Blood absolute neutrophil count was calculated automatically using the CELL-DYN Emerald cell counter (Abbott, Wiesbaden, Germany).

This study was performed with the approval of the Mansoura Faculty of Medicine Institutional Review Board “MFM-IRB”. All patients provided written informed consent prior to participation in any protocol-specific procedure.

Statistical analysis

Data were analyzed with SPSS version 21. The normality of data was first tested with one-sample Kolmogorov-Smirnov test. Qualitative data were described using numbers and percentage. Association between categorical variables was tested using Chi-square. Continuous variables were presented as mean $\pm$ SD (standard deviation) for parametric data, but Median and IQR was used for non-parametric data. The two groups were compared with Student t test (parametric data) and Mann–Whitney test (non-parametric data). Clinical variables were considered statistically significant when ($p < 0.05$). Sensitivity and specificity of absolute neutrophil count for SBP diagnosis were estimated by receiver operator characteristic curve. Significant parameters in univariate binary logistic regression analysis were entered into a stepwise multivariate logistic regression analysis to identify the independent predictors for the occurrence of SBP and culture positive SBP.

Results

Table 1 shows the clinical, laboratory and demographic details of the studied groups. The two groups were matched for age, sex and etiology of cirrhosis (viral hepatitis, NASH and cryptogenic cirrhosis). Significant differences were found between both groups as regards, fever, abdominal pain, bilirubin, AST, ALT, creatinine, prothrombine time (PT), CRP, platelet count, blood WBC, ANC. No significant differences were found in serum albumin, blood lymphocyte and hemoglobin.

To determine predictive parameters for the diagnosis of SBP, variables with significant associations on univariate analysis were subjected to a logistic regression analysis. Increased ANC, WBC, CRP and creatinine and decreased platelets were independent predictors of SBP development (**Table 2**).

Among the 300 patients with SBP, 184 (61.3%) patients were culture positive SBP and 116 (38.6%) were culture negative SBP. No significant differences were found between both groups regarding, abdominal pain, fever, albumin, bilirubin, AST, ALT, INR, blood lymphocyte and haemoglobin. There was a significant increase in age and platelet count in patients with culture-negative compared to culture-positive SBP. A significant increase in, serum creatinine, CRP, blood WBC, blood neutrophil count was found in patients with culture-positive SBP vs. culture-negative SBP (**Table 3**).

To determine reliable predictive parameters for the diagnosis of culture positive SBP, variables with significant associations on univariate analysis were subjected to a logistic regression analysis. Decreased platelets and increased ANC and WBC and were independent predictors of culture positive SBP (**Table 4**).

The ROC curve analysis demonstrated that, absolute blood neutrophil count at a cut-off value of >2.804 had 84% sensitivity and 78% specificity for diagnosis of SBP with 79.4% PPV, 83.6% NPV and AUC equal to 0.88. (**Figure 1**). At cut-off point > 5.6 , absolute neutrophil count was capable of differentiating culture positive from culture negative SBP cases with 62.07% sensitivity and 60.87 % specificity with an AUC equal to 0.59. (**Figure 2**).

Discussion

In the present study, we investigated the use of absolute blood neutrophil count as a noninvasive diagnostic test for spontaneous bacterial peritonitis in cirrhotic ascitic patients. The advantages of this test are simple, low cost and reliability. It is known that, white blood cells (leukocytes) are a defense line against bacterial infection. The most abundant leukocytes are neutrophil, which are the first line of defense against microbial invasion.²⁸ Moreover, absolute neutrophil count can be used as a predictive factor for

infections.²⁹In general, bacterial infection is suspected if neutrophil in the blood are increased.³⁰

The diagnostic utility of the absolute neutrophil count for the diagnosis of spontaneous bacterial peritonitis is rarely found in research. Our study showed that, absolute neutrophil count is significantly higher in patients with spontaneous bacterial peritonitis versus non-SBP patients. In addition, absolute neutrophil count at a cut-off point > 2.804 is capable of differentiating spontaneous bacterial peritonitis from non-SBP with 84% sensitivity and 78% specificity with positive and negative predictive values of 79.4% and 83.6 respectively and AUC equal to 0.88.

The diagnostic utility of the blood neutrophil count as non-invasive test for the diagnosis of several infections has been previously suggested.³¹⁻³⁴ Mousa et al. evaluated the clinical utility of combined blood neutrophil to lymphocyte ratio (NLR) and C-reactive protein (CRP) as non-invasive tests for diagnosis of spontaneous bacterial peritonitis and concluded that, NLR was significantly higher in spontaneous bacterial peritonitis versus non-SBP, in addition, for SBP diagnosis, a blood NLR of >2.89 had a sensitivity of 80.3% and a specificity of 88.9%.³³

Many laboratory tests have been introduced as predictive parameters for SBP, including CRP level, platelet count, impaired prothrombin time and serum creatinine level.^{33,35,36} In this study, logistic regression analysis demonstrated that, increased blood neutrophil, WBC CRP and creatinine and decreased platelets were independent predictors of SBP development.

Even with the use of sensitive methods, ascites culture is negative in nearly 60% of patients with clinical manifestations suggestive of SBP and increased ascites neutrophil count.³⁷As culturing bacteria needs time, delays in antibacterial treatment can be fatal and result in the death of the patient from overwhelming infection. Mousa et al, in his study found no significant difference in patients with culture-positive SBP versus culture-negative SBP regarding NLR.³³ However, in our study, we found that, absolute neutrophil count is significantly higher in culture positive SBP versus culture negative SBP and at a cut-off point >5.6 , neutrophil count is capable of differentiating culture positive from culture negative SBP cases with 62.07% sensitivity and 60.87 % specificity with an AUC equal to 0.59.

In this study, logistic regression analysis demonstrated that, increased blood absolute neutrophil, WBC, CRP and decreased platelet count were independent predictors for culture positive SBP. Schwabl et al in his study found that, WBC were elevated in patients presenting with SBP versus cirrhotic ascitic patients without SBP.³⁶

Conclusion. Based on current evidence, the absolute blood neutrophil count, which requires only routine laboratory test could be used as a new, simple, cheap, non-invasive test for SBP diagnosis allowing rapid diagnosis and treatment of SBP so that reducing its complications.

Acknowledgements. The authors would like to express their appreciation for all staff in Tropical and Internal Medicine Departments, Mansoura University, Department of Tropical Medicine, Menoufia University, and Damietta Cardiology and Gastroenterology Center.

List of abbreviations.

SBP, spontaneous bacterial peritonitis; ANC, absolute neutrophil count; WBC, white blood cell; CRP, C reactive protein; PMN polymorph nuclear neutrophil; CNNA, culture negative neutrocytic ascites.

Declaration

Ethical Approval. This study was performed with the approval of the Mansoura Faculty of Medicine Institutional Review Board “MFM-IRB”.

Consent to participate. All patients provided written informed consent prior to participation in any protocol-specific procedure. All authors approve the publication of this work.

Availability of data and materials. Not applicable

Competing interests. The authors declare that they have no competing interests.

Funding. we have no funding

Authors' contributions.

Nasser Mousa, Ahmed Abdel-Razik, Tarek Sheta, developed the original idea and the protocol; Ahmed Deiab, Khaled Taha, Rasha Mahmoud, Ahmed Yassen, Alaa Habib analyzed data; Mohmoud Awad, Amany Amer, Eman Mousa, wrote the main manuscript text ; Niveen El-Wakeel, Amr El-Sabbagh, Narmen effat and Ola El-Emam, done laboratory work; Muhammad Diasty, Aya Fathy statistical

analysis; Mostafa Abdelsalam, Dina Elhammady, Ayman Elgamal prepared
figures, Sherif Elbaz, Essam Elmahdi revision of manuscript.

References

1. Ripoll C, Groszmann R, Garcia-Tsao G, Burroughs A, Planas R, Escorsell A, et al. Hepatic venous pressure gradient predicts clinical decompensation in patients with compensated cirrhosis. *Gastroenterology* 2007;133 (2):481-488.
2. Elhadidy A, Farid K. Eosinophilic ascites: Uncommon Presentation of Eosinophilic Gastroenteritis in children. *Med J Viral Hepatitis* 2019; 3 (2) - pp. 37-40.
3. Pericleous M, Sarnowski A, Moore A, Fijten R, Zaman M. The clinical management of abdominal ascites, spontaneous bacterial peritonitis and hepatorenal syndrome: a review of current guidelines and recommendations. *Eur J Gastroenterol Hepatol* 2016; 28(3):e10-18.
4. Elmoghazy M, El Shabrawi A and Mousa N. Portal Hypertension, an Overview. *Med J Viral Hepatitis* 2019;4 (1): 15-21.
5. Lucena MI, Andrade RJ, Tognoni G, Hidalgo R, De La Cuesta FS. Multicenter hospital study on prescribing patterns for prophylaxis and treatment of complications of cirrhosis. *Eur J Clin Pharmacol* 2002; 58:435-40.
6. Ibrahim I, Elbadrawy E, Nasr D. Frequency and characterization of mixed ascites among cirrhotic patients admitted to Zagazig University hospital. *Med J Viral Hepatitis* 2020;4 (2): 75-80.
7. Arvaniti V, D'Amico G, Fede G, Manousou P, Tsochatzis E, Pleguezuelo M. Infections in patients with cirrhosis increase mortality four-fold and should be used in determining prognosis. *Gastroenterology* 2010;139: 1246–1256.
8. El Shabrawi A, Mohammed Abdelaziz A, Mousa N. Infections in cirrhotic patients. *Med J Viral Hepatitis* 2019; 4 (1) - pp. 5-14.
9. Elbatae H, Ahmed Abdel-Razik A, Mousa E. Periodontal disease as predictor of chronic liver diseases. *Med J Viral Hepatitis* 2020; 4 (2):57-61.
10. Bonnel AR, Bunchorntavakul C, Reddy KR. Immune dysfunction and infections in patients with cirrhosis. *Clin Gastroenterol Hepatol* 2011;9:727-738.
11. Mandorfer M, Bota S, Schwabl P, Bucsics T, Pfisterer N, Kruzik M, et al. Nonselective β blockers increase risk for hepatorenal syndrome and death in patients with cirrhosis and spontaneous bacterial peritonitis. *Gastroenterology*. 2014 Jun. 146(7):1680-90.e

- 297 12. Amr S, Ahmed A, Hoda E. Spontaneous Bacterial Peritonitis: An Overview. *Med J*
298 *Viral Hepatitis* 2018; **3** (1) - pp. 13-17.
- 299 13. Jalan R, Fernandez J, Wiest R, Schnabl B, Moreau R, Angeli P. Bacterial infections
300 in cirrhosis: a position statement based on the EASL Special Conference 2013. *J Hepatol*
301 2014;**60**:1310–1324.
- 302 14. Garcia-Tsao G. Current Management of the Complications of Cirrhosis and Portal
303 Hypertension: Variceal Hemorrhage, Ascites, and Spontaneous Bacterial Peritonitis. *Dig*
304 *Dis*2016;34 (4): 382–386.
- 305 15. Lutz P, Nischalke HD, Strassburg CP, Spengler U. Spontaneous bacterial peritonitis:
306 The clinical challenge of a leaky gut and a cirrhotic liver. *World J Hepatol* 2015;7 (3):
307 304–314.
- 308 16. Fernández J, Bert F, Nicolas-Chanoine MH. The challenges of multi-drug-resistance
309 in hepatology. *J Hepatol* 2016;65(5):1043–1054.
- 310 17. Lee JH, Song S, Yoon SY, Lim CS, Song JW, Kim HS. Neutrophil to lymphocyte
311 ratio and platelet to lymphocyte ratio as diagnostic markers for pneumonia severity. *Br J*
312 *Biomed Sci* 2016;73(3):140–142.
- 313 18. Iliaz R, Ozpolat T, Baran B, Demir K, Kaymakoglu S, Besisik F. Predicting mortality
314 in patients with spontaneous bacterial peritonitis using routine inflammatory and
315 biochemical markers. *Eur J Gastroenterol Hepatol* 2018;30(7):786-791.
- 316 19. Strauss E, de Sá Ribeiro MDFG. Bacterial infections ssociated with hepatic
317 encephalopathy: prevalence and outcome. *Ann Hepatol*. 2003;2:41–50.
- 318 20. Abdel-Razik A, Mousa N, Elbaz S, Eissa M, Elhelaly R, Eldars W. Diagnostic utility
319 of interferon gamma-induced protein 10 kDa in spontaneous bacterial peritonitis: single-
320 center study. *Eur J Gastroenterol Hepatol* 2015;27(9):1087-1093.
- 321 21. Lee SS, Min HJ, Choi JY, Cho HC, Kim JJ, Lee JM, et al. Usefulness of ascitic fluid
322 lactoferrin levels in patients with liver cirrhosis. *BMC Gastroenterol* 2016;**13**;16(1):132.
- 323 22.Su DH, Zhuo C, Liao K, Cheng WB, Cheng H, Zhao XF. Value of serum
324 procalcitonin levels in predicting spontaneous bacterial
325 peritonitis. *Hepatogastroenterology* 2013;60(124):641-6.

- 326 23. Luo J, Wu X, Zhang Y, Huang W, Jia B. Role of ascitic prostaglandin E2 in diagnosis
327 of spontaneous bacterial peritonitis and prediction of in-hospital mortality in patients with
328 decompensated cirrhosis. *Medicine (Baltimore)* 2019;98(26):e16016.
- 329 24. Abdel-Razik A, Mousa N, Abdel-Aziz M, Elsherbiny W, Zakaria S, Shabana W, et al.
330 Mansoura simple scoring system for prediction of spontaneous bacterial peritonitis:
331 lesson learnt. *Eur J Gastroenterol Hepatol* 2019;31(8):1017-1024.
- 332 25. Nasser Mousa, Sahar Zakaria, Mohamed Abd EL Maksoud. The Predictive Factors
333 for Recurrence of Spontaneous Bacterial Peritonitis. *Med J Viral Hepatitis* 2018; 2 (2) -
334 pp. 25-29.
- 335 26. EASL Clinical Practice Guidelines for the management of patients with
336 decompensated cirrhosis. *Journal of Hepatology* 2018; 69: 406–460.
- 337 27. Runyon BA, AASLD. Introduction to the revised American Association for the study
338 of liver diseases practice guideline management of adult patients with ascites due to
339 cirrhosis 2012. *Hepatol Baltim Md.* (2013) ;57:1651–1653
- 340 28. Jenne CN, Liao S, Singh B. Neutrophils: multitasking first responders of immunity
341 and tissue homeostasis. *Cell Tissue Res* 2018;371(3):395-397.
- 342 29. Gibson C, Berliner N. How we evaluate and treat neutropenia in adults. *Blood*
343 2014;124(8):1251-8.
- 344 30. Shapiro MF, Greenfield S. The complete blood count and leukocyte differential
345 count: an approach to their rational application. *Ann Intern Med* 1987;106:65–74.
- 346 31. Alfar M, Abd El-Maksoud M, Mousa N. B-cell monoclonal lymphocytosis in chronic
347 hepatitis C virus infection. *Med J Viral Hepatitis* 2018;3:27–35.
- 348 32. Lee JH, Song S, Yoon SY, Lim CS, Song JW, Kim HS.. Neutrophil to lymphocyte
349 ratio and platelet to lymphocyte ratio as diagnostic markers for pneumonia severity. *Br J*
350 *Biomed Sci* 2016;73(3):140–142.
- 351 33. Mousa N, Besheer T, Abdel-Razik A, Hamed M, Deiab AG, Sheta T, et al. Can
352 combined blood neutrophil to lymphocyte ratio and C-reactive protein be used for
353 diagnosis of spontaneous bacterial peritonitis?. *Br J Biomed Sci* 2018;75(2):71-75.

354 34. Abdel-Razik A, Mousa N, Besheer TA, Eissa M, Elhelaly R, Arafa M, et al.
355 Neutrophil to lymphocyte ratio as a reliable marker to predict insulin resistance and
356 fibrosis stage in chronic hepatitis C virus infection. *Acta Gastroenterol Belg* 2015;
357 **78**:386–392.

358 35. Shi KQ, Fan YC, Ying L, Lin XF, Song M, Li LF, et al. Risk stratification of
359 spontaneous bacterial peritonitis in cirrhosis with ascites based on classification and
360 regression tree analysis. *Mol Biol Rep* 2012;**39**:6161–6169.

361 36. Schwabl P, Bucsics T, Soucek K, Mandorfer M, Bota S, Blacky A, et al. Risk factors
362 for development of spontaneous bacterial peritonitis and subsequent mortality in cirrhotic
363 patients with ascites. *Liver Int* 2015; 35 (9): 2121-8.

364 37. Rimola A, Garcia-Tsao G, Navasa M, Piddock LJ, Planas R, Bernard B, et al.
365 Diagnosis, treatment and prophylaxis of spontaneous bacterial peritonitis: a consensus
366 document. International Ascites Club. *J Hepatol* 2000;**32**:142–153.

380 **Table 1.** Clinical and biochemical characteristic of studied patient.

381

382

Parameters	SBP (Case) (N=300)	Non-SBP (Control) (N=300)	P value
Age	55.60±8.44	55.36±7.78	0.29
Sex (M/F)	207/93	194/106	0.14
HCV (N/%)	273 (91%)	275 (91.6%)	0.66
HBV (N/%)	15 (5%)	17 (5.6%)	0.71
NASH (N/%)	8 (2.6%)	5 (1.6%)	0.4
Cryptogenic (N/%)	4 (1.3%)	3 (1%)	0.7
Fever (N/%)	126 (37.3%)	0	<0.001
Abdominal pain (N%)	201(67%)	115/300	<0.001
Albumin (g/dl)	2.36±0.61	2.41±0.62	0.25
Bilirubin (mg/dL)	3.30 (1.80-6.30)	0.90 (0.70-4.10)	<0.001
ALT (U/L)	30.0 (20.25-42.00)	36.0 (20.0-56.0)	0.006
AST (U/L)	61.0 (37.0 -96.0)	31.0 (22.0-95.0)	<0.001
Creatinine (mg/dl)	1.5 (0.9-2.4)	1.4 (0.8-1.6)	0.003
PT (seconds)	13.9 ± 2.8	13.4 ± 2.6	0.023
CRP (mg/dl)	48.0 (44.0-48.0)	7.0 (5.0-9.0)	<0.001
Platelets (10 ⁹ /L)	79 (46-139)	93 (59-112)	0.006
Blood WBC (10 ³ /cmm)	8.13±3.23	4.88±1.60	<0.001
ANC (10 ³ /cmm)	5.98±2.86	2.25±0.6	<0.001
Blood lymphocyte (10 ³ /cmm)	1.66±0.79	1.64±0.89	0.77
Haemoglobin (g/dl)	10.10±1.87	10.05±2. 11	0.57

383

384 HCV, hepatitis C virus; HBV, hepatitis B virus; NASH, non-alcoholic steatohepatitis ;

385 ALT, alanine aminotransferase; AST, aspartate aminotransferase; PT, prothrombin

386 time;WBC, white blood cells; CRP, C reactive protein; ANC,Absolute neutrophil count.

387

388

389

390

391

392 **Table 2.** Logistic regression analysis of risk factors associated with SBP.

Parameters			95% CI		Sig.
	Exp (B)		Lower-Upper		
Bilirubin	1.002	0.911	1.102		0.969
AST	0.993	0.983	1.002		0.137
ALT	1.003	0.990	1.015		0.701
Creatinie	0.320	0.182	0.563		0.000
Platelets	1.05	0.975	1.533		0.011
CRP	0.761	0.000	1.03		0.009
Blood WBC	1.9	0.999	2.65		0.001
ANC	0.946	0.913	0.980		0.002

393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423

Table 3. Comparison between culture positive and culture negative patients with SBP.

Parameters	Culture Positive SBP (N=184)	Culture Negative SBP (N=116)	P value
Age	54.81±7.53	56.86±9.82	0.02
Abdominal pain (N/%)	128 (69.5.9%)	73 (62.9%)	0.23
Fever (N/%)	73 (39.67%)	53 (45.6%)	0.30
Albumin (g/dl)	2.42±.65	2.34±.59	0.5
Bilirubin (mg/dl)	3.5 (2.0-5.05)	3.3 (1.80-7.1)	0.36
ALT (U/L)	48 (33-74)	48 (36.5-80.5)	0.53
AST (U/L)	33 (20-37)	29 (21-43)	0.17
Creatinine (mg/dl)	1.5 (0.9-2.4)	1.4 (0.8-1.6)	0.003
INR	1.73±0.53	1.60±0.51	0.14
CRP (mg/dl)	48 (12-96)	47 (24-48)	0.02
Platelets (10 ⁹ /L) Min-max	72.5 (46-128) (12-210)	97 (45.7-157.2) (10-41)	0.001
Blood WBC (10 ³ /cmm)	8.79±3.7	7.80±3.28	0.008
ANC(10 ³ /cmm)	6.54±3.0	5.64±2.6	0.003
Blood lymphocytes (10 ³ /cmm)	1.4±0.19	1.6±0.23	0.5
Haemoglobin (g/dl)	10.24±2.15	10.05±1.67	0.20

CRP, C reactive protein; ANC, Absolute neutrophil count.

447 **Table 4.** Logistic regression analysis of risk factors associated with culture positive SBP.

Parameters	Exp (B)	95% CI		Sig.
		Lower	Upper	
Age	1.030	0.972	1.091	0.314
Creatinie	.976	0.672	1.418	0.898
Platelets	0.921	0.901	1.77	0.016
WBCs	.999	0.922	1.45	0.001
ANC	1.081	1.036	1.127	0.001

448
449
450
451
452
453
454
455
456
457
458

Figure 1. Receiver operator characteristics curve for blood absolute neutrophil count in cases of spontaneous bacterial peritonitis (case control).

466

467

468 **Figure 2.** Receiver operator characteristics curve for blood absolute neutrophil count in
469 culture-positive and culture negative spontaneous bacterial peritonitis.

470

471