

The predictive value of lncRNA MIR31HG expression on clinical outcomes in patients with solid malignant tumors

Chao Tu

The Second Xiangya Hospital, Central South University

Xiaolei Ren

The Second Xiangya Hospital, Central South University

Jieyu He

The Second Xiangya Hospital, Central South University

Shuangqing Li

The Second Xiangya Hospital, Central South University

Lin Qi

The Second Xiangya Hospital, Central South University

Zhixi Duan

The Second Xiangya Hospital, Central South University

Wanchun Wang

The Second Xiangya Hospital, Central South University

Zhihong Li (✉ lizhihong@csu.edu.cn)

The Second Xiangya Hospital, Central South University <https://orcid.org/0000-0002-1944-9671>

Primary research

Keywords: lncRNA MIR31HG; sarcoma; tumor; prognosis; overall survival

Posted Date: February 6th, 2020

DOI: <https://doi.org/10.21203/rs.2.22773/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Cancer Cell International on April 7th, 2020. See the published version at <https://doi.org/10.1186/s12935-020-01194-y>.

Abstract

Background Emerging studies have explored the prognostic value of MIR31HG in cancers, but its role remains elusive. Herein, we aimed to summarize the prognostic potential of MIR31HG in this study. **Methods** Several databases were searched for literature retrieval on Dec 5, 2019. Overall and subgroup analyses were conducted to measure the relationship between MIR31HG expression and clinical outcomes. Moreover, GEPIA was applied for validation of prognostic value of MIR31HG in tumor patients in TCGA dataset. **Results** Overall, seventeen studies with 2,573 patients were enrolled. Compared to counterparts, those patients with high MIR31HG expression tended to have shorter RFS. Notably, MIR31HG overexpression predicted unfavorable OS in lung cancer. By contrast, gastrointestinal cancer patients with elevated MIR31HG expression predicted better OS and disease-free survival. Additionally, MIR31HG overexpression was significantly associated with worse clinicopathological features including advanced tumor stage and LNM in lung cancer, but favorable clinical characteristics in gastrointestinal cancer. Moreover, the positive association between MIR31HG and OS in lung cancer was further confirmed in TCGA dataset. **Conclusion** Overexpression of MIR31HG suggested remarkable association with poor prognosis in terms of OS, tumor stage, and LNM in lung cancer, but favorable prognosis in gastrointestinal cancer. Therefore, MIR31HG may serve as a promising prognostic biomarker in multiple cancers.

Background

Cancer is the major cause of death and public health problem worldwide (1). In spite of notable continuous advances in cancer treatment over the last decades, the patients' survival rate remains disappointed (2). Nowadays, only a minority of patients in early clinical stage could benefit from curative approaches, while the rest patients, particularly those with advanced stage, may be resistant to conventional treatments (3). Against this backdrop, it is necessary to explore reliable biomarkers harboring sufficient sensitivity and specificity for early diagnosis and attractive targets for therapeutic intervention before reaching to advanced stages (4).

Long non-coding RNAs (lncRNAs), longer than 200 nucleotides (nt), are defined as highly conserved non-coding RNAs with limited or no protein-coding capacity (5). LncRNA are involved in diverse physiological and pathological cellular processes (6) by promoting or restraining the protein-coding genes expression (7). Accumulating evidences have showed that aberrant expression of lncRNA could drive tumor phenotypes, including initiation, invasion, and metastasis, via interacting with other cellular macromolecules (8). More recently, emerging lncRNAs have been identified and characterized as crucial factors in tumorigenesis, with certain reports highlighting them as biomarkers and targets in various cancers, such as breast cancer anti-estrogen resistance 4 (BCAR4) (9), small nucleolar RNA host gene 16 (SNHG16) (10), and TTN antisense RNA 1 (TTN-AS1) (11).

LncRNA microRNA-31 host gene (MIR31HG), also known as LOC554202 (12), is a newly identified lncRNA with a length of 2,166 nt (2, 4). Meanwhile, MIR31HG was also named as LncHIFCAR (long noncoding HIF-1 α co-activating RNA) since it is a hypoxia-inducible lncRNA (4). P16 (INK4A) is a well-known tumor suppressor, MIR31HG was reported to negatively regulate p16-dependent senescence phenotype (13), suggesting a potential role in carcinogenesis.

Recently, an increasing number of studies have been exploring the prognostic association between MIR31HG and cancers. Abnormal expression of MIR31HG has been reported in a diverse range of cancer types, including osteosarcoma (14), chordoma (15), colorectal cancer (CRC) (16–18), hepatocellular carcinoma (HCC) (5), lung cancer (2, 19–23), gastric cancer (24), bladder cancer (25), pancreatic ductal adenocarcinoma (PDAC) (6), oral squamous cell carcinoma (OSCC) (4), breast cancer (12, 26), cervical cancer (27), vulvar squamous cell carcinoma (VSCC) (28), and esophageal squamous cell carcinoma (ESCC) (29, 30). Furthermore, MIR31HG expression was remarkably associated with survival time, tumor-node metastasis (TNM) stage and carcinogenesis of certain cancers. However, the prognostic significance of MIR31HG may be jeopardized by controversial results, ethnic or geographical limitations, and limited sample size. For example, Sun K et al. (29) reported that MIR31HG expression was significantly elevated in ESCC tissues and plasma, and MIR31HG could promote ESCC cell proliferation, migration and invasion (29). On the contrary, another study revealed that MIR31HG appeared to have lower expression in ESCC tissues and patients with reduced MIR31HG were correlated with malignant clinical features. The

contradictory expression patterns of MIR31HG were also found in breast cancer (12, 26), CRC (18, 31, 32) and gastric cancer (24, 33). In view of these conflicting data, for the first time, we performed this comprehensive meta-analysis of the current evidences in order to determine the predictive value of MIR31HG in various carcinomas.

Materials And Methods

Literature search strategy

Studies in relevant with MIR31HG and cancer prognosis were thoroughly searched via PubMed, Web of Science, and Cochrane library (up to Dec 5, 2019). The search strategy was used as following: ("Cancer" OR "Malignancy" OR "Sarcoma" OR "Tumor" OR "Neoplasm" OR "Carcinoma") AND ("MIR31HG" OR "LOC554202" OR "LncHIFCAR") AND ("Long non-coding RNA" OR "LncRNA" OR "long untranslated RNA" OR "lincRNA") AND ("Prognosis" OR "Prognostic"). The reference lists of enrolled studies were further reviewed to identify relevant articles.

Study selection criteria and data extraction

Studies were included based on the following inclusion criteria: (1) original studies; (2) the diagnosis of cancer was validated by pathology and histology; (3) studies exploring the correlation between MIR31HG and cancers; (4) MIR31HG expression levels were measured and divided into high or low expression group; (5) reporting of at least one of the endpoints: overall survival (OS), disease-free survival (DFS) and relapse-free survival (RFS), et al; (6) sufficient data: the hazard ratios (HRs) or odds ratios (ORs) with 95% confidence intervals (CIs) or K-M survival curves were shown; (7) the more recent article was selected if duplicated researches were reported.

On contrary, studies were excluded due to following exclusion criteria: (1) article type: meta-analysis, reviews, editorial material, or case reports; (2) in vivo or in vitro studies without clinical data; (3) sufficient survival data were not provided or cannot be extracted; (4) full text was not accessible; (5) duplicates;

Two independent investigators (CT and XLR) screened and extracted the data from selected studies according to the criteria abovementioned. Any divergence views were resolved by another senior researcher (WCW) to reach a consensus. The following items were provided: first author, year of publication, country, sample size, tumor type, detection methods, follow-up time, and prognostic outcomes. HRs (ORs) and 95% CIs were obtained from the original texts in eligible studies as provided. If not, Engauge Digitizer (version 4.1) would be used to calculate the HR through K-M survival curve as previously described (10).

Quality assessment

Two investigators (JYH and SQL) assessed the quality of studies independently in accordance with the Newcastle-Ottawa scale (NOS). Usually, studies with $NOS \geq 6$ were considered as high-quality. The scores of all eligible studies in this study ranged from 6 to 8, which indicated high quality.

Prognosis analysis in GEPIA using TCGA dataset

Gene Expression Profiling Interactive Analysis (GEPIA) (<http://gepia.cancer-pku.cn/index.html>) based on The Cancer Genome Atlas (TCGA) data (34) was further searched for cross-validation of the prognostic significance of MIR31HG in cancers.

Publication bias and sensitivity analysis

Sensitivity analysis was performed by omitting individual study consecutively to evaluate their impact on this study. Begg's test and Egger's test were both utilized to evaluate the publication bias.

Statistical analysis

Data analysis was conducted by using STATA (version 12.0) and RevMan (Version 5.3). A two-tailed P value less than 0.05 was considered statistically significant. HRs and corresponding 95% CIs were adopted to evaluate the relation between MIR31HG expression and cancer prognosis as well as other clinicopathological characteristics. Heterogeneity of the included studies was

assessed by Cochran's Q statistic and I^2 test. If $p \leq 0.05$, or $I^2 \geq 50\%$, a random-effects model shall be applied to calculate the pooled results due to substantial heterogeneity. Otherwise, a fixed-effects model would be used. Subgroup analysis was performed as per the tumor type and follow-up months in case of obvious heterogeneity among the selected studies.

Results

Characteristics of the included studies

According to the screening criteria, seventeen articles comprising 2,573 patients were included in our study (1, 2, 4, 5, 14, 16, 18, 22, 24, 25, 27–32, 35). All the included studies were related to lncRNA MIR31HG in cancer and comprised clinical data. The study screening flow diagram was shown in Fig. 1. Among the included seventeen articles, twelve types of cancers were identified, including osteosarcoma, cervical cancer, gastric cancer, bladder cancer, CRC, lung adenocarcinoma (LUAD), non-small-cell lung carcinoma (NSCLC), HCC, VSCC, ESCC, OSCC, and laryngeal squamous cell carcinoma (LSCC). The included studies were performed between 2015 and 2019, and almost all of them were originated in China except one from Norway (16). The sample size ranged from 16 to 1265. Survival analysis regarding OS, DFS and RFS were measured in twelve studies with follow-up time ranging from 25 to 100 months. Quantitative reverse transcription polymerase chain reaction (qRT-PCR) was utilized for analysis of the relative expression level of MIR31HG in tumor tissues except one used RNA sequencing (4). The cut-off values of MIR31HG expression were median value. NOS scores were calculated for the quality evaluation with value ranging from 6 to 8, indicating that the enrolled studies were of high quality. Details of the abovementioned characteristics of the studies were presented in Table 1.

Table 1
Baseline characteristics of eligible studies.

Authors	Publication year	Country	Cancer type	Sample size	Follow-up months	Detection method	Clinical outcomes	Analysis method	NOS
Chen J, et al. [27]	2017	China	Cervical cancer	120	60	qRT-PCR	OS	Multivariate	7
Ding J, et al. [18]	2015	China	CRC	48	-	qRT-PCR	-	Multivariate	6
Eide, PW, et al. [16]	2019	Norway	CRC	1265	60	qRT-PCR	RFS	Multivariate	7
He A, et al. [25]	2016	China	Bladder cancer	55	-	qRT-PCR	-	Multivariate	6
Li Y, et al. [31]	2018	China	CRC	157	100	qRT-PCR	OS, DFS	Multivariate	7
Ni S, et al. [28]	2016	China	VSCC	16	-	qRT-PCR	-	Multivariate	6
Nie F, et al. [24]	2016	China	Gastric cancer	42	40	qRT-PCR	OS	Multivariate	6
Qin J, et al. [22]	2018	China	LUAD	112	85	qRT-PCR	OS	Multivariate	7
Ren Z, et al. [30]	2017	China	ESCC	185	60	qRT-PCR	OS	Multivariate	6
Shih J, et al. [4]	2017	China	OSCC	42	40	RNA-seq	OS, RFS	Multivariate	7
Sun K, et al. [29]	2018	China	ESCC	53	-	qRT-PCR	-	Multivariate	6
Sun Y, et al. [14]	2019	China	Osteosarcoma	40	-	qRT-PCR	-	Multivariate	6
Wang R, et al. [35]	2018	China	LSCC	60	85	qRT-PCR	OS, RFS	Multivariate	7
Wu D, et al. [40]	2019	China	NSCLC	50	85	qRT-PCR	OS	Multivariate	8
Yan S, et al. [5]	2018	China	HCC	42	25	qRT-PCR	OS	Multivariate	7

Abbreviation: CRC: colorectal cancer; DFS: disease-free survival; ESCC: esophageal squamous cell carcinoma; HCC: hepatocellular carcinoma; LSCC: laryngeal squamous cell cancer; LUAD; lung adenocarcinoma; NOS: Newcastle-Ottawa Scale; NSCLC: Nonsmall cell lung cancer; OS: overall survival; OSCC: oral squamous cell carcinoma; qRT-PCR: real-time quantitative reverse transcription polymerase chain reaction; RFS: relapse-free survival; VSCC: vulvar squamous cell carcinoma

Authors	Publication year	Country	Cancer type	Sample size	Follow-up months	Detection method	Clinical outcomes	Analysis method	NOS
Yang L, et al. [32]	2016	China	CRC	178	70	qRT-PCR	OS, DFS	Multivariate	7
Zheng S, et al. [2]	2019	China	NSCLC	88	50	qRT-PCR	OS	Multivariate	8

Abbreviation: CRC: colorectal cancer; DFS: disease-free survival; ESCC: esophageal squamous cell carcinoma; HCC: hepatocellular carcinoma; LSCC: laryngeal squamous cell cancer; LUAD; lung adenocarcinoma; NOS: Newcastle-Ottawa Scale; NSCLC: Non-small cell lung cancer; OS: overall survival; OSCC: oral squamous cell carcinoma; qRT-PCR: real-time quantitative reverse transcription polymerase chain reaction; RFS: relapse-free survival; VSCC: vulvar squamous cell carcinoma

Expression of lncRNA MIR31HG as a prognosis marker in various cancers

In this study, the association between prognosis and MIR31HG expression were reported in twelve studies. Eleven studies were included for OS meta-analysis. As the result of meta-analysis exhibited obvious heterogeneity ($P < 0.001$, $I^2 = 88.1\%$), subgroup analysis was further conducted to calculate the pooled HRs or ORs and 95% CIs in which the patients were stratified by cancer type (gastrointestinal cancer, lung cancer or others), or follow-up time (more or less than 50 months), as shown in Fig. 2A and Figure S1, respectively. The gastrointestinal cancer included ESCC, gastric cancer, HCC, and CRC, while the lung cancer consisted of NSCLC and LUAD. Besides, OSCC, LSCC and cervical cancer were grouped in other cancers. The random-effect model was applied to reduce the possible bias. All the pooled HRs and 95% CIs of the overall and subgroup analysis were shown in Table 2. The results demonstrated that elevated expression of the MIR31HG predicted favorable OS in gastrointestinal cancer (HR = 0.56, 95% CI: 0.41–0.77), but unfavorable OS in lung cancer (HR = 1.64, 95% CI: 1.15–2.34) and other cancers (HR = 3.37, 95% CI: 2.69–4.23) (Fig. 2A and Table 2). Interestingly, the pooled results also showed that overexpression of MIR31HG predicted shorter OS in studies with follow-up months less than 50, while no correlation was found in studies with follow-up more than 50 months (Figure S1).

Table 2
The pooled results of association between clinical prognosis and MIR31HG expression.

Outcomes		Studies	Participants	HRs and 95% CIs	Model	Heterogeneity	
							P, I ²
OS	Total	11	1096	1.10 (0.60, 2.03)	Random	< 0.001, 88.1%	
	Gastrointestinal cancer	5	604	0.56 (0.41, 0.77)	Random	0.825, 0%	
	Lung cancer	3	270	1.64 (1.15, 2.34)	Random	0.615, 0%	
	Other cancers	3	222	3.37 (2.69, 4.23)	Random	0.746, 0%	
DFS		2	335	0.44 (0.27, 0.72)	Fixed	0.551, 0%	
RFS		3	1367	2.54 (1.91, 3.38)	Fixed	0.610, 0%	

Abbreviation: CI: confidence interval; DFS: disease-free survival; HR: hazard ratio; OS: overall survival; RFS: relapse-free survival.

Moreover, two studies on CRC for DFS and three studies on various cancers reported available data for RFS were further pooled for measurement. The HR and 95% CI (HR = 0.44, 95% CI: 0.27–0.72) indicated overexpression of MIR31HG predicted favorable

DFS in CRC (Fig. 2B). However, overexpression of MIR31HG was correlated with poor RFS (HR = 2.54, 95% CI: 1.91–3.38) (Fig. 2C).

Association between MIR31HG expression and other clinicopathologic parameters

The association between MIR31HG and clinicopathologic features, including clinical stage, lymph node metastasis (LNM), distant metastasis (DM), and tumor size were further analyzed by ORs and their 95% CIs. There was no significant difference in MIR31HG expression detected in DM (OR = 0.75, 95% CI: 0.31–1.85) and tumor size (OR = 0.95, 95% CI: 0.58–1.56), even if the stratified analysis were performed by cancer type (Fig. 3C and 3D). Although, the pooled overall OR and 95% CI showed no significant difference in MIR31HG expression analysis in various clinical stage (OR = 1.41, 95% CI: 0.77–2.56) and LNM (OR = 1.65, 95% CI: 0.93–2.91), after subgroups analysis by cancer type, the results suggested that upregulated expression of MIR31HG indicated the advanced clinical stage (OR = 6.26, 95% CI: 3.53–11.11) and higher risk of LNM in lung cancer (OR = 2.08, 95% CI: 1.04–4.15) (Fig. 3A and 3B).

Sensitivity analysis

Sensitivity analysis was performed to the studies which had more than two articles. All results showed that this meta-analysis was stable except the analysis of clinical stage (Fig. 5B) and LNM (Fig. 5C). After dropping the research performed by Sun. K. et al, the pooled ORs and 95% CIs showed that elevated expression of MIR31HG were negatively associated with clinical stage (OR = 0.30, 95% CI: 0.20–0.44) and LNM risk in gastrointestinal cancer (OR = 0.55, 95% CI: 0.32–0.95) (Fig. 4A and 4B).

Publication bias

Publication bias was measured by using the funnel plots, Begg's and Egger's test. No significant publication was observed as shown in Fig. 5A.

Prognostic analysis in GEPIA using TCGA dataset

Furthermore, the relationships between expression of MIR31HG and OS/ DFS in lung cancers were validated in TCGA dataset by using GEPIA as previously described (10). The results indicated that high MIR31HG expression was correlated with significant shorter OS (HR = 1.3, P = 0.012), but not DFS (HR = 0.99, P = 0.96) (Fig. 6).

Discussion

Recently, with the development of next generation sequencing, lncRNAs are revealed as multifaceted regulators in a wide range of cellular homeostasis (36) via participating in epigenetic, transcriptional and posttranscriptional regulation of genes (25). Increasing evidences have established a strong relationship between dysfunction of lncRNAs and cell fate determination as well as disease pathogenesis, such as aging (37), arthritis (38), and cancer (12). LncRNA MIR31HG, locating at human chromosome 9p21.3 (39) with transcription regulated by methylation of the promoter region (26), was found aberrantly expressed in several types of cancers. Specifically, MIR31HG was found elevated expressed in breast cancer (12), cervical cancer (27), chordoma (15), osteosarcoma (14), lung cancer (2, 22, 40), OSCC (4), PDAC (6), VSCC (28) and LSCC (35, 41, 42). Nevertheless, the expression levels of MIR31HG was reported down-regulated in triple-negative breast cancer (TNBC) cell lines of basal subtype (26), bladder cancer (25), gastric cancer (24), CRC (31, 32), and HCC (5). Besides, abnormal expression of MIR31HG was deemed as potential biomarker to reflect the clinicopathological characteristics and prognostic outcomes of cancer patients (27). For instance, increased MIR31HG expression was found both in LUAD/ NSCLC tissues when compared with counterparts, and positively associated with unfavorable TNM stage, metastasis, histological differentiated degree, and forecasted shorter OS (2, 22, 40). Notably, MIR31HG expression was even higher in gefitinib-resistant NSCLC cells, and knockdown of MIR31HG could promote cell apoptosis and cell cycle arrest, and subsequently induce gefitinib sensitivity (19). More recently, He J et al. further verified that MIR31HG was also remarkably increased in gefitinib-resistant NSCLC patients, and MIR31HG overexpression contributed to reduced sensitivity of NSCLC cell to gefitinib in vitro (43). By contrast, Yan S et al. reported that overexpression of MIR31HG significantly inhibited HCC proliferation and metastasis in vitro and impeded

tumorigenesis *in vivo* (5). In CRC, the MIR31HG expression was negatively correlated with unfavorable prognosis, as indicated by advanced pathologic stage, and larger tumor size (18). Consistently, Li Y et al. found that high expression of MIR31HG predict high OS and DFS in CRC patients treated with oxaliplatin (31).

Given the prognostic significance of MIR31HG in cancers, in the current study, we firstly explored the association between MIR31HG level and clinical outcomes by performing a meta-analysis containing seventeen literatures with 2,573 patients. The pooled results of subgroup analysis as per the tumor types demonstrated that high MIR31HG expression predicted unfavorable OS in patients with lung cancer and other cancers, and poor RFS in all selected studies, respectively. On the contrary, overexpression of MIR31HG was closely associated with favorable OS and DFS in gastrointestinal cancer, indicating a tissue-specific predictive prognosis of MIR31HG in multiple human cancers. Besides, the relationship between expression of MIR31HG and other parameters were analyzed. Consistent with the predictive value of MIR31HG in OS, high level of MIR31HG was also significantly correlated with advanced clinical stage and LNM potential in lung cancer. Conversely, MIR31HG expression was negatively associated with clinical stage and LNM risk in gastrointestinal cancer. However, no correlation between MIR31HG expression and other clinical characteristics, such as DM and tumor size, were noted. Beside, in order to increase the credibility of the results, we performed cross-validation of the results by using TCGA dataset. The results showed the elevated MIR31HG was significantly correlated with poor OS rather than DFS in lung cancer.

Mechanistically, lncRNA could execute versatile functions in diseases through alternative splicing, epigenetic modulation, chromatin modification, scaffolding/decoy function or acting as molecular sponge (44). To begin with, given the critical role of MIR31HG in hypoxia-associated malignant progression, MIR31HG was observed to promote tumorigenesis by serving as a HIF-1 α co-activator and regulating HIF-1 transcriptional network (4, 41). For instance, in LSCC, MIR31HG could facilitate cell proliferation, cell cycle progression and suppress apoptosis via HIF-1 α and p21 (35). In OSCC, MIR31HG knockdown impaired the HIF-1 α transactivation, sphere-forming ability, metabolic shift and metastatic cascade both *in vitro* and *in vivo* (4). Besides, lncRNA could involve in the competing endogenous RNA (ceRNA) regulatory work to function as endogenous miRNA sponge (Table 3) (5). As the host gene of miR-31, MIR31HG was firstly identified to co-express or modulate the expression of miR-31 in certain cancers (43). For example, MIR31HG could enhance proliferation, migration and invasion by up-regulating EZH2/miR-31 and then indirectly activating the oncogene RNF144B in chordoma (15). Yang S et al. demonstrated that elevated MIR31HG could inhibit miR-31 expression, and increase RhoA in LSCC, and via which promote cell growth, cell cycle and invasion (42). In addition to miR-31, other miRNAs were also identified to interact with MIR31HG in carcinogenesis. It was reported that MIR31HG could directly sponge to tumor suppressor- miR-361, rather than miR-31, and in turn regulate cell proliferation, cell cycle arrest, and apoptosis via targeting VEGF, FOXM1 and Twist in osteosarcoma (14). In NSCLC, MIR31HG behaved as an oncogene by inhibiting miR-214 expression, thereby facilitating cancer cell migration and invasion (40). In HCC cancer, MIR31HG could competitively bind miR-575, and positively regulate ST7L to impair cell proliferation and metastasis (5). Moreover, MIR31HG could regulate tumorigenesis by targeting a broad spectrum of target genes or pivotal signaling pathways (Table 3). In gastric cancer, MIR31HG may suppress cell proliferation and hinder tumorigenesis partly by regulation of E2F1 and p21 expression (24). In NSCLC, MIR31HG was found to enhance Wnt/ β -catenin pathway, and induce epithelial-mesenchymal transition (EMT) phenotype (2). Jin C et al. identified that MIR31HG was induced by nuclear translocation of NF- κ B, and in turn directly bind to I κ B α and participated in NF- κ B activation, revealing an interaction between MIR31HG and NF- κ B in osteogenic differentiation (45). Of note, MIR31HG could modulate gefitinib resistance partly through EGFR/PI3K/Akt (19, 43) or RAF-MEK-ERK pathways (43). Furthermore, Zheng S and colleagues discovered that down-regulation of MIR31HG inhibited NSCLC cell proliferation, invasion and EMT phenotype via up-regulating E-cadherin expression, inhibiting Wnt/ β -catenin cascade, and down-regulating expression of Twist1 and vimentin (2).

Table 3

Expression pattern of MIR31HG in cancers and its interaction with miRNAs and target genes

Tumor type	Expression	Signaling pathways	miRNAs and target genes	Biological functions	References
Osteosarcoma	↑	/	miR-361, VEGF, FOXM1, Twist	Cell proliferation, migration, apoptosis, G1/G arrest, EMT	[14]
Chordoma	↑	/	miR-31, EZH2, RNF144B	Cell proliferation, migration, invasion, EMT	[15]
OSCC	↑	/	HIF-1 α , p300	Sphere-forming ability, metabolic shift and metastasis	[4]
HNSCC/LSCC	↑	/	HIF-1 α , p21	Cell proliferation, cell cycle progression, apoptosis	[35]
LSCC	↑	/	miR-31, RhoA	Cell growth, cell cycle, invasion	[42]
Breast cancer	↑	/	/	Cell proliferation, apoptosis, migration, invasion	[12]
Cervical cancer	↑	/	/	Cell proliferation, apoptosis	[27]
VSCC	↑	/	/	/	[28]
NSCLC	↑	Wnt/ β -catenin	GSK3 β , Twist1, Vimentin, E-cadherin	Cell proliferation, invasion, EMT	[2]
NSCLC	↑	PI3K/Akt	EGFR	Cell proliferation, apoptosis, G0/G1 arrest, gefitinib resistance	[19]
NSCLC	↑	RAF/MEK/ERK, PI3K/Akt	miR-31, FIH-1, RASA1	Cell proliferation, clonogenic growth, gefitinib resistance	[43]
NSCLC	↑	/	miR-214	Cell migration, invasion	[40]
LUAD	↑	/	/	Cell proliferation, cell cycle arrest	[22]
PDAC	↑	/	miR-193b, CCND1, Mcl-1, NT5E, KRAS, uPA, ETS1	Cell growth, apoptosis, G1/S arrest, invasion	[6]
ESCC	↑	/	Furin, MMP1	Cell proliferation, migration, invasion	[29]
ESCC	↓	/	/	/	[30]
Gastric cancer	↑	/	p21, E-cadherin	Cell proliferation, migration	[33]
Gastric cancer	↓	/	E2F1	Cell proliferation	[24]

Abbreviation: ↑up-regulated; ↓down-regulated; Akt: protein kinase B; CCND1: cyclin D1; CRC: colorectal cancer; E2F1: E2F transcription factor 1; EGFR: epidermal growth factor receptor; EMT: Epithelial-mesenchymal transition; ERK: extracellular regulated protein kinases; ESCC: esophageal squamous cell carcinoma; ETS1: ETS proto-oncogene 1, transcription factor; EZH2: enhancer of zeste 2 polycomb repressive complex 2 subunit; FIH-1: hypoxia inducible factor 1 subunit alpha inhibitor; FOXM1: forkhead box M1; GSK-3 β : glycogen synthase kinase-3 β ; HCC: hepatocellular carcinoma; HIF-1 α : hypoxia inducible factor 1 subunit alpha; HNSCC: head and neck squamous cell carcinoma; IFN- α/γ : interferon- α/γ ; LSCC: laryngeal squamous cell cancer; LUAD; lung adenocarcinoma; MMP1: matrix metalloproteinase 1; NSCLC: Non-small cell lung cancer; NT5E: 5'-nucleotidase ecto; OSCC: oral squamous cell carcinoma; PDAC: pancreatic ductal adenocarcinoma; PI3K: phosphoinositide 3-kinase; RASA1: RAS p21 protein activator 1; RhoA: ras homolog gene family, member A; ST7L: suppression of tumorigenicity 7 like; TGF- β : transforming growth factor- β ; TNF: tumor necrosis factor; VEGF: vascular endothelial growth factor; VSCC: vulvar squamous cell carcinoma.

Tumor type	Expression	Signaling pathways	miRNAs and target genes	Biological functions	References
CRC	↑	/	miR-31-5p, MYC, TNF- α /NF- κ B, TGF- β , IFN- α / γ	EMT	[16]
CRC	↓	/	/	Cell proliferation, apoptosis	[18]
HCC	↓	/	miR-575, ST7L	Cell proliferation, metastasis	[5]
Bladder cancer	↓	/	/	/	[25]

Abbreviation: ↑up-regulated; ↓down-regulated; Akt: protein kinase B; CCND1: cyclin D1; CRC: colorectal cancer; E2F1: E2F transcription factor 1; EGFR: epidermal growth factor receptor; EMT: Epithelial-mesenchymal transition; ERK: extracellular regulated protein kinases; ESCC: esophageal squamous cell carcinoma; ETS1: ETS proto-oncogene 1, transcription factor; EZH2: enhancer of zeste 2 polycomb repressive complex 2 subunit; FIH-1: hypoxia inducible factor 1 subunit alpha inhibitor; FOXM1: forkhead box M1; GSK-3 β : glycogen synthase kinase-3 β ; HCC: hepatocellular carcinoma; HIF-1 α : hypoxia inducible factor 1 subunit alpha; HNSCC: head and neck squamous cell carcinoma; IFN- α / γ : interferon- α / γ ; LSCC: laryngeal squamous cell cancer; LUAD; lung adenocarcinoma; MMP1: matrix metalloproteinase 1; NSCLC: Non-small cell lung cancer; NT5E: 5'-nucleotidase ecto; OSCC: oral squamous cell carcinoma; PDAC: pancreatic ductal adenocarcinoma; PI3K: phosphoinositide 3-kinase; RASA1: RAS p21 protein activator 1; RhoA: ras homolog gene family, member A; ST7L: suppression of tumorigenicity 7 like; TGF- β : transforming growth factor- β ; TNF: tumor necrosis factor; VEGF: vascular endothelial growth factor; VSCC: vulvar squamous cell carcinoma.

Despite an effort to make a comprehensive analysis, limitations remained inevitably in this study. First, most of the studies were among Asian populations though we do not impose restrictions on region during literature retrieval, which may introduce possible geographical bias. However, since these results have been further confirmed by TCGA dataset (20), it could be more credible to generalize the data to other populations. Second, some HRs with corresponding 95% CIs were indirectly extracted from the Kaplan–Meier (K-M) survival curves, which may introduce possible bias. Third, the cut-off value of MIR31HG levels varied among the selected studies, which may contribute to potential limitation to the clinical use.

Conclusion

Taken together, our study provided evidence that MIR31HG overexpression suggested remarkable association with poor prognosis in terms of OS, tumor stage, and LNM in lung cancer, but favorable prognosis in gastrointestinal cancer. Therefore, MIR31HG may serve as a novel promising prognostic biomarker in patients with multiple cancers. However, more rigorously designed prospective studies with large sample size and complete clinical information are still requested for further confirmation of the results.

Abbreviations

Akt: protein kinase B; BCAR4: breast cancer anti-estrogen resistance 4; CCND1: cyclin D1; ceRNA: competing endogenous RNA; CI: confidence interval; CRC: colorectal cancer; DFS: disease-free survival; DM: distant metastasis; E2F1: E2F transcription factor 1; EGFR: epidermal growth factor receptor; EMT: epithelial-mesenchymal transition; ERK: extracellular regulated protein kinases; ESCC: esophageal squamous cell carcinoma; ETS1: ETS proto-oncogene 1, transcription factor; EZH2: enhancer of zeste 2 polycomb repressive complex 2 subunit; FIGO: FOXM1: forkhead box M1; GEPIA: Gene Expression Profiling Interactive Analysis; GSK-3 β : glycogen synthase kinase-3 β ; HCC: hepatocellular carcinoma; HIF-1 α : hypoxia inducible factor 1 subunit alpha; HNSCC: head and neck squamous cell carcinoma; HR: hazard ratio; IFN- α / γ : Interferon- α / γ ; K-M: Kaplan-Meier; lncRNA: long noncoding RNA; LncHIFCAR: long noncoding HIF-1 α co-activating RNA; LNM: lymph node metastasis; LSCC: laryngeal squamous cell cancer; LUAD; lung adenocarcinoma; MIR31HG: microRNA-31 host gene; MMP1: matrix metalloproteinase 1; NOS: Newcastle-Ottawa Scale; NSCLC: non-small cell lung cancer; NT5E: 5'-nucleotidase ecto; OS: overall survival; OSCC: oral squamous cell carcinoma; PDAC: pancreatic ductal adenocarcinoma; PFS: progression-free survival; PI3K: phosphoinositide 3-kinase; qRT-PCR: real-time quantitative reverse transcription polymerase chain reaction; RFS: relapse-free survival; RhoA: ras homolog gene family, member A; SNHG16: small nucleolar RNA host gene 16; ST7L: suppression of tumorigenicity 7 like;

TCGA: The Cancer Genome Atlas; TGF- β : transforming growth factor- β ; TTN-AS1: TTN antisense RNA 1; TNF: tumor necrosis factor; VEGF: vascular endothelial growth factor; VSCC: vulvar squamous cell carcinoma.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests in this work.

Funding

This work was supported by the National Natural Science Foundation of China (No. 81902745), Natural Science Foundation of Hunan Province, China (2018JJ3716, 2018JJ3762), China Scholarship Council (201806375067, 201806375068), and the Fundamental Research Funds for the Central Universities of Central South University (2017zzts231).

Authors' contributions

CT and ZHL conceptualized the study and designed the methodology of the study. CT and XLR performed the literature retrieval, data analysis and interpretation. JYH and SQL participated in the quality assessment, and assisted in the statistical analysis. LQ and ZXD participated in the data analysis, and assisted in creating the figures. WCW participated in the data interpretation and assisted in manuscript drafting. ZHL and WCW critically reviewed and revised the manuscript. All authors read and approved the final manuscript as submitted.

Acknowledgements

We would like to thank the researchers and study participants for their contributions.

References

1. Wu Y, Hou Y, Xu P, Deng Y, Liu K, Wang M, et al. The prognostic value of YAP1 on clinical outcomes in human cancers. *Aging*. 2019;11(19):8681-700.
2. Zheng S, Zhang X, Wang X, Li J. MIR31HG promotes cell proliferation and invasion by activating the Wnt/beta-catenin signaling pathway in non-small cell lung cancer. *Oncol Lett*. 2019;17(1):221-9.
3. Huo X, Han S, Wu G, Latchoumanin O, Zhou G, Hebbard L, et al. Dysregulated long noncoding RNAs (lncRNAs) in hepatocellular carcinoma: implications for tumorigenesis, disease progression, and liver cancer stem cells. *Mol Cancer*. 2017;16(1):165.
4. Shih JW, Chiang WF, Wu ATH, Wu MH, Wang LY, Yu YL, et al. Long noncoding RNA lncHIFCAR/MIR31HG is a HIF-1 α co-activator driving oral cancer progression. *Nat Commun*. 2017;8:15874.

5. Yan S, Tang Z, Chen K, Liu Y, Yu G, Chen Q, et al. Long noncoding RNA MIR31HG inhibits hepatocellular carcinoma proliferation and metastasis by sponging microRNA-575 to modulate ST7L expression. *J Exp Clin Cancer Res*. 2018;37(1):214.
6. Yang H, Liu P, Zhang J, Peng X, Lu Z, Yu S, et al. Long noncoding RNA MIR31HG exhibits oncogenic property in pancreatic ductal adenocarcinoma and is negatively regulated by miR-193b. *Oncogene*. 2016;35(28):3647-57.
7. Atianand MK, Caffrey DR, Fitzgerald KA. Immunobiology of Long Noncoding RNAs. *Annu Rev Immunol*. 2017;35:177-98.
8. Schmitt AM, Chang HY. Long Noncoding RNAs in Cancer Pathways. *Cancer Cell*. 2016;29(4):452-63.
9. Tu C, Ren X, He J, Zhang C, Chen R, Wang W, et al. The Value of LncRNA BCAR4 as a Prognostic Biomarker on Clinical Outcomes in Human Cancers. *J Cancer*. 2019;10(24):5992-6002.
10. Zhang C, Ren X, He J, Wang W, Tu C, Li Z. The prognostic value of long noncoding RNA SNHG16 on clinical outcomes in human cancers: a systematic review and meta-analysis. *Cancer Cell Int*. 2019;19:261.
11. Fu D, Lu C, Qu X, Li P, Chen K, Shan L, et al. LncRNA TTN-AS1 regulates osteosarcoma cell apoptosis and drug resistance via the miR-134-5p/MBTD1 axis. *Aging*. 2019;11(19):8374-85.
12. Shi Y, Lu J, Zhou J, Tan X, He Y, Ding J, et al. Long non-coding RNA Loc554202 regulates proliferation and migration in breast cancer cells. *Biochem Biophys Res Commun*. 2014;446(2):448-53.
13. Montes M, Nielsen MM, Maglieri G, Jacobsen A, Hojfeldt J, Agrawal-Singh S, et al. The lncRNA MIR31HG regulates p16(INK4A) expression to modulate senescence. *Nat Commun*. 2015;6:6967.
14. Sun Y, Jia X, Wang M, Deng Y. Long noncoding RNA MIR31HG abrogates the availability of tumor suppressor microRNA-361 for the growth of osteosarcoma. *Cancer Manag Res*. 2019;11:8055-64.
15. Ma X, Qi S, Duan Z, Liao H, Yang B, Wang W, et al. Long non-coding RNA LOC554202 modulates chordoma cell proliferation and invasion by recruiting EZH2 and regulating miR-31 expression. *Cell Prolif*. 2017;50(6).
16. Eide PW, Eilertsen IA, Sveen A, Lothe RA. Long noncoding RNA MIR31HG is a bona fide prognostic marker with colorectal cancer cell-intrinsic properties. *Int J Cancer*. 2019;144(11):2843-53.
17. Zhang H, Wang Z, Wu J, Ma R, Feng J. Long noncoding RNAs predict the survival of patients with colorectal cancer as revealed by constructing an endogenous RNA network using bioinformation analysis. *Cancer Med*. 2019;8(3):863-73.
18. Ding J, Lu B, Wang J, Wang J, Shi Y, Lian Y, et al. Long non-coding RNA Loc554202 induces apoptosis in colorectal cancer cells via the caspase cleavage cascades. *J Exp Clin Cancer Res*. 2015;34:100.
19. Wang B, Jiang H, Wang L, Chen X, Wu K, Zhang S, et al. Increased MIR31HG lncRNA expression increases gefitinib resistance in non-small cell lung cancer cell lines through the EGFR/PI3K/AKT signaling pathway. *Oncol Lett*. 2017;13(5):3494-500.
20. Liu B, Chen Y, Yang J. LncRNAs are altered in lung squamous cell carcinoma and lung adenocarcinoma. *Oncotarget*. 2017;8(15):24275-91.
21. Ning P, Wu Z, Hu A, Li X, He J, Gong X, et al. Integrated genomic analyses of lung squamous cell carcinoma for identification of a possible competitive endogenous RNA network by means of TCGA datasets. *Peer J*. 2018;6:e4254.
22. Qin J, Ning H, Zhou Y, Hu Y, Yang L, Huang R. LncRNA MIR31HG overexpression serves as poor prognostic biomarker and promotes cells proliferation in lung adenocarcinoma. *Biomed Pharmacother*. 2018;99:363-8.
23. Sui J, Yang S, Liu T, Wu W, Xu S, Yin L, et al. Molecular characterization of lung adenocarcinoma: A potential four-long noncoding RNA prognostic signature. *J Cell Biochem*. 2019;120(1):705-14.
24. Nie FQ, Ma S, Xie M, Liu YW, De W, Liu XH. Decreased long noncoding RNA MIR31HG is correlated with poor prognosis and contributes to cell proliferation in gastric cancer. *Tumour Biol*. 2016;37(6):7693-701.
25. He A, Chen Z, Mei H, Liu Y. Decreased expression of LncRNA MIR31HG in human bladder cancer. *Cancer Biomark*. 2016;17(2):231-6.
26. Augoff K, McCue B, Plow EF, Sossey-Alaoui K. miR-31 and its host gene lncRNA LOC554202 are regulated by promoter hypermethylation in triple-negative breast cancer. *Mol Cancer*. 2012;11:5.

27. Chen J, Zhu J. Elevated Expression Levels of Long Non-Coding RNA, Loc554202, Are Predictive of Poor Prognosis in Cervical Cancer. *Tohoku J Exp Med.* 2017;243(3):165-72.
28. Ni S, Zhao X, Ouyang L. Long non-coding RNA expression profile in vulvar squamous cell carcinoma and its clinical significance. *Oncol Rep.* 2016;36(5):2571-8.
29. Sun K, Zhao X, Wan J, Yang L, Chu J, Dong S, et al. The diagnostic value of long non-coding RNA MIR31HG and its role in esophageal squamous cell carcinoma. *Life Sci.* 2018;202:124-30.
30. Ren ZP, Chu XY, Xue ZQ, Zhang LB, Wen JX, Deng JQ, et al. Down-regulation of lncRNA MIR31HG correlated with aggressive clinicopathological features and unfavorable prognosis in esophageal squamous cell carcinoma. *Eur Rev Med Pharmacol Sci.* 2017;21(17):3866-70.
31. Li Y, Xin S, Wu H, Xing C, Duan L, Sun W, et al. High expression of microRNA31 and its host gene LOC554202 predict favorable outcomes in patients with colorectal cancer treated with oxaliplatin. *Oncol Rep.* 2018;40(3):1706-24.
32. Yang L WH, Xiao HJ. Long non-coding RNA Loc554202 expression as a prognostic factor in patients with colorectal cancer. *Eur Rev Med Pharmacol Sci.* 2016;20(20):4243-7.
33. Lin Y, Zhang CS, Li SJ, Li Z, Sun FB. LncRNA LOC554202 promotes proliferation and migration of gastric cancer cells through regulating p21 and E-cadherin. *Eur Rev Med Pharmacol Sci.* 2018;22(24):8690-7.
34. Tang Z, Kang B, Li C, Chen T, Zhang Z. GEPIA2: an enhanced web server for large-scale expression profiling and interactive analysis. *Nucleic Acids Res.* 2019;47(W1):W556-W60.
35. Wang R, Ma Z, Feng L, Yang Y, Tan C, Shi Q, et al. LncRNA MIR31HG targets HIF1A and P21 to facilitate head and neck cancer cell proliferation and tumorigenesis by promoting cell-cycle progression. *Mol Cancer.* 2018;17(1):162.
36. Huarte M. The emerging role of lncRNAs in cancer. *Nat Med.* 2015;21(11):1253-61.
37. He J, Tu C, Liu Y. Role of lncRNAs in aging and age-related diseases. *Aging Med (Milton).* 2018;1(2):158-75.
38. Zou Y, Xu S, Xiao Y, Qiu Q, Shi M, Wang J, et al. Long noncoding RNA LERFS negatively regulates rheumatoid synovial aggression and proliferation. *J Clin Invest.* 2018;128(10):4510-24.
39. Rajbhandari R, McFarland BC, Patel A, Gerigk M, Gray GK, Fehling SC, et al. Loss of tumor suppressive microRNA-31 enhances TRADD/NF-kappaB signaling in glioblastoma. *Oncotarget.* 2015;6(19):17805-16.
40. Dandan W, Jianliang C, Haiyan H, Hang M, Xuedong L. Long noncoding RNA MIR31HG is activated by SP1 and promotes cell migration and invasion by sponging miR-214 in NSCLC. *Gene.* 2019;692:223-30.
41. Feng L, Wang R, Lian M, Ma H, He N, Liu H, et al. Integrated Analysis of Long Noncoding RNA and mRNA Expression Profile in Advanced Laryngeal Squamous Cell Carcinoma. *PLoS one.* 2016;11(12):e0169232.
42. Yang S, Wang J, Ge W, Jiang Y. Long non-coding RNA LOC554202 promotes laryngeal squamous cell carcinoma progression through regulating miR-31. *J Cell Biochem.* 2018;119(8):6953-60.
43. He J, Jin S, Zhang W, Wu D, Li J, Xu J, et al. Long non-coding RNA LOC554202 promotes acquired gefitinib resistance in non-small cell lung cancer through upregulating miR-31 expression. *J Cancer.* 2019;10(24):6003-13.
44. Bar C, Chatterjee S, Thum T. Long Noncoding RNAs in Cardiovascular Pathology, Diagnosis, and Therapy. *Circulation.* 2016;134(19):1484-99.
45. Jin C, Jia L, Huang Y, Zheng Y, Du N, Liu Y, et al. Inhibition of lncRNA MIR31HG Promotes Osteogenic Differentiation of Human Adipose-Derived Stem Cells. *Stem Cells.* 2016;34(11):2707-20.

Supplementary File Legend

Fig. S1. Forest plots describing subgroup analysis of HRs for association between MIR31HG expression and OS based on follow-up months. Abbreviations: CI: confidence interval; HR, hazard ratio; OS, overall survival.

Figures




Figure 1

Figure 1

The PRISMA flow diagram of study selection.




Figure 2

Figure 2

Forest plots describing the overall and subgroup analysis of HRs for association between MIR31HG expression and OS (A), DFS (B), and RFS (C). Abbreviations: CI: confidence interval; DFS: disease-free survival; HR, hazard ratio; OS, overall survival; RFS: relapse-free survival.




Figure 3

Figure 3

Forest plot describing overall and stratified analysis of relationship between MIR31HG expression and clinicopathological parameters, including advanced clinical stage (A), LNM (B), DM (C), and tumor size (D). Abbreviations: DM: distant metastasis; LNM: lymph node metastasis.




Figure 4

Figure 4

Overall and subgroup analysis of association between MIR31HG expression and advanced clinical stage (A), LNM (B) after deletion according to the sensitive analysis. Abbreviations: LNM: lymph node metastasis.




Figure 5

Figure 5

Begg's funnel plot for included studies reporting OS (A) and sensitivity analysis of each study reporting clinical stages (B) and LNM (C). Abbreviations: LNM: lymph node metastasis; OS, overall survival.




Figure 6

Figure 6

The prognostic value of MIR31HG in lung cancer patients with tumors in TCGA dataset, including OS (A), DFS (B). Abbreviation: DFS: disease-free survival; HR, hazard ratio; OS, overall survival; TCGA: The Cancer Genome Atlas.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS1.pdf](#)