Association Between Ambient Fine Particulate Matter and Maternal Vitamin D Levels During The Second Trimester of Pregnancy

Chong Li

Affiliated Kunshan Hospital of Jiangsu University

Ya-qin Gong

Affiliated Kunshan Hospital of Jiangsu University

Yun-yu Xia

Meteorological Bureau of Kunshan City

Xiao-chun Wang

Meteorological Bureau of Kunshan City

Lin Chen

Ecology and Environment Bureau of Kunshan City

Shan-jun Yan

Ecology and Environment Bureau of Kunshan City

Ke Lu (✉ sgu8434@sina.com)

Affiliated Kunshan Hospital of Jiangsu University

Research Article

Keywords: Vitamin D, Air Pollution, Particulate Matter, Pregnancy, Meteorology

Posted Date: February 14th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1342874/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Evidence for an association between the amount of ambient fine particulate matter (PM) in the atmosphere and vitamin D status of pregnant women is limited. We aimed to examine the independent association between ambient fine PM and maternal levels of serum 25-hydroxyvitamin D (25OHD) during the second trimester and to explore possible modifications to the association by meteorological factors.

Methods: 27,768 pregnant women presenting for prenatal examination who were tested for serum 25OHD concentration during the second trimester between January 1, 2016, and 31 December, 2020, were included in this retrospective analysis. Exposure to PM was evaluated based on daily average PM with an aerodynamic diameter of $\leq 2.5 \mu\text{m}$ (PM_{2.5}) and PM with an aerodynamic diameter of $\leq 10 \mu\text{m}$ (PM₁₀). Corresponding meteorological data for daily average atmospheric temperature, atmospheric pressure, relative humidity, sunshine duration, and wind speed, were collected.

Results: Vitamin D deficiency ($<12 \text{ ng/mL}$), inadequacy ($12\text{--}20 \text{ ng/mL}$), and adequacy ($\geq 20 \text{ ng/mL}$) were respectively present in 23.5%, 41.3%, and 35.2% of pregnant women during their second trimester. The maximum cumulative effects of PM_{2.5} occurred at lag 45 days, and the maximum cumulative effects of PM₁₀ occurred at lag 60 days. In crude models, 45-day moving daily average PM_{2.5} concentrations were negatively associated with 25OHD levels (β , -0.200 ; 95% CI, -0.206 to -0.193), as were 60-day moving daily average PM₁₀ concentrations (β , -0.142 ; 95% CI, -0.146 to -0.137). After adjusting for temporal and meteorological factors, the effect values were drastically reduced (adjusted β of PM_{2.5}, -0.032 ; 95% CI, -0.046 to -0.018 ; adjusted β of PM₁₀, -0.039 ; 95% CI, -0.049 to -0.028).

Conclusions: Our study showed an independent negative association between ambient fine PM in the atmosphere and maternal serum 25OHD levels during the second trimester of pregnancy. However, the effect values were small after adjusting for temporal and/or meteorological factors, which indicates that ambient fine PM may have a limited influence on maternal serum 25OHD levels.

1. Introduction

The key fat-soluble nutrient vitamin D has multiple functions, and its deficiency is thought to be a risk factor in skeletal health and various non-skeletal conditions, such as schizophrenia, skin disorders, certain types of cancer, type 2 diabetes, and infections [1]. For pregnant women, sufficient vitamin D stores need to be maintained to provide for both themselves and their fetus [2]. Research continues to provide evidence of an association between maternal Vitamin D deficiency (VDD) and a higher risk of numerous undesirable pregnancy outcomes, such as a low body weight, respiratory tract infections, and neurocognitive developmental problems in newborns and preeclampsia and high blood pressure in mothers [3-8]. Therefore, VDD is a key clinical issue for pregnant women and deserves research attention. Throughout the world at present, VDD during pregnancy is still a frequent occurrence. In one study of pregnant women in Finland, 77.4% were found to suffer from VDD during their first trimester [9]. While in,

the USA, Ginde et al. [10] reported 46%, 32%, and 18% of women with VDD in the first, second, and third trimesters, respectively, and there were comparable observations among British [11], Indian [12], and Chinese [13] women.

For the most part, factors determining the vitamin D status during pregnancy do not differ from those important for the general population, such as the number of melanocytes in the skin, the amount of exposure of the skin to the sun, the amount of adipose tissue in the body, geographical latitude, dietary factors, and intake of supplementary vitamins [14-18]. An indoor lifestyle can also increase the risk of developing a VDD [19]. Moreover, the amount of PM in the atmosphere has been associated with circulating levels of 25-hydroxyvitamin D (25OHD), which is a key marker for vitamin D status, in humans [20-23], including pregnant women [24].

It is not yet clear what mechanisms coordinate the links between maternal vitamin D and air pollution levels. Because vitamin D is a relatively scarce resource in food, humans generally rely on the vitamin D produced by cells in the skin on exposure to natural solar ultraviolet (UV) B radiation [25, 26]. Some scholars have hypothesized that the reduction in the amount of solar UVB radiation reaching ground level by ambient fine particulate matter (PM) is the main mechanism linking PM to circulating 25OHD levels in humans [23, 24]. In an investigation into the relationship between certain pollutants and ground-level UVB intensity, PM with an aerodynamic diameter of $\leq 10 \mu\text{m}$ (PM_{10}) was shown to be a significant negative predictor of solar UVB radiation, but the impact of PM_{10} was very small [27]. In addition, vitamin D status correlated with temporal and meteorological factors. During their second trimester, the 25OHD levels of women in eastern China demonstrated variation with season and air temperature [28]. Another study reported 25OHD levels were positively correlated with the prior month's temperature [29]. Thus, when exploring the independent association between ambient fine PM and vitamin D status, meteorological factors may need to be taken into account.

Thus, our study objective was to examine the independent association between ambient fine PM and maternal serum 25OHD levels during the second trimester in pregnant women and to explore if meteorological factors have an impact on this association.

2. Methods

2.1 Study Population

We designed a single-hospital-based cross-sectional retrospective observational study to test our hypotheses. All prenatal examination data were collected from the hospital medical database in Affiliated Kunshan Hospital of Jiangsu University located in Kunshan, eastern China, at 31.2°N latitude and approximately 30 kilometers from Shanghai. This is the largest hospital that provides prenatal care in all the districts of Kunshan. Consecutively, 35,476 pregnant women at the 15-20th week gestation (during the second trimester) who visited our institution for prenatal examination and were tested for serum 25OHD concentrations between January 1, 2016, and 31 December, 2020, were included in this study. We

excluded 7,708 women for the following reasons: those (1) with a chronic metabolic disease with consequences for the metabolism of vitamin D ($n = 5,389$); (2) with a high-risk pregnancy ($n = 1,087$), or (3) who had lived in Kunshan for less than 1 year ($n = 1,232$). Finally, 27,768 pregnant women were included in this analysis (Additional file 1: Figure S1).

The study protocol, submitted for review by the ethical committee at the Affiliated Kunshan Hospital of Jiangsu University (approval No. 2020-03-046-K01), was approved, and it complied with the Declaration of Helsinki. Patient information was initially documented for hospital's quality improvement purposes. Data analyzers were blinded to the identity of patients. Written informed consent was not required due to the anonymous data gathering and observational nature of the study.

2.2 Measurement of Vitamin D

In humans, the most abundant form of vitamin D in the blood is 25OHD, and serum levels are used to reliably estimate a patient's vitamin D status. All pregnant women in this study were in their 15-20th week of gestation (during the second trimester of pregnancy) and were asked to fast before blood samples were taken. Serum 25OHD concentrations were measured immediately using an automated electrochemiluminescence immunoassay on a Roche Cobas 8000/e602 analyzer (Roche Diagnostics, Mannheim, Germany). There is no medical consensus regarding the status cut-off values of 25OHD concentrations. However, the Institute of Medicine (National Academy of Sciences, Washington, DC, USA) and the National Osteoporosis Society (Bath, England) concur that, for bone health, a serum 25OHD level ($2.5 \text{ nmol/L } 25\text{OHD} = 1 \text{ ng/mL } 25\text{OHD}$) of $< 30 \text{ nmol/L}$ (12 ng/mL) is deficient, $30\text{--}50 \text{ nmol/L}$ ($12\text{--}20 \text{ ng/mL}$) is insufficient for certain individuals, while $>50 \text{ nmol/L}$ (20 ng/mL) is sufficient for most people [30, 31]. The time of blood collection was included in the analysis. Seasons were defined as: Spring, March–May; Summer, June–August; Autumn, September–November; Winter, December–February.

2.3 PM Exposure and Meteorological Data Assessment

We received data for the daily average concentration of PM with an aerodynamic diameter of $\leq 2.5 \mu\text{m}$ ($\text{PM}_{2.5}$; $\mu\text{g}/\text{m}^3$) and PM_{10} ($\mu\text{g}/\text{m}^3$) from the Meteorological Bureau of Kunshan City. Meteorological data (daily average atmospheric temperature [$^{\circ}\text{C}$], atmospheric pressure [hPa], relative humidity [%], sunshine duration [hour], and wind speed [m/s]) were obtained from the Ecology and Environment Bureau of Kunshan City. The distances between the environmental and meteorological monitoring sites and the hospital are approximately 8 and 3 kilometers, respectively. Kunshan City covers an area of 927.7 square kilometers; this distance is shorter than that used for sensitivity analysis in the study by Di et al. in 2017 [32].

Because the serum 25OHD half-life is about 15 days [33], the cumulative effects of $\text{PM}_{2.5}$ and PM_{10} on maternal serum 25OHD levels during the second trimester of pregnancy were approximated using the moving-average lag structures (blood collection day [day 0] was not included) as follows: 0-3 days (3-day moving daily average PM concentration), 0-7 days (7-day moving daily average PM concentration), 0-15

days (15-day moving daily average PM concentration), 0-30 days (30-day moving daily average PM concentration), 0-45 days (45-day moving daily average PM concentration), 0-60 days (60-day moving daily average PM concentration), 0-75 days (75-day moving daily average PM concentration), and 0-90 days (90-day moving daily average PM concentration). When adjusting for the influence of meteorological factors, the corresponding moving average lag structures for the meteorological factors were calculated.

2.4 Statistics

The summary statistics for the characteristics of pregnant women were expressed as frequencies (proportions) for categorical variables and as the means (standard deviation [SD]) or medians (Q1-Q3) for continuous variables. We also conducted univariate logistic regression analysis to evaluate the association between the characteristics of pregnant women and maternal serum 25OHD concentrations during the second trimester. The daily average meteorological variables and daily average PM_{2.5}, PM₁₀, and 25OHD concentrations were screened for correlations by Pearson's.

Using generalized estimating equations (GEE), we evaluated the relationships between the cumulative effects of PM on maternal serum 25OHD levels. The cumulative effects of PM were divided into four levels on the basis of quartiles. After adjusting for year and age at blood collection (adjust I) or year, season, and age at blood collection (adjust II), β -values (95% confidence interval [CI]) of the maternal serum 25OHD levels were calculated based on group trends. As part of the sensitivity analysis, we also calculated the β -values (95% CI) for the 25OHD levels when using the concentrations of PM_{2.5} and PM₁₀ as continuous variables.

Controlling for the influence of meteorological covariances by applying multivariate linear regression analysis allowed us to evaluate independent associations between the maximum cumulative effects of PM and maternal serum 25OHD. We calculated the results of the unadjusted or minimally adjusted analysis and those from fully adjusted analysis. First, collinearity diagnosis of covariances was performed using variance inflation factor (VIF) analysis (the variable average atmospheric temperature was first removed due to VIF >10). Then, a judgement on whether to adjust covariances was made using the following principles: Criteria 1, the covariate is added to the basic model or removed from the full model and the matched odds ratio (OR) is changed by at least 10%; Criteria 2: Criteria 1 or a covariate P -value of <0.1 in the univariate model [34]. As part of the sensitivity analysis, we transformed 25OHD quantitative variables into dichotomous qualitative variables (1 = vitamin D deficiency and inadequacy [<20 ng/mL]; 0 = adequacy [≥ 20 ng/mL]), then the OR and 95% CI for maternal vitamin D deficiency and inadequacy (<20 ng/mL) associated with a 10 mg/m^3 increase in PM_{2.5} or PM₁₀ was determined.

Non-linear relationships were additionally identified via a generalized additive model (GAM), and on finding a non-linear correlation, the threshold effect in terms of the smoothing curve was calculated using a two-piecewise linear regression model. When a clear ratio was apparent in the smoothing curve, the recursive method was applied to automatically calculate the turning point at which to use the maximum

likelihood model [35]. In addition, to test the robustness and potential variation in the different subgroups, we repeated the subgroup analyses while stratifying by season, age, and meteorological factors. The age threshold was derived from the turning point calculated by the GAM followed by an inspection of the modification and interaction of the subgroups with the likelihood ratio test.

All statistical analyses were performed using the Empower Stats (www.empowerstats.com, X&Y solutions, Inc., Boston, MA, USA) and R software version 3.6.3 (<http://www.r-project.org>). A P -value <0.05 was set as the significance threshold.

3. Results

Table 1 displays the data on the pregnant women and their vitamin D statuses. This analysis included 27,768 individuals with a mean age of 28.86 (SD, 4.27) years. The mean (SD) and median (Q1-Q3) values of maternal serum 25OHD concentrations during the second trimester of pregnancy were 17.7 (7.9) ng/mL and 16.0 (12.0-22.0) ng/mL, respectively. Vitamin D deficiency, inadequacy, and adequacy were present in 23.5%, 41.3%, and 35.2% of women, respectively. Univariate logistic regression analysis revealed that maternal 25OHD concentration was positively associated with maternal age and showed seasonal variation, with the peak in September and the nadir in February.

Pearson's correlation analysis was conducted to compare serum 25OHD concentration, meteorological variables, and air pollutant exposure. Figure S2 (Additional file 1) presents that, except PM_{2.5} vs. sunshine duration (P -value = 0.821), all correlations among the variables were statistically significant (P -value < 0.001). There was a strong positive correlation between the daily average PM_{2.5} concentration and PM₁₀ concentration, and the Pearson coefficient was 0.924. There was also a strong negative correlation between daily average atmospheric temperature and atmospheric pressure, and the Pearson coefficient was -0.892. There was a moderate negative correlation between daily sunshine duration and relative humidity, and the correlation coefficient was -0.683.

Figure 1 shows periodic changes in the above indicators in terms of monthly average values over time. It can be seen that the variation in the average temperature of the last month was most consistent with the periodic variation of monthly average serum 25OHD concentrations. Similar periodic changes were seen for sunshine duration and relative humidity. However, monthly average PM_{2.5} and PM₁₀ concentrations and atmospheric pressure showed periodic changes that were diametrically opposite to those of the monthly average serum 25OHD concentration.

Figure 2 and Table S1 (Additional file 1) show the cumulative effects of PM_{2.5} and PM₁₀ on maternal serum 25OHD levels during the second trimester of pregnancy. After we adjusted for year, season, and age at blood collection, the maximum cumulative effects occurred at lag 0-45 days of PM_{2.5} and lag 0-60 days of PM₁₀. When PM concentrations were used as continuous variables, similar results were observed, and these are provided in Table S2 (Additional file 1).

The independent associations between the maximum cumulative effects of PM and serum 25OHD levels were further investigated, and Table 2 shows the results adjusted for different covariances. In the crude models, the effect sizes of PM_{2.5} (β , -0.200; 95% CI: -0.206 to -0.193; P -value <0.00001) and PM₁₀ (β , -0.142; 95% CI: -0.146 to -0.137; P -value <0.00001) were relatively high. After adjustments for year, season, and age at blood collection (Model 1), the effect sizes of PM_{2.5} (β , -0.113; 95% CI: -0.125 to -0.101; P -value <0.00001) and PM₁₀ (β , -0.090; 95% CI: -0.099 to -0.081; P -value <0.00001) were reduced. On the basis of the Model 1 adjustment, the effect values of PM_{2.5} (β , -0.042; 95% CI: -0.055 to -0.028; P -value <0.00001) and PM₁₀ (β , -0.039; 95% CI: -0.049 to -0.030; P -value <0.00001) were further reduced after adding atmospheric pressure adjustment (Model 2). After further adjustment for sunshine duration, the effect values of Model 3 were similar to those of Model 2. In the fully adjusted Model 4 (adjusted for year, age, season, daily average atmospheric pressure, sunshine duration, relative humidity, and wind speed), there was a negative relationship between the 45-day moving daily average PM_{2.5} concentration and the women's 25OHD levels (β , -0.032; 95% CI, -0.046 to -0.018; P -value <0.00001), and the 60-day moving daily average PM₁₀ concentration was negatively associated with 25OHD levels (β , -0.039; 95% CI, -0.049 to -0.028; P -value <0.00001). As part of the sensitivity analysis, the crude and adjusted OR for PM exposure's association with maternal vitamin D deficiency and inadequacy (<20 ng/mL) were determined, which are given in Table S3 (Additional file 1). We observed greater odds of maternal vitamin D deficiency and inadequacy (<20 ng/mL) with higher PM levels. In the fully adjusted Model 4, an increase in the PM_{2.5} 45-day moving daily average by 10 mg/m³ was associated with a 12.4% (OR, 1.124; 95% CI, 1.071 to 1.180, P -value <0.00001) increase in the odds for maternal vitamin D deficiency and inadequacy (<20 ng/mL). In the fully adjusted Model 4, a 10 mg/m³ increase in the PM₁₀ 60-day moving daily average was associated with a 11.3% (OR, 1.113; 95% CI, 1.074 to 1.153, P -value <0.00001) increase in the odds of maternal vitamin D deficiency and inadequacy (<20 ng/mL).

In the subgroup analyses stratified by season, we further investigated the role of season on the association between the maximum cumulative effects of PM and serum 25OHD levels. For PM_{2.5} (Table 3), both linear and nonlinear effect values were higher in summer and autumn and lower in winter and spring (P -values for interaction <0.001). Figure 3A shows the different nonlinear associations between the 45-day moving daily average PM_{2.5} concentration and maternal serum 25OHD levels stratified by season. In addition, we calculated, using the two-piecewise linear regression model, the turning point of the adjusted smoothed curve. Specifically, the difference between the two slopes was at its maximum in autumn, and the turning point (45-day moving daily average PM_{2.5} concentration) was 31.11 $\mu\text{g}/\text{m}^3$. For PM₁₀ (Table 3), linear regression analysis showed a significant interaction for season (interaction P -value <0.001), but the non-linear model showed this interaction was not significant (interaction P -value = 0.383). Figure 3B shows the different nonlinear associations between the 60-day moving daily average PM₁₀ concentration and maternal serum 25OHD levels stratified by season. It is worth mentioning that, in autumn, the relationship between PM₁₀ and 25OHD levels was nonlinear, and the turning point was 38.10 $\mu\text{g}/\text{m}^3$. Specifically, when the 60-day moving daily average PM₁₀ concentration ranged from 32.32 $\mu\text{g}/\text{m}^3$

to $38.10 \mu\text{g}/\text{m}^3$, a stronger negative relationship was found between PM_{10} and serum 25OHD levels (β , -1.121 ; 95% CI, -1.513 to -0.728 ; P -value <0.0001 ; number of pregnant women, 878).

The threshold effect analysis used to examine the associations between maternal age and serum 25OHD levels during the second trimester of pregnancy revealed a stronger positive relationship between age and serum 25OHD levels when the women were 15 to 25 years of age and a weaker positive relationship when they were 25 to 47. The results of four different adjusted models were robust (Additional file 1: Figure S3 and Table S4). We then categorized the pregnant women using a threshold of 25 years according to the results of the threshold effect analysis and further investigated the modification effect of age on the association between the maximum cumulative effects of PM and serum 25OHD levels. Table S5 (Additional file 1) shows that the interaction between age and $\text{PM}_{2.5}$ was not significant (linear interaction P -value = 0.083 and non-linear interaction P -value = 0.058), while Table S6 (Additional file 1) shows the interaction between age and PM_{10} had a marginally significant effect on serum 25OHD levels (linear P -value for interaction = 0.045 and non-linear P -value for interaction = 0.038).

In the subgroup analyses stratified by meteorological factors, Table 4 showed the associations between PM and serum 25OHD levels during the second trimester of pregnancy were generally modified by meteorological factors, although wind speed had no modification effect on $\text{PM}_{2.5}$. Of particular interest, there was a positive association between PM concentration and 25OHD levels under low relative humidity. The result of this stratification was the opposite to the final conclusion.

4. Discussion

Based on our knowledge of the literature, this epidemiological study is the first to focus on the independent relationship between PM and maternal serum 25OHD levels during the second trimester of pregnancy after adjusting for meteorological factors. We found vitamin D deficiency, inadequacy, and adequacy in 23.5%, 41.3%, and 35.2% of pregnant women during the second trimester of pregnancy, respectively. Prenatal exposure to higher levels of PM was associated with decreased maternal serum 25OHD concentrations. We also found that the maximum cumulative effects of $\text{PM}_{2.5}$ occurred at lag 45 days and the maximum cumulative effects of PM_{10} occurred at lag 60 days. However, the effect values were drastically reduced after adjusting for temporal and/or meteorological factors. The results indicated that ambient fine PM has a limited influence on maternal serum 25OHD levels.

Our conclusions are similar to those of Zhao et al., who reported an association between prenatal exposure to higher $\text{PM}_{2.5}$ and PM_{10} levels and a decrease in circulating 25OHD concentrations in women in later stages of pregnancy [24]. In contrast to the analysis they conducted, we first investigated the time window for the largest cumulative effect of PM at the individual level and then adjusted for temporal and meteorological factors. Although both effect sizes of $\text{PM}_{2.5}$ and PM_{10} in the fully adjusted models were small (β -value of $\text{PM}_{2.5}$ = -0.032 ; β -value of PM_{10} = -0.039 , respectively), they were highly significant (P -value < 0.00001). The β -values showed two large drops, one after adjusting for season and the other after

adjusting for atmospheric pressure. Due to the collinearity of atmospheric pressure and temperature, it can be inferred that season and atmospheric pressure/temperature may have been important confounders in the regression model. Thus, in some correlation analyses of PM and vitamin D levels, the independent role of PM may have been overestimated when there was no adjustment for meteorological factors.

Indeed, in areas with distinct seasons, 25OHD concentrations in the population fluctuate over time [29, 36, 37]. Many studies showed season to be the primary factor affecting serum 25OHD levels [15, 29, 38-40]. Periodic changes in the seasons affect periodic changes in meteorological factors, such as temperature, humidity, and sunshine duration. Hence, unlike other studies that only adjusted for season, our study precisely adjusted for meteorological factor exposure at the individual level.

On the other hand, meteorological factors are also related to periodic changes in PM concentrations. Weather and climate are the most influential forces affecting the chemistry and atmospheric residence time of PM [41]. In a study in China, the geographical and temporal variations in PM levels and coinciding meteorological conditions were analyzed for 366 cities, and peak PM concentrations occurred in winter in most regions, and there were negative correlations between PM levels and precipitation, relative humidity, air temperature, and wind speed but a positively correlation was found between PM levels and surface pressure [42]. Guan et al. [43] reported the spatiotemporal variability of PM in three cities in western China and the influence of meteorological factors on PM. Crawford et al. [44], who analyzed the impact of meteorology on particulate source types in Lucas Heights, Australia, reported that temperature significantly affected PM_{2.5} levels. An Indian study reported that PM displayed substantial seasonal variations and a strong negative association with temperature, with considerable dependency levels [41].

Over 90% of the vitamin D in the body is synthesized in human skin after the exposure of precursor 7-dehydrocholesterol to UVB (290-315 nm) radiation from the sun [45]. The strength of the UVB during sun exposure, and therefore the amount of vitamin D synthesized, is affected by solar zenith angle (SZA)—the angle between the local vertical and the sun's position. Hence, the most intense solar radiation occurs when the SZA is small, i.e., which at lower altitudes, is around 11:00-15:00h in the summer, at which time the synthesis of vitamin D by the skin is most active [46]. At latitudes >50°, human skin participates in very little vitamin D synthesis during winter and spring for all skin types and ethnicities [47]. In addition, the amount of vitamin D produced is dependent on local weather conditions, such as the percentage cloud cover, which filters UVB radiation, and has an impact in all seasons and hours of the day [37]. Thus, because PM can absorb and diffuse solar irradiation, some scholars have hypothesized that PM can indirectly reduce vitamin D formation by reducing UVB exposure [23, 24]. However, other scholars have argued that there is significant spatiotemporal variation in the morphology, chemical makeup, density of PM, and it is difficult to determine its effect on UV radiation [48]. More recently, PM₁₀ was found to be a significant negative predictor of solar UVB radiation, but the effect of PM₁₀ was miniscule [27]. This conclusion can be used to further deduce that PM₁₀ has an effect on vitamin D levels, but the effect is small, which is consistent with the results of our study. Combined with the above views, we propose a hypothesis (see Additional file 1: Figure S4) that PM and meteorological factors indirectly influence the

cyclical changes of vitamin D in pregnant women by impacting the level of personal UVB exposure. For example, in winter, when the temperature is lower and there are fewer hours of sunshine, the PM concentration is higher, but pregnant women spend less time outdoors and wear more clothes, which lead to lower solar UVB radiation exposure, reducing the synthesis of vitamin D.

In the subgroup analysis, with the exception of the low relative humidity, a negative effect between PM and 25OHD was evident in all subgroups considered and after careful adjustments. In the interaction analysis, season and most meteorological factors interacted with the association between PM and 25OHD. However, the mechanisms of their interactions were unclear and need to be further investigated in the future.

There were some limitations regarding our study. First, our results were obtained from a Chinese population of pregnant women during the second trimester and cannot be extrapolated to other populations. Second, the study was an analytical retrospective study and hence provides limited evidence that PM exposure and vitamin D outcomes were related, and the difference between cause and effect is uncertain. Third, demographic information was lacking (education level, vitamin D/calcium supplement use, outdoor activities, use of sun protection, and BMI, etc.) for the individuals whose test results were used. However, valuable insights can be gleaned from the study, as it involved the retrospective analysis of a large dataset from a prenatal examination population. We used precise adjustments at the individual level to reveal whether PM levels are independently associated with 25OHD concentration.

5. Conclusion

Our study revealed an independent negative association between ambient fine PM and maternal serum 25OHD levels during the second trimester of pregnancy. However, the effect values were drastically reduced after adjusting for temporal and/or meteorological factors, which indicates that ambient fine PM may have a limited influence on maternal serum 25OHD levels.

Declarations

Funding

This study was funded by National Natural Science Foundation of China (CN) [82172441], Clinical Medical Science and Technology Development Fund of Jiangsu University (CN) (JLY2021048) and Suzhou Key Clinical Diagnosis and Treatment Technology Project (CN) (LCZX202024).

Competing interests

The authors declare that they have no competing interests.

Ethics approval

This study complies with the Declaration of Helsinki and has been approved by the Ethics Committee of the First People's Hospital of Kunshan (no. 2020-03-046-K01).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

Ke Lu contributed to the study conception and design. Ya-qin Gong, Yun-yu Xia, Xiao-chun Wang, Lin Chen and Shan-jun Yan contributed to the acquisition of data. Ke Lu and Ya-qin Gong contributed to the analysis and interpretation of data. Chong Li drafted the manuscript, and Ke Lu revised it. All authors critically revised the manuscript, agree to be fully accountable for ensuring the integrity and accuracy of the work, and read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. Holick MF: **Vitamin D: evolutionary, physiological and health perspectives.** *Curr Drug Targets* 2011, **12**(1):4-18.
2. Hatun S, Ozkan B, Orbak Z, Doneray H, Cizmecioglu F, Toprak D, Calikoglu AS: **Vitamin D deficiency in early infancy.** *J Nutr* 2005, **135**(2):279-282.
3. Achkar M, Dodds L, Giguère Y, Forest JC, Armson BA, Woolcott C, Agellon S, Spencer A, Weiler HA: **Vitamin D status in early pregnancy and risk of preeclampsia.** *Am J Obstet Gynecol* 2015, **212**(4):511.e511-517.
4. Arnold DL, Enquobahrie DA, Qiu C, Huang J, Grote N, VanderStoep A, Williams MA: **Early pregnancy maternal vitamin D concentrations and risk of gestational diabetes mellitus.** *Paediatr Perinat Epidemiol* 2015, **29**(3):200-210.
5. Wang H, Xiao Y, Zhang L, Gao Q: **Maternal early pregnancy vitamin D status in relation to low birth weight and small-for-gestational-age offspring.** *J Steroid Biochem Mol Biol* 2018, **175**:146-150.
6. Murthi P, Davies-Tuck M, Lappas M, Singh H, Mockler J, Rahman R, Lim R, Leaw B, Doery J, Wallace EM *et al*: **Maternal 25-hydroxyvitamin D is inversely correlated with foetal serotonin.** *Clin Endocrinol (Oxf)* 2017, **86**(3):401-409.
7. Meems LM, Mahmud H, Buikema H, Tost J, Michel S, Takens J, Verkaik-Schakel RN, Vreeswijk-Baudoin I, Mateo-Leach IV, van der Harst P *et al*: **Parental vitamin D deficiency during pregnancy is associated with increased blood pressure in offspring via Panx1 hypermethylation.** *Am J Physiol Heart Circ Physiol* 2016, **311**(6):H1459-h1469.

8. Weinert LS, Silveiro SP: **Maternal-fetal impact of vitamin D deficiency: a critical review.** *Matern Child Health J* 2015, **19**(1):94-101.
9. Viljakainen HT, Saarnio E, Hytinantti T, Miettinen M, Surcel H, Mäkitie O, Andersson S, Laitinen K, Lamberg-Allardt C: **Maternal vitamin D status determines bone variables in the newborn.** *J Clin Endocrinol Metab* 2010, **95**(4):1749-1757.
10. Ginde AA, Sullivan AF, Mansbach JM, Camargo CA, Jr.: **Vitamin D insufficiency in pregnant and nonpregnant women of childbearing age in the United States.** *Am J Obstet Gynecol* 2010, **202**(5):436.e431-438.
11. McAree T, Jacobs B, Manickavasagar T, Sivalokanathan S, Brennan L, Bassett P, Rainbow S, Blair M: **Vitamin D deficiency in pregnancy - still a public health issue.** *Matern Child Nutr* 2013, **9**(1):23-30.
12. Vijayendra Chary A, Hemalatha R, Seshacharyulu M, Vasudeva Murali M, Jayaprakash D, Dinesh Kumar B: **Vitamin D deficiency in pregnant women impairs regulatory T cell function.** *J Steroid Biochem Mol Biol* 2015, **147**:48-55.
13. Song SJ, Zhou L, Si S, Liu J, Zhou J, Feng K, Wu J, Zhang W: **The high prevalence of vitamin D deficiency and its related maternal factors in pregnant women in Beijing.** *PloS one* 2013, **8**(12):e85081.
14. Bodnar LM, Catov JM, Roberts JM, Simhan HN: **Prepregnancy obesity predicts poor vitamin D status in mothers and their neonates.** *J Nutr* 2007, **137**(11):2437-2442.
15. Bodnar LM, Simhan HN, Powers RW, Frank MP, Cooperstein E, Roberts JM: **High prevalence of vitamin D insufficiency in black and white pregnant women residing in the northern United States and their neonates.** *J Nutr* 2007, **137**(2):447-452.
16. Brembeck P, Winkvist A, Olausson H: **Determinants of vitamin D status in pregnant fair-skinned women in Sweden.** *Br J Nutr* 2013, **110**(5):856-864.
17. Jensen CB, Petersen SB, Granström C, Maslova E, Mølgaard C, Olsen SF: **Sources and determinants of vitamin D intake in Danish pregnant women.** *Nutrients* 2012, **4**(4):259-272.
18. Richard A, Rohrmann S, Quack Lötscher KC: **Prevalence of Vitamin D Deficiency and Its Associations with Skin Color in Pregnant Women in the First Trimester in a Sample from Switzerland.** *Nutrients* 2017, **9**(3).
19. Sowah D, Fan X, Dennett L, Hagtvedt R, Straube S: **Vitamin D levels and deficiency with different occupations: a systematic review.** *BMC Public Health* 2017, **17**(1):519.
20. Agarwal KS, Mughal MZ, Upadhyay P, Berry JL, Mawer EB, Puliyel JM: **The impact of atmospheric pollution on vitamin D status of infants and toddlers in Delhi, India.** *Arch Dis Child* 2002, **87**(2):111-113.
21. Kelishadi R, Moeini R, Poursafa P, Farajian S, Yousefy H, Okhovat-Souraki AA: **Independent association between air pollutants and vitamin D deficiency in young children in Isfahan, Iran.** *Paediatr Int Child Health* 2014, **34**(1):50-55.
22. Feizabad E, Hossein-Nezhad A, Maghbooli Z, Ramezani M, Hashemian R, Moattari S: **Impact of air pollution on vitamin D deficiency and bone health in adolescents.** *Arch Osteoporos* 2017, **12**(1):34.

23. Yang C, Li D, Tian Y, Wang P: **Ambient Air Pollutions Are Associated with Vitamin D Status.** *Int J Environ Res Public Health* 2021, **18**(13).
24. Zhao Y, Wang L, Liu H, Cao Z, Su X, Cai J, Hua J: **Particulate Air Pollution Exposure and Plasma Vitamin D Levels in Pregnant Women: A Longitudinal Cohort Study.** *J Clin Endocrinol Metab* 2019, **104**(8):3320-3326.
25. Glerup H, Mikkelsen K, Poulsen L, Hass E, Overbeck S, Thomsen J, Charles P, Eriksen EF: **Commonly recommended daily intake of vitamin D is not sufficient if sunlight exposure is limited.** *J Intern Med* 2000, **247**(2):260-268.
26. Chen TC, Chimeh F, Lu Z, Mathieu J, Person KS, Zhang A, Kohn N, Martinello S, Berkowitz R, Holick MF: **Factors that influence the cutaneous synthesis and dietary sources of vitamin D.** *Arch Biochem Biophys* 2007, **460**(2):213-217.
27. Rahman A, Elmi A: **Air pollutants are negatively associated with vitamin D-synthesizing UVB radiation intensity on the ground.** *Sci Rep* 2021, **11**(1):21480.
28. Xiao JP, Zang J, Pei JJ, Xu F, Zhu Y, Liao XP: **Low maternal vitamin D status during the second trimester of pregnancy: a cross-sectional study in Wuxi, China.** *PloS one* 2015, **10**(2):e0117748.
29. Ding F, Nie X, Li X, He Y, Li G: **Data mining: Biological and temporal factors associated with blood parathyroid hormone, vitamin D, and calcium concentrations in the Southwestern Chinese population.** *Clin Biochem* 2021, **90**:50-57.
30. Aspray TJ, Bowring C, Fraser W, Gittoes N, Javaid MK, Macdonald H, Patel S, Selby P, Tanna N, Francis RM: **National Osteoporosis Society vitamin D guideline summary.** *Age Ageing* 2014, **43**(5):592-595.
31. Institute of Medicine Committee to Review Dietary Reference Intakes for Vitamin D, Calcium: **The National Academies Collection: Reports funded by National Institutes of Health.** In: *Dietary Reference Intakes for Calcium and Vitamin D.* Edited by Ross AC, Taylor CL, Yaktine AL, Del Valle HB. Washington (DC): National Academies Press (US) Copyright © 2011, National Academy of Sciences.; 2011.
32. Di Q, Wang Y, Zanobetti A, Wang Y, Koutrakis P, Choirat C, Dominici F, Schwartz JD: **Air Pollution and Mortality in the Medicare Population.** *N Engl J Med* 2017, **376**(26):2513-2522.
33. Jones KS, Assar S, Vanderschueren D, Bouillon R, Prentice A, Schoenmakers I: **Predictors of 25(OH)D half-life and plasma 25(OH)D concentration in The Gambia and the UK.** *Osteoporos Int* 2015, **26**(3):1137-1146.
34. Kernan WN, Viscoli CM, Brass LM, Broderick JP, Brott T, Feldmann E, Morgenstern LB, Wilterdink JL, Horwitz RI: **Phenylpropanolamine and the risk of hemorrhagic stroke.** *N Engl J Med* 2000, **343**(25):1826-1832.
35. Liu S, Wang X, Lu Y, Li T, Gong Z, Sheng T, Hu B, Peng Z, Sun X: **The effects of intraoperative cryoprecipitate transfusion on acute renal failure following orthotopic liver transplantation.** *Hepatol Int* 2013, **7**(3):901-909.

36. Dovnik A, Mujezinović F, Treiber M, Pečovnik Balon B, Gorenjak M, Maver U, Takač I: **Determinants of maternal vitamin D concentrations in Slovenia : A prospective observational study.** *Wien Klin Wochenschr* 2017, **129**(1-2):21-28.
37. Saraff V, Shaw N: **Sunshine and vitamin D.** *Arch Dis Child* 2016, **101**(2):190-192.
38. Yuan Y, Liu H, Ji C, Guo X, Hu L, Wen J, Cai M: **Association of Maternal Serum 25-hydroxyvitamin D Concentrations in Second Trimester with Delivery Mode in A Chinese Population.** *Int J Med Sci* 2017, **14**(10):1008-1014.
39. Krieger JP, Cabaset S, Canonica C, Christoffel L, Richard A, Schröder T, von Wattenwyl BL, Rohrmann S, Lötscher KQ: **Prevalence and determinants of vitamin D deficiency in the third trimester of pregnancy: a multicentre study in Switzerland.** *Br J Nutr* 2018, **119**(3):299-309.
40. Li H, Ma J, Huang R, Wen Y, Liu G, Xuan M, Yang L, Yang J, Song L: **Prevalence of vitamin D deficiency in the pregnant women: an observational study in Shanghai, China.** *Arch Public Health* 2020, **78**:31.
41. Murari V, Kumar M, Mhawish A, Barman SC, Banerjee T: **Airborne particulate in Varanasi over middle Indo-Gangetic Plain: variation in particulate types and meteorological influences.** *Environ Monit Assess* 2017, **189**(4):157.
42. Li X, Song H, Zhai S, Lu S, Kong Y, Xia H, Zhao H: **Particulate matter pollution in Chinese cities: Areal-temporal variations and their relationships with meteorological conditions (2015-2017).** *Environ Pollut* 2019, **246**:11-18.
43. Guan Q, Cai A, Wang F, Yang L, Xu C, Liu Z: **Spatio-temporal variability of particulate matter in the key part of Gansu Province, Western China.** *Environ Pollut* 2017, **230**:189-198.
44. Chambers, Scott, Williams, Alastair, Crawford, Jagoda, Stelcer, Eduard, Dyer, Leisa: **Impact of meteorology on fine aerosols at Lucas Heights, Australia.** *Atmospheric Environment* 2016.
45. Diehl JW, Chiu MW: **Effects of ambient sunlight and photoprotection on vitamin D status.** *Dermatol Ther* 2010, **23**(1):48-60.
46. Webb AR: **Who, what, where and when-influences on cutaneous vitamin D synthesis.** *Prog Biophys Mol Biol* 2006, **92**(1):17-25.
47. Webb AR, Kline L, Holick MF: **Influence of season and latitude on the cutaneous synthesis of vitamin D3: exposure to winter sunlight in Boston and Edmonton will not promote vitamin D3 synthesis in human skin.** *J Clin Endocrinol Metab* 1988, **67**(2):373-378.
48. Zegarska B, Pietkun K, Zegarski W, Bolibok P, Wiśniewski M, Roszek K, Czarnecka J, Nowacki M: **Air pollution, UV irradiation and skin carcinogenesis: what we know, where we stand and what is likely to happen in the future?** *Postepy Dermatol Alergol* 2017, **34**(1):6-14.

Tables

Table 1. Characteristics of study participants (N =27,768)

Characteristics	Statistics	25OHD, Mean (SD) , ng/mL	β (95%CI) ^a	<i>P</i> - value ^a
Maternal serum 25OHD concentrations during pregnancy continuous, Mean (SD) Median (Q1-Q3), ng/mL	17.7 (7.9) 16.0 (12.0-22.0)			
Maternal serum 25OHD concentrations during pregnancy categorical, N (%)				
Deficiency (< 12 ng/mL)	6532 (23.5%)			
Inadequacy (>= 12, < 20 ng/mL)	11465 (41.3%)			
Adequacy (>= 20 ng/mL)	9771 (35.2%)			
Age tertile, N (%)				
Tertile 1 (15 - 26 y)	8605 (30.99%)	17.18 (7.70)	Reference	-
Tertile 2 (27 - 29 y)	7948 (28.62%)	17.57 (7.85)	0.39 (0.15, 0.63)	0.0013
Tertile 3 (30 - 47 y)	11215 (40.39%)	18.08 (8.03)	0.90 (0.68, 1.12)	<0.0001
<i>P</i> for trend				<0.0001
Year of blood collection , N (%)				
2016	5943 (21.40%)	16.28 (7.45)	Reference	-
2017	7410 (26.69%)	18.10 (7.65)	1.82 (1.55, 2.08)	<0.0001
2018	5616 (20.22%)	17.68 (8.52)	1.40 (1.11, 1.68)	<0.0001
2019	5204 (18.74%)	18.38 (7.96)	2.10 (1.81, 2.39)	<0.0001
2020	3595	17.94	1.65	<0.0001

	(12.95%)	(7.65)	(1.33, 1.98)	
<i>P</i> for trend				< 0.0001
Month of blood collection, N (%)				
January	2161 (7.78%)	13.83 (7.02)	Reference	-
February	1549 (5.58%)	13.65 (6.69)	-0.18 (-0.66, 0.29)	0.4535
March	2285 (8.23%)	14.37 (6.91)	0.54 (0.11, 0.97)	0.0133
April	2676 (9.64%)	16.19 (7.24)	2.36 (1.95, 2.77)	<0.0001
May	2578 (9.28%)	15.76 (6.80)	1.93 (1.52, 2.35)	<0.0001
June	2505 (9.02%)	18.98 (7.23)	5.15 (4.73, 5.56)	<0.0001
July	2617 (9.42%)	20.05 (7.39)	6.22 (5.81, 6.64)	<0.0001
August	2381 (8.57%)	22.10 (7.97)	8.27 (7.85, 8.70)	<0.0001
September	2276 (8.20%)	22.45 (8.09)	8.62 (8.19, 9.05)	<0.0001
October	2183 (7.86%)	20.65 (7.78)	6.82 (6.38, 7.25)	<0.0001
November	2273 (8.19%)	17.53 (7.27)	3.70 (3.27, 4.13)	<0.0001
December	2284 (8.23%)	14.79 (6.74)	0.96 (0.53, 1.39)	<0.0001
<i>P</i> for trend				< 0.0001

Season of blood collection, N (%)				
Spring (March, April and May)	7539 (27.15%)	15.49 (7.03)	Reference	-
Summer (June, July and August)	7503 (27.02%)	20.34 (7.64)	4.85 (4.61, 5.09)	<0.0001
Autumn (September, October and November)	6732 (24.24%)	20.20 (7.98)	4.71 (4.47, 4.95)	<0.0001
Winter (December, January and February)	5994 (21.59%)	14.15 (6.85)	-1.34 (-1.60, -1.09)	<0.0001
<i>P</i> for trend				< 0.0001

Abbreviations: 25OHD, 25-hydroxy vitamin D; OR, odds ratio; CI, confidence interval; SD, standard deviation.

^aCrude associations with maternal serum 25OHD concentrations during pregnancy continuous.

Table 2. Associations of the maximum cumulative effects of PM and maternal serum 25OHD levels during the second trimester in models adjusted for different temporal and meteorological factors

Model	β (95% CI)	
	<i>P</i> -value	
	45-day moving daily average PM _{2.5} concentration	60-day moving daily average PM ₁₀ concentration
Crude Model ^a	-0.200 (-0.206, -0.193) <0.00001	-0.142 (-0.146, -0.137) <0.00001
Model 1 (Age + Year + Season) ^b	-0.113 (-0.125, -0.101) <0.00001	-0.090 (-0.099, -0.081) <0.00001
Model 2 (Model 1 + Atmospheric Pressure) ^c	-0.042 (-0.055, -0.028) <0.00001	-0.039 (-0.049, -0.030) <0.00001
Model 3 (Model 2 + Sunshine Duration) ^d	-0.039 (-0.053, -0.026) <0.00001	-0.046 (-0.055, -0.036) <0.00001
Model 4 (Model 3 + Relative Humidity +Wind Speed) ^e	-0.032 (-0.046, -0.018) <0.00001	-0.039 (-0.049, -0.028) <0.00001

^aNo adjusted.

^bAdjusted for year, age and season.

^cAdjusted for year, age, season, and corresponding-day moving daily average atmospheric pressure.

^dAdjusted for year, age, season, the corresponding-day moving daily average atmospheric pressure and sunshine duration.

^eAdjusted for year, age, season, the corresponding-day moving daily average atmospheric pressure, sunshine duration, relative humidity and wind speed.

Abbreviations: PM, particulate matter; PM_{2.5}, particulate matter with an aerodynamic diameter of ≤ 2.5 μm ; PM₁₀, particulate matter with an aerodynamic diameter of ≤ 10 μm ; 25OHD, 25-hydroxy vitamin D; CI, confidence interval.

Table 3A. Threshold effect analysis examining associations between 45-day moving daily average PM_{2.5} levels and maternal serum 25OHD levels during second trimester in subgroups stratified by season of blood collection

	Spring	Summer	Autumn	Winter	Total
Model A ^a					<i>P</i> -interaction: <0.001
One line slope, β (95%CI) <i>P</i> -value	-0.027 (-0.069, 0.014) 0.1919	-0.187 (-0.247, -0.128) <0.0001	-0.173 (-0.250, -0.097) <0.0001	-0.029 (-0.051, -0.006) 0.0136	-0.032 (-0.046, -0.018) <0.0001
Model B ^b					<i>P</i> -interaction: <0.001
Turning point (K), $\mu\text{g}/\text{m}^3$	38.62	35.4	31.11	46.31	20.07
< K, β (95%CI) <i>P</i> -value	-0.229 (-0.331, -0.127) <0.0001	-0.348 (-0.427, -0.268) <0.0001	0.020 (-0.086, 0.125) 0.7130	-0.130 (-0.192, -0.068) <0.0001	-1.798 (-2.065, -1.532) <0.0001
> K, β (95%CI) <i>P</i> -value	0.080 (0.016, 0.145) 0.0150	0.169 (0.038, 0.300) 0.0115	-0.552 (-0.713, -0.390) <0.0001	-0.013 (-0.038, 0.011) 0.2825	-0.022 (-0.036, -0.008) 0.0018
Slope 2 – Slope 1, β (95%CI) <i>P</i> -value	0.309 (0.166, 0.452) <0.0001	0.516 (0.347, 0.685) <0.0001	-0.571 (-0.787, -0.356) <0.0001	0.117 (0.050, 0.183) 0.0006	1.776 (1.509, 2.044) <0.0001
Predicted 25OHD levels at K (95% CI), ng/mL	15.977 (15.655, 16.298)	18.303 (17.982, 18.625)	19.469 (19.137, 19.802)	14.284 (14.024, 14.543)	20.635 (20.488, 20.782)
LRT ^c , <i>P</i> -value	<0.001	<0.001	<0.001	<0.001	<0.001

Adjusted for year, age, 45-day moving daily average atmospheric pressure, sunshine duration, relative humidity and wind speed.

^aLinear analysis, *P*-value <0.05 indicates a linear relationship.

^bNon-linear analysis.

^c*P* <0.05 means Model B is significantly different from Model A, which indicates a non-linear relationship.

Abbreviations: PM_{2.5}, particulate matter with an aerodynamic diameter of $\leq 2.5 \mu\text{m}$; 25OHD, 25-hydroxy vitamin D; CI, confidence interval; LRT, logarithmic likelihood ratio test.

Table 3B. Threshold effect analysis examining associations between 60-day moving daily average PM₁₀ level and maternal serum 25OHD levels during second trimester in subgroups stratified by season of

blood collection

	Spring	Summer	Autumn	Winter	Total
Model A ^a					<i>P</i> -interaction: <0.001
One line slope, β (95%CI) <i>P</i> -value	-0.102 (-0.142, -0.061) <0.0001	-0.058 (-0.098, -0.019) 0.0040	-0.003 (-0.062, 0.057) 0.9227	-0.023 (-0.044, -0.002) 0.0313	-0.039 (-0.049, -0.028) <0.0001
Model B ^b					<i>P</i> -interaction: 0.383
Turning point (K), $\mu\text{g}/\text{m}^3$	93.15	45.07	38.10	113.18	38.28
< K, β (95%CI) <i>P</i> -value	-0.080 (-0.122, -0.038) 0.0002	-0.310 (-0.430, -0.189) <0.0001	-1.121 (-1.513, -0.728) <0.0001	-0.012 (-0.034, 0.010) 0.2946	-0.711 (-0.841, -0.580) <0.0001
> K, β (95%CI) <i>P</i> -value	-0.530 (-0.734, -0.325) <0.0001	-0.030 (-0.071, 0.012) 0.1648	-0.046 (-0.108, 0.015) 0.1383	-0.379 (-0.576, -0.182) 0.0002	-0.038 (-0.049, -0.027) <0.0001
Slope 2 – Slope 1, β (95%CI) <i>P</i> -value	-0.450 (-0.661, -0.239) <0.0001	0.280 (0.154, 0.407) <0.0001	1.074 (0.702, 1.447) <0.0001	-0.367 (-0.570, -0.165) 0.0004	0.673 (0.542, 0.803) <0.0001
Predicted 25OHD levels at K (95% CI), ng/mL	14.927 (14.625, 15.230)	20.666 (20.345, 20.988)	20.828 (20.475, 21.180)	12.769 (12.366, 13.172)	20.950 (20.789, 21.112)
LRT ^c , <i>P</i> -value	<0.001	<0.001	<0.001	<0.001	<0.001

Adjusted for year, age, 60-day moving daily average atmospheric pressure, sunshine duration, relative humidity and wind speed.

^aLinear analysis, *P*-value <0.05 indicates a linear relationship.

^bNon-linear analysis.

^c*P* <0.05 means Model B is significantly different from Model A, which indicates a non-linear relationship.

Abbreviations: PM₁₀, particulate matter with an aerodynamic diameter of $\leq 10 \mu\text{m}$; 25OHD, 25-hydroxy vitamin D; CI, confidence interval; LRT, logarithmic likelihood ratio test.

Table 4A. Associations between 45-day moving daily average PM_{2.5} levels and maternal serum 25OHD levels during second trimester of pregnancy in subgroups stratified by meteorological factors

	N	β (95% CI) <i>P</i> -value	<i>P</i> -value for interaction
45-day moving daily average atmospheric pressure			<0.0001
Tertile 1 (1002.65 - 1010.69 hPa)	9247	-0.191 (-0.250, -0.132) <0.0001	
Tertile 2 (1010.74 - 1020.73 hPa)	9260	-0.042 (-0.077, -0.007) 0.0193	
Tertile 3 (1020.75 - 1028.82 hPa)	9261	-0.023 (-0.041, -0.005) 0.0141	
45-day moving daily average relative humidity			<0.0001
Tertile 1 (62.58 - 71.11 %)	9220	0.140 (0.105, 0.175) <0.0001	
Tertile 2 (71.13 - 77.04 %)	9275	-0.086 (-0.111, -0.060) <0.0001	
Tertile 3 (77.07 - 94.33 %)	9273	-0.135 (-0.169, -0.102) <0.0001	
45-day moving daily average sunshine duration			<0.0001
Tertile 1 (1.28 - 4.10 hours)	9136	-0.044 (-0.064, -0.024) <0.0001	
Tertile 2 (4.12 - 5.14 hours)	9303	-0.030 (-0.063, 0.003) 0.0717	
Tertile 3 (5.15 - 8.68 hours)	9329	-0.190 (-0.242, -0.137) <0.0001	
45-day moving daily average wind speed			0.2390
Tertile 1 (1.36 - 1.87 m/s)	9138	-0.068 (-0.088, -0.049) <0.0001	
Tertile 2 (1.88 - 2.12 m/s)	9117	-0.040 (-0.069, -0.011) 0.0066	
Tertile 3 (2.13 - 2.65 m/s)	9513	-0.047 (-0.091, -0.004) 0.0331	

Adjusted for year, season, 45-day moving daily average atmospheric pressure, sunshine duration, relative humidity and wind speed except the subgroup variable.

Abbreviations: PM_{2.5}, particulate matter with an aerodynamic diameter of $\leq 2.5 \mu\text{m}$; 25OHD, 25-hydroxy vitamin D; CI, confidence interval.

Table 4B. Associations between 60-day moving daily average PM₁₀ levels and maternal serum 25OHD levels during second trimester of pregnancy in subgroups stratified by meteorological factors

	N	β (95% CI) <i>P</i> -value	<i>P</i> -value for interaction
60-day moving daily average atmospheric pressure			<0.0001
Tertile 1 (1003.24 - 1010.76 hPa)	9241	-0.154 (-0.190, -0.118) <0.0001	
Tertile 2 (1010.78 - 1020.92 hPa)	9270	-0.132 (-0.157, -0.108) <0.0001	
Tertile 3 (1020.97 - 1028.09 hPa)	9257	-0.021 (-0.037, -0.005) 0.0100	
60-day moving daily average relative humidity			<0.0001
Tertile 1 (63.32 - 71.30 %)	9189	0.071 (0.044, 0.099) <0.0001	
Tertile 2 (71.32 - 77.18 %)	9289	-0.053 (-0.071, -0.035) <0.0001	
Tertile 3 (77.23 - 92.27 %)	9290	-0.074 (-0.096, -0.051) <0.0001	
60-day moving daily average sunshine duration			<0.0001
Tertile 1 (1.84 - 4.07 hours)	9237	-0.026 (-0.043, -0.008) 0.0034	
Tertile 2 (4.08 - 5.15 hours)	9234	-0.057 (-0.078, -0.035) <0.0001	
Tertile 3 (5.16 - 7.90 hours)	9297	-0.128 (-0.157, -0.099) <0.0001	
60-day moving daily average wind speed			0.0032
Tertile 1 (1.34 - 1.89 m/s)	8993	-0.043 (-0.059, -0.027) <0.0001	
Tertile 2 (1.90 - 2.11 m/s)	9306	-0.078 (-0.102, -0.054) <0.0001	
Tertile 3 (2.12 - 2.58 m/s)	9469	-0.094 (-0.124, -0.065) <0.0001	

Adjusted for year, season, 60-day moving daily average atmospheric pressure, sunshine duration, relative humidity and wind speed except the subgroup variable.

Abbreviations: PM₁₀, particulate matter with an aerodynamic diameter of $\leq 10 \mu\text{m}$; 25OHD, 25-hydroxy vitamin D; CI, confidence interval.

Figures

Figure 1

Temporal trends in monthly average serum 25OHD concentration vs. monthly average PM_{2.5} concentration (A), PM₁₀ concentration (B), atmospheric pressure (C), temperature (D), relative humidity (E), sunshine duration (F), and wind speed (G) from January 2016 to December 2020. 25OHD, 25-hydroxy vitamin D; PM_{2.5}, particulate matter with an aerodynamic diameter of $\leq 2.5 \mu\text{m}$; PM₁₀, particulate matter with an aerodynamic diameter of $\leq 10 \mu\text{m}$.

Figure 2

Association among exposure to PM_{2.5} (A) or PM₁₀ (B) and maternal serum 25OHD levels during pregnancy. The cumulative effects of PM were divided into four levels based on quartiles. After adjusting for year and age at blood collection (adjust I) or year, season, and age at blood collection (adjust II), the β -values (95% CI) of maternal serum 25OHD levels during pregnancy were calculated based on the group trend. The dashed vertical line represents lag days corresponding to the maximum cumulative effects. 25OHD, 25-hydroxy vitamin D; PM_{2.5}, particulate matter with an aerodynamic diameter of $\leq 2.5 \mu\text{m}$; PM₁₀, particulate matter with an aerodynamic diameter of $\leq 10 \mu\text{m}$; CI, confidence interval.

Figure 3

Adjusted smoothed curves for 45-day moving daily average PM_{2.5} concentration (A), 60-day moving daily average PM₁₀ concentration (B), and maternal serum 25OHD levels during pregnancy stratified by seasons. Thresholds were nonlinear associations between PM and 25OHD, as evidenced in generalized additive models. Adjustment factors included year, age, corresponding-day moving daily average atmospheric pressure, sunshine duration, relative humidity, and wind speed. 25OHD, 25-hydroxy vitamin D; PM_{2.5}, particulate matter with an aerodynamic diameter of ≤ 2.5 μm; PM₁₀, particulate matter with an aerodynamic diameter of ≤ 10 μm.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [FigureS1.jpg](#)
- [FigureS2.jpg](#)
- [FigureS3.jpg](#)
- [FigureS3.jpg](#)
- [FigureS4.jpg](#)
- [TableS1.docx](#)
- [TableS2.docx](#)
- [TableS3.docx](#)
- [TableS4.docx](#)
- [TableS5.docx](#)
- [TableS6.docx](#)