

# Acute pancreatitis in 60 Iranian children: do pediatricians follow the new guidelines in diagnosis and management of acute pancreatitis?

**Mahsa soti khiabani**

Children's medical center, Tehran University of medical sciences, Tehran, Iran

**Mahya Sadat Mohammadi**

Children's medical center, Tehran University of medical sciences, Tehran, Iran

**Seyyed amirreza Ghoreyshi**

Children's medical center, Tehran University of medical sciences, Tehran, Iran

**Pejman Rohani**

Pediatric Gastroenterology and Hepatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

**Hosein Alimadadi** (✉ [hoseinalimadadi@yahoo.com](mailto:hoseinalimadadi@yahoo.com))

Pediatric Gastroenterology and Hepatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

**Mohammad Hassan Sohoul**

Pediatric Gastroenterology and Hepatology Research Center, Tehran University of Medical Sciences, Tehran, Iran

---

## Research Article

**Keywords:** pancreatitis, pediatric, etiology, fluid management, antibiotic, imaging, COVID-19

**Posted Date:** March 15th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1347854/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** the incidence of acute pancreatitis in children is increasing but causes and diagnostic and therapeutic methods are different in different centers. The aim of this study was to investigate the common causes and routine diagnostic and therapeutic methods of acute pancreatitis in children in a pediatric gastrointestinal referral center and its compliance with existing guidelines.

**Methods:** In this retrospective, cross-sectional study, a total of 60 children with a diagnosis of acute pancreatitis, were studied.

**Results:** The most common causes of acute pancreatitis were systemic and metabolic diseases and medications. CT scan was performed for 36% of patients, but 31% of patients, for whom a CT scan was performed had no clear indication of CT scan. Only half of the patients received fluid 1.5 times their maintenance in the first 24 hours. Antibiotic therapy was performed for 48% of patients but medical indications for antibiotic treatment were found in only 34% of cases. During the COVID-19 pandemic, the relative incidence of acute pancreatitis was increased.

**Conclusions:** In children with systemic and metabolic disease and using anticonvulsant drugs, it is important to consider the incidence of this disease. In clinical education, the risks of radiation due to unnecessary CT scans and inappropriate prescription of antibiotics need to be emphasized. More research should be done to study the association between COVID-19 and acute pancreatitis.

## Background

The incidence of acute pancreatitis, as the most common pancreatic disorder in children, has increased in recent decade [1–3]. In children, the most common etiologies include disorders or obstructions of the biliary system, impenetrable abdominal injuries, multisystem diseases, metabolic diseases, drug poisoning, and genetic predisposing factors [4–7]. Moreover, the most common systemic diseases associated with acute pancreatitis in children include autoimmune pancreatitis, Crohn's disease, diabetes mellitus (diabetic ketoacidosis), Henoch-Schoenlein purpura, and hemolytic uremic syndrome [8, 9]. Therefore, it is important to find the etiological causes of acute pancreatitis for disease management in children.

According to the International Study Group of Pediatric Pancreatitis in Search for a CuRE (INSPPIRE) Consortium, diagnosis of pancreatitis in children is defined as having two of the following three signs: 1) abdominal pain associated with pancreatitis; 2) amylase or lipase activity at least three times higher than normal; and 3) imaging findings suggesting pancreatitis. Although contrast-enhanced computed tomography (CT) is the gold standard modality in the diagnosis of acute pancreatitis, ultrasonography is recommended for the initial examination; however, CT scan is not recommended for all patients [10].

The goal of medical treatment for acute pancreatitis is to control pain and maintain metabolic balance. Water, electrolyte, and mineral modification is an important part of patient management[10]. Evidence

suggests that an earlier onset of mouth feeding reduces the rate of complications and the length of hospital stay [11]. In severe acute pancreatitis, antibiotics are used to treat infectious necrosis, whereas prophylactic antibiotics are not recommended[9]. Besides, suppression of gastric acid with proton-pump inhibitors may be beneficial[8].

As only few original studies have investigated acute pancreatitis in children in Iran, in the present study, the clinical manifestations and etiological factors of this disease, as well as diagnostic and therapeutic measures, were investigated in children.

## Methods

In this retrospective, cross-sectional study, a total of 60 children with a diagnosis of acute pancreatitis, admitted to Children's Medical Center Hospital in Tehran, Iran, from 2013 to 2020, were studied. The patients' data were collected by reviewing the hospital data archive. Based on the evaluation of medical records, patients under the age of 18 years, who met the diagnostic criteria for acute pancreatitis according to the INSPPIRE criteria, were included in the study. On the other hand, patients diagnosed with another disease during hospitalization or those without a definite diagnosis of acute pancreatitis were excluded. If a patient had recurrent pancreatitis, only the first occurrence was considered in this study.

The patients' demographic information, signs, and symptoms were collected from their medical history. To find the predisposing factors for pancreatitis in children, the patients were surveyed for hepatobiliary disease, infectious diseases, systemic diseases, history of trauma, and medication history. Diagnostic data, including laboratory and radiological findings, were also collected and reviewed. The treatment measures were determined based on the physician orders documented in the medical files. All **diagnostic measures** and **therapeutic measures** compared with the North American Society for Pediatric Gastroenterology Hepatology and Nutrition Pancreas Committee guideline for management of Acute Pancreatitis in the Pediatric Population[10].

### *Statistical analysis methods:*

Data analysis was performed using SPSS software version 22. The normality of data distribution was assessed using the Shapiro-Wilk test. Mean (standard deviation) and frequency (percentage) were used to describe quantitative and qualitative data, respectively. Independent sample T-Test or, if necessary, its non-parametric equivalent (Mann-Whitney test) was used to compare the mean of quantitative variables. The  $\chi^2$  test was used for comparisons of categorical variables. In this study,  $P < 0.05$  was considered significant. This study was approved by the Tehran University of Medical Sciences (ethics code: IR.TUMS.CHMC.REC.1399.118).

## Results

Among 60 children with acute pancreatitis, 33 were boys (55%), and 27 were girls (45%). The mean age of the patients was  $8.25 \pm 4.4$  years (range: 4 months-16 years).

**Clinical manifestations:** Regarding the patients' clinical manifestations, abdominal pain (91%) and nausea and vomiting (80%) were the most common symptoms. Some patients also had fever, diarrhea, back pain, and jaundice. In the physical examination at the time of admission, the most common sign was epigastric tenderness (50%). Other common signs included tachycardia, tachypnea, and hyperthermia (Table 1).

**Table 1: Clinical manifestations**

	<b>Total number:60</b>
Abdominal pain	55 (91.7%)
Nausea/vomiting	48 (80%)
Epigastric tenderness	30 (50%)
Tachycardia	19 (31%)
Tachypnea	15 (25%)
Temperature>38	9 (15%)
Diarrhea	7 (11.3%)
Jaundice	3 (5%)
Back pain	3 (5%)
Irritability	3 (5%)

**Etiologies:** The evaluation of predisposing factors for pancreatitis in children showed systemic diseases (21%), drugs (13%) , hepatobiliary abnormalities (10%), infectious diseases (10%) and recent trauma history (8%) as common causes of acute pancreatitis. However, the study of 17 (28%) patients did not indicate the specific cause of acute pancreatitis (Table 2). corticosteroid, valproic acid and carbamazepine were the most common causes of drug-induced pancreatitis in this study. All children with acute pancreatitis during infancy had predisposing factors, such as hyperlipidemia.

The North American Society for Pediatric Gastroenterology Hepatology and Nutrition Pancreas Committee, has published a guideline for management of Acute Pancreatitis in the Pediatric Population in 2018.(9) In this study all **diagnostic measures** and **therapeutic measures compared** with **this guideline to determine whether our pediatricians follow the guideline recommendations or not.**

**Table 2. Acute pancreatitis etiologies**

**Total number:60**

Hepato biliary system anomaly	6 (10%)	<i>Choledocal cyst: 4(6.6%)</i> <i>Common bile duct stone: 1 (1.7%)</i> <i>Pancreatic duct stenosis: 1(1.7%)</i>
Infection disease	6 (10%):	<i>Urinary tract infection: 1 (1.7%)</i> <i>Upper respiratory tract infection: 1(1.7%)</i> <i>Spontaneous bacterial peritonitis (SBP):1(1.7%)</i> <i>Pneumonia: 2 (3.3%)</i> <i>Covid-19 infection (positive PCR): 1(1.7%)</i>
Systemic disease	13 (21%)	<i>Hyperlipidemia: 5 (8.3%)</i> <i>Minor thalassemia: 2 (3.3%)</i> <i>Systemic lupus erythematosus: 1(1.7%)</i> <i>Hypothyroidism: 1 (1.7%)</i> <i>End stage renal disease: 1(1.7%)</i> <i>Metabolic disease: 1 (1.7%)</i> <i>Cystic fibrosis: 1 (1.7%)</i> <i>Diabetes mellitus: 1(1.7%)</i>
Trauma history	5(8.3%)	<i>Blunt trauma: 3(5%)</i> <i>Post-surgery: 2(3%)</i>
Drugs associated with acute pancreatitis	8(13.3%)	<i>Sodium valproate: 3 (5%)</i> <i>Corticosteroid: 4(6.6%)</i> <i>Carbamazepine + sodium valproate: 1 (1.7%)</i>

**Diagnostic measures:** elevated serum amylase and lipase activity was the most common finding in the tests. The lowest level of amylase was 13 U/L, while its highest level was 6,117 U/L. Besides, the lowest lipase level was 20 U/L, while the highest level was 4,510 U/L. Normal amylase and lipase levels were observed in only four patients with hyperlipidemia during medical assessments. In addition, increased inflammatory markers and leukocytosis were common findings in the patients. Elevated hepatic transaminases and increased levels of gamma-glutamyl transferase (GGT) were also observed in some patients. Hypocalcemia, hyperglycemia, and hypoglycemia were detected in a small number of patients. Also, regarding the patients' lipid profile, four patients were diagnosed with hyperlipidemia (Table 3). Considering the imaging procedures, ultrasonography was performed for 53 (88%) patients; findings

consistent with pancreatitis were found in only 27 (50%) cases. Moreover, CT scan was performed for 22 (36%) patients, 17 (77%) of whom showed evidence of acute pancreatitis. In the clinical case study of 7 (31%) patients, for whom a CT scan was performed, no specific indication was found for CT scan, based on the medical records (Table 4).

**Table 3. Laboratory data analysis**

Leukocytosis	<b>32 (53%)</b>	Raised amylase	<b>54 (90%)</b>
Anemia	10 (16.6%)	Raised lipase	<b>55 (91%)</b>
Thrombocytosis	8 (13%)	Hyperlipidemia	<b>4 (6.6%)</b>
Thrombocytopenia	7 (11.6%)	Raised transaminases	<b>8 (13.3%)</b>
Abnormal renal function test	2 (3.3%)	Raised gamma GT	<b>6 (10%)</b>
Hypoglycemia	2 (3.3%)	Raised ALK P	<b>6 (10%)</b>
Hyperglycemia	5 (8.3%)	Raised ESR/CRP	<b>15 (25%)</b>
Hypocalcemia	3 (5%)		

**Table 4. Imaging data analysis.**

	<b>Total number: 60 patients</b>
Abdominal sonography	Performed: 53 (88%) Positive for acute pancreatitis: 27 (45%)
Abdominal computed tomography (CT scan)	Performed: 22 (36%) Positive for acute pancreatitis: 17 (28%) No clear indication for CT scan: 7 (11.6%)
Magnetic resonance cholangiopancreatography (MRCP)	Performed: 5 (8%)

**Therapeutic measures:** The therapeutic measures taken for the patients were examined in this study (table 5). For 11 (18%) patients, an isotonic intravenous bolus fluid was prescribed to treat dehydration. Only half of the patients received fluid 1.5 times their maintenance in the first 24 hours. Analgesia was prescribed for only 28 (46%) patients. Fifteen patients (25%) kept NPO for more than 24 hours. Antibiotic therapy was performed for 28 (48%) patients. Medical indications for antibiotic treatment (e.g., infectious necrosis) were found in only 10 (34%) cases, based on the medical records. One of the patients initially underwent surgery due to a diagnosis of acute appendicitis; however, during surgery, it was found that the appendix was normal and that the patient had acute pancreatitis.

**Table 5. Therapeutic measures**

<b><i>Therapeutic measures</i></b>	<b><i>Total number:60</i></b>
Resuscitation fluid	11(18%)
Correct fluid therapy (1.5-2* maintenance)	30(50%)
Analgesic treatment	28 (46%)
Acetaminophen	16(26%)
Petidine	12(20%)
Proton pump inhibitor	54(90%)
<b><i>Start oral feeding:</i></b>	20 (33%)
0-12hours	25 (41%)
12-24 hours	15 (25%)
later than 24hours	
Antibiotic	29 (48%)
	<i>No clear indication for antibiotic therapy :19(31.6%)</i>
TPN	1 (1.6%)
Surgery	4 (6%)

### ***Outcomes:***

Four patients required surgery after being diagnosed with acute pancreatitis. The causes of surgery were *choledochal* cyst excision, pancreatic pseudocysts, and pancreatic lesions due to trauma, followed by recurrent pancreatitis and pancreatic divisum (Table 5).

One patient with a history of cerebral palsy, presenting with fever, seizures, sepsis, and acute pancreatitis, expired after 15 days of intensive care unit (ICU) admission. It should be noted that in the last six months of this study (i.e., the first half of 2020), the COVID-19 pandemic occurred. During these six months, 11 cases of acute pancreatitis were documented in our hospital. Four cases had clinical, imaging, and laboratory evidence of COVID-19 and were admitted to the COVID-19 ward; one of them had positive nasopharyngeal PCR results for COVID-19.

## **Discussion**

In the present study, which was done on 60 children with acute pancreatitis, the number of boys (55%) was slightly higher than girls. In previous studies, the male-to-female ratio ranged from 0.9 to 1.2, which is not significantly different from the present study [2, 12–15]. The mean age of children was 8.2 years; also, other studies reported an age range of 6.9–12.5 years [2, 12–15]. In the present study, the most common clinical symptom was abdominal pain in the epigastric region (91%), followed by nausea and

vomiting (80%). In other previous studies, abdominal pain (95 – 80%) was also the most common clinical symptom [10, 16, 17].

Importantly, the clinical signs of acute pancreatitis may be atypical during infancy (e.g., nausea and vomiting, fever, irritability, and abdominal distension)[18]. Therefore, it is essential to consider the diagnosis of acute pancreatitis, especially in infants with the above-mentioned unexplained symptoms. In this study, the most common causes of acute pancreatitis were systemic and metabolic diseases (35%), medications (13%), hepatobiliary diseases (10%), and infections (10%). In other studies, systemic diseases, especially diabetic ketoacidosis, sepsis, and hemolytic uremic syndrome, were the etiologies of 10–50% of cases [10, 16, 17]. In the present study, the most common systemic diseases associated with pancreatitis was hyperlipidemia.

In previous studies, medications were the cause of 10–30% of acute pancreatitis cases [10, 16–18]. The most common medications reported in previous studies were valproic acid, L-asparaginase, 6-mercaptopurine, azathioprine, steroids, mesalamine, and metronidazole [10, 15, 16, 19]. Additionally, the most common biliary cause of acute pancreatitis in the present study was choledochal cyst, while in other studies, pancreas divisum, biliary stones, and biliary stenosis were the most common anatomical causes [10, 16, 17]; the delayed diagnosis of choledochal cysts in Iran may be the cause of this difference. Corticosteroid, valproic acid, and carbamazepine were the most common causes of drug-induced pancreatitis in this study.

In the present study, no specific cause was found for acute pancreatitis in 28% of cases. The corresponding rate was estimated at 20–24% in similar studies [10, 16, 17]; this slight difference might be due to limitations in the diagnostic accuracy of modalities. Ultrasonography was performed in 88% of patients in our study; in half of the cases, the findings were consistent with acute pancreatitis. Although ultrasound is recommended as the first diagnostic method for the diagnosis of acute pancreatitis, it is operator-dependent [10, 18, 19]. An abdominal CT scan was performed for more than one-third of cases in this study. The most important indications for CT scan were necrosis, abscess, and other complications [10, 18, 19]. Given the risk of radiation exposure, the need to perform a CT scan must be carefully considered. Also, in one-third of CT scans performed in this study, no specific reason was found for the CT scans, based on the hospital records.

Adequate fluid therapy (bolus fluid infusion and intravenous fluid therapy with 1.5-2 times maintenance in the first 24–48 hours of disease) and early onset of oral nutrition play important roles in reducing the length of hospitalization and the need for ICU admission and decreasing the likelihood of disease progression into its severe form [10, 18, 19]. Only 18% of patients in the current study received bolus intravenous fluid therapy, and only half of the patients obtained more fluid than the maintenance requirement in the first 48 hours of disease; therefore, fluid therapy should be emphasized in clinical education due to its great importance in these patients.

Moreover, one-quarter of patients remained NPO after the first 24 hours, which also increased the need for further training of physicians. Almost half of the patients in this study were treated with antibiotics, and

according to the hospital records, only one-third of them had indications for antibiotic treatment. According to the North American Society for Pediatric Gastroenterology Hepatology and Nutrition Pancreas Committee guidelines, administration of antibiotics for acute pancreatitis is only necessary in the case of infectious necrosis or disease progression into necrotic pancreatitis[10]. Besides, the inappropriate use of antibiotics, along with the side effects of medications, increases the likelihood of antibiotic resistance. In this study, only one child with cerebral palsy expired, with signs of sepsis, seizures, and acute pancreatitis. According to previous studies, the mortality rate of acute pancreatitis in children is less than 5%, which is consistent with our study [10, 16, 17].

The relatively increased incidence of acute pancreatitis in our center during the COVID-19 pandemic may be related to this infection; before the pandemic, 4–5 cases of acute pancreatitis were registered every six months according to the hospital records, while after the pandemic in Iran, 11 cases were registered during six months. It seems that several viruses play an etiological role in acute pancreatitis, including mumps, measles, Epstein-Barr virus (EBV), hepatitis A virus (HAV), hepatitis E virus (HEV), and coxsackievirus [20, 21]; therefore, an association between SARS-CoV-2 and pancreatitis is probable.

Previous studies have confirmed the gastrointestinal involvement of SARS-CoV-2 [22]. However, limited case reports have been published on the association between COVID-19 and acute pancreatitis in adult or pediatric populations during this pandemic [23–25]. The proposed pathophysiology of pancreatic involvement in COVID-19 is the expression of angiotensin-converting enzyme 2 (ACE2) in both islet cells and the exocrine portion of the pancreas[26, 27]. Pancreatic injury during an acute SARS-CoV-2 infection can be also related to indirect systemic inflammatory and immune-mediated cellular responses. Besides, antipyretics that are commonly used for COVID-19 patients can cause drug-related pancreatic damages[26, 27]. Further research is required to determine the definite effects of SARS-CoV-2 on pancreatic function and regulation.

## **Conclusion**

Considering the increasing prevalence of acute pancreatitis in children, if symptoms, such as vomiting or abdominal pain, occur in a patient with systemic or metabolic diseases, acute pancreatitis should be considered. Medications play an important role in the development of acute pancreatitis. In children using anticonvulsants, immunosuppressant drugs, and chemotherapy, it is important to consider the incidence of this disease. Moreover, in clinical education, the risks of radiation due to unnecessary CT scans for children with acute pancreatitis need to be emphasized. Besides, inappropriate prescription of antibiotics, insufficient fluid therapy, and delayed initiation of oral feeding in these patients should be highlighted in therapeutic and educational programs. Also, healthcare providers should consider COVID-19 as a differential diagnosis when managing patients with gastrointestinal symptoms.

## **Declarations**

### ***Acknowledgements***

We appreciate the contributions of study participants to this study.

### ***Authors' contributions***

The authors confirm contribution to the paper as follows: Study conception and design: Mahsa soti khiabani and hosein alimadadi. Data collection: Seyyed amirreza Ghoreyshi and Mahya Sadat Mohammadi. Analysis and interpretation of results: Mahsa soti khiabani. Draft manuscript preparation: Mahsa soti khiabani and hosein alimadadi and pejman rohani and Mohammad Hassan Sohoul. All authors reviewed the results and approved the final version of the manuscript.

### ***Funding***

No founding

### ***Availability of data and materials***

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### ***Declarations Ethics approval and consent to participate***

This study was approved by the Tehran University of Medical Sciences (ethics code: IR.TUMS.CHMC.REC.1399.118). The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The requirement for written informed consent was waived owing to the retrospective nature of the study by the ethics committees of Tehran University of Medical Sciences.

### ***Consent for publication***

Not applicable.

### ***Competing interests***

The authors declare that they have no competing interests.

## **References**

1. Bornman P: **The Pancreas: An Integrated Textbook of Basic Science, Medicine and Surgery. 2nd Edn H. Beger, A. Warshaw, M. Büchler, R. Kozarek, M. Lerch, J. Neoptolemos, K. Shiratori, D. Whitcomb (eds). 230× 283 mm. Pp. 1006. Illustrated. 2008. Blackwell Publishing: Oxford.£ 245· 00. In.: Oxford University Press; 2008.**
2. Lopez MJ: **The changing incidence of acute pancreatitis in children: a single-institution perspective. *The Journal of pediatrics* 2002, **140**(5):622-624.**
3. Morinville VD, Barmada MM, Lowe ME: **Increasing incidence of acute pancreatitis at an American pediatric tertiary care center: is greater awareness among physicians responsible? *Pancreas* 2010,**

39(1):5-8.

4. Garipey CE, Heyman MB, Lowe ME, Pohl JF, Werlin SL, Wilschanski M, Barth B, Fishman DS, Freedman SD, Giefer MJ: **The causal evaluation of acute recurrent and chronic pancreatitis in children: consensus from the INSPPIRE group.** *Journal of pediatric gastroenterology and nutrition* 2017, **64**(1):95.
5. Jamer T: **Etiologia ostrego zapalenia trzustki–niedoceniony problem w pediatrii.** *Developmental Period Medicine* 2015:341-346.
6. Alabdulkareem A, Almahmoud T, Al-Tahan H, Javad S, Al Hatlani M: **Etiology and clinical characteristics of pediatric acute pancreatitis in Saudi Arabia: a 20-year experience from a single tertiary center.** *International Journal of Pediatrics and Adolescent Medicine* 2018, **5**(1):13-17.
7. Kandula L, Lowe ME: **Etiology and outcome of acute pancreatitis in infants and toddlers.** *The Journal of pediatrics* 2008, **152**(1):106-110. e101.
8. Larino-Noia J, Lindkvist B, Iglesias-Garcia J, Seijo-Ríos S, Iglesias-Canle J, Domínguez-Muñoz J: **Early and/or immediately full caloric diet versus standard refeeding in mild acute pancreatitis: a randomized open-label trial.** *Pancreatology* 2014, **14**(3):167-173.
9. Raizner A, Phatak UP, Baker K, Patel MG, Husain SZ, Pashankar DS: **Acute necrotizing pancreatitis in children.** *The Journal of pediatrics* 2013, **162**(4):788-792.
10. Abu-El-Haija M, Lin TK, Nathan JD: **Management of acute pancreatitis in children.** *Current Opinion in Pediatrics* 2017, **29**(5):592-597.
11. Abu-El-Haija M, Wilhelm R, Heinzman C, Siqueira BNF, Zou Y, Fei L, Cole CR: **Early enteral nutrition in children with acute pancreatitis.** *Journal of Pediatric Gastroenterology and Nutrition* 2016, **62**(3):453-456.
12. Werlin SL, Kugathasan S, Frautschy BC: **Pancreatitis in children.** *Journal of pediatric gastroenterology and nutrition* 2003, **37**(5):591-595.
13. DeBanto JR, Goday PS, Pedroso MR, Iftikhar R, Fazel A, Nayyar S, Conwell DL, DeMeo MT, Burton FR, Whitcomb DC: **Acute pancreatitis in children.** *The American journal of gastroenterology* 2002, **97**(7):1726-1731.
14. Benifla M, Weizman Z: **Acute pancreatitis in childhood: analysis of literature data.** *Journal of clinical gastroenterology* 2003, **37**(2):169-172.
15. Calatayud GA, Bermejo F, Morales J, Claver E, Huber L, Abunaji J, Canete A, Boixeda D: **Acute pancreatitis in childhood.** *Revista española de enfermedades digestivas: organo oficial de la Sociedad Española de Patología Digestiva* 2003, **95**(1):40-48.
16. Husain SZ, Srinath AI: **What's unique about acute pancreatitis in children: risk factors, diagnosis and management.** *Nature Reviews Gastroenterology & Hepatology* 2017, **14**(6):366-372.
17. Uc A, Husain SZ: **Pancreatitis in children.** *Gastroenterology* 2019, **156**(7):1969-1978.
18. Wyllie R, Hyams JS: **Pediatric Gastrointestinal and Liver Disease E-Book:** Elsevier Health Sciences; 2010.

19. Kleinman RE, Goulet O-J, Mieli-Vergani G, Sanderson IR, Sherman PM, Shneider BL: **Walker's pediatric gastrointestinal disease: physiology, diagnosis, management**. PMPH USA, Ltd; 2018.
20. Párniczky A, Abu-El-Haija M, Husain S, Lowe M, Oracz G, Sahin-Tóth M, Szabó FK, Uc A, Wilschanski M, Witt H: **EPC/HPSG evidence-based guidelines for the management of pediatric pancreatitis**. *Pancreatology* 2018, **18**(2):146-160.
21. Alloway BC, Yaeger SK, Mazzaccaro RJ, Villalobos T, Hardy SG: **Suspected case of COVID-19-associated pancreatitis in a child**. *Radiology case reports* 2020, **15**(8):1309-1312.
22. Patel KP, Patel PA, Vunnam RR, Hewlett AT, Jain R, Jing R, Vunnam SR: **Gastrointestinal, hepatobiliary, and pancreatic manifestations of COVID-19**. *Journal of Clinical Virology* 2020, **128**:104386.
23. Samies NL, Yarbrough A, Boppana S: **Pancreatitis in pediatric patients with COVID-19**. *Journal of the Pediatric Infectious Diseases Society* 2021, **10**(1):57-59.
24. de-Madaria E, Capurso G: **COVID-19 and acute pancreatitis: examining the causality**. *Nature Reviews Gastroenterology & Hepatology* 2021, **18**(1):3-4.
25. Aloysius MM, Thatti A, Gupta A, Sharma N, Bansal P, Goyal H: **COVID-19 presenting as acute pancreatitis**. *Pancreatology* 2020, **20**(5):1026-1027.
26. Liu F, Long X, Zhang B, Zhang W, Chen X, Zhang Z: **ACE2 expression in pancreas may cause pancreatic damage after SARS-CoV-2 infection**. *Clinical Gastroenterology and Hepatology* 2020, **18**(9):2128-2130. e2122.
27. Wang F, Wang H, Fan J, Zhang Y, Wang H, Zhao Q: **Pancreatic injury patterns in patients with coronavirus disease 19 pneumonia**. *Gastroenterology* 2020, **159**(1):367-370.