

Prevalence, antimicrobial susceptibility pattern, and associated factors of *Salmonella* and *Shigella* among food handlers in Adigrat University student's cafeteria, Northern Ethiopia, 2018

Haftom Legese (✉ legesehaftom2@gmail.com)

Adigrat university <https://orcid.org/0000-0002-6280-1116>

Tsega Kahsay

Adigrat University College of Health Sciences

Aderajew Gebrewahd

Adigrat University College of Health Sciences

Brhane Berhe

Adigrat University College of Health Sciences

Berhane Fseha

Adigrat University College of Health Sciences

Senait Tadesse

Bahir Dar University College of Medical and Health Sciences

Guesh Gebremariam

Adigrat University College of Health Sciences

Hadush Negash

Adigrat University College of Health Sciences

Fitsum Mardu

Adigrat University college of Medicine and health Sciences

Kebede Tesfay

Adigrat University College of Health Sciences

Gebre Adhanom

Adigrat University College of Health Sciences

Research

Keywords: Antimicrobial susceptibility, Food handlers, Salmonella, Shigella, Ethiopia

Posted Date: July 30th, 2020

DOI: <https://doi.org/10.21203/rs.2.22938/v5>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Tropical Diseases, Travel Medicine and Vaccines on September 11th, 2020. See the published version at <https://doi.org/10.1186/s40794-020-00119-x>.

Abstract

Background: Food handlers play a significant role in the transmission of foodborne infections. *Salmonella* and *Shigella* are the most common foodborne pathogens and their infections are a major public health problem globally. Thus, this study aimed to determine the prevalence, antimicrobial susceptibility patterns, and associated factors of *Salmonella* and *Shigella* among food handlers.

Methodology: A cross-sectional study was conducted from March to August 2018 at Adigrat University student cafeteria, Northern Ethiopia. Data on socio-demographic and associated factors were collected using a structured questionnaire. Fresh stool samples were collected from 301 food handlers and transported to Adigrat University Microbiology Laboratory. Bacterial isolation and antimicrobial susceptibility test were performed using standard bacteriological methods. Data analysis was performed using SPSS version 22 and $P < 0.05$ with a corresponding 95% confidence interval was considered statistically significant.

Results: A total of 301 food handlers were included in this study. The majority of study participants were females 265 (88.0 %). About 22 (7.3%) and 11 (3.7%) of food handlers were found to be positive for *Salmonella* and *Shigella* respectively. Hand washing after using a bathroom with water only, no hand washing after using the bathroom, no hand washing after touching dirty materials, no hand washing before food handling, and untrimmed fingernails were significant associated factors identified. None of the *Salmonella* and *Shigella* isolates were sensitive to ampicillin. On the other hand, low resistance against chloramphenicol, ceftriaxone, and ciprofloxacin was found.

Conclusion: The present study revealed that the prevalence of *Salmonella* and *Shigella* among food handlers was 22 (7.3%) and 11 (3.7%) respectively. Such colonized food handlers can contaminate food, drinks and could serve as a source of infection to consumers via the food chain. This indicates that there is a need for strengthened infection control measures to prevent *Salmonella* and *Shigella* transmission in the students' cafeteria.

Background

Foodborne diseases are a major public health problem globally. The severity is higher among developing countries due to low hygienic food handling practices, lack of environmental sanitation, and poor access to safe drinking water [1]. In developing countries, approximately 70% of cases of diarrheal diseases are associated with the consumption of contaminated food [2].

Salmonella remains a major cause of foodborne infection in humans [3], which leads to approximately 93 million infections every year [4, 5]. The World Health Organization (WHO) estimates that there are around 16 million new cases and 600,000 deaths due to typhoid fever each year worldwide [6]. It causes bacterial bloodstream infections with a fatality rate of 20-25% [7]. The widespread nature of salmonellosis increases antibiotic resistance which in turn increases the treatment cost, hospitalization, morbidity, and mortality [8].

These bacteria are transmitted directly and indirectly through contaminated objects such as food, water, nails, and fingers. This indicates those microorganisms can be spread by fecal-oral human-to-human transmission [9, 10]. Compared to other parts of the hand, fingernails harbor the most microorganisms and are difficult to clean.

Shigella continues to play a major role in the etiology of inflammatory diarrhea and dysentery in food handlers [11]. The annual incidence of *Shigella* is estimated to be 164.7 million people, with 69% of all deaths attributable to shigellosis worldwide [12, 13]. The highest prevalence of shigellosis is observed in tropical and subtropical parts of the world [14].

Salmonella and *Shigella* are a significant cause of severe post-diarrheal complications such as reactive arthritis, sepsis, Reiter's syndrome, myocarditis, inflammatory bowel diseases, irritable bowel syndrome, and peritonitis [15, 8, 16]. The emergence of antimicrobial-resistant *Salmonella* and *Shigella* becomes a significant threat to deliver reliable therapies [17, 18].

In Ethiopia, it is difficult to estimate the severity of salmonellosis and shigellosis as well as their antibiotic resistance due to the limited scope of studies, lack of coordinated epidemiological surveillance system, poor reporting system, and limited availability of culture facilities [19].

Determining the prevalence and antimicrobial susceptibility pattern of *Salmonella* and *Shigella* is very important for the proper selection of antimicrobial agents to control the spread of infection. However, in the study area, there was a scarcity of data on the carriage of *Salmonella* and *Shigella* among food handlers. Therefore, this study aims to assess the prevalence, antimicrobial

susceptibility patterns, and associated factors of *Salmonella* and *Shigella* among food handlers in Adigrat University, Tigray, Northern Ethiopia.

Materials And Methods

Study Design, Area and Period

A cross-sectional study was conducted among food handlers who participated in food preparation, dispatch and storage of Adigrat University student's cafeteria from March to August 2018. The annual rainfall ranges from 400-600mm and the minimum and maximum temperature range from 6-21.8°C. Currently, the university enrolled more than 15,000 students who are getting dining services in the student cafeteria. There are six cafeterias, and a total of 700 food handlers are working in the student's cafeteria (Adigrat University human resource management and registrar office). A large number of food handlers in the university were neither well-trained nor graduated by food handling, preparation and serving. Although there was a periodic medical checkup in the university, it was not consistent. However, those food handlers employed in serving the public and tourists are well-trained and graduated in food preparation and handling practice from college or universities.

Sample size determination and sampling technique

Sample size determination

The sample size was determined by using a single population proportion formula.

$$n = \frac{(Z_{\alpha/2})^2 \cdot P \cdot (1-P)}{d^2}$$

The sample size was determined based on the prevalence of *Salmonella* among university food handlers done by Mama and Alemu (2016) at Arba Minch University, South Ethiopia (6.9%) [14]. Then with a margin of error (5%), (d=0.03) and 95% level of confidence (z=1.96), the sample size was calculated as follows:

$$n = \frac{(1.96)^2 \cdot 0.069(0.931)}{(0.03)^2} = 274, \text{ with } 10\% \text{ non response rate} = 301$$

Therefore, a total of 301 food handlers were included in the study from all cafeteria of the university. A simple random sampling technique was employed. The lottery method was used to select the study subjects after a complete list of food handlers was obtained from a roster of cafeteria staff Adigrat University.

Eligibility criteria

Inclusion criteria

Food handlers working in Adigrat University student's cafeterias were included in the study.

Exclusion criteria

Food handlers who have taken antibiotics and/or antihelminthics within one week and those with clinical signs of typhoid fever were excluded from the study.

Data collection and Sample processing

Socio-demographic and Specimen Collection, Handling and Transportation

A structured questionnaire was used to collect the data regarding socio-demographic and associated factors. Questionnaires were checked for accuracy and completeness. After proper instruction, about 2 g of fresh stool specimens were collected from food handlers with a labeled wide-mouthed plastic container and a clean wooden applicator stick. Specimens were immediately transported to the laboratory using an icebox.

Isolation and identification

The stool specimen was collected and transported to Adigrat University Medical Microbiology laboratory within one hour of collection. Stool specimens were immediately inoculated in Selenite F enrichment broth and incubated at 37°C for 24 hours, and then subculture onto selective media of xylose-lysine desoxycholate agar (XLD) and Hektoen enteric medium agar at 37°C for 18-24 hours. The isolated colonies were differentiated and identified based on Gram stain, colonial morphology and pigmentation, hemolysis on blood agar, catalase test, oxidase test, carbohydrate fermentation, H₂S production, motility, indole formation and urease production, citrate utilization and incubated for 24 to 48 hours at 37°C. Then colonies producing an alkaline slant with acid butt and hydrogen sulfide production on Triple Sugar Iron Agar, positive for lysine, negative for urea hydrolysis, negative for indole test, positive for citrate utilization and motility test were considered to be *Salmonella*. Colonies which were urease negative, indole positive/negative, produce a pink-red slope and yellow butt with no blackening on Triple Sugar Iron agar, lysine decarboxylase negative and citrate negative is identified as *Shigella* species. Finally, all of the confirmed *Salmonella* and *Shigella* isolates were examined for antimicrobial susceptibility.

Antimicrobial susceptibility tests

Antimicrobial susceptibility testing was performed using the modified Kirby- Bauer disc diffusion method according to Clinical and Laboratory Standards Institute (CLSI) guidelines, 2016 [20]. Using a sterile wire loop, 3-5 well-isolated colonies of the test organism was emulsified into a tube of 3-4 ml sterile physiological saline to get bacterial inoculums equivalent to 0.5 McFarland turbidity standards. Then the standardized suspension (test organisms) were uniformly swabbed within 15 minutes using a sterile cotton swab into Muller-Hinton agar and allowed to dry. After that, the antibiotic discs were placed manually on the medium and incubated at 37°C for about 18 hours and the zones of inhibition were measured using a caliper. The interpretation of the results was made based on the CLSI criteria as sensitive, intermediate and resistant [20]. The following antimicrobials are prioritized by considering local prescription; gentamicin (10 µg), ampicillin (30 µg), amoxicillin (30 µg), ciprofloxacin (5 µg), clarithromycin (30 µg), chloramphenicol (30 µg), cotrimoxazole (25 µg), amoxicillin-clavulanic acid (30 µg), and ceftriaxone (30 µg) [20].

Data Quality Assurance

Data quality was ensured at various activities of the study by following a prepared standard operating procedure (SOP). Questionnaires were prepared in a clear and precise way and translated into the local language and back-translated to English to ensure the consistency of the questionnaires. The pretest was done on 5% of food handlers and modifications were made accordingly. To ensure general safety; universal bio-safety precautions were followed. American Type Culture Collection (ATCC) strains *P.aeruginosa* (ATCC 27853), *E. coli* (ATCC-25922) were used as control strains for the culture and antimicrobial susceptibility testing.

Statistical Analysis

After the collection of socio-demographic characteristics, associated factors and laboratory data using a structured questionnaire and laboratory report format, data were edited, cleaned, entered and analyzed using statistical package for social sciences (SPSS) version 22. Descriptive statistics, bivariate, and multivariate logistic regressions were performed. Bivariate logistic regression was employed to determine the association between the outcome variable and each independent variable. A binary logistic regression analysis was used to calculate the odds ratios (OR); crude odds ratio (COR) and adjusted odds ratio (AOR) to ascertain the degree of association between dependent and independent variables. All variables with p-value 0.20 in the bivariate logistic regression were transferred to multivariate logistic regression analysis to compute AOR. The regression model was first examined by the goodness of fit test using the Hosmer-Lemeshow test to determine whether the fitted model adequately described the study. In this study, multi-collinearity among independent variables was detected using the standard errors for regression coefficients. Finally, variables with a p-value < 0.05 with a corresponding 95% confidence interval (CI) were considered as statistically significant.

Operational Definition

Dirty materials: Dirty materials are soiled objects, unclean or impure

Fingernail status: Status of the nail was either trimmed or untrimmed fingernails that serve as a vehicle for the transmission of food contaminating pathogens.

Results

Socio-demographic characteristics

A total of 301 food handlers were included in the study. Out of the total respondents, 265 (88.0 %) were females. The age of study participants ranged from 19-38 years (23.51 ± 3.186 years). The majority of 241 (80.1%) of the participants were between the ages of 21 and 30 years. One hundred fifty-six (51.8%) were enrolled in secondary school with an average of 3.7 years of work experience in the cafeteria. Out of the total study participants, 37 (12.3%) were certified for training in food handling and 265 (88.0%) had previously undergone a medical checkup stool microscopy examination (**Table 1**).

Table 1 Socio-demographic characteristics of food handlers in Adigrat University student cafeteria, Tigray, North Ethiopia March to August 2018 (N=301)

Demographic characteristics	Characteristics	Frequency	Percent
Sex	Male	37	12.3
	Female	264	87.7
Age	≤ 20	54	17.9
	21-40	247	80.1
Marital status	Single	127	42.2
	Married	174	57.8
Educational level	Secondary school	187	62.12
	Higher than secondary school	114	37.9
Years of service	≤1	36	12.0
	1-2	42	14.0
	>2	223	74.1
Certified in food preparation and handling	No	264	87.7
	Yes	37	12.3
Medical check-up	No	36	12.0
	Yes	265	88.0

Prevalence and associated factors of *Salmonella* and *Shigella* carriers

The prevalence of *Salmonella* and *Shigella* in this study was 22 (7.3%) and 11 (3.7%) respectively. In the current study, 13 independent variables were considered during the analysis of associated factors for *Salmonella* and *Shigella* carriers (**Table 2**). Accordingly in the multivariate analysis, hand washing after using the bathroom with water only (AOR=23.239, 95%CI: 2.125-254.17, P<0.01), no handwashing after using the bathroom (AOR=2.25, 95%CI: 5.11-77.34, P<0.001), no hands washing after touching dirty materials (AOR=37.19, 95%CI: 5.66-244.45, P<0.001), no handwashing before food handling (AOR=33.1, 95%CI: 4.958-220.52, P<0.001), untrimmed fingernail (AOR=13.97, 95%CI: 3.404-57.362, P<0.001) were significant factors associated with *Salmonella* and *Shigella* carriers. However, no significant association was found between years of service, medical check-up, workers blowing their noses, touching food with bare hands, and certification in food preparation and handling, with *Salmonella* and *Shigella* colonization in this study (**Table 3**).

Table 2 Bivariate logistic regression analysis of factors associated with *Salmonella* and *Shigella* infections among food handler's working at Adigrat University Students' Cafeteria, Tigray, Northern Ethiopia, March to August 2018 (N=301)

Variables		Growth of <i>Salmonella</i> and <i>Shigella</i>		COR (95% CI)	P-value
		Positive N (%)	Negative N (%)		
Sex	Male	1(2.7)	36(97.3)	0.22(0.03-1.58)	0.20
	Female	32(12.1)	232(87.9)	1	
Age	≤ 20	2(3.7)	52(96.3)	0.3(0.07-1.2)	0.081
	21-40	31(12.55)	216(87.45)	1	
Marital status	Single	16(12.6)	111(87.4)	1.29(.68-2.45)	0.341
	Married	17(9.78)	157(90.23)	1	
Educational status	Lower than Secondary school	16(8.56)	171(91.44)		0.507
	Higher than secondary school	17(14.9)	97(85.1)	1.74(0.92-3.31)	
Hand washing after using the bathroom	Yes with water and soap	1(0.9)	112(99.1)	1	0.006
	Yes only with water	17(13.6)	108(86.4)	17.630(2.306-134.7)	
	No	15(23.8)	48(76.2)	35.000(4.495-272.49)	
Hand washing After touching dirty materials	Yes with water and soap	5(3.4)	141(96.6)	1	0.020
	Yes only with water	14(11.0)	113(89.0)	3.494(1.222- 9.991)	
	No	14(50.0)	14(50.0)	28.200(8.845-89.906)	
Handwashing before food handling	Yes with water and soap	2(1.8)	112(98.2)	1	0.025
	Yes only with water	13(9.2)	129(90.8)	5.643(1.247- 25.548)	
	No	18(40.0)	27(60.0)	37.333(8.164 -170.714)	
Fingernail status	Trimmed	4(2.2)	174(97.8)	1	0.001
	Untrimmed	29(23.6)	94(76.4)	13.420(4.580-39.324)	
After blowing nose	Yes with water and soap	7(8.0)	81(92.0)	1	0.842
	Yes only with water	11(11.7)	83(88.3)	0.919(0.401-2.107)	
	No	15(12.6)	104(87.4)	0.599(0.233-1.538)	
Touch food with bare hands	No	16(14.0)	98(86.0)	1	0.207
	Yes	17(9.1)	170(90.9)	1.597(0.772-3.302)	
Years of service	≤1	9(25.0)	27(75.0)	3.72(1.76-78.50)	0.001
	1-2	9(21.4)	33(78.6)	0.818(0.285- 2.349)	
	>2	15(6.7)	208(93.3)	1	
Certified in food preparation and handling	No	29(11.5)	224(88.5)	1	0.527
	Yes	4(8.3)	44(91.7)	0.702(0.235-2.097)	
Medical check-up	No	10(18.5)	44(81.5)	1	0.04
	Yes	23(9.3)	224(90.7)	1.99(1.01-3.93)	

Key: ^a(COR=Crude odds ratio); ^b(CI=Confidence interval);1(referent).

Table 3 Multivariate logistic regression analysis of factors associated with *Salmonella* and *Shigella* isolates among food handler's working at Adigrat University Students' Cafeteria, Tigray, Northern Ethiopia, March to August 2018 (N=301)

Variables	Growth of <i>Salmonella</i> and <i>Shigella</i>		COR ^a (95% CI ^b)	P-value	AOR ^c (95%CI)	P-value
	Negative N (%)	Positive N (%)				
Hand washing using after using the bathroom						
Yes with water and soap	112(99.1)	1(0.9)	1		1	
Yes only with water	108(86.4)	17(13.6)	17.630(2.306,134.7)	0.006	23.239(2.125-254.17)	0.010*
No	48(76.2)	15(23.8)	35.000(4.495,272.49)	0.001	62.917(5.11-77.34)	0.001*
Hand washing after touching dirty materials						
Yes with water and soap	141(96.6)	5(3.4)	1		1	
Yes only with water	113(89.0)	14(11.0)	3.494(1.222, 9.991)	0.020	1.089(0.187-6.345)	0.924
No	14(50.0)	14(50.0)	28.200(8.845, 89.906)	0.000	37.189(5.658-244.45)	0.001*
Hand washing before food handling						
Yes with water and soap	112(98.2)	2(1.8)	1		1	
Yes only with water	129(90.8)	13(9.2)	5.643(1.247, 25.548)	0.025	5.972(0.899-39.677)	0.064
No	27(60.0)	18(40.0)	37.333(8.164, 170.714)	0.000	33.065(4.958-220.52)	0.001*
Finger nail status						
Trimmed	174(97.8)	4(2.2)	1		1	
Untrimmed	94(76.4)	29(23.6)	13.420(4.580, 39.324)	0.000	13.973(3.404-57.362)	0.001*

Key: ^a(COR=Crude odds ratio); ^b(CI=Confidence interval); ^c(AOR=Adjusted odds ratio); 1(referent).

Antimicrobial Susceptibility Patterns of *Salmonella* and *Shigella* isolates

Antimicrobial susceptibility patterns were performed for 22 *Salmonella* and 11 *Shigella* isolates against 9 antimicrobial agents. 100% *Salmonella* and *Shigella* isolates were resistant to ampicillin. 22(100%) and 10(90.90%) *Salmonella* and *Shigella* isolates were resistant to gentamicin, respectively. 21(95.50%), and 11(100%) *Salmonella* and *Shigella* isolates were resistant to amoxicillin, respectively. However, all isolates were susceptible to ciprofloxacin. Additionally, low resistance was observed for ceftriaxone, chloramphenicol and cotrimoxazole for *Salmonella* and *Shigella* respectively. None of the isolates showed intermediate resistance (Table 4). Multidrug resistance in this study is defined as resistance to at least three classes of antimicrobial agents and out of the thirty-three isolates 12 (54.5%) *Salmonella* and 10(90.90%) *Shigella* species were multi-drug resistant isolates (Table 5).

Table 4 Antimicrobial susceptibility patterns of *Salmonella* and *Shigella* isolated from food handlers Of Adigrat University Students' Cafeteria, Tigray, Northern Ethiopia, March to August 2018 (N=33)

Sensitivity pattern n(%)	Antimicrobial agents N (%)									
	GM	AML	AMP	CIP	CLR	TS	AMC	CHL	CRO	
n(22)	S	22(100)	1(4.5)	0(0.00)	22(100)	18(81.8)	20(90.9)	13(59.1)	22(100)	22(100)
	R	0(0.00)	21(95.5)	22(100)	0(0.00)	4(18.2)	2(9.1)	9(40.9)	0(0.00)	0(0.00)
(11)	S	10(90.9)	0(0.00)	0(0.00)	11(100)	2(18.2)	10(90.9)	6(54.5)	9(81.2)	10(90.9)
	R	1(9.1)	11(100)	11(100)	0(0.00)	9(81.2)	1(9.1)	5(45.5)	2(18.2)	1(9.1)

Key: S = Sensitive R = Resistant, GM Gentamicin, AML Amoxicillin, AMP Ampicillin, CIP=Ciprofloxacin, CLR Clarithromycin, TS Cotrimoxazole, AMC Amoxicillin-clavulanic acid CHL Chloramphenicol, CRO Ceftriaxone

Table 5 Multidrug-resistant of *Salmonella* and *Shigella* isolated from food handler's working at Adigrat University Students' Cafeteria, Tigray, Northern Ethiopia, March to August 2018 N=33

		Resistant isolates no. (%)	
	Antimicrobials	<i>Salmonella</i>	<i>Shigella</i>
For three	AML,AMP,CLR	2(16.67)	4(40)
	AML,AMP,CHL	-	1(10)
	AML,CLR,AMC	7(58.34)	-
	AMP,CLR,AMC	1(8.33)	-
	AML,AMP,TS	1(8.33)	-
FOR FOUR	AML,AMP,CLR,AMC	1(8.33)	3(30)
FOR FIVE	AML,AMP,CLR,AMC,CHL	-	1(10)
FOR SIX	GM,AML,AMP,CLR,TS,AMC	-	1(10)
		12(100)	10(100)

Key: GM Gentamicin, AML Amoxicillin, AMP Ampicillin, CIP=Ciprofloxacin, CLR Clarithromycin, TS Cotrimoxazole, AMC Amoxicillin-clavulanic acid CHL Chloramphenicol, CRO, Ceftriaxone

MDR multidrug resistant; MDR definition for *Salmonella* and *Shigella* percent is computed from total number of *Salmonella* and *Shigella*

Discussion

In this study, the prevalence of *Salmonella* among food handlers was 22 (7.3%). This was similar to studies carried out in Southern Ethiopia, Arba Minch University (6.9%) [14], and Nigeria, Abeokuta (5.5%) [21]. However, it was higher than the studies reported from Ethiopia, Addis Ababa (3.5%) [22], Bahir Dar (1.6%) [23], and Gondar (3.1%) [24]. On the other hand, our result was lower than the studies reported from Ethiopia, Addis Ababa (10.5%) [25], and Nigeria (42.3%) [26]. The variation might be attributed to poor personal hygiene and environmental sanitation differences among the study areas.

The prevalence of *Shigella* (3.7%) in our study is consistent with studies done in Southern Ethiopia, Arba Minch University (3%) [14], Ethiopia, Addis Ababa (4.5%) [25], and Gondar (3.1%) [26]. However, our finding was lower than a study conducted in Nigeria (15.5%) [27]. These might be due to the differences in inconsistent training on food preparation and handling and hygiene practices of the food handlers.

In the present study, not practicing handwashing after using the bathroom among food handlers was significantly associated with *Salmonella* and *Shigella* carriers. Food handlers who hadn't washed their hands after using the bathroom were more likely to be colonized with *Salmonella* and *Shigella* compared to those who washed with water and soap after using the bathroom. This finding was similar to a study conducted in Ethiopia, Mekelle [28], Gondar [29], and Bahir Dar [30]. The acquisition of *Salmonella* and *Shigella* is due to poor sanitary conditions, poor toilet facilities, and scarcity of availability of facilities used for handwashing practice. A majority of food handlers of the university reported that they washed their hands with only water and some of them not wash their hands neither with water only nor with soap and water after the bathroom.

Our finding also revealed that there is a statistically significant difference in handwashing after touching dirty materials among food handlers with *Salmonella* and *Shigella* carriers. Food handlers who did not wash their hands after touching dirty materials are twenty-eight fold more likely to be colonized with *Salmonella* and *Shigella* than those who washed with water and soap after touching dirty materials. This finding is consistent with a study conducted in Ethiopia, Bahir Dar [23]. This might be due to the absence of handwashing facilities within proximity of the food handler's workplace.

Our study showed that food handlers who washed their hand with soap and water before touching food were less likely to be colonized with *Salmonella* and *Shigella* than food handlers who did not wash their hand with soap and water before food preparation. This is in line with the finding of a similar study reported from Ethiopia, Yebu Town [31]. In the majority of food handlers, handwashing before handling food was practiced. However, a very large proportion (42.8%) washed their hands only with water. There are food handlers who apply some hygiene practice, though many of them do not use soap nor do they appreciate or understand the need for handwashing [32].

Furthermore, in this finding, untrimmed fingernail was significantly associated with *Salmonella* and *Shigella* colonization among food handlers. This study is similar to studies conducted in Ethiopia, Yebu Town [31], and Arba Minch [14]. This result might be due to the lifestyle of food handlers. Examination of fingernail contents of food handlers for *Salmonella* or *Shigella* is one way of indicating a source of possible food contamination [31]. However, the current study did not assess the *Salmonella* and *Shigella* carriage of fingernail contents.

Salmonella and *Shigella* carriers who are preparing and handling food daily can act as sources of infection to the community the university via the food chain. Therefore, regular training, medical check-up programs and accessibility of personal hygiene guidelines with intensive health education could be important to prevent and control the carriage.

Antimicrobial susceptibility pattern data showed that ciprofloxacin, ceftriaxone, gentamicin, chloramphenicol, and cotrimoxazole were effective against the *Shigella* isolates. Our finding was comparable with studies reported from Ethiopia, Haramaya University on ceftriaxone (16.7%) [33], Jimma on gentamicin (1.3%) [34], and Harar on gentamicin (3.6%) [35]. Whereas our result showed lower resistance patterns compared to the studies conducted in Ethiopia, Addis Ababa on gentamicin (75.6%) [36], and Gondar on ciprofloxacin (8.9%) and cotrimoxazole (73.4%) [37,38]. This increase of resistance from those reports indicates that there are differences in the geographical area, study period and study design. Increased resistance was observed in our finding which is in line with a study reported from Harar on ampicillin (100%) [35], Arba Minchon amoxicillin (100%) [14].

In the current study, isolates of *Salmonella* species were sensitive to gentamicin, ciprofloxacin, chloramphenicol, ceftriaxone, cotrimoxazole, and clarithromycin. This is consistent with reports from Gondar University, Ethiopia [24, 25, 38]. Increased resistance was observed in our findings for amoxicillin-clavulanic, amoxicillin, and ampicillin which were supported by studies reported from Ethiopia, Arba Minch, Jimma, and Bahir Dar [14,12,23,35]. This might be due to misuse or inappropriate use of antibiotics and the use of clinical diagnosis for treatment by physicians may lead to emerging of drug-resistant bacterial strains and replacement of sensitive strains by resistant strains.

The prevalence of multidrug resistance towards *Salmonella* and *Shigella* were observed. Out of all the isolates, 54.54% *Salmonella* and 90.9% *Shigella* species were resistant to at least three antimicrobials. One isolate of *Shigella* was resistant to six classes of antimicrobial agents. This study is supported by a study conducted in Ethiopia Butajira [25], Addis Ababa [22], Haramaya University [33], and Gondar [24]. This increased multidrug resistance might be due to genetic variation by mutations, irrational use of antimicrobials, and less hygienic practice of the food handlers. *Salmonella* and *Shigella* species are becoming resistant to most antimicrobials, indicates that there might be easy availability, irrational use common antimicrobials from different governmental and private pharmacies.

Limitation

Fingernail content examination could not be identified. This may be supportive to know either the contamination is due to poor fingernail hygiene or poor food handling practices. Despite this limitation, the methods used to isolate and characterize the antimicrobial susceptibility pattern of *Salmonella* and *Shigella* spp. are comprehensive. In the current study, because of the self-reported nature of the study, recall bias was a limitation but it is not reflected in the findings. We also do not know what biases have the greatest impact on self-reports. In addition to that, even though this study used a random sampling technique to select study participant's it was facility-based. Then Hosmer-Lemeshow test is used for overall calibration error but not used for a particular lack of fitness, so it does not properly take overfitting into account. Therefore generalizability might be hardly achieved. Additionally, Since there is variation in seasons, geography, and in the definition of antimicrobial resistance guidelines among different studies and across regions we couldn't infer the external validity.

Conclusion

The overall prevalence of *Salmonella* and *Shigella* in the study area was found to be 7.3% and 3.7% respectively. The *Salmonella* and *Shigella* carriage was significantly associated with not washing hands after touching dirty materials, not washing hands after using the bathroom, untrimmed fingernails. No resistance against chloramphenicol, ceftriaxone, and ciprofloxacin was identified. The majority of *Salmonella* and *Shigella* were multidrug-resistant. Regular medical checkups improved personal hygiene, and environmental sanitation should be practiced. Consistent training on food preparation and handling for the food handlers of Adigrat University is very important to prevent the risk of infection for the university community having close contact with those carriers.

Additionally, this study suggested that physicians should prescribe based on the laboratory result. drug dispensing by different governmental and private pharmacies should be according to the prescription of the physicians

Abbreviation

ATCC: American Type Culture Collection; **CI:** confidence interval; **CLSI:** Clinical and Laboratory Standards Institute; **MDR:** Multi-drug resistant Multi-Drug Resistant; **SOPs:** Standard Operating Procedures **USA:** United States of America; **WHO:** World health organization;

Declarations

Ethics approval and consent to participate

The study was approved by the College of medicine and health sciences Research ethical review committee of Adigrat University, Ethiopia (Consent Ref Number AGU/CMHS/044/2018 approval dated 07/04/2018 Official letter was obtained from Adigrat University (Consent Ref Number AGU/CMHS/RCSH/19/2018 approval dated 25/04/2018. Written informed consent was sought from each study participant before sample collection and maintained throughout the study. All participants were given code numbers to keep their identity confidential.

Consent for publication

Not applicable

Availability of data and materials

All data collected and analyzed during this study were included in the manuscript. But if the full paper is needed, it will be shared upon request by the editor from the corresponding author.

Competing interests

The authors' declared that there were no competing interests

Funding

Not applicable

Authors' contributions

HL designed the study, collection, analysis, and interpretation of data, and drafted the manuscript. TK, AG, BB, and BF designed the study, supervised data collection both on-field and in the laboratory, and prepared the manuscript. ST, GG, HN, and GA read and approved the final manuscript.

Acknowledgments

The authors gratefully acknowledge the food handlers of Adigrat University for their participation in the study. We also wish to extend our deep appreciation to Adigrat University, College of Medicine and Health Sciences for providing us with the opportunity to do this research and allowing the finance.

References

1. Centers for Disease Control and Prevention (CDC). Surveillance for foodborne disease outbreaks-United States, 2008. *MMWR* 2010; **59**(31):1277-1280.
2. World Health Organization (WHO). (2007). Fact Sheet Number 237: Food safety and foodborne illness. Geneva, Switzerland: World Health Organization. Available at <http://www.who.int/media/centre/factsheets/fs237/en/>. accessed June 26, 2017
3. Coburn, G. A. Grassl, and B. B. Finlay, "Salmonella, the host and disease: a brief review," *Immunology and Cell Biology* 2007;**85**:2:112-118

4. Majowicz SE, Musto J, Scallan E, Angulo FJ, O'Brien SJ, Jones TF, The global burden of nontyphoidal Salmonella gastroenteritis. *Clin Infect Dis.* 2010; **50**:882-9
5. Centers for Disease Control and Prevention. Surveillance for foodborne disease outbreaks the United States, 2008. *MMWR Morb Mortal Wkly Rep* 2011; **60**(35):1197- 1202
6. World Health Organization (WHO) laboratory identification and antimicrobial susceptibility testing of bacterial pathogens of public health concern in the developing world Geneva, Switzerland: World Health Organization; 2003
7. Smith SI, Fowora M.A, Goodluck H.A, Nwaokorie F.O, Aboaba O.O, Opere B. Molecular typing of *Salmonella* spp isolated from food handlers and animals in Nigeria *Int J Mol Epidemiol Genet* 2011; **2**(1):73-77
8. Feasey NA, Dougan G, Kingsley RA, Heyderman RS, and Gordona MA. Invasive non-typhoidal salmonella disease: an emerging and neglected tropical disease in Africa. *Lancet* 2012;**379**(9835): 2489-2499
9. Pala K, Ozakin C, Akis N, Sinirtas M, Gediko S, Aytekin H: Asymptomatic carriage of bacteria in food workers in Nilüfer district, Bursa, Turkey. *Turk J Med Sci* 2010; **40**(1):133-139.
10. Khurana S, Taneja N, Thapar R, Sharma M, Malla N: Intestinal bacterial and parasitic infections among food handlers in a tertiary care hospital of North India. *Trop Gastroenterol* 2008; **29**:207-209.
11. Ao TT, Feasey NA, Gordon MA, Keddy KH, Angulo FJ, Crump JA. Global Burden of Invasive Nontyphoidal *Salmonella* Disease *Infect Dis* 2015 ;**21**(6):441-449 DOI: <http://dx.doi.org/10.3201/eid2106.140999>
12. Beyene G, Tasew H. Prevalence of intestinal parasite, Shigella and Salmonella species among diarrheal children in Jimma health center, Jimma southwest Ethiopia: a cross-sectional study. *Ann Clin Microbiol Antimicrob.* 2014; **13**: Published 2014 Feb 5. doi:10.1186/1476-0711-13-10
13. Mokhtari W, Nsaibia S, Majouri D, Ben Hassen A, Gharbi A, and Aouni M, Detection and characterization of Shigella species isolated from food and human stool samples in Nabeul, Tunisia, by molecular methods and culture techniques *J Applied Microbiol* 2012;**113**: 209-222 doi: 10.1111/j.1365-2672.2012.05324
14. Mama M, Getaneh A. Prevalence, antimicrobial susceptibility patterns and associated risk factors of *Shigella* and *Salmonella* among food handlers in Arba Minch University, South Ethiopia. *BMC Infect Dis.* 2016; **16**:686
15. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. Foodborne illness acquired in the United States-major pathogens. *Emerg Infect Dis.* 2011;**17**(1):7-15
16. Bonkougou IJO, Haukka K, Österblad M, Hakanen AJ, Traoré AS, Barro N, et al. Bacterial and viral etiology of childhood diarrhoea in Ouagadougou, Burkina Faso. *BMC Pediat.* 2013; **13**(36):1-6
17. Afeworki G, Lirneh Y. Multiple drug resistance within Shigella serogroups. *Ethiop Med J.* 1980; **18**:7-11.
18. Roma B, Worku S, T/Mariam S, Langeland N. Antimicrobial susceptibility pattern of Shigella isolates in Awassa. *Ethiop J of Health Dev.* 2000; **14**:149-54.
19. Beyene G., Asrat D., Mengistu Y., Aseffa A. and Wain J. Typhoid fever in Ethiopia. *J Infect Developing Countries* 2008; **2**:448-453
20. Clinical and Laboratory Standards Institute (CLSI): Performance standards for antimicrobial susceptibility testing; Twenty-sixth Informational Supplement. *Wayne PA* 19087 USA 2016; **36**(1).
21. Mobolaji OA, Olubunmi OF. Assessment of the hygienic practices and the incidence of enteric bacteria in food handlers in small businesses in an urban area in Abeokuta. *Int J Microbiol Res.* 2014;**5**(3):41-9
22. Aklilu A, Kahase D, Dessalegn M, Tarekegn N, Gebremichael S, Zenebe S. et al. Prevalence of intestinal parasites, salmonella and shigella among apparently health food handlers of Addis Ababa University student's cafeteria, Addis Ababa, Ethiopia. *BMC Res Notes* 2015; **8**; 17:1-6
23. Abera B, Biadegelgen F, Bezabih B. Prevalence of Salmonella typhi and intestinal parasites among food handlers in Bahir Dar Town, Northwest Ethiopia. *Ethiop J Health Dev.* 2010;**24**(1):46-50
24. Garedew-Kifelew L, Wondafrash N, Feleke A. Identification of drug-resistant Salmonella from food handlers at the University of Gondar, Ethiopia. *BMC Res Notes.* 2014;**7**(1):545
25. Mengistu G, Mulugeta G, Lema T, Aseffa A. Prevalence and Antimicrobial Susceptibility Patterns of *Salmonella serovars* and *Shigella* *J Microb Biochem Technol.*2014: S2: 006. doi:10.4172/1948-5948.S2-006

26. Ifeadike C, Ironkwe O, Adogu P, Nnebue C, Emelumadu O, Nwabueze S, Ubajaka C. Prevalence and pattern of bacteria and intestinal parasites among food handlers in the Federal Capital Territory of Nigeria. *Niger Med J.* 2012;**53**(3):166
27. Andargie G, Kassu A, Moges F, Tiruneh M, Huruy K: Prevalence of bacteria and intestinal parasites among food handlers in Gondar Town, Northwest Ethiopia. *J Health Popul Nutri* 2008;**26**(4):451-455.
28. Nigusse D and Kumie A. Food hygiene practices and prevalence of intestinal parasites among food handlers working in Mekelle university student's cafeteria, Mekelle. *Global Adv Res J of Soc Sci (GARJSS)*,2012;**1**(4):065-071
29. Dagne M, Tiruneh M, Moges F, Gizachew M. Bacterial Profile and Antimicrobial Susceptibility Pattern among Food Handlers at Gondar University Cafeteria, Northwest Ethiopia. *J Infect Dis Ther.*2013;**1**: 105. doi:10.4172/2332-0877.1000105
30. Abera B, Yitayew G, Amare H. *Salmonella* serotype Typhi, *Shigella*, and intestinal parasites among food handlers at Bahir Dar University, Ethiopia *J Infect Dev Ctries* 2016; **10**(2):121-126
31. Tefera T, Mebrie G. Prevalence and Predictors of Intestinal Parasites among Food Handlers in Yebu Town, Southwest Ethiopia. *PLoS ONE*.2014; **9** (10): e110621. doi:10.1371/journal.pone.0110621
32. Zain MM, Naing NN. Sociodemographic characteristics of food handlers and their knowledge, attitude and practice towards food sanitation: A preliminary report. *Southeast Asian J Trop Med Public Health* 2002; **33**(2): 410-7.
33. Marami D, Hailu K, and Tolera M. Prevalence and antimicrobial susceptibility pattern of *Salmonella* and *Shigella* species among asymptomatic food handlers working in Haramaya University cafeterias, Eastern Ethiopia. *BMC Res Notes.*2018; **11**:74
34. Mache A. *Salmonella* serogroups and their antibiotic resistance patterns isolated from diarrhoeal stools of pediatric out-patients in Jimma Hospital and Jimma Health Center, South West Ethiopia. *Ethiop J Health Sci.*2002; **37**: 37-45
35. Reda AA, Seyoum B, Yimam J, Andualem G, Fiseha S, Vandeweerd J-M. Antibiotic susceptibility patterns of *Salmonella* and *Shigella* isolates in Harar, Eastern, Ethiopia. *J Infect Dis Immun.* 2011; **3**: 134-139
36. Asrat D. *Shigella* and *Salmonella* serogroups and their antibiotic susceptibility patterns in Ethiopia. *East Mediterr Heal J.* 2008;**14**:760-7
37. Yismaw O, Negeri C, Kassu A. A five-year antimicrobial resistance pattern observed in *Shigella* species isolated from stool samples in Gondar University Hospital, northwest Ethiopia. *Ethiop J Heal Dev.* 2006;**20**:194-8
38. Tiruneh M Serodiversity and antimicrobial resistance pattern of *Shigella* isolates at Gondar University teaching hospital, Northwest Ethiopia. *Jpn J Infect Dis.* 2009; **62**(2):93-7

Tables

Table 1

Demographic characteristics	Characteristics	Frequency	Percent
Sex	Male	37	12.3
	Female	264	87.7
Age	≤ 20	54	17.9
	21-40	247	80.1
Marital status	Single	127	42.2
	Married	174	57.8
Educational level	Secondary school	187	62.12
	Higher than secondary school	114	37.9
Years of service	≤1	36	12.0
	1-2	42	14.0
	>2	223	74.1
Certified in food preparation and handling	No	264	87.7
	Yes	37	12.3
Medical check-up	No	36	12.0
	Yes	265	88.0

Table 2

Variables		Growth of <i>Salmonella</i> and <i>Shigella</i>		COR (95% CI)	P-value
		Positive N (%)	Negative N (%)		
Sex	Male	1(2.7)	36(97.3)	0.22(0.03-1.58)	0.20
	Female	32(12.1)	232(87.9)	1	
Age	≤ 20	2(3.7)	52(96.3)	0.3(0.07-1.2)	0.081
	21-40	31(12.55)	216(87.45)	1	
Marital status	Single	16(12.6)	111(87.4)	1.29(.68-2.45)	0.341
	Married	17(9.78)	157(90.23)	1	
Educational status	Lower than Secondary school	16(8.56)	171(91.44)		0.507
	Higher than secondary school	17(14.9)	97(85.1)	1.74(0.92-3.31)	0.862
Hand washing after using the bathroom	Yes with water and soap	1(0.9)	112(99.1)	1	
	Yes only with water	17(13.6)	108(86.4)	17.630(2.306-134.7)	0.006
	No	15(23.8)	48(76.2)	35.000(4.495-272.49)	0.001
Hand washing after touching dirty materials	Yes with water and soap	5(3.4)	141(96.6)	1	
	Yes only with water	14(11.0)	113(89.0)	3.494(1.222-9.991)	0.020
	No	14(50.0)	14(50.0)	28.200(8.845-89.906)	0.001
Handwashing before food handling	Yes with water and soap	2(1.8)	112(98.2)	1	
	Yes only with water	13(9.2)	129(90.8)	5.643(1.247-25.548)	0.025
	No	18(40.0)	27(60.0)	37.333(8.164-170.714)	0.001
Fingernail status	Trimmed	4(2.2)	174(97.8)	1	
	Untrimmed	29(23.6)	94(76.4)	10.490(3.780-29.09)	0.001
After blowing nose	Yes with water and soap	7(8.0)	81(92.0)	1	
	Yes only with water	11(11.7)	83(88.3)	0.919(0.401-2.107)	0.842
	No	15(12.6)	104(87.4)	0.599(0.233-1.538)	0.287
Touch food with bare hands	No	16(14.0)	98(86.0)	1	
	Yes	17(9.1)	170(90.9)	1.597(0.772-3.302)	0.207
Years of service	≤1	9(25.0)	27(75.0)	3.72(1.76-78.50)	0.001
	1-2	9(21.4)	33(78.6)	0.818(0.285-	0.709

				2.349)	
	>2	15(6.7)	208(93.3)	1	
Certified in food preparation and handling	No	29(11.5)	224(88.5)	1	
	Yes	4(8.3)	44(91.7)	0.702(0.235-2.097)	0.527
Medical check-up	No	10(18.5)	44(81.5)	1	
	Yes	23(9.3)	224(90.7)	1.99(1.01-3.93)	0.04

Key: ^a(COR=Crude odds ratio); ^b(CI=Confidence interval);1(referent).

Variables	Growth of <i>Salmonella</i> and <i>Shigella</i>		COR ^a (95% CI ^b)	P-value	AOR ^c (95%CI)	P-value
	Positive N (%)	Negative N (%)				
Hand washing using after using the bathroom						
Yes with water and soap	1(0.9)	112(99.1)	1		1	
Yes only with water	17(13.6)	108(86.4)	17.630(2.306,134.7)	0.006	23.239(2.125-254.17)	0.010*
No	15(23.8)	48(76.2)	35.000(4.495,272.49)	0.001	62.917(5.11-77.34)	0.001*
Hand washing after touching dirty materials						
Yes with water and soap	5(3.4)	141(96.6)	1		1	
Yes only with water	14(11.0)	113(89.0)	3.494(1.222, 9.991)	0.020	1.089(0.187-6.345)	0.924
No	14(50.0)	14(50.0)	28.200(8.845, 89.906)	0.000	37.189(5.658-244.45)	0.001*
Hand washing before food handling						
Yes with water and soap	2(1.8)	112(98.2)	1		1	
Yes only with water	13(9.2)	129(90.8)	5.643(1.247, 25.548)	0.025	5.972(0.899-39.677)	0.064
No	18(40.0)	27(60.0)	37.333(8.164, 170.714)	0.000	33.065(4.958-220.52)	0.001*
Finger nail status						
Trimmed	4(2.2)	174(97.8)	1		1	
Untrimmed	29(23.6)	94(76.4)	10.490(3.780, 29.09)	0.000	13.973(3.404-57.362)	0.001*

Table 3

Key: ^a(COR=Crude odds ratio); ^b(CI=Confidence interval); ^c(AOR=Adjusted odds ratio); 1(referent).

Table 4

Isolates(n)	Sensitivity pattern n(%)	Antimicrobial agents N (%)								
		GM	AML	AMP	CIP	CLR	TS	AMC	CHL	CRO
Salmonella n(22)	S	22(100)	1(4.5)	0(0.00)	22(100)	18(81.8)	20(90.9)	13(59.1)	22(100)	22(100)
	R	0(0.00)	21(95.5)	22(100)	0(0.00)	4(18.2)	2(9.1)	9(40.9)	0(0.00)	0(0.00)
Shigella n (11)	S	10(90.9)	0(0.00)	0(0.00)	11(100)	2(18.2)	10(90.9)	6(54.5)	9(81.2)	10(90.9)
	R	1(9.1)	11(100)	11(100)	0(0.00)	9(81.2)	1(9.1)	5(45.5)	2(18.2)	1(9.1)

Key: S = Sensitive R = Resistant, GM Gentamicin, AML Amoxicillin, AMP Ampicillin, CIP=Ciprofloxacin, CLR Clarithromycin, TS Cotrimoxazole, AMC Amoxicillin-clavulanic acid CHL Chloramphenicol, CRO Ceftriaxone

Table 5

		Resistant isolates no. (%)	
	Antimicrobials	<i>Salmonella</i>	<i>Shigella</i>
For three	AML,AMP,CLR	2(16.67)	4(40)
	AML,AMP,CHL	-	1(10)
	AML,CLR,AMC	7(58.34)	-
	AMP,CLR,AMC	1(8.33)	-
	AML,AMP,TS	1(8.33)	-
FOR FOUR	AML,AMP,CLR,AMC	1(8.33)	3(30)
FOR FIVE	AML,AMP,CLR,AMC,CHL	-	1(10)
FOR SIX	GM,AML,AMP,CLR,TS,AMC	-	1(10)
		12(100)	10(100)

Key: GM Gentamicin, AML Amoxicillin, AMP Ampicillin, CIP=Ciprofloxacin, CLR Clarithromycin, TS Cotrimoxazole, AMC Amoxicillin-clavulanic acid CHL Chloramphenicol, CRO, Ceftriaxone

MDR multidrug resistant; MDR definition for *Salmonella* and *Shigella* percent is computed from total number of *Salmonella* and *Shigella*