Transcriptomic analysis reveals the role of *Oryza sativa* FOUR LIPS in response to salt stress in rice

Chunxia Zhang (✉ spring1984@163.com)

CAS Institute of Botany: Institute of Botany Chinese Academy of Sciences <https://orcid.org/0000-0001-8525-8868>

Jie Zhang

Institute of Botany Chinese Academy of Sciences

Huichao Liu

Institute of Botany Chinese Academy of Sciences

Xiaoxiao Qu

Institute of Botany Chinese Academy of Sciences

Junxue Wang

Institute of Botany Chinese Academy of Sciences

Qixiumei He

Institute of Botany Chinese Academy of Sciences

Junjie Zou

Institute of Botany Chinese Academy of Sciences

Kezhen Yang

Institute of Botany Chinese Journal of Plant Ecology: Institute of Botany Chinese Academy of Sciences

Jie Le

Institute of Botany Chinese Academy of Sciences <https://orcid.org/0000-0001-8854-3031>

Research Article

Keywords: RNA-seq, ChIP-seq, OsFLP, OsCDKB1;1, Rice, Salt stress

Posted Date: February 23rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1354294/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Title: Transcriptomic analysis reveals the role of *Oryza sativa* FOUR LIPS in response to salt stress in rice

Authors:

Chunxia Zhang^{1*}, Jie Zhang^{1,2*}, Huichao Liu^{1,2}, Xiaoxiao Qu^{1,2}, Junxue Wang^{1,2,§}, Qixiumei He^{1,2}, Junjie Zou^{1,3}, Kezhen Yang¹ and Jie Le^{1,2†}

Addresses:

¹Key Laboratory of Plant Molecular Physiology, CAS Center for Excellence in Molecular Plant Sciences, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China

²University of Chinese Academy of Sciences, Beijing 100049, China

³Biotechnology Research Institute, Chinese Academy of Agricultural Sciences, Beijing 100081, China

[†]Correspondence: Jie Le (lejie@ibcas.ac.cn), Tel: +86 10 6283 6673

Orcid-ID: Jie Le, <https://orcid.org/0000-0001-8854-3031>

*These authors contributed equally to this work.

Present address:

[§]Wenbo School, Jinan 250100, China

Author e-mail addresses:

Chunxia Zhang: spring1984@163.com

Jie Zhang: jieer1027@163.com

Huichao Liu: liuhuichao@ibcas.ac.cn

Xiaoxiao Qu: quxiaoxiao721@sina.com

Junxue Wang: wangjunxues@163.com

Qixiumei He: heqixiumei@ibcas.ac.cn

Junjie Zou: zoujunjie@caas.cn

Kezhen Yang: ykzdsp@163.com

Jie Le: lejie@ibcas.ac.cn

Key message: An R2R3-MYB transcription factor *Oryza sativa* FOUR LIPS associated with *Oryza sativa* B-type Cyclin-Dependent Kinase 1;1 confers salt tolerance in rice

Keywords: RNA-seq, ChIP-seq, *OsFLP*, *OsCDKB1;1*, Rice, Salt stress

Abstract

The Arabidopsis FOUR LIPS (AtFLP), an R2R3 MYB transcription factor, acts as an important stomatal development regulator. Only one orthologue protein of AtFLP, *Oryza sativa* FLP (OsFLP), was identified in rice. However, the function of OsFLP is largely unknown. In this study, we conducted RNA-seq and ChIP-seq to investigate the potential role of OsFLP in rice. Our results reveal that OsFLP is probably a multiple functional regulator involved in many biological processes in growth development and stress responses in rice. However, we mainly focus on the role of OsFLP in salt stress response. Consistently, phenotypic analysis under salt stress conditions showed that *Osflp* exhibited significant sensitivity to salt stress, while *OsFLP* over-expression lines displayed obvious salt tolerance. Additionally, Yeast one-hybrid assay and electrophoretic mobility shift assay (EMSA) showed that OsFLP directly bond to the promoter region of *Oryza sativa* B-type Cyclin-Dependent Kinase 1;1 (*OsCDKB1;1*), and the expression of *OsCDKB1;1* was repressed in *Osflp*. Disturbing the expression of *OsCDKB1;1* remarkably enhanced the tolerance to salt stresses. Taken together, our findings reveal a crucial function of OsFLP regulating *OsCDKB1;1* in salt tolerance and largely extend the knowledge about the role of OsFLP in rice.

Introduction

MYB (myeloblastosis) transcription factor genes family have a wide function in plant adaptive responses to adverse environment and growth development (El-Kereamy et al. 2012; Lv et al. 2017; Yang et al. 2012; Ye et al., 2015). MYB genes family are classified into four groups, including 1R-MYB, 2R-MYB, 3R-MYB, and 4R-MYB. Among them, R2R3-MYB genes are more prevalent in plants compared to those in animals (Martin and Paz-Ares 1997; Dubos et al. 2010). In Arabidopsis, FOUR LIPS (AtFLP) is an atypical R2R3-MYB transcription factor as a regulator in stomatal development (Yang and Sack 1995; Lai et al. 2005). Besides this, AtFLP/AtMYB88 play important role in abiotic stress responses (Xie et al. 2010b), root gravitropism by regulating the transcription of *PIN* genes (Wang et al. 2015), as well as regulating female reproductive development (Makkena et al. 2012). In addition, AtFLP/AtMYB88 are widely expressed in various tissues (Lei et al. 2015). These reports indicate that AtFLP serves as a pleiotropic regulator participating in several aspects of biological processes in Arabidopsis. Recently, the

homolog of AtFLP in rice OsFLP was knocked out via the genome editing system. In contrast to the role in Arabidopsis, OsFLP is required for the proper orientation of symmetric division of guard mother cells in rice (Wu et al. 2019). However, little is known about the function of OsFLP and the downstream targets.

Salinity stress as a crucial environmental factor affects plant growth and development and leads to reduce crop yield (Kreps et al. 2002; Gupta and Huang 2014; Roy et al. 2014). To adapt to unfavorable environmental conditions, plants have evolved molecular mechanisms to protect themselves against stresses (Matsui et al. 2008). Thus, unraveling the salt stress-responsive mechanisms will be beneficial for improving plant salinity tolerance and subsequently increasing crop production to ensure food security for the rapid growth of the global population. Salinity stress causes several aspects of detrimental effects on plants. One of the most severe effects is the accumulation of Na^+ and Cl^- , which leads to ion imbalance and inhibits the uptake of K^+ resulting in reduced productivity and even death (James et al. 2011). Hence, maintaining Na^+/K^+ homeostasis in cells is vital for plants to survive under salt stress conditions. On the other hand, the production of ROS is enhanced and further leads to oxidative damages to the membranes, lipids, proteins (Mittler 2002). Many enzyme proteins such as superoxide dismutase (SOD), peroxidase (POD), catalase (CAT), glutathione peroxidase (GPX), ascorbate peroxidase (APX), are involved in ROS scavenging, and the activity of these proteins positively correlates with salt tolerance in plants (Mäkelä et al. 1999; Xia et al. 2014).

Besides *AtFLP/AtMYB88*, several genes have functions not only in controlling stomatal development but also in response to abiotic stress. For example, HIC, a negative regulator of stomatal development, controls stomatal density in response to elevated CO_2 (Gray et al. 2000). The cold-responsive gene *ICE1* participates in stomatal differentiation (Kanaoka et al. 2008). The *Arabidopsis* receptor-like kinase ERECTA (ER) controls stomatal patterning (Shpak et al. 2005). Over-expression of ER remarkably increased heat tolerance in plants (Shen et al. 2015). Hence, these data implicate the cross-talk of the signal transduction pathways between stomatal development and abiotic stress.

Cyclin-dependent kinases (CDKs) mainly play roles in regulating cell division in plants (De Veylder et al. 2003; Dewitte and Murray 2003). In Arabidopsis, 12 CDKs have been identified with six categories (CDKA to CDKF) (Vandepoele et al. 2002) (28). B-type CDKs including B1-type (CDKB1;1 and CDKB1;2) and B2-type (CDKB2;1 and CDKB2;2) subgroups are unique in plants (Vandepoele et al. 2002; Boudolf et al. 2001). In Arabidopsis, CDKB1;1 and CDKB1;2 act redundantly in regulating the

symmetric division of Guard Mother Cells (GMCs). In addition, FLP/MYB88 could directly bind to the promoter region of *CDKB1;1* to regulate its expression during stomatal development (Xie et al. 2010a). Moreover, CDKB1;1 together with the RPA2a, which is a subunit of replication protein A complexes, restricts stomatal proliferation by CDKB1;1-mediated phosphorylation (Yang et al. 2019). *OsCDKB1;1* could restore the impaired GMC division and stomatal production in *Atcdkb1;1 1;2* mutants (Qu et al. 2018). In addition, salt stress affects the cell cycle regulation at the transcriptional level, ultimately leading to adaptive growth responses (BursSENS et al. 2000). Salt stress responses involve the decline of cell cycle activity in the meristem of the root (West et al. 2004). However, the involvement of *OsCDKB1;1* in response to abiotic stress, especially salt stress, is still unknown yet.

In this study, we isolated an *Osflp* mutant and generated the complementary transgenic lines. Using these materials, we performed the integrative analysis of RNA-seq and ChIP-seq. We found that the potential role of OsFLP was involved in several aspects of biological processes, especially in salt stress responses. *Osflp* was more sensitive to salt stresses with more ROS accumulation in plants compared with wild-type ZH11. Additionally, OsFLP could directly target the promoter region of *OsCDKB1;1* *in vitro* and repress its transcriptional expression. Consistently, *OsCDKB1;1*-RNAi plants exhibited significant salt tolerance with less ROS accumulation. Collectively, we conclude that upon salt stresses, OsFLP represses the expression of *OsCDKB1;1* to confer salt tolerance in rice. These findings provide new functions of *OsCDKB1;1* out of cell cycle and novel insights into the role of OsFLP in rice.

Materials and Methods

Plant Materials and growth conditions

Rice plants, *Oryza sativa* L. ssp. *Japonica* cv. Zhonghua 11 (ZH11, wild type), were cultivated in the field of the Institute of Botany, Chinese Academy of Sciences, Beijing for propagating seeds.

Plant growth chambers were also used for rice cultivating with the following setting: 12 h day at 30 °C ± 2 °C, 12 h night at 22 °C ± 2 °C, and 60% humidity.

Osflp mutant was isolated by TILLING (Targeting Induced Local Lesions IN Genomes) through an in-house TILLING platform (www.croptilling.org), which had a G→A point mutant in the sixth exon at the 2126 bp position downstream of ATG position which led to the substitution of Ala with Thr.

To generate *OsFLP* and *OsCDKB1;1* overexpression lines, the full-length cDNA fragments were amplified by PCR and were inserted into TOPO vector, afterward were sub-cloned into pH7WG2D.1

vector via LR reaction of gateway cloning with 35S promoter. To obtain the *Osflp* complementation line, 4733bp genomic fragment of *OsFLP* was amplified and was sub-cloned into the destination vector pCAMBIA1301. For making the GUS line, the promoter region of *OsFLP* was fused to the vector pGUS1301. These constructs were introduced into rice via an *Agrobacterium*-mediated transformation (Hiei et al. 1994). The homozygous lines were selected for further analysis.

Germination rates assay under salt stress

Salt tolerance phenotype assay on half-strength Murashige & Skoog medium (MS) solid medium plates. Three-day-old rice seedlings grown on 1/2 MS solid medium vertical plates were transferred to new 1/2 MS solid medium vertical plates supplemented with or without 200 mM NaCl, 5 days later taking pictures and performing measurements.

Salt tolerance phenotype assay using hydroponic culture. 14-day-old rice seedlings cultivated with 1/2 MS hydroponic culture were transferred to liquid medium supplemented with or without 150 mM NaCl for 3 days or 10 days, changing liquid medium every 3 days, afterward transferred to fresh liquid medium for 3 days recovery.

Salt tolerance phenotype assay with rice seedlings grown on soil. 14-day-old rice seedlings grown on 1/2 MS solid medium were transferred to soil growing for 2 weeks and stopped watering for one week, afterward watering plants with tap water supplemented with or without 200 mM NaCl for 15 days and 21 days.

Survival rates and soluble sugar content measurement.

Survival rates were calculated after 21 days of salt stress treatment. No green shoots were counted as dead plants and the survival rate was shown as the percentage of alive seedlings.

Soluble sugar content was determined referring to the previous report (Tang et al. 2013). 500 mg leaves were ground in 2 mL 10% trichloroacetic acid (TCA), and the mixture was centrifuged at 4000 rpm for 10 min. 2 mL extracts were taken and added 2 ml 0.6% thiobarbituric acid (TBA). Reacted in boiled water for 15 min and quickly cooled on ice, afterward centrifuged at 4000 rpm for 15 min. The spectral absorbance of the supernatant was determined at 450 nm, 532 nm, and 600 nm with an ultraviolet spectrophotometer. Soluble concentration (mM)= $11.71 \times D_{450}$.

SOD, POD, and CAT enzyme activities assays

500 mg rice leaves were ground with pre-cooled mortar and pestle in liquid nitrogen. Afterward, 50 mM

potassium phosphate buffer (pH 7.8) containing 1% polyvinylpyrrolidone was added and mixed thoroughly. Then the mixture was centrifuged at 15000 g at 4 °C for 20 min. The supernatant was the crude enzyme extraction. The activities of SOD, POD, and CAT were determined following the previously described protocols (Miao et al. 2010). The final enzyme activities were calculated according to the following formulations. SOD activity (U/g·min) = $(A_0 - A_s) \times V_t \div (A_0 \times W \times 0.5 \times V_s)$, (A_0 , absorbance value of the control; A_s , absorbance value of the samples; V_t , the total volume of the sample; W , the fresh weight of the sample; V_s , the volume of the sample for measurement). POD activity (U/g·min) = $(\Delta A_{470} \times V_t) \div (W \times V_s \times 0.01 \times t)$, (ΔA_{470} , the change of absorbance value at 470nm wavelength per minute; V_t , the total volume of the sample; W , the fresh weight of the sample; V_s , the volume of the sample for measurement; t , interval time). CAT activity (U/g·min) = $(\Delta A_{240} \times V_t) \div (W \times V_s \times 0.1 \times t)$, ($\Delta A_{240} = A_{S0} - (A_{S1} + A_{S2}) / 2$, (ΔA_{240} , the change of absorbance value at 240 nm wavelength per minute; V_t , the total volume of the sample; W , the fresh weight of the sample; V_s , the volume of the sample for measurement; t , interval time; A_{S0} , absorbance value of the control; A_{S1} and A_{S2} , absorbance value of the samples).

Na⁺ and K⁺ content assay

Shoots and roots of rice seedlings were harvested separately. Samples were dried at 104 °C for 15 min and 55 °C for 3 days. Dry samples were digested in nitric acid overnight in the dark and then transferred to 120 °C for 8-12 h. The content of Na⁺ and K⁺ in the extractions was measured with an ICP emission spectrometer (iCAP6300, Thermo scientific, US) according to the methods described previously (Shen et al. 2015b).

ROS content determination

The content of H₂O₂ was determined referring to the protocol described previously (53). The commercial H₂O₂ detection kit (SolarBio Inc.) was used. In brief, approximately 100 mg rice leaves were ground with liquid nitrogen and 1 ml lysate buffer was added and mixed gently. The extraction was centrifuged at 8000 g for 10 min at 4 °C. The reaction buffer was added to the supernatant. Then mixed them gently, and reacted for 5 min at room temperature. The mixture was measured with an ultraviolet spectrophotometer at the 415 nm wavelength, and the H₂O₂ content was calculated according to the formulation in the manufacturer's protocol. We also used 3,3-diaminobenzidine (DAB) staining for the detection of H₂O₂ and nitroblue tetrazolium (NBT) staining to determine O₂^{·-} according to the previously

described protocol (Zhang *et al.*, 2014a). In brief, for the DAB staining assay, the rice leaves were cut into around 1cm length, immersed in 1 mg/mL DAB solution, incubated for 8 h at room temperature. Afterward, samples were transferred into ethanol to remove the chlorophyll heating with a water bath at 100 °C for 2 h. For NBT staining assay, around 1 cm length of rice leaves was immersed in 6 mM NBT solution and incubated for 8 hours at room temperature. Ethanol was used to remove the chlorophyll and pictures were taken for analysis.

Yeast one-hybrid assay

In the yeast one-hybrid assay, the activated domain AD vector and the LacZ reporter BD vector were used. *OsFLP* was fused to the AD vector (*OsFLP*-AD). The 2266 bp promoter of *OsCDKB1;1* was truncated into nine fragments (-2266 bp to -1132 bp, -2266 bp to -1683 bp, -1701 bp to -1132 bp, -1208 bp to -537 bp, -632 bp to -1 bp, -536 bp to -413 bp, -418 bp to -275 bp, -274 bp to -138 bp, -137 bp to -1 bp), and fused with LacZ reporter BD vector, respectively. Afterward, the fused AD and BD vectors were co-transformed into the EGY48 yeast strain and performed X-gal staining.

RNA-sequencing, ChIP-sequencing, and RT-qPCR analysis

For RNA-sequencing analysis, ZH11 and *Osflp* mutant plants were grown under 30 °C ± 2 °C, 12 h day / 22 °C ± 2 °C, 12 h night conditions for 7 days. The shoots of the seedlings were harvested and frozen in liquid nitrogen. RNA was extracted with TRIzol reagent (Invitrogen). The libraries were sequenced based on the Illumina platform and were performed by BioMarker Inc. (Beijing, China). The clean reads were used for the subsequent analysis. Genes with adjusted p-value < 0.05 and log2 fold- change >1 were taken as the differential expressed genes (DEGs). The GO enrichment of DEGs analysis was conducted with AgriGO V2.0 (<http://systemsbiology.cau.edu.cn/agriGOv2/>).

To generate *OsFLP:gOsFLP*-MYC plants, the genomic sequence of *OsFLP* was amplified with the gOsFLP-F primer (ATGGCGACCGGACCGATCT) and gOsFLP-R primer (AAGCTTTTATACCGAGTTCAAGTCT), and sub-cloned into a pSUPER:1300-MYC vector driven with *OsFLP* native promoter. Chromatin immunoprecipitation(ChIP) was performed with 7-day-old rice seedlings. The sequencing was performed by Wuhan Igenebook Biotechnology Co., Ltd (Wuhan, China).

For RT-qPCR analysis, RNA was extracted with HiPure Plant RNA Mini Kit (Angen Biotech CO., Ltd). The first-strand cDNA was synthesized with a PrimeScript® RT Reagent Kit (Takara). Real-time PCR was performed in a 96-well plate with the LightCycler96 system. Each reaction with 20 µl final

volume contained the following ingredients: 10 µl Hieff® qPCR SYBR® Green Master Mix (No Rox), 0.8 µl gene-specific primers (10 µM), 2 µl cDNA which had been diluted 10 times from the original cDNA solution. Three steps amplification program was used as follows: 95 °C for 5 min, 40 cycles with 95 °C for 10 sec, 55 °C for 20 sec, and 72 °C for 20 sec, finally following with melting curve step. *OsACT1* was used as the internal reference gene for normalization. The primers are listed in Table S3.

Electrophoretic mobility shift assays (EMSA)

The coding sequence of *OsFLP* was amplified by PCR inserted into the pET28a vector and transformed into *E. coli* DE3 (BL21) cells for expressing the recombinant protein. The OsFLP protein was purified with His tag. Oligonucleotide probes were synthesized and labeled with Biotin 3' End DNA Labeling Kit (PIERCE). For EMSA assay, 20 µL volume of reactions included the following components: His-OsFLP recombinant protein, labeled probes, 10×binding buffer, 100 mM MgCl₂, 50% (v/v) glycerol, 1 µg/µL poly (dI-dC), and 1% (v/v) Nonidet P-40. Unlabeled probes were added for the competition assay. The reaction was incubated at room temperature for 20 min. The reaction mixtures were analyzed by electrophoresis. Te LightShift Chemiluminescent EMSA kit (Thermo Scientific, 20148) was used to detect the Biotin-labeled probes.

Results

The phylogenetic relationship analysis of FLP in plants

The FLP encodes an R2R3-MYB transcription factor protein (Lai et al. 2005; Wu et al. 2019). To explore the evolutionary relationship of the FLP in different species, including algae, lower plants, and higher plants, we conducted phylogenetic tree analysis with OsFLP amino acids as a probe. The results revealed that the OsFLP orthologous proteins were not only present in higher plants (*Panicum miliaceum*, *Brachypodium distachyon*, *Setaria viridis*, *Setaria italic*, *Zea mays*, *Sorghum bicolor*, *Oryza sativa*, *Arabidopsis thaliana*, *Glycine max*, *Medicago truncatula*, and *Medicago sativa*) with stomata but also in the lower plants (*Zostera marina*) without stomata, even in the single-cell green algae (*Coccomyxa subellipsoidea* and *Chlamydomonas reinhardtii*) (Fig.S1). Thus, these data indicate that besides the function in regulating stomatal development, FLP may be also involved in other functions in plant development or adaptation to unfavorable environmental conditions.

Transcriptomic analysis of the *Osflp* mutant

To facilitate functional analysis of OsFLP, we isolated an *Osflp* mutant line based on TILLING system.

The substitution of Ala with Thr in the sixth exon in *Osflp* mutant led to the abnormal GCs development in the mature leaves. The genomic region of *OsFLP* driven with the native promoter was introduced into the *Osflp*. Several complementation lines were obtained and the abnormal GCs phenotype was rescued in these lines. One complementation line was used in this study and indicated as COM (Supplementary Fig. 1A-E). This phenotype is consistent with the previous report on *Osflp* created by CRISPR (Wu et al. 2019).

To explore the potential functions of OsFLP, we performed the RNA-sequencing (RNA-seq) analysis with the shoots of seven-day-old seedlings of *Osflp* and the wild-type ZH11 under normal growth conditions. In total, we identified 755 differentially expressed genes (DEGs), including 328 up-regulated genes and 427 down-regulated genes in *Osflp* compared with ZH11 (Fig. 2A, Dataset S1). In these DEGs, several types of genes might be regulated by OsFLP including “transcription factors” “transferase proteins” “cell wall-related genes” “environmental stress-responsive genes” “metabolism pathway” and “hormone-related genes”, etc. (Fig. 2B, Dataset S2). These results indicate that OsFLP may be involved in various aspects of plant development and environmental stress responses.

To further unravel the function of OsFLP, we performed Gene Ontology (GO) analysis with up-regulated and down-regulated DEGs, respectively. Intriguingly, GO analysis revealed that GO terms related to peroxidase activity, oxidative stress reactive oxygen species metabolic process, and so on were significantly enriched with down-regulated DEGs other than up-regulated DEGs (Fig. 2C, Dataset S3). Several class III peroxidases (PRXs) were identified according to the GO terms and most of these PRXs were downregulated in *Osflp* (Supplementary Table S1). Furthermore, to confirm the RNA-seq data, the expression of four randomly selected genes was detected by RT-qPCR. The results are consistent with RNA-seq (Supplementary Fig. S2A). Considering PRXs responsible for H₂O₂ metabolism (Almagro et al. 2009), OsFLP may play a role in controlling ROS-related pathways. Thus, the expression of *POD1*, *POD2*, and *CAT*, which is involved in ROS scavenging (Miller et al. 2010; Mittler 2002) (Noctor and Foyer 1998), were detected. The results showed that the expression of three genes significantly decreased in *Osflp* compared with ZH11 (Supplementary Fig. S2B).

To further identify the direct targets of OsFLP, the ChIP-seq analysis was conducted with *pOsFLP::OsFLP-MYC* in *Osflp*. After removal of the background peaks, a total of 12899 peaks were identified (Dataset S4). KEGG pathway analysis showed that the potential direct targets of OsFLP related

to various pathways, including “spliceosome”, “mRNA surveillance pathway”, “N-Glycan biosynthesis”, “endocytosis”, “RNA transport” and “oxidative phosphorylation”, etc. (Supplementary Fig. S3, Dataset S5). Among them, oxidative phosphorylation is closely related to ROS metabolism.

Through the integrative analysis of RNA-seq data and ChIP-seq data, 125 genes were found to be present in both RNA-seq and ChIP-seq profiles (Fig. 2D, Dataset S6). KEGG pathway analysis showed that peroxisome related pathway was included (Supplementary Fig. S4, Dataset S7). Peroxisome mainly controlling ROS scavenging was enriched. This result implied that *OsFLP* may play a role in ROS metabolism.

In addition, 10 genes out of 125 genes were related to salt stress responses, Na⁺ and K⁺ transports, as well as oxidoreductase activity (Supplementary Table S2). In addition, to further confirm the salt stress-responsive function of *OsFLP*, the expression levels of four salt stress-responsive genes including *OsRab16A*, *OsNAC5*, *OsDREB2A*, and *OsDREB1F* (Song et al. 2011; Ponce et al. 2021; Yang et al. 2012), were examined in ZH11, *Osflp*, COM, and OE with or without NaCl stress. With salt stress conditions, the expression of these four genes significantly decreased in *Osflp*, whereas increased in OE, compared with ZH11 and COM in rice (Fig. 2E-H). Taken together, these data show that *OsFLP* affects the expression of salt stress-responsive genes as well as the ROS metabolic genes, suggesting the possible involvement of *OsFLP* in salt stress responses.

***OsFLP* plays a vital role in response to salt stress**

To verify the role of *OsFLP* in response to salt stress, the expression pattern of *OsFLP* under salt stresses was determined. The expression of *OsFLP* was increased after 2 h salt stress treatment and reached the maximal level after 6 h salt stress treatment (Supplementary Fig. S5A). Consistently, the shoots and leaves of the *pOsFLP::GUS* transgenic plants also showed increased GUS activity after 6 h salt stress treatment (Supplementary Fig. S5B).

The increased expression of *OsFLP* induced by salt stress encouraged us to examine the phenotype of *Osflp* under salt stresses. Upon 150 mM NaCl stress, the germination rate of ZH11 and COM were almost 100%, while the germination rate of *Osflp* was much lower, around 53% after 5 days (Supplementary Fig. S6A, B, and C). Furthermore, under 200 mM NaCl stress conditions, compared to ZH11 and COM, *Osflp* had shorter roots and shoots. In contrast, significant increases in shoot and root length were observed in the *OsFLP* over-expression plants (OE) (Fig. 3A – C). To further confirm the

salt responsive function of *OsFLP*, we performed the phenotypic analysis with plants cultivated in soil. Three-week-old seedlings were treated with 200 mM NaCl for 15 days and 21 days, respectively. *Osflp* exhibited more sensitivity with the significantly decreased survival rate (about 18.75%) compared with ZH11 (about 87.5%), and COM plants could completely rescue the salt-sensitive phenotype with around 81.25% of plants survival rate, whereas OE plants behaved more resistance to salt stress with 100% survival rate (Fig. 3D and E). Taken together, these data indicate that *OsFLP* serves as a positive regulator in salt stress response in rice.

***Osflp* accumulates more ROS under salt stress conditions**

Salt stress usually causes ROS accumulation in plants (Xiong et al. 2002; Yang and Guo 2018). A high concentration of ROS leads to oxidative damage to membranes (lipid peroxidation), proteins, RNA, and DNA molecules (Mittler 2002). Several enzymes such as superoxide (SOD), peroxidase (POD), and catalase (CAT) contribute to the ROS detoxification in plants (Mittler et al. 2004). Our transcriptomic analysis showed that many genes, involved in ROS metabolism, were enriched (Fig 2C.). Hence, we investigated if ROS homeostasis was affected in *Osflp* and OE plants under salt stress conditions. Considering hydroponic cultivation is more stable and easier for stress treatment, we performed the following measurements with hydroponic cultivation. Two-week-old seedlings of ZH11, *Osflp*, COM, and OE were treated with or without 150 mM NaCl for 3 days and 10 days, respectively, afterward following 3 days recovery (Fig. 4A). Salt stress generally increased soluble sugar concentration (Dubey and Singh 1999). Thus we performed measurements with survival rate and the soluble sugar content. It showed that the survival rate and the soluble sugar content were significantly lower in *Osflp* but higher in OE plants, compared with ZH11 and COM under salt stresses (Fig. 4B-C). Then, we checked the enzyme activities of SOD, POD, and CAT in ZH11, *Osflp*, OE, and COM. Under non-salt treatment conditions, the activities of these enzymes in *Osflp* and OE were very similar to those in ZH11 and COM. However, upon salt stresses, the activities of these enzymes significantly reduced in *Osflp* but enhanced in OE (Fig. 4D-F). In addition, with salt treatment conditions, more H₂O₂ accumulated in *Osflp* but less in OE (Fig. 4G). Consistently, diaminobenzidine (DAB) and nitro blue tetrazolium (NBT) staining also confirmed the higher contents of ROS in *Osflp* relative to ZH11 and COM (Fig. 4H and I). Collectively, these results suggest that *OsFLP* affects ROS accumulation in response to salt stress.

***OsFLP* confers salt tolerance by reducing sodium accumulation**

Under salt stress conditions, intracellular Na⁺ and K⁺ homeostasis is essential for cell activities and is crucial for salt tolerance in plants (Zhu 2003; Horie et al. 2001). To test if Na⁺ and K⁺ homeostasis was changed in *Osflp* mutant and OE lines with salt stress, we measured Na⁺ and K⁺ contents in both shoots and roots. Under normal growth conditions, no significant differences in Na⁺ and K⁺ contents were observed in both *Osflp* and OE lines compared to ZH11 and COM. However, with 150mM NaCl treatment, Na⁺ contents were significantly higher in both shoots and roots of *Osflp* mutant compared to ZH11 and COM. In contrast, in OE lines, Na⁺ contents were much lower in both shoots and roots (Fig. 5A and D). Whereas, K⁺ contents behaved oppositely (Fig. 5B and E). Thus, with NaCl treatment, the ratio of Na⁺/K⁺ in both shoots and roots was increased in *Osflp* and decreased in OE plants (Fig. 5C and F). This result indicated that the excessive Na⁺ was accumulated in *Osflp* under the salt stress condition.

***OsCDKB1;1* is the direct target of OsFLP**

In Arabidopsis, AtFLP/AtMYB88 directly controlled *AtCDKB1;1* expression to regulate symmetric division in stomatal development (Xie et al. 2010a). *OsCDKB1;1* shares 88.5% amino acid identity with *AtCDKB1;1* and *AtCDKB1;2* (Qu et al. 2018). To identify the target of OsFLP, *OsCDKB1;1* was taken as one potential candidate. The expression of *OsCDKB1;1* was significantly induced in *Osflp* but repressed in OE, compared with that in ZH11 and COM (Fig. 6A), suggesting that OsFLP negatively regulates the transcriptional expression of *OsCDKB1;1*. To examine whether OsFLP can directly bind to the promoter region of *OsCDKB1;1*, a yeast one-hybrid assay was conducted. The 2266 bp upstream of the transcriptional start site (ATG) was truncated into several fragments. The results showed that OsFLP is directly bound to the region of 137 bp upstream of ATG, but not to the other regions of *OsCDKB1;1* promoter (Fig. 6B). To determine the exact sequence which was bound by OsFLP, we scanned the *OsCDKB1;1* promoter region 137 bp upstream of ATG (-137 bp to -1bp) and found six potential binding sites including -130 bp to -126bp (TTCCC), -125 bp to -121 bp (TTCCC), -114 bp to -110 bp (ATCCC), -78 bp to -74 bp (ATCCC), -18 bp to -14 bp (ATCGG) and -10 bp to -6 bp (TTCGG) (Fig. 6C). To confirm the binding specificity, we divided this 137 bp promoter region into three fragments with -137 bp to -95 bp, -95bp to -45 bp and -45 bp to -1 bp, respectively, and performed electrophoretic mobility shift assay (EMSA) using recombinant OsFLP. Ultimately, we found that -125 bp to -121 bp (TTCCC) and -78 bp to -74 bp (ATCCC) were required for OsFLP binding with obviously shifted bands. By contrast, the mutant versions and the excess of the unlabeled probe as competitors, failed to form the

shifted bands (Fig. 6D-F).

***OsCDKB1;1* negatively regulates salt stress responses**

As the expression of *OsCDKB1;1* was directly regulated by OsFLP (Fig. 6A), we were wondering if *OsCDKB1;1* was also involved in the salt stress responses in rice. Then, we were encouraged to examine whether *OsCDKB1;1* was required for salt tolerance. We exposed two-week-old seedlings of *OsCDKB1;1* over-expression plant (*OsCDKB1;1*-OE) and the *OsCDKB1;1* knockdown plant (*OsCDKB1;1*-RNAi) (Qu et al. 2018) to 150 mM NaCl stress conditions, with ZH11 as control. After 12 days of salt stress and 3 days recovery, *OsCDKB1;1*-OE plants exhibited more sensitivity to salt stress, while *OsCDKB1;1*-RNAi behaved more tolerance to salt stress compared with ZH11 (Fig. 7A). Consistently, under salt stresses, the survival rate was significantly lower in *OsCDKB1;1*-OE plants but much higher in *OsCDKB1;1*-RNAi plants compared with ZH11 (Fig. 7B). It suggested that *OsCDKB1;1* had a negative function on salt stress tolerance in rice.

To investigate whether the ROS accumulation was impacted in *OsCDKB1;1*-RNAi plants, we measured the H₂O₂ content with or without salt stresses. Intriguingly, upon salt stress, compared with ZH11, a higher level of H₂O₂ was accumulated in *OsCDKB1;1*-OE plants, while the lower level of H₂O₂ was accumulated in *OsCDKB1;1*-RNAi plants (Fig. 7C), suggesting that *OsCDKB1;1* influenced the content of H₂O₂ under salt stress condition. Collectively, these results indicate that *OsCDKB1;1* plays a negative role in response to salt stress by increasing ROS production.

Discussion

It has been implicated that *AtFLP* as a multiple functional regulator involves several aspects of plant physiological processes, including stomatal development, abiotic stress response, root gravitropism, and female reproductive development (Xie et al. 2010a; Xie et al. 2010b; Wang et al. 2015; Makkena et al. 2012). In rice, it has been reported that OsFLP controls the orientation of the GMC symmetric division in stomatal development (Wu et al. 2019), but the downstream targets are unknown (Lai et al. 2005). It seems that the function of FLP is conserved and divergent in monocots and dicots plants. However, the function of OsFLP in rice is still largely unknown. In this study, by transcriptomic analysis, the role of OsFLP was involved in many aspects, while we mainly focused on the salt stress responses. We also found that the *Osflp* mutant was sensitive to salt stress with more ROS accumulation and the OE lines behaved conversely. It indicates that OsFLP acts as a key regulator in salt tolerance in rice.

The Na⁺/K⁺ ratio in the cytosol is important for keeping the normal cellular activities in plants. Under salt stress conditions, the uptake of Na⁺ competes with K⁺ and even the salt stress blocks the K⁺ specific transporters (Zhu 2003). Our findings showed that more Na⁺ was taken up into the cells of *Osflp* under salt stress conditions. In this process, some sodium transporters may be involved (Fig. 7D). Interestingly, the integrative analysis of RNA-seq and ChIP-seq identified the Ca²⁺/Na⁺ antiporter (Os03g0411300) with increased expression, and the potassium channel (Os01g0846300) with decreased expression (Supplementary Table S2). Hence, we proposed that OsFLP may regulate the activity of Ca²⁺/Na⁺ antiporter and potassium channels to control the intracellular concentration of Na⁺. In Arabidopsis, AtFLP together with *AtCDKB1;1* and *AtCDKB1;2* controls the stomatal division. However, in rice, no obvious phenotype in stomatal development in *Oscdkb1;1* was found. *OsCDKB1;1* may have functional redundancy with *OsCDKB1;2* in regulating the stomatal development (Qu et al. 2018; Menges et al. 2005; Porceddu et al. 2001; Andersen et al. 2008). However, in Arabidopsis, the response to salt stress involves a rapid decline of cell cycle activity (Burssens et al. 2000; West et al. 2004), and the expression of CDKB1;1 and CDKB1;2 was downregulated under salt stress (Mahapatra and Roy 2021). Intriguingly, in our findings, *Oscdkb1;1* exhibited significantly salt tolerance with less ROS accumulation under salt stress conditions, suggesting that *OsCDKB1;1* had a negative function in salt tolerance. Considering OsFLP directly binding to the promoter region of *OsCDKB1;1* and repressing its transcription, we conclude that OsFLP associated with *OsCDKB1;1* confers salt tolerance in rice. Thus, our findings uncover a novel OsFLP-*OsCDKB1;1* module conferring salt tolerance by involving ROS accumulation in rice.

In addition, by RNA-seq analysis, many PRXs genes were identified in *Osflp*. PRXs mainly function in catalyzing H₂O₂ scavenging and ROS production (Bhatt and Tripathi 2011; Almagro et al. 2009), indicating that OsFLP might directly or indirectly regulate PRXs to influence the ROS contents in plants. Hence, in the future, it will be interesting to explore how OsFLP mediates the transcription of PRXs to affect the ROS accumulation and will be beneficial for uncovering the mechanism of *OsFLP* involved in physical processes of salt tolerance.

Based on our findings, we proposed a working model for OsFLP-*OsCDKB1;1* mediated salt tolerance in rice (Fig. 7D). Under salt stress conditions, in wild-type plant ZH11, OsFLP protein binds to the promoter region of *OsCDKB1;1* and represses the expression of *OsCDKB1;1* leading to a decrease in

the ROS accumulation resulting in salt tolerance in rice. In *Osflp* mutant plant, the expression of *OsCDKB1;1* is activated and the ROS accumulation is increased, ultimately, enhancing salt sensitivity in rice. Together, our findings extend the knowledge about the role of OsFLP in rice, especially in salt stress response, indicating the potential benefits for salt-tolerant rice cultivars' improvement.

Accession numbers

Sequences used in this study were download from the Rice Genome Annotation Project Database (<http://rice.plantbiology.msu.edu/>) with the following accession numbers: *OsFLP* LOC_Os07g43420; *OsCDKB1;1* LOC_Os01g67160; *OsPRX107* LOC_Os07g44550; *OsPRX109* LOC_Os07g47990; *OsPRX129* LOC_Os10g41720; *OsPRX112* LOC_Os07g48030; *OsRab16A* LOC_Os02g04680; *OsNAC5* LOC_Os07g37920; *OsDREB2A* LOC_Os01g07120; *OsDREB1F* LOC_Os01g73770; *OsACTIN2* LOC_Os03g50885.

Acknowledgements

This work was supported by grants from the Strategic Priority Research Program of the Chinese Academy of Sciences (grant number. XDA24010303) and the National Natural Science Foundation of China (grant numbers 31970804 and 32170722 to J. L, grant number 31470362 to J.-J. Z and grant number 31871377 and 32070723 to K.-Z. Y).

References

- Almagro L, Gómez Ros LV, Belchi-Navarro S, Bru R, Ros Barceló A, Pedreño MA (2009) Class III peroxidases in plant defence reactions. *J Exp Bot* 60: 377-390. [https://10.1093/jxb/ern277](https://doi.org/10.1093/jxb/ern277)
- Andersen SU, Buechel S, Zhao Z, Ljung K, Novák O, Busch W, Schuster C, Lohmann JU (2008) Requirement of B2-type cyclin-dependent kinases for meristem integrity in *Arabidopsis thaliana*. *Plant Cell* 20: 88-100. [https://10.1105/tpc.107.054676](https://doi.org/10.1105/tpc.107.054676)
- Bhatt I, Tripathi BN (2011) Plant peroxiredoxins: catalytic mechanisms, functional significance and future perspectives. *Biotechnol Adv* 29: 850-859. [https://10.1016/j.biotechadv.2011.07.002](https://doi.org/10.1016/j.biotechadv.2011.07.002)
- Boudolf V, Rombauts S, Naudts M, Inzé D, De Veylder L (2001) Identification of novel cyclin-dependent kinases interacting with the CKS1 protein of *Arabidopsis*. *J Exp Bot* 52: 1381-1382. <https://doi.org/10.1093/jexbot/52.359.1381>
- Burssens S, Himanen K, van de Cotte B, Beeckman T, Van Montagu M, Inzé D, Verbruggen N (2000) Expression of cell cycle regulatory genes and morphological alterations in response to salt stress in *Arabidopsis thaliana*. *Planta* 211: 632-640. [https://10.1007/s004250000334](https://doi.org/10.1007/s004250000334)
- De Veylder L, Joubès J, Inzé D (2003) Plant cell cycle transitions. *Curr Opin Plant Biol* 6: 536-543. [https://10.1016/j.pbi.2003.09.001](https://doi.org/10.1016/j.pbi.2003.09.001)
- Dewitte W, Murray JA (2003) The plant cell cycle. *Annu Rev Plant Biol* 54: 235-264. [https://10.1146/annurev.arplant.54.031902.134836](https://doi.org/10.1146/annurev.arplant.54.031902.134836)
- Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L (2010) MYB transcription factors in *Arabidopsis*. *Trends Plant Sci* 15: 573-581. [https://10.1016/j.tplants.2010.06.005](https://doi.org/10.1016/j.tplants.2010.06.005)

447 Dubey RS, Singh AK (1999) Salinity induces accumulation of soluble sugars and alters the
 448 activity of sugar metabolizing enzymes in rice plants. *Biol Plant* 42:233-9.
 449 <https://doi.org/10.1023/A:1002160618700>
 450 El-Kereamy A, Bi YM, Ranathunge K, Beatty PH, Good AG, Rothstein SJ (2012) The rice R2R3-MYB
 451 transcription factor OsMYB55 is involved in the tolerance to high temperature and modulates amino acid
 452 metabolism. *PLoS One* 7: e52030. <https://10.1371/journal.pone.0052030>
 453 Gray JE, Holroyd GH, van der Lee FM, Bahrami AR, Sijmons PC, Woodward FI, Schuch W,
 454 Hetherington AM (2000) The HIC signalling pathway links CO₂ perception to stomatal development.
 455 *Nature* 408: 713-716. <https://10.1038/35047071>
 456 Gupta B, Huang B (2014) Mechanism of salinity tolerance in plants: physiological, biochemical, and
 457 molecular characterization. *Int J Genomics* 2014: 701596. <https://10.1155/2014/701596>
 458 Hiei Y, Ohta S, Komari T, Kumashiro T (1994) Efficient transformation of rice (*Oryza sativa* L.) mediated
 459 by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J* 6: 271-282.
 460 <https://10.1046/j.1365-313x.1994.6020271.x>
 461 Horie T, Yoshida K, Nakayama H, Yamada K, Oiki S, Shinmyo A (2001) Two types of HKT transporters
 462 with different properties of Na⁺ and K⁺ transport in *Oryza sativa*. *Plant J* 27: 129-138.
 463 <https://10.1046/j.1365-313x.2001.01077.x>
 464 James RA, Blake C, Byrt CS, Munns R (2011) Major genes for Na⁺ exclusion, Nax1 and Nax2 (wheat
 465 HKT1;4 and HKT1;5), decrease Na⁺ accumulation in bread wheat leaves under saline and waterlogged
 466 conditions. *J Exp Bot* 62: 2939-2947. <https://10.1093/jxb/err003>
 467 Kanaoka MM, Pillitteri LJ, Fujii H, Yoshida Y, Bogenschutz NL, Takabayashi J, Zhu JK, Torii KU (2008)
 468 SCREAM/ICE1 and SCREAM2 specify three cell-state transitional steps leading to arabidopsis stomatal
 469 differentiation. *Plant Cell* 20: 1775-1785. <https://10.1105/tpc.108.060848>
 470 Kreps JA, Wu Y, Chang HS, Zhu T, Wang X, Harper JF (2002) Transcriptome changes for Arabidopsis
 471 in response to salt, osmotic, and cold stress. *Plant Physiol* 130: 2129-2141. <https://10.1104/pp.008532>
 472 Lai LB, Nadeau JA, Lucas J, Lee EK, Nakagawa T, Zhao L, Geisler M, Sack FD (2005) The Arabidopsis
 473 R2R3 MYB proteins FOUR LIPS and MYB88 restrict divisions late in the stomatal cell lineage. *Plant*
 474 *Cell* 17: 2754-2767. <https://10.1105/tpc.105.034116>
 475 Lei Q, Lee E, Keerthisinghe S, Lai L, Li M, Lucas JR, Wen X, Ren X, Sack FD (2015) The FOUR LIPS
 476 and MYB88 transcription factor genes are widely expressed in Arabidopsis thaliana during development.
 477 *Am J Bot* 102: 1521-1528. <https://10.3732/ajb.1500056>
 478 Lv Y, Yang M, Hu D, Yang Z, Ma S, Li X, Xiong L (2017) The OsMYB30 Transcription Factor
 479 Suppresses Cold Tolerance by Interacting with a JAZ Protein and Suppressing β -Amylase Expression.
 480 *Plant Physiol* 173: 1475-1491. <https://10.1104/pp.16.01725>
 481 Mahapatra K, Roy S (2021) SOG1 transcription factor promotes the onset of endoreduplication under
 482 salinity stress in Arabidopsis. *Sci Rep* 11: 11659. <https://10.1038/s41598-021-91293-1>
 483 Mäkelä P, Kontturi M, Pehu E, Somersalo S (1999) Photosynthetic response of drought- and salt-stressed
 484 tomato and turnip rape plants to foliar-applied glycinebetaine. 105: 45-50.
 485 <https://doi.org/10.1034/j.1399-3054.1999.105108.x>
 486 Makkena S, Lee E, Sack FD, Lamb RS (2012) The R2R3 MYB transcription factors FOUR LIPS and
 487 MYB88 regulate female reproductive development. *J Exp Bot* 63: 5545-5558. <https://10.1093/jxb/ers209>
 488 Martin C, Paz-Ares J (1997) MYB transcription factors in plants. *Trends Genet* 13: 67-73.
 489 [https://10.1016/s0168-9525\(96\)10049-4](https://10.1016/s0168-9525(96)10049-4)
 490 Matsui A, Ishida J, Morosawa T, Mochizuki Y, Kaminuma E, Endo TA, Okamoto M, Nambara E,

491 Nakajima M, Kawashima M, Satou M, Kim JM, Kobayashi N, Toyoda T, Shinozaki K, Seki M (2008)
 492 Arabidopsis transcriptome analysis under drought, cold, high-salinity and ABA treatment conditions
 493 using a tiling array. *Plant Cell Physiol* 49: 1135-1149. <https://10.1093/pcp/pcn101>
 494 Menges M, de Jager SM, Gruissem W, Murray JA (2005) Global analysis of the core cell cycle regulators
 495 of Arabidopsis identifies novel genes, reveals multiple and highly specific profiles of expression and
 496 provides a coherent model for plant cell cycle control. *Plant J* 41: 546-566. <https://10.1111/j.1365-313X.2004.02319.x>
 497
 498 Miao BH, Han XG, Zhang WH (2010) The ameliorative effect of silicon on soybean seedlings grown in
 499 potassium-deficient medium. *Ann Bot* 105: 967-973. <https://10.1093/aob/mcq063>
 500 Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R (2010) Reactive oxygen species homeostasis and
 501 signalling during drought and salinity stresses. *Plant Cell Environ* 33: 453-467. <https://10.1111/j.1365-3040.2009.02041.x>
 502
 503 Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7: 405-410.
 504 [https://10.1016/s1360-1385\(02\)02312-9](https://10.1016/s1360-1385(02)02312-9)
 505 Mittler R, Vanderauwera S, Gollery M, Van Breusegem F (2004) Reactive oxygen gene network of plants.
 506 *Trends Plant Sci* 9: 490-498. <https://10.1016/j.tplants.2004.08.009>
 507 Noctor G, Foyer CH (1998) ASCORBATE AND GLUTATHIONE: Keeping Active Oxygen Under
 508 Control. *Annu Rev Plant Physiol Plant Mol Biol* 49: 249-279. <https://10.1146/annurev.arplant.49.1.249>
 509 Ponce KS, Meng L, Guo L, Leng Y, Ye G (2021) Advances in Sensing, Response and Regulation
 510 Mechanism of Salt Tolerance in Rice. *Int J Mol Sci* 22. <https://10.3390/ijms22052254>
 511 Porceddu A, Stals H, Reichheld JP, Segers G, De Veylder L, Barroco RP, Casteels P, Van Montagu M,
 512 Inzé D, Mironov V (2001) A plant-specific cyclin-dependent kinase is involved in the control of G2/M
 513 progression in plants. *J Biol Chem* 276: 36354-36360. <https://10.1074/jbc.M011060200>
 514 Qu X, Yan M, Zou J, Jiang M, Yang K, Le J (2018) A2-type cyclin is required for the asymmetric entry
 515 division in rice stomatal development. *J Exp Bot* 69: 3587-3599. <https://10.1093/jxb/ery158>
 516 Roy SJ, Negrão S, Tester M (2014) Salt resistant crop plants. *Curr Opin Biotechnol* 26: 115-124.
 517 <https://10.1016/j.copbio.2013.12.004>
 518 Shen H, Zhong X, Zhao F, Wang Y, Yan B, Li Q, Chen G, Mao B, Wang J, Li Y, Xiao G, He Y, Xiao H,
 519 Li J, He Z (2015a) Overexpression of receptor-like kinase ERECTA improves thermotolerance in rice
 520 and tomato. *Nat Biotechnol* 33: 996-1003. <https://10.1038/nbt.3321>
 521 Shen Y, Shen L, Shen Z, Jing W, Ge H, Zhao J, Zhang W (2015b) The potassium transporter OsHAK21
 522 functions in the maintenance of ion homeostasis and tolerance to salt stress in rice. *Plant Cell Environ*
 523 38: 2766-2779. <https://10.1111/pce.12586>
 524 Shpak ED, McAbee JM, Pillitteri LJ, Torii KU (2005) Stomatal patterning and differentiation by
 525 synergistic interactions of receptor kinases. *Science* 309: 290-293. <https://10.1126/science.1109710>
 526 Song SY, Chen Y, Chen J, Dai XY, Zhang WH (2011) Physiological mechanisms underlying OsNAC5-
 527 dependent tolerance of rice plants to abiotic stress. *Planta* 234: 331-345. <https://10.1007/s00425-011-1403-2>
 528
 529 Tang L, Cai H, Ji W, Luo X, Wang Z, Wu J, Wang X, Cui L, Wang Y, Zhu Y, Bai X (2013) Overexpression
 530 of GsZFP1 enhances salt and drought tolerance in transgenic alfalfa (*Medicago sativa* L.). *Plant Physiol*
 531 *Biochem* 71: 22-30. <https://10.1016/j.plaphy.2013.06.024>
 532 Vandepoele K, Raes J, De Veylder L, Rouzé P, Rombauts S, Inzé D (2002) Genome-wide analysis of
 533 core cell cycle genes in Arabidopsis. *Plant Cell* 14: 903-916. <https://10.1105/tpc.010445>
 534 Wang HZ, Yang KZ, Zou JJ, Zhu LL, Xie ZD, Morita MT, Tasaka M, Friml J, Grotewold E, Beeckman

T, Vanneste S, Sack F, Le J (2015) Transcriptional regulation of PIN genes by FOUR LIPS and MYB88 during Arabidopsis root gravitropism. *Nat Commun* 6: 8822. <https://10.1038/ncomms9822>

West G, Inzé D, Beemster GT (2004) Cell cycle modulation in the response of the primary root of Arabidopsis to salt stress. *Plant Physiol* 135: 1050-1058. <https://10.1104/pp.104.040022>

Wu Z, Chen L, Yu Q, Zhou W, Gou X, Li J, Hou S (2019) Multiple transcriptional factors control stomata development in rice. *New Phytol* 223: 220-232. <https://10.1111/nph.15766>

Xia XJ, Gao CJ, Song LX, Zhou YH, Shi K, Yu JQ (2014) Role of H₂O₂ dynamics in brassinosteroid-induced stomatal closure and opening in *Solanum lycopersicum*. *Plant Cell Environ* 37: 2036-2050. <https://10.1111/pce.12275>

Xie Z, Lee E, Lucas JR, Morohashi K, Li D, Murray JA, Sack FD, Grotewold E (2010a) Regulation of cell proliferation in the stomatal lineage by the Arabidopsis MYB FOUR LIPS via direct targeting of core cell cycle genes. *Plant Cell* 22: 2306-2321. <https://10.1105/tpc.110.074609>

Xie Z, Li D, Wang L, Sack FD, Grotewold E (2010b) Role of the stomatal development regulators FLP/MYB88 in abiotic stress responses. *Plant J* 64: 731-739. <https://10.1111/j.1365-313X.2010.04364.x>

Xiong L, Schumaker KS, Zhu JK (2002) Cell signaling during cold, drought, and salt stress. *Plant Cell* 14 Suppl: S165-183. <https://10.1105/tpc.000596>

Yang A, Dai X, Zhang WH (2012) A R2R3-type MYB gene, OsMYB2, is involved in salt, cold, and dehydration tolerance in rice. *J Exp Bot* 63: 2541-2556. <https://10.1093/jxb/err431>

Yang K, Zhu L, Wang H, Jiang M, Xiao C, Hu X, Vanneste S, Dong J, Le J (2019) A conserved but plant-specific CDK-mediated regulation of DNA replication protein A2 in the precise control of stomatal terminal division. *Proc Natl Acad Sci U S A* 116: 18126-18131. <https://10.1073/pnas.1819345116>

Yang M, Sack FD (1995) The too many mouths and four lips mutations affect stomatal production in Arabidopsis. *Plant Cell* 7: 2227-2239. <https://10.1105/tpc.7.12.2227>

Yang Y, Guo Y (2018) Unraveling salt stress signaling in plants. *J Integr Plant Biol* 60: 796-804. <https://10.1111/jipb.12689>

Ye Y, Liu B, Zhao M, Wu K, Cheng W, Chen X, Liu Q, Liu Z, Fu X, Wu Y (2015) CEF1/OsMYB103L is involved in GA-mediated regulation of secondary wall biosynthesis in rice. *Plant Mol Biol* 89: 385-401. <https://10.1007/s11103-015-0376-0>

Zhang H, Liu Y, Wen F, Yao D, Wang L, Guo J, Ni L, Zhang A, Tan M, Jiang M (2014) A novel rice C₂H₂-type zinc finger protein, ZFP36, is a key player involved in abscisic acid-induced antioxidant defence and oxidative stress tolerance in rice. *J Exp Bot* 65: 5795-5809. <https://10.1093/jxb/eru313>

Zhu JK (2003) Regulation of ion homeostasis under salt stress. *Curr Opin Plant Biol* 6: 441-445. [https://10.1016/s1369-5266\(03\)00085-2](https://10.1016/s1369-5266(03)00085-2)

Statements and Declarations

Funding

This work was supported by grants from the Strategic Priority Research Program of the Chinese Academy of Sciences (grant number. XDA24010303) and the National Natural Science Foundation of China (grant numbers 31970804 and 32170722 to J. L, grant number 31470362 to J.-J. Z and grant number 31871377 and 32070723 to K.-Z. Y).

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

C.-X. Z and J. Z performed the experiments and analyzed the data. H.-C. L, X.-X. Q, J.-X. W, Q.-X.-M. H and J.-J. Z participated in parts of experiments and data analysis. J. L, K.-Z. Y, C.-X. Z and J. Z designed the experiments. C.-X. Z and J.L wrote the manuscript. All authors read and approved the final manuscript.

Data Availability

All data generated during this study are included in this published article and its supplementary information files. The RNA-seq datasets generated during the current study are available in the NCBI Sequence Read Archive (SRA) repository, <https://dataview.ncbi.nlm.nih.gov/object/PRJNA805061?reviewer=m847fdbhc5h8hlajg783fodski> <https://dataview.ncbi.nlm.nih.gov/object/PRJNA805087?reviewer=gj1raljgoss270nmmf2d8dgmfu> The ChIP-seq datasets generated during the current study are available in the NCBI Sequence Read Archive (SRA) repository, <https://dataview.ncbi.nlm.nih.gov/object/PRJNA803610?reviewer=8ark21mh1c3mrblaoj1fnpek34>

Figure captions

Fig. 1 The characterization of *Osflp*

A. *Osflp* mutant line was generated by TILLING system. A point mutation nucleotide G to A was identified in the sixth exon at position 2126 bp downstream of ATG which led to the substitution of amino acid Ala with Thr

B. Multiple alignments of amino acid sequences of OsFLP, AtFLP, AtMYB88, Bradi1g20586, Glyma01g05190, Solyc04g005710, Solyc05g007160, Zm24064630 showed the mutation site of *Osflp* located in the highly conserved region of R3 domain

C-E. Normal stomata in ZH11 and complementation line. Abnormal division and differentiation stomata in *Osflp* was indicated with red arrowhead, Scale bars=20 μ m

Fig. 2 Identification of potential targets of OsFLP by transcriptomic analysis

A. Volcano Plot shows the up-and down-regulated genes in *Osflp* mutant compared with ZH11

B. The potential proteins or biological processes which are related to OsFLP, are identified by RNA-seq analysis

C. GO term enrichment of differentially expressed genes (DEGs) for biological process with RNA-seq analysis

D. The integrative of RNA-seq and ChIP-seq. In total, 125 genes were identified as the potential targets of OsFLP. Among them, 71 genes were induced and 54 genes were repressed

E-H. The expression of salt stress-related genes *OsRab16A*, *OsNAC5*, *OsDREB2A*, *OsDREB1F* were detected using RT-qPCR with or without salt treatment in ZH11, *Osflp*, COM and OE lines, respectively.

OsActin was always taken as the reference gene. Data are means $\pm$ SD. The student's *t*-test was performed.

Statistically significant differences are indicated with asterisks:*, $P<0.05$, **, $P<0.01$

Fig. 3 *Osflp* mutants are more sensitive to salt stress

A. Salt stress tolerance phenotype assay of ZH11, *Osflp*, COM and OsFLP overexpression line (OE) seedlings with or without NaCl treatment. Seedlings were grown on 1/2 MS plates for three days and transferred to 1/2 MS plates supplemented with or without 200 mM NaCl for 5 days. Scale bars=1cm

B and C. The root length and the shoot length of ZH11, *Osflp*, COM and OE seedlings were quantified, respectively, in D. Data are means $\pm$ SD. The student's *t*-test was performed. Statistically significant differences are indicated with asterisks: **, $P<0.01$

D. Salt stress tolerance phenotype assay of ZH11, *Osflp*, COM, and OE seedlings were grown on soil. 7-day-old seedlings cultivated on soil irrigated with water or water supplement with 200 mM NaCl for 15 days and 21 days, respectively. Scale bars=10 cm

E. Survival rates of ZH11, *Osflp*, COM, and OE plants after 21 days of stress treatment were analyzed. Data are means $\pm$ SD. The student's *t*-test was performed. Statistically, significant differences are indicated with asterisks: **, $P<0.01$

Fig. 4 More ROS was accumulated in *Osflp* rice mutant under salt stress conditions

A. Salt stress phenotype assay of ZH11, *Osflp*, COM, and OE seedlings cultivated with hydroponic culture. Two-week-old seedlings were treated with or without 150 mM NaCl in hydroponic culture after 3 days, 10 days, and 3 days of restored growth.

B. Survival rate was analyzed after 3 days of recovery.

C. Soluble sugar content was measured after 3 days of recovery.

D-G. SOD activity (D), POD activity (E), CAT activity (F), and H₂O₂ content (G) were assayed after 3 days of recovery. Data are means $\pm$ SD from three biological replicates. Student's *t*-test was performed.

Student's *t* - test was performed. Statistically significant differences are indicated with asterisks: *, $P < 0.05$, **, $P < 0.01$

H-I. Both DAB (H) and NBT(I) staining showed more ROS accumulation in *Osflp* mutant leaves treated with 150 mM NaCl treatment after 24 hours. Scale bars=0.5 cm

Fig. 5 *Osflp* mutant accumulates more Na^+ and less K^+ in shoots and roots under salt stress

Na^+ and K^+ contents assay in root and shoot of ZH11, *Osflp*, COM and OE lines of rice plants. Two-week-old seedlings were treated with or without 150mM NaCl for 4 days

A. and B. Na^+ and K^+ contents were measured in the shoot

C. The ratio of Na^+ to K^+ in the shoot was calculated according to the measurements in A. and B

D. and E. Na^+ and K^+ contents were measured in root

F. The ratio of Na^+ to K^+ in root was calculated according to the measurements in D. and E. Data are means $\pm$ SD from three biological replicates. The student's *t*-test was performed. Statistically, significant differences were indicated with asterisks: *, $p < 0.05$, **, $p < 0.01$

Fig. 6 OsFLP directly binds to the promoter region of *OsCDKB1;1* and regulates its transcriptional expression

A. The transcriptional expression of *OsCDKB1;1* was negatively regulated by OsFLP. The transcriptional level of *OsCDKB1;1* was detected by RT-qPCR in ZH11, *Osflp*, COM, and OE lines of rice plants

B. OsFLP binds to the promoter region of *OsCDKB1;1* in yeast one-hybrid assay. The full length of the promoter region of *OsCDKB1;1* was truncated into nine fragments. The area from -137 bp to -1 bp was identified as the minimal promoter region, which is the region for the binding of OsFLP

C. Schematic diagram of *OsCDKB1;1* promoter region showing potential binding sites of OsFLP in three different regions including P1, P2, and P3

D-F. Electrophoretic mobility shift assays (EMSA) indicated OsFLP specifically bound to the promoter region of the *OsCDKB1;1*. The arrows indicate the protein-DNA complexes. The asterisks indicate the non-specific binding. The competitor was the probe without biotin label

Fig. 7 *OsCDKB1;1* negatively regulates the response to salt stress in rice

A. Salt stress phenotype analysis of ZH11, *OsCDKB1;1*-OE, *OsCDKB1;1*-RNAi. Two-week-old seedlings were cultivated in hydroponic culture with or without 150 mM NaCl for 12 days and then restored growth for 3 days

B. The survival rates of ZH11, *OsCDKB1;1*-OE, *OsCDKB1;1*-RNAi were analyzed after salt stress recovery

C. The content of H₂O₂ was monitored in ZH11, *OsCDKB1;1*-OE, *OsCDKB1;1*-RNAi after salt stress recovery. Data in B-E are means $\pm$ SD from three biological replicates. The student's *t*-test was performed. Statistically, significant differences are indicated with asterisks: **, *P*<0.01

D. Proposed working model for OsFLP-*OsCDKB1;1* transcriptional regulation module mediated salt stress response in rice. Under salt stress conditions, in ZH11, OsFLP binds to the promoter region of *OsCDKB1;1* to repress the transcriptional expression of *OsCDKB1;1* resulting in less ROS accumulation, ultimately conferring salt tolerance (in the left panel). However, in *Osflp* mutant plants, the expression of *OsCDKB1;1* is enhanced, and promotes the ROS accumulation in plants, thereby leading to salt sensitivity in rice (in the right panel). OsFLP_m indicates mutation of OsFLP

Supplementary Information

Fig. S1 Phylogenetic tree shows the relationship of FLP proteins from algae, lower plants and higher plants

Fig. S2 RT-qPCR was performed to verify the results of the RNA-seq

Fig. S3 KEGG enrichment analysis with ChIP-seq

Fig. S4 KEGG enrichment analysis with genes identified by the integrative analysis of RNA-seq data and ChIP-seq data

Fig. S5 The expression of *OsFLP* upon salt stress

Fig. S6 *Osflp* mutants are more sensitive to salt stress

Table. S1 Genes are related to ROS metabolism enriched by RNA-seq analysis

Table. S2 Genes are related to salt stress enriched by the integrative analysis of RNA-seq and ChIP-seq

Table. S3 Primers were used in this study

Dataset. S1 RNA-seq *Osflp_vs_ZH11*_annotation

Dataset. S2 RNA-seq data analysis with categories

Dataset. S3 GO analysis with RNA-seq data

Dataset. S4 ChIP-seq original data_annotation

Dataset. S5 KEGG Enrichment with ChIP-seq data

Dataset. S6 RNA-seq and ChIP-seq integrative analysis annotation_Total

692 **Dataset. S7** KEGG Enrichment genes from RNA-seq and ChIP-seq

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [DatasetS1.RNAseqOsflpvsZH11annotation.xlsx](#)
- [DatasetS2.RNAseqdataanalysiswithcategories.xls](#)
- [DatasetS3.GOanalysiswithRNAseqdata.xlsx](#)
- [DatasetS4.ChIPseqoriginaldataannotation.xls](#)
- [DatasetS5.KEGGENrichmentwithChIPseqdata.xls](#)
- [DatasetS6.RNAseqandChIPseqintegrativeanalysis.xlsx](#)
- [DatasetS7.KEGGENrichmentgenesfromRNAseqandChIPseq.xlsx](#)
- [Supplementaryfiguresandtables.pdf](#)