

Validation of Perceived Physical Fatigability Using the Simplified-Chinese Version of the Pittsburgh Fatigability Scale

Yixin Hu (✉ chhyxcn@126.com)

Chinese PLA General Hospital

Hangming Zhang

Peking University Shougang Hospital <https://orcid.org/0000-0003-2373-2207>

Weihao Xu

Chinese PLA General Hospital

Ming Zhao

Haidian 37th Ex-Cadre Rest and Recuperation Center

Juan Liu

Beijing North Hospital

Linna Wu

Chinese PLA General Hospital

Lin Zou

Chinese PLA General Hospital

Jing Zuo

Chinese PLA General Hospital

Yunxia Liu

Chinese PLA General Hospital

Woei-Nan Bair

Department of Physical Therapy, University of the Sciences in Philadelphia

Yujia(Suanna) Qiao

University of Pittsburgh Department of Epidemiology

Nancy W. Glynn

University of Pittsburgh Department of Epidemiology

Research article

Keywords: older adults, fatigue, validity, physical performance, ADL

Posted Date: December 29th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-135784/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Validation of Perceived Physical Fatigability using the Simplified-Chinese Version of the Pittsburgh Fatigability Scale

Yixin Hu^{1*}, Hangming Zhang², Weihao Xu¹, MingZhao³, Juan Liu⁴, Linna Wu⁵, Lin Zou¹, Jing Zuo¹, Yunxia Liu¹, LiFan¹, Woei-Nan Bair⁶, Yujia (Susanna) Qiao⁷, Nancy W. Glynn⁷

Author details (addresses)

1. Geriatric Health Care Department of The Second Medical Center & National Clinical Research Center for Geriatric Diseases, Chinese PLA General Hospital, Beijing, China.
2. Department of Orthopedics, Peking University Shougang Hospital, Beijing, China
3. Outpatient Department, Haidian 37th Ex-Cadre Rest and Recuperation Center, Beijing, P.R. China
4. Geriatric Department of Beijing North Hospital of Ordnance Industry, Beijing, China
5. Geriatric Emergency Department of The Second Medical Center & National Clinical Research Center for Geriatric Diseases, Chinese PLA General Hospital, Beijing, China
6. Department of Physical Therapy, University of the Sciences in Philadelphia, Philadelphia, PA, USA
7. Department of Epidemiology, Graduate School of Public Health, Center for Aging and Population Health, University of Pittsburgh, Pittsburgh, PA, USA

Acknowledgments We acknowledge all participants involved in the present study, and thank Shuangshuang Du, Jinxia Li, Longqin Zhang and Guangji Wang for their thoughtful contribution for this research, and Jinhui Ju for help with the translation procedure of the simplified-Chinese version of the PFS.

Authors' contributions

Yixin Hu involved in the study design, interpreting the results, wrote the first draft of the manuscript.

Hangming Zhang performed statistical analyses, amend the first draft of the manuscript, Weihao Xu, MingZhao, Lin Zou, Juan Liu and Linna Wu, Jing Zuo and Yunxia Liu performed the investment in community and prepare the data. LiFan, Woei-Nan Bair, Yujia (Susanna) Qiao, and

Nancy W. Glynn aided in interpreting the results and revised the manuscript draft for important intellectual content. All authors read and approved the final manuscript, thereby taking full responsibility for the work and manuscript content.

Funding sources: The study was supported by Opening Foundation of National Clinical Research Center for Geriatric Diseases (NCRCG-PLAGH-2017010), Healthcare Grants (15BJZ41), Healthcare Grants (20BJZ30) and Healthcare Grants of The Second medical center (ZXD2010). Additionally, the Claude D. Pepper Older Americans Independence Center, Research Registry and Developmental Pilot Grant (NIH P30 AG024827), and the Intramural Research Program, National Institute on Aging supported Nancy W. Glynn, PhD to develop the Pittsburgh Fatigability Scale.

Availability of data and materials

The datasets used during the current study are available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

The ethics committee of the Chinese PLA General Hospital approved this study, which complies with the current laws of China (Ethic number: S2018-102-01). Written informed consent was obtained from all individual participants included in the study.

Consent for publication

Not applicable.

Competing interests

The authors declare they have no competing interests.

*Corresponding author: Yixin Hu, Email:chhyxcn@126.com

Abstract

Background: The Pittsburgh Fatigability Scale (PFS) was developed to capture fatigue and demand in a single tool, filling a gap that no validated questionnaire existed to measure perceived fatigability. Since fatigability is a more sensitive measure of a person's susceptibility to fatigue, we validated the simplified-Chinese version of the PFS among Chinese community-dwelling older adults.

Methods: The PFS was translated into the simplified-Chinese by the translation, retro-translation method. Internal consistency of the Physical subscale of the PFS was evaluated by Cronbach's alpha. Convergent validity and discriminant validity were evaluated against physical performance measures (i.e., Short Physical Performance Battery & Timed Up and Go Test) and daily living performance (i.e., Barthel Index & Instrumental activity of daily living).

Results: The simplified-Chinese version of PFS Physical scores showed strong internal consistency (Cronbach's alpha=0.81). Higher PFS Physical scores were associated with worse physical performance, and daily living performance (|correlation coefficient| range: 0.36-0.56, p<.001). Age- and sex-adjusted PFS Physical scores had moderate to good overall discrimination for correctly classifying people by their physical performance and daily living performance (AUCs range 0.70-0.87, p<.001)

Conclusions: The PFS simplified-Chinese version is a valid instrument to assess perceived physical fatigability in Chinese-speaking older adults with good convergent validity. Thus, the PFS, with low

cost and greater feasibility, is a desired tool to measure fatigability in large population studies.

Keywords: older adults, fatigue, validity, physical performance, ADL

Background

In developing countries, people aged 65 years and older represent the fast-growing proportion and absolute number^[1]. The 2010 China Census revealed that 13.3% (i.e., 177.6 million) of the total population was represented by older persons^[2]. Fatigability is a common and distressing self-perceived symptom while performing usual mental and physical activities, especially among older adults, and strongly associated with adverse health-related events^[3]. Perceived fatigability measures an individual's whole-body tiredness anchored to the standardized activity of a specific intensity and duration, which may help to overcome the limitation of current fatigue scales by providing a less-biased and more sensitive measure^[3-8].

Since there was no questionnaire-based tool to measure perceived fatigability, Glynn et al. developed the Pittsburgh Fatigability Scale (PFS) in English, a 10-item validated tool evaluating both physical and mental fatigability in separate subscales^[6]. The PFS Physical subscale has good convergent validity against performance measures of mobility, physical function and fitness, as well as good concurrent validity against performance-based measures of fatigability^[6]. Greater perceived physical fatigability using the PFS has been shown to be higher in women than men, associated with poorer mobility, meaningful decline in physical function, and other important health indicators^[3, 4, 7, 9, 10].

The decline in physical function with aging is usually defined in 4 stages: active pathology, impairment, functional limitation, ultimately disability.^[11] Given the importance of measuring fatigability in older adults^[8], and the lack of a validated tool in China, it is important to evaluate the

validity of the PFS in Chinese community-dwelling older adults with different physical function assessment tools. This study aimed to validate simplified-Chinese version of the PFS Physical scores by assessing convergent validity against physical performance and daily living performance among Chinese older adults.

Methods

Data and Study Participants

A cross-sectional study of older adults from a senior community was performed in Beijing, China. This study was approved by the Research Ethics Committee of Chinese PLA General Hospital (Ethic number: S2018-102-01) and has been registered in Chinese Clinical Trial Register (ChiCTR1900022576). All participants signed the written informed consent.

A total of 457 participants were recruited from a senior community. At enrollment, the inclusion criteria was voluntary participation in this study, and participants were excluded if they were: (1) unable to perform Timed Up and Go Test (TUG) test or gait speed test; (2) had severe cognitive impairment, dementia, or severe hearing problem; (3) had an implanted electronic device or orthopedic metal implantations; and (4) had terminal diseases. All of the interviewers were well-trained before the study inception. Participants' data were collected by face-to-face interview and real-time physical and cognitive measurements at the community medical center or their home.

Linguistic Translation of the Simplified-Chinese Version of PFS

The validated method of translation, Retro-translation of the Pittsburgh Fatigability Scale was followed to obtain the final simplified-Chinese version of the PFS^[12]. Two Chinese-speaking researchers independently translated the PFS into simplified-Chinese. Discrepancies were discussed and an agreement was reached by consensus. We administered the simplified-Chinese version of the

PFS to two Chinese older community-dwellers to evaluate the proper understanding of items used in the scale. Then, the simplified-Chinese version of the PFS was retro-translated into English by a bilingual researcher and then checked for accuracy by another independent bilingual individual. For culturally relevancy some example activities (dusting, straightening up, baking; raking; aerobic machines, Zumba) from the original PFS were removed and a new example, Ping-Pong was added to the high-intensity activity item. The final simplified-Chinese version of the PFS is available upon request by the developer: <https://publichealth.pitt.edu/epidemiology/research-practice/faculty-research/pittsburgh-fatigability-scale>.

Measurement of Perceived Physical Fatigability

The 10-item Pittsburgh Fatigability Scale was used to assess perceived physical fatigability. Participants were asked to rate their level of physical tiredness that they expected or imagined they would feel after completing each activity on a scale from 0 (no fatigue) to 5 (extreme fatigue). These activities ranged in intensity from low (e.g. watching TV for 2 h) to high (e.g. high-intensity activity for 30 min). Responses to each item were summed to create PFS Physical scores ranging from 0 to 50 (higher scores = greater perceived fatigability). A PFS Physical Score ≥ 15 denotes greater perceived physical fatigability.^[3, 7] Additionally, PFS Physical scores were categorized into six groupings according to previously published work,^[3, 7] and then collapsed into three groupings due to distribution of the sample to examine severity of perceived physical fatigability.

Physical Performance Measures

The Short Physical Performance Battery (SPPB) includes three objective tests: 1) two-timed 4-m walks at a usual pace; 2) rising 5 times from a seated position to an upright position as quickly as possible with the arms folded across the chest; and 3) three individual standing balance tests,

including 3 progressively more difficult positions for 10s each: side-by-side stand, semi-tandem stand, and tandem stand^[13]. Lower limb strength was measured by the total time of repeated chair stands (\geq 16.7 sec means poor lower limb strength)^[14]. Each of the 3 tests was scored 0 (worst performance) to 4 (best performance). The three tests were summed, and a total SPPB score ranged from 0 to 12, with higher scores reflecting better physical function.

Daily Living Performance

The Barthel Index was used to measure the Activities of Daily Living (ADL)^[15], ranging between 0-100 with a score of 0-80 indicating functional dependency. Instrumental Activity of Daily Living (IADL) was measured by the 8-item scale developed by Lawton and Brody^[16]. IADL dependence was defined as self-reported inability to perform any of the following tasks independently: using a telephone, shopping, preparing food, doing laundry, administering medications, going places, housekeeping, or personal finances.

Covariates

Covariates were measured from questionnaires which included age, sex, education level, marital status (married vs. others including divorce, widowed, and never married), smoking status, body mass index (BMI), nutritional status, and self-reported physician-diagnosed diseases including hypertension, coronary artery disease (CAD), chronic obstructive pulmonary diseases (COPD), diabetes mellitus, and chronic kidney disease (CKD).

Statistical Analyses

The descriptive statistics for continuous variables were by mean \pm standard deviation and

categorical variables were by N(%). Cronbach's alpha was used to measure the internal consistency of the simplified-Chinese version of the PFS Physical scores. Commonly, a Cronbach's alpha coefficient is preferably above 0.8, and 0.7 to 0.8 is considered acceptable^[17].

Spearman's rank-order correlations were used to examine the correlations of the PFS Physical scores with physical performance and daily living performance. The strength of correlations reflected via Spearman correlation coefficients^[18]: $|r| \leq 0.3$: low or weak correlation; $|r|=0.3-0.5$: moderate correlation; and $|r| \geq 0.5$: strong correlation.

When using the dichotomous measures of distinguishing different levels of physical performance and daily living performance, the least-square means and standard error adjusted for age and sex of the two groups were calculated. The discriminant validity was assessed by the Area Under the Curve (AUC): AUC: 0.5-0.7, poor or low; AUC: 0.7-0.8, fair or moderate; and AUC: 0.8-1.0, good or excellent. All statistical analyses were performed using Statistical Package for Social Science (SPSS) software, version 22.

Results

Sample Description

The average age of the 457 participants was 84.5 ± 5.8 years, 60.2% women, 27.7% had lower educational status, and 61.3% were married. The average BMI of the participants was 24.0 ± 3.5 kg/m². Mean PFS Physical score was 25.5 ± 9 points, and 403 (88.2%) of them had a PFS score ≥ 15 , and 271 (59.3%) of them ≥ 25 (Table 1). Additionally, women had higher PFS Physical scores than men, 26.4 ± 8.4 versus 24.1 ± 9.4 , respectively, $P = .007$ (Figure 1). PFS Physical scores were distributed across the range of potential values, and no sign of ceiling or floor effects. The Cronbach's alpha of the simplified-Chinese version of the PFS was 0.81, which showed good

internal consistency.

Convergent Validity

Higher PFS Physical scores were correlated with both lower extremity and hand function.

Specifically, for SPPB score ($r=-0.51$), gait speed ($r=-0.55$), the time to complete 5 chair stands ($r=0.40$) and TUG test ($r=0.28$). For daily living performance, higher PFS Physical scores were also associated with ADL ($r=-0.56$) and IADL ($r=-0.48$), all $p<.001$ (Table 2).

Age- and sex-adjusted PFS Physical scores were significantly higher in subgroups of worse physical and daily living performance, indicating greater perceived physical fatigability (differences 2.2 to 10.7 points, Table 3). Moreover, AUC values (Table 3) indicated PFS Physical scores had moderate to excellent overall discrimination for correctly classifying people by their physical and daily living performance status (AUCs: 0.70–0.87).

Discussion

The present study was the first to translate the PFS into simplified-Chinese, modify according to cultural context and validate it among mainland Chinese older adults. This study showed that the simplified-Chinese version of the PFS Physical subscale had good content validity, indicating that it is a valid measure of perceived physical fatigability in Chinese older adults. Our findings indicated that PFS Physical scores were moderate- to strongly associated with physical performance (SPPB, gait speed, TUG) and daily living performance (ADL and IADL). Greater perceived physical fatigability was significantly associated with worse physical and daily living performance. Overall, the simplified-Chinese version of the PFS exhibited moderate to excellent convergent validity and strong internal consistency.

The original English PFS has been translated into 12 languages, with several versions already validated in their associated populations. The English version of the PFS^[6] presents high concurrent

and convergent validity. Like the original and Spanish versions of the PFS, the simplified-Chinese version was able to distinguish those exhibiting high versus low fatigability against physical performance measures.^[6, 19] Additionally, the English and Dutch versions of the PFS had good reliability for the Physical subscale with intraclass correlations ≥ 0.80 ^[6, 18]. The PFS Physical scores of our study were higher than others, likely because our population was older. In our population of older adults from Beijing, 88.2% of this population (mean age around 85 years) were categorized as having greater perceived physical fatigability comparable to the oldest-old (≥ 90 years) from the Long Life Family Study (LLFS), a generally healthier population enriched for exceptional longevity.^[10] In the LLFS^[10], mean PFS Physical scores of those 80-89 year and ≥ 90 years old were 19.1 and 28.6 points, respectively, indicating that our sample had greater fatigability, with a mean of 25.5 points. Similar to LaSorda et al.,^[10] we found that women had significantly higher PFS Physical scores than men.

ADL and Instrument ADLs are common questionnaires for assessment of disability, but are less sensitive for their “ceiling effects”,^[16, 20] especially for the individual’s healthy status modifications. Further, it is of importance to identify those with physical limitations in clinical settings, since they may benefit more from early interventions to reduce physical fatigability. Earlier functional limitations are commonly measured with the SPPB and its components.^[11, 21] Total SPPB score is a strong predictor of mobility disability and ADL disability.^[13] Gait speed, which could be easily measured in clinical setting and communities, has been reported by many studies, and independently predicts several adverse health outcomes.^[22, 23] In the present study, we found that the PFS not only had good concurrent and convergent validity against ADL and IADL, but also was positively associated with total SPPB score and gait speed. Our findings concur with findings from the Long Life Family Study^[10] showing that having difficulty with ADLs (physical disability) was a

risk factor associated with greater perceived physical fatigability. The current study also indicated that perceived physical fatigability was a risk factors associated with physical limitation in this very old Chinese group.

The PFS includes commonly performed activities from low to high-intensity of older adults. Even persons who are unable to perform particular activities can complete the PFS by imaging their perception of tiredness, thus we are still able to obtain a valid measure of their perceived physical fatigability. Additionally, the PFS is a brief, simple tool which is easy and inexpensive to use, especially for large population-based studies. Moreover, using the PFS can better elucidate how fatigability fits in the disablement pathway and functional decline, and identify older adults at risk of dysfunction in clinical and research settings instead of using performance-based fatigability measures that some older adults are unable to do.

The strengths of this study include the relatively large sample size and the assessment of convergent validity of PFS with a wide variety of performance-based measurements and daily living performance. Further, the linguistic translation of the simplified-Chinese version of PFS was used the validated method of translation, retro-translation. A limitation of this study is that the sample came from one community of retired older adults, limiting generalizability due to potential selection bias.

Conclusions

The PFS, simplified-Chinese version, is a brief and easy to use tool to measure perceived physical fatigability in Chinese older adults. The PFS Physical subscale has high concurrent and convergent validity against physical performance measurements and daily living performance. This is of great significance for improving the evaluation of fatigability impact and identifying older adults at risk

of mobility limitation in clinical and research settings. This study also provides the possibility of using perceived physical fatigability as clinical indicator or predictor variable in further studies, as it is an important prognostic measure of phenotypic aging.

References

- [1] United Nations DoEaSA. World Population Prospects: The 2012 Revision .
- [2] China NBoS. Communiqué on the main data of the sixth national census in 2010. http://www.gov.cn/guoqing/2012-04/20/content_2582698.htm .
- [3] Schrack JA, Simonsick EM, Glynn NW. Fatigability: A Prognostic Indicator of Phenotypic Aging. *J Gerontol A Biol Sci Med Sci.* 2020. 75(9): e63-e66.
- [4] Eldadah BA. Fatigue and fatigability in older adults. *PM R.* 2010. 2(5): 406-13.
- [5] Wasson E, Rosso AL, Santanasto AJ, et al. Neural correlates of perceived physical and mental fatigability in older adults: A pilot study. *Exp Gerontol.* 2019. 115: 139-147.
- [6] Glynn NW, Santanasto AJ, Simonsick EM, et al. The Pittsburgh Fatigability scale for older adults: development and validation. *J Am Geriatr Soc.* 2015. 63(1): 130-5.
- [7] Simonsick EM, Schrack JA, Santanasto AJ, Studenski SA, Ferrucci L, Glynn NW. Pittsburgh Fatigability Scale: One-Page Predictor of Mobility Decline in Mobility-Intact Older Adults. *J Am Geriatr Soc.* 2018. 66(11): 2092-2096.
- [8] Simonsick EM, Glynn NW, Jerome GJ, Shardell M, Schrack JA, Ferrucci L. Fatigued, but Not Frail: Perceived Fatigability as a Marker of Impending Decline in Mobility-Intact Older Adults. *J Am Geriatr Soc.* 2016. 64(6): 1287-92.
- [9] Alexander NB, Taffet GE, Horne FM, et al. Bedside-to-Bench conference: research agenda for idiopathic fatigue and aging. *J Am Geriatr Soc.* 2010. 58(5): 967-75.
- [10] LaSorda KR, Gmeli T, Kuipers AL, et al. Epidemiology of Perceived Physical Fatigability in Older Adults: The Long Life Family Study. *J Gerontol A Biol Sci Med Sci.* 2019 .
- [11] Nagi SZ. An epidemiology of disability among adults in the United States. *Milbank Mem Fund Q Health Soc.* 1976. 54(4): 439-67.
- [12] Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine (Phila Pa 1976).* 2000. 25(24): 3186-91.
- [13] Guralnik JM, Simonsick EM, Ferrucci L, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol.* 1994. 49(2): M85-94.
- [14] Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med.* 1995. 332(9): 556-61.
- [15] MAHONEY FI, BARTHEL DW. FUNCTIONAL EVALUATION: THE BARTHEL INDEX. *Md State Med J.* 1965. 14: 61-5.
- [16] Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental activities of daily living. *Gerontologist.* 1969. 9(3): 179-86.
- [17] Sousa VD, Rojjanasrirat W. Translation, adaptation and validation of instruments or scales for use in cross-cultural health care research: a clear and user-friendly guideline. *J Eval Clin Pract.* 2011. 17(2): 268-74.
- [18] Feenstra M, Smidt N, van Munster BC, Glynn NW, de Rooij SE. Translation and validation of the Dutch Pittsburgh Fatigability Scale for older adults. *BMC Geriatr.* 2020. 20(1): 234.
- [19] Pérez LM, Roqué M, Glynn NW, et al. Validation of the Spanish version of the Pittsburgh Fatigability Scale for older adults. *Aging Clin Exp Res.* 2019. 31(2): 209-214.
- [20] KATZ S, FORD AB, MOSKOWITZ RW, JACKSON BA, JAFFE MW. STUDIES OF ILLNESS IN THE AGED. THE INDEX

- OF ADL: A STANDARDIZED MEASURE OF BIOLOGICAL AND PSYCHOSOCIAL FUNCTION. JAMA. 1963. 185: 914-9.
- [21] Justice JN, Cesari M, Seals DR, Shively CA, Carter CS. Comparative Approaches to Understanding the Relation Between Aging and Physical Function. *J Gerontol A Biol Sci Med Sci*. 2016. 71(10): 1243-53.
 - [22] Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *JAMA*. 2011. 305(1): 50-8.
 - [23] Studenski S, Perera S, Wallace D, et al. Physical performance measures in the clinical setting. *J Am Geriatr Soc*. 2003. 51(3): 314-22.
 - [24] Porto JM, Nakaishi A, Cangussu-Oliveira LM, Freire Júnior RC, Spilla SB, Abreu D. Relationship between grip strength and global muscle strength in community-dwelling older people. *Arch Gerontol Geriatr*. 2019. 82: 273-278.

Table 1. Participant Characteristics of the Simplified-Chinese version PFS Validation Sample

Characteristics	Total sample N=457	PFS Physical Score Severity		
		0-14 54 (11.8)	15-24 132 (28.9)	≥25 271 (59.3)
Age, years	84.8±5.8	80.2±8.7	84.5±5.8	85.8±4.5
≥ 80 years	404 (92.3)	35 (64.8)	116 (87.9)	253 (93.4)
Women	275 (60.2)	26 (48.1)	65 (49.2)	184 (67.9)
Married	280 (61.3)	35 (64.8)	93 (70.5)	152 (56.1)
Education level				
< Senior middle school	127 (27.8)	16 (29.6)	25 (18.9)	86 (31.7)
Senior middle school	109 (23.9)	15 (27.8)	28 (21.2)	66 (24.4)
> Senior middle school	221 (48.4)	23 (42.6)	79 (59.8)	119 (43.9)
Monthly income, RMB				
<10000	201 (44.0)	23 (42.6)	51 (38.6)	127 (46.9)
≥10000	256 (56.0)	31 (57.4)	81 (61.4)	144 (53.1)
BMI, kg/m ²	24.0±3.5	24.3±3.3	24.2±3.1	23.9±3.7
Smoking status				
Current	8 (1.8)	42 (77.8)	98 (74.2)	218 (80.4)
Previous ^a	91 (19.9)	10 (18.5)	31 (23.5)	50 (18.5)
Never	358 (78.3)	2 (3.7)	3 (2.3)	3 (1.1)
MNA score, 0-14 range	12.9±1.6	13.3±1.0	13.2±1.1	12.7±1.8
≤ 11	64 (14.2)	4 (7.4)	12 (9.2)	48 (18.1)
Health Conditions				
Hypertension	320 (70.6)	35 (64.8)	88 (67.2)	197 (73.5)
CAD	227 (50.3)	23 (42.6)	60 (45.8)	144 (54.1)
COPD	82 (18.2)	3 (5.6)	27 (20.8)	52 (19.5)
Diabetes	132 (28.9)	10 (18.5)	33 (25.2)	89 (32.8)
CKD	45 (10.0)	6 (11.1)	12 (9.2)	27 (10.1)
Number of comorbidities				
0-1	10 (19.2)	20 (15.9)	24 (9.4)	10 (19.2)
2-4	32 (61.5)	62 (49.2)	124 (48.6)	32 (61.5)
≥5	10 (19.2)	44 (34.9)	107 (42)	10 (19.2)
Total SPPB score, 0-12 range	8.4±2.8	10.3±1.9	9.1±2.3	7.5±2.8
10-12	141 (39.9)	37 (71.2)	53 (48.6)	51 (26.6)
7-9	116 (32.9)	13 (25.0)	42 (38.5)	61 (31.8)
≤ 6	96 (27.2)	2 (3.8)	14 (12.8)	80 (41.7)
Gait speed, m/s	0.9±0.3	1.1±0.3	1.0±0.2	0.8±0.2

<1m/s	273 (67.9)	18 (34.0)	74 (57.8)	181 (81.9)
Chair stand test time, s	14.2±4.9	12.5±4.8	12.8±3.1	15.8±5.4
≥16.7s or incapable ^b	133 (35.2)	3 (5.9)	30 (28.6)	100 (45.0)
TUG time, s	12.9±4.9	11.4±5.4	12.1±4.0	14.0±5.3
≥14.0s	112 (33.0)	10 (19.6)	33 (26.8)	69 (41.8)
ADL≤80	48 (10.5)	1 (1.9)	0 (0.0)	47 (17.4)
IADL ≤7	284 (62.1)	18 (33.3)	67 (50.8)	199 (73.4)

Values are presented in mean ± standard deviation and N (%). BMI: Body Mass Index. MNA: Mini Nutritional Assessment. CAD: Coronary Artery Disease. COPD: Chronic Obstructive Pulmonary Disease. CKD: Chronic Kidney Disease. PFS: Pittsburgh Fatigability Score. SPPB: Short Physical Performance Battery. TUG: Timed Up and Go Test. Barthel index: Instrumental Activity of Daily Living (Lawton and Brody)^[16].

^a Defined as used to smoke and has quit smoking for more than 6 months.

^b “Incapable” means the participants did not complete the 5 times chair stand test for physical reasons or safety reasons.




Figure 1. Mean PFS physical scores by age strata and sex.

Table 2. Convergent Validity of the PFS Physical Scores -Simplified-Chinese version

Characteristics	Correlation Coefficient (r) ^a with the PFS Physical scores	P-Value
Total SPPB score	-0.51	< .001
Gait speed, m/s	-0.55	< .001
Chair stand test time, s	0.40	< .001
TUG, s	0.28	< .001
Barthel Index	-0.56	< .001
IADL	-0.48	< .001

SPPB: Short Physical Performance Battery. TUG: Timed Up and Go Test. ADL: Activity of Daily Living. IADL: Instrumental Activity of Daily Living.

^a Spearman's rank-order correlations

Table 3. Concurrent and convergent validity of the PFS Physical scores: Simplified-Chinese Version with physical performance and physical ability measures

Measures	LS means (SE) ^a		Difference		AUC (95% CI)
	Yes	No	Mean (SE)	P-Value	
Low SPPB ^b	27.0 (3.4)	19.4 (3.0)	7.7 (0.4)	< .001	0.77 (0.72, 0.81)
Slower gait speed ^c	26.9 (3.3)	19.0 (4.0)	7.8 (0.4)	< .001	0.76 (0.72, 0.80)
Poor lower limb strength ^d	26.8 (2.4)	24.6 (4.2)	2.3 (0.4)	< .001	0.78 (0.74, 0.82)
High TUG ^e	25.7 (4.2)	21.7 (4.1)	4.3 (0.9)	< .001	0.77 (0.72, 0.81)
ADL dependence ^f	35.1 (5.7)	24.4 (3.9)	10.7 (0.6)	< .001	0.87 (0.84, 0.90)
IADL dependence ^g	28.1 (4.1)	21.3 (3.3)	6.8 (0.4)	< .001	0.81 (0.77, 0.84)

LS means: Least squares means, SE: Standard Error, AUC: Area Under the Curve. SPPB: Short Physical Performance Battery. ADL: Activities of Daily Living. IADL: Instrumental Activities of Daily Living.

^a Adjusted for age and sex

^b Defined as SPPB <10 points

^c Defined as gait speed: gait speed <1.0 m/sec

^d Defined as the time to performed chair stand test ≥16.7 seconds or not able to do 5 stands

^e Defined as Timed Up and Go Test time ≥14.0 seconds

^f Defined as Barthel Index ≤80

^g Defined as IADL ≤7

Figures




Figure 1

Mean PFS physical scores by age strata and sex.