

Trends of Follow-Up Clinic Visits and Admissions Three-Months Before and During COVID-19 Pandemic at Tikur Anbessa Specialized Hospital, Ethiopia: an Interrupted Time Series Analysis

Workeabeba Abebe (✉ workeabebasol@gmail.com)

Addis Ababa University

Alemayehu Worku

Addis Ababa University

Tamirat Moges

Addis Ababa University

Nuhamin Tekle

Addis Ababa University

Wondowossen Amogne

Addis Ababa University

Tewodros Haile

Addis Ababa University

Desalew Mekonen

Addis Ababa University

Abebe Habtamu

Addis Ababa University

Wakgari Deressa

Addis Ababa University

Research Article

Keywords: Admission, COVID-19, Ethiopia, Follow-up visit, Health-care utilization, TASH, Trends.

Posted Date: January 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-135977/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Health Services Research on July 23rd, 2021. See the published version at <https://doi.org/10.1186/s12913-021-06730-8>.

1 **Trends of follow-up clinic visits and admissions three-months before and during COVID-19**
2 **pandemic at Tikur Anbessa specialized hospital, Ethiopia: an interrupted time series**
3 **analysis**

4 Workeabeba Abebe^{1*}, Alemayehu Worku², Tamirat Moges¹, Nuhamin Tekle³, Wondowossen
5 Amogne⁴, Tewodros Haile⁴, Desalew Mekonen⁴, Abebe Habtamu¹, Wakgari Deressa².

6

7

8

9 1. Department of Pediatrics and Child Health, College of Health Sciences, Addis Ababa
10 University, Addis Ababa, Ethiopia

11 2. School of Public Health, College of Health Sciences, Addis Ababa University, Addis
12 Ababa, Ethiopia

13 3. Department of Family Medicine, College of Health Sciences, Addis Ababa University,
14 Addis Ababa, Ethiopia

15 4. Department of Internal Medicine, College of Health Sciences, Addis Ababa University,
16 Addis Ababa, Ethiopia

17

18

19

20

21

22

23 * Corresponding author: Workeabeba Abebe (WA)

24 E-mail: workeabebasol@gmail.com

25 **Abstract**

26 **Background:** Following the first report of the COVID-19 case in Ethiopia on the 13th of
27 March 2020, the country adopted a lockdown policy to contain the spread of the virus.
28 Responding to the health-care burden imposed by the COVID-19 pandemic had to be coupled
29 with ensuring essential health care services. This study assessed the impact of COVID-19 on the
30 trends of non-COVID follow-up visits and admissions at Tikur Anbessa Specialized Hospital,
31 Addis Ababa, Ethiopia.

32 **Methods:** A retrospective, time-series study with the 1st case of COVID-19 report as a
33 reference time examined the trend in follow-up visits and admissions between December 1st, 2019
34 and May 31st, 2020. A comparison of health care utilization between December 2019 to May 2020
35 and its equivalent period in 2018/19 was also done. A data abstraction tool was used to collect
36 secondary data from the hospital's electronic medical recordings and logbooks of each unit.

37 **Results:** A total of 7,717 visits from eight follow-up clinics and 3,310 admissions were
38 collected during three months before the onset of COVID-19. During the following three months
39 after the pandemic, 4,597 visits and 2,383 admissions were collected. Overall, a 40.4% decrease
40 in follow-up visits and a 28% decline in admissions were observed during the COVID-19. The
41 drop in the daily follow-up visits was observed for both genders. The number of visits in all
42 follow-up clinics in 2019/2020 decreased when compared to the same months in
43 2018/19($p<0.05$). Follow-up visits were substantially lower for renal patients (-68%), patients
44 with neurologic problems (-53.9%), antiretroviral treatment clinics (-52.3%), cardiac patients
45 (-51.4%). Although pediatric emergency admission was significantly lower (-54.1%) from the

46 baseline ($p=0.04$), admissions from the general pediatric and adult wards did not show a
47 significant difference.

48 **Conclusions:** Significant decreases in-hospital follow-up clinic visits were observed during
49 the first months of the COVID-19 pandemic. Public health guidance on how best to access care,
50 more for patients with serious illnesses are required. Promoting self-care, alternatives health-care
51 services like home-based care, and phone clinics might be considered for patients with mild
52 symptoms. Further studies needed to track the long-term effect of the pandemic among non-
53 COVID-19 patients.

54 **Keywords:** Admission, COVID-19, Ethiopia, Follow-up visit, Health-care utilization, TASH,
55 Trends.

56

57

58

59

60

61

62

63

64

65

66

67

68

69 **Background**

70 In Ethiopia, the first case of the novel coronavirus disease 2019 (COVID-19) was reported on 13th
71 of March 2020. Ethiopia adopted a lockdown policy to contain the spread of the virus and schools
72 were closed on March 16, 2020 and travel bans were put in place on March 20, 2020.
73 Subsequently stay at home orders were recommended to the public and the country declared a
74 state of emergency on April 8, 2020 (1, 2). Despite efforts to prevent the spread of COVID-19 in
75 Ethiopia, new cases continued to emerge among individuals with travel history and even without
76 clear exposure. The community transmission has been already established and the virus has been
77 distributed to every part of the country. As of November 5th, 2020, the total number of confirmed
78 COVID-19 cases in Ethiopia was 98391, of which 1508 died and 57000 recovered (3, 4).
79 Although the virus was detected in all age groups, severe symptoms were reported mainly in older
80 age groups. People with pre-existing medical conditions such as hypertension, diabetes, and heart
81 disease appear to be more vulnerable to becoming severely ill with COVID-19 (5).

82 The COVID-19 pandemic continues to challenge the healthcare systems around the world.
83 Responding to the healthcare burden imposed by the pandemic required intense resources. Even
84 more, this had to be coupled with ensuring essential healthcare services (6). Countries struggled
85 to fairly allocate resources for these needs. Hospitals operated with less capacity due to the
86 distribution of resources for COVID-19 prevention, detection, and case management. Some of the
87 care provided in hospitals such as elective surgery and other non-critical medical services were
88 purposefully curtailed (7). In addition, following the initial public health messaging that
89 discouraged unnecessary healthcare use, many patients delayed or canceled follow-up
90 appointments in hospitals (8, 9). The low utilization of emergency services was also evident (10).
91 In Singapore, doctor visits decreased by up to 30% while a massive drop of 63.8% in pediatric

92 emergency health-care utilization was observed in Germany and a comparable 63.5% reduction in
93 emergency department visits were reported in the U.S. (11-13).

94 The delayed access or provision of care affected both acute and chronic care (14, 15), and resulted
95 in increased adverse outcomes from non-COVID-19-related illnesses (16, 17). The main reason
96 for the reduced hospital visits was fear of contracting COVID-19 in health-care settings and
97 restricted transports due to social distancing measures (18, 19). However, a reduction in some
98 acute infections that have a person-to-person transmission risk and illnesses related to
99 environmental exposure such as allergens and air pollution as well as traumas was also observed
100 in some settings (14, 20). Furthermore, the application of telemedicine created alternative access
101 to care; partly explaining the reduced number of in-person care (21).

102 In this study, we assessed the impact of COVID-19 on the trends in hospital visits and admissions
103 at Tikur Anbessa Specialized Hospital (TASH) in Ethiopia. We compared the rate of follow-up
104 clinic visits and admissions for the three months before and after the first report of the COVID-19
105 case in 2020. In addition, we compared the trends in hospital visits from December 2019 to May
106 2020 to the equivalent period in the previous year (Dec 2018 to May 2019).

107

108

109

110

111

112

113

114

115

116 **Methods**

117 **Setting**

118 This study was conducted at TASH located in Addis Ababa, Ethiopia. TASH serves as a public
119 institute training center with about 700 beds. The hospital serves a large population of patients
120 referred for outpatient and inpatient care in all core specialties (internal medicine, emergency,
121 critical care, surgery, pediatrics and obstetrics/gynecology). The hospital also provides specialized
122 and sub-specialized clinical services to the patients referred from other hospitals and health
123 institutions across all parts of the country. Of the total beds, about 150 are medical (including 12
124 intensive care unit [ICU] beds), 172 beds are in pediatrics (including 4 pediatric ICU beds). Both
125 adult and pediatric emergency rooms (ER) are open to patient's 24 hours a day and seven days a
126 week. Clinical care teams composed of nursing staff; medical students and residents supervised
127 by attending staff provide all care for in-patients. TASH started to have functional electronic
128 database recording system for most of the adult clinics since 2018; however, the health records
129 from pediatrics and some clinics from adult side were still paper-based recordings.

130 **Study design and period**

131 A retrospective time series study was conducted to collect daily follow-up visits, and monthly
132 admissions data at TASH between December 1st, 2019 and May 31st, 2020. A before-after study
133 was done to analyze the trend in daily follow-up visits and monthly admissions with the 1st case
134 of COVID-19 report (March 13th 2020) as a cut of point. We also compared health care utilization,
135 between December 2019 to May 2020 and its equivalent period in 2018/19. A quantitative
136 method using structured data abstraction tools was used to collect the data. Data abstraction was
137 conducted from May to June 30, 2020.

138 **Data collection and procedures**

139 A study team comprised of clinicians and public health experts developed a data abstraction tool
140 after reviewing the available literature. We collected secondary data during the selected periods
141 from the hospital's Health Management Information System (HMIS) and logbooks in each
142 outpatient clinic, emergency/casualty department, and wards of TASH. A one-day training to data
143 collectors was conducted on the data collection tools and procedures. Trained nurses from each
144 unit collected data that were collected as part of the routine hospital service. Eight outpatient
145 follow-up clinics (4 adult and 4 pediatric clinics) were included. These clinics were selected based
146 on the availability of the documentation of the required information. In addition, both adult and
147 pediatric inpatient wards were included. The main outcome variables were daily follow-up visits
148 and monthly hospital admissions of adult and pediatric wards.

149 **Inclusion/Exclusion**

150 Follow-up clinics visits, and admissions, between December 1st, 2019, and May 31st, 2020 as well
151 as previous year follow-up clinics visits, and admissions between December 1st, 2018 and May
152 31st 2019 were included in this study. Clinics with incomplete documentation were excluded

153 **Statistical analysis**

154 All data were entered into a password-secured database using Microsoft Excel. After cleaning the
155 data were exported to STATA version 14. We analyzed the trend of daily follow-up visits and
156 monthly hospital admissions. Data were summarized using graphs; mean (standard error of mean
157 [SEM])). Wilcoxon Signed-Ranks test was used to detect mean differences of 2018/19 and
158 2019/20 trend of the follow up visits and ward admission differences. P-value below 0.05 was
159 considered as significant.

160 **Ethical consideration**

161 The study was granted ethical approval from the Institutional Review Board (IRB) of the College
162 of Health Sciences, Addis Ababa University (Protocol Number:042/20/SPH). Informed
163 consent was waived by the IRB, which approved the study. All personnel involved in this study
164 trained in infection prevention and control procedures (standard contact, droplet or airborne
165 precautions). Data were completely anonymous. All methods were carried out in accordance with
166 relevant guidelines and regulations.

167 **Results**

168 Overall, 12,314 follow-up visits and 5,693 hospital admissions solely non-COVID-19 data were
169 collected. Taking 1st case of COVID-19 report in country as a cut of, 7,717 follow-up visits
170 happened before (3 months) and the rest 4,597 follow-up visits after (3 months). Separately,
171 3,310 hospital admissions before (3 months) and 2,383 admissions after (3 months) of the 1st case
172 of COVID-19 report was collected (**Table 1**). Similarly, 15, 583 follow-up visits and 6,639
173 hospital admissions solely non-COVID-19 data of last year of the same time also collected. The
174 decrease in follow-up clinic visits and admissions after the onset of COVID-19 was 40.4% and
175 28%, respectively. We observed variations in the number of visits and admissions across the
176 follow-up clinics and inpatient wards. Follow-up visits were substantially lower for renal patients
177 (-68% below baseline), patients with neurologic problems (-53.9%), Anti-retroviral treatment
178 clinics (-52.3%), cardiac patient (-51.4%). Although, pediatric emergency admission was
179 significantly lower (-54.1%) from the baseline, admissions from the general pediatric (-14.7%)
180 and adult (-6.9%) wards were not that much affected.

181

182 **Table 1. Number of follow-up visits and admissions before and during COVID-19, and**
 183 **percentage reduction, TASH, 2020**

Characteristics	Pre-COVID-19			Total	During-COVID-19			Total	*Percent (%) reduction
	Dec	Jan	Feb		Mar	Apr	May		
Follow-up clinics									
Neurology	599	643	467	1709	275	265	247	787	-53.9
Cardiac	511	618	422	1551	288	260	206	754	-51.4
Hema. - Oncology	481	525	582	1588	449	460	465	1374	-13.5
Pediatric - ART	296	337	264	897	127	152	149	428	-52.3
Endocrine	251	238	269	758	222	205	139	566	-25.3
HRIC	226	110	196	532	125	130	133	388	-27.1
Renal	123	128	124	375	38	40	42	120	-68.0
Gastro-intestinal	110	71	126	307	88	58	34	180	-41.4
2019/20 (Overall visit) Total	2597	2670	2450	7717	1612	1570	1415	4597	-40.4
2019/20 (Ped. ER admission)	406	421	398	1225	200	212	150	562	-54.1
2019/20 (Ped. Ward admission)	532	535	469	1536	460	480	370	1310	-14.7
2019/20 (Adult ward admission)	173	181	195	549	185	164	162	511	-6.9
Overall admission	1111	1137	1062	3310	845	856	682	2383	-28.0

184 *Percentage Changes in follow-up clinics visits & admission

185

186 In total, 131 observation (daily working days hospital follow up visits in six months) occurred at
 187 TASH during the analyzed period with 66 before and 65 after first COVID-19 case seen in
 188 Ethiopia. We described the daily mean and standard error of mean separately for male and female
 189 who visited outpatient clinics three months before and after the first report of COVID-19 case
 190 reported in Ethiopia. There were statistically significant declines in the number of daily visits in
 191 all follow-up clinics except the Endocrine and GI clinics (**Table 2**).

192

193

194 **Table 2. Comparison of number of outpatient visits by gender before and after the first case**
 195 **of COVID-19, TASH, 2020**

Clinics	Average daily visits 3months before 1 st COVID case		Average 3months Daily (±SEM)	Average daily visits 3months after 1 st COVID		P-value
	Daily (±SEM)	mean 95% CI		Daily (±SEM)	mean 95% CI	
*ART clinic						
Male	13.8 (0.7)	12.3-15.2	12.2(0.8)	10.5-13.8	0.07	
Female	23.6(1.0)	21.5-25.6	19.7(1.3)	17.1-22.3	0.01	
Cardiac clinic						
Male	24.5(2.4)	19.8-29.3	11.8(1.7)	8.3-15.2	0.00	
Female	37.2 (3.6)	30.0-44.6	17.1(2.4)	12.3-21.8	0.00	
Chest clinic						
Male	13.2 (1.5)	10.1-16.2	7.0(1.0)	5.0-9.0	0.00	
Female	12.8 (1.6)	9.6-15.9	8.5(1.3)	6.0-11.0	0.02	
Endocrine clinic						
Male	19.0 (3.1)	12.7-25.3	15.1(2.5)	9.9-20.1	0.16	
Female	23.9 (3.9)	16.0-31.7	17.5(2.9)	11.6-23.4	0.09	
# GI clinic						
Male	2.0(0.4)	1.3-2.8	1.5(0.3)	0.8-2.1	0.13	
Female	1.2(0.2)	0.8-1.7	0.7(0.2)	0.4-1.1	0.05	
\$ HRIC clinic						
Male	3.3 (0.2)	2.9-3.8	2.0(0.2)	1.7-2.3	0.00	
Female	2.8 (0.2)	2.3-3.2	1.9(0.2)	1.6-2.4	0.00	
Neurology clinic						
Male	18.7 (1.0)	16.7-20.7	10.7(0.7)	9.3-12.1	0.00	
Female	11.8 (0.7)	10.5-13.2	6.2(0.4)	5.3-7.0	0.00	

196 ***ART=Anti Retroviral Treatment, #GI=Gastro-Intestinal, \$HRIC=High Risk Infant Clinic**

197

198 The trend for six-month daily attendance by male and female patients from two follow-up clinics
 199 (Cardiac and ART) at TASH is illustrated in (**Fig 1**). The clinics were selected since these clinics
 200 had daily follow-up throughout the week over the observation period. There was a sharp drop in
 201 the daily hospital visits for both genders after mid-March, when the first COVID-19 case was
 202 reported. Hospital visit did not rebound to the baseline observation period by the end of May
 203 2020.

204

205 **Figure 1. Trends in the number of daily follow-up visits from selected clinics by gender from**
206 **December 2019 to May 2020**

207 **Figure 2** shows the total number of follow-up visits and admissions for six months of Dec 2019-
208 May 2020 compared with the same period of last year (2018/19). The number of visits at the
209 follow up clinics decreased when compared to the same time period in 2018/19. Ward admissions
210 at TASH decreased after the month of February, however the trend is not similar with the follow-
211 up clinics.

212

213 **Figure 2. Trends in the number of daily follow-up visits from selected clinics by gender from**
214 **December 2019 to May 2020**

215 **Figure 3** depicted follow up clinics visits separately. In 2019/20, the number of visits at the
216 follow up clinics decreased when compared to the same time period in 2018/19. The relative
217 reduction in monthly follow-up visits was observed in all clinics, except endocrine and
218 hematology- oncology clinics.

219

220 **Figure 3: Monthly follow-up visits, December 2019 to May 2020 relative to equivalent**
221 **period in 2018/19.**

222 The trend of admissions in the pediatric emergency between December 1st, 2019, and May 31st,
223 2020 is illustrated in **Figure 4**. The monthly report of the number of children who were admitted
224 to the pediatrics Emergency room (ER) showed a significant drop in the month of March and
225 there was no rebound in the month of May (**p=0.04**).

226

227 **Figure 4: Pediatric ER Admission, December 2019 to May 2020 relative to equivalent period**
228 **in 2018/19.**

229 **Figure 5a and 5b** illustrate the monthly pediatrics and adult ward admissions and deaths
230 respectively. Pediatrics ward admissions at TASH decreased since the month of February, with
231 the highest drop observed in the month of May 2020. However the difference was not statistically
232 significant ($p=0.34$). Adult medical admissions did not show any difference in 2020 compared to
233 the previous year ($P=0.12$).

234

235 **Figure 5a**

Figure 5b

236 **Figure 5a and 5b: Pediatrics and adult ward admissions, December 2019 to May 2020**
237 **relative to equivalent period in 2018/19.**

238 **Discussion**

239 This study demonstrated a significant decrease in follow-up clinics visits and emergency
240 admissions during the first three months of the COVID-19 pandemic in a tertiary care center in
241 Ethiopia. Furthermore, the case numbers in follow-up clinics during the observed period
242 decreased when compared to the same time in 2018/19. However, the number of children who
243 were admitted to the pediatrics and adult wards didn't show significant difference in the months
244 following March.

245 The decline in the number of overall visits was consistent with the reports from a study in the
246 U.S. where a decline of 0.7 million non- COVID admissions in April 2020 compared to April
247 2019 was demonstrated in hospitals across the nation (20). The decline in non-COVID admissions
248 ranged from 39.5% to 50% in the minimally affected to greatly affected regions respectively.

249 Similar studies revealed that the decrease was observed in both primary health care and
250 emergency department visits (17, 22). Moreover, these findings were true for pediatric and adult
251 populations across specialty clinics (23-25).

252 Ethiopia, compared to other countries that had significant surge of COVID-19 cases in the first
253 months of 2020, reported fewer cases. However, as a measure of prevention of the spread of the
254 COVID-19 virus in the hospital and effective case management, hospitals were designated as
255 COVID-19 and non-COVID-19 case management centers. TASH was among the designated non-
256 COVID-19 case management centers, patients who were suspected to have COVID-19, were
257 isolated in a ward and those whose tests confirmed the diagnosis of COVID-19 were transferred
258 out to COVID-19 centers. However, considering the overcrowding and the limited use of personal
259 protective equipment at TASH the anxiety associated with being at an increased risk in this
260 setting was understandable. The fear of contracting COVID-19 in the health-care settings was the
261 commonest explanation for the drop in hospital visits (7, 13). Moreover, the testing capacity at
262 institutions and in the country at large was poor to support prevention of transmission. The
263 communication of alarming medical information to the public and the announcement of stay at
264 home order, have also impacted health care seeking behavior.

265 Being a referral hospital, it was believed that patients who were on follow-up and those visiting
266 the emergency department of TASH often have a real need that requires hospital management. In
267 such situations, people opted for a conservative, ‘watch and wait’ approach (7) while some might
268 have resorted to traditional medicine, as it is a common cultural alternative in Ethiopia. The delay
269 in seeking appropriate health care will have deleterious health outcomes (12). Few outpatient
270 departments had phone clinics that provided an alternative for accessing health care. A
271 compelling contrary argument for the decreased hospital visit was a reduced incidence of illnesses

272 such as community-acquired pneumonia, which was a reason for most of the pediatric ED visits
273 and in adult patients with co-morbidities (13, 23).

274 Looking at the number of children who were admitted to the pediatrics and adult wards showed
275 overall decline –28, which is lower with the report from U.S where all medical admissions in
276 April declined by 34.1 percent (95% CI: –34.6, –33.6) (20). We can still consider the drop in
277 admission is significant since TASH is a big hospital/ward where usually all beds were occupied.
278 The impact of COVID-19 on health care resource is huge in the world and more so in low-income
279 countries. Ethiopia being among the low-income countries struggles to adequately allocate human
280 resource, support community-based testing capacity, and provide equipment needed to manage
281 COVID- 19 cases. It is wise to consider that the attempt to respond to the COVID-19 pandemic
282 has compromised the attention given for non-COVID illnesses, which might be a reason for
283 admission declines to reserve beds for possible anticipated surge (13).

284 **Limitations**

285 We abstracted a secondary data, and the data were not crosschecked. We took data from the
286 clinics not from individuals therefore we did not do subgroup analysis except by sex. The data in
287 the hospital was not collected for scientific purposes and poor documentation was the reality in
288 our setting so there may be cases that were not documented (missed). TASH is a referral center,
289 and the service was structured according to sub-specialty/specialty clinics so these findings might
290 not be generalized to primary health care and other health care systems.

291 **Conclusion**

292 A significant decrease in follow-up clinics visit and emergency admissions was observed during
293 the first months of the COVID-19 pandemic. Appropriate public health guidance on how best to
294 access care emphasizing on the importance of continuing to visit the health-care facilities for

295 serious illnesses is required. Promoting of self-care, implementing of home care and
296 strengthening of phone clinics might be considered for patient with mild symptoms. Further
297 studies needed to track the long-term effect of the pandemic among non-COVID-19 patients.

298 **Declarations**

299 **Consent for publication**

300 Not applicable

301 **Availability of data and material:** The datasets used and/or analyzed during the current study
302 are available from the corresponding author on reasonable request.

303 **Competing interests:** We declare that we have no competing interest

304 **Funding**

305 Addis Ababa University (AAU) funded this study.

306 **Author Contributions**

307 WA, AW, W. Amogne and WD conceived and designed the study. WA, TM, TH, DM and AH
308 supervised data collection. WA and NT prepared the final dataset for analysis. WA and AW
309 analyzed and interpreted the data. WA and NT drafted the manuscript. WD, AW and TM
310 provided major roles in revising the manuscript. All authors read and approved the final
311 manuscript.

312 **Acknowledgments**

313 The authors are grateful to the research staff at the College of Health Sciences. The authors would
314 also like to thank the data collectors and study participants for their time and contributing to the
315 research.

316

317 **References**

- 318 1. Ethiopian Public Health Institute. National Public Health Emergency Operation Center.
319 COVID-19 Pandemic Preparedness and Response. Bulletin No. 1, May 03, 2020. Addis
320 Ababa, Ethiopia.
- 321 2. Ethiopian Public Health Institute. National Public Health Emergency Operation Center.
322 COVID-19 Pandemic Preparedness and Response. Bulletin No. 7, June 15, 2020. Addis
323 Ababa, Ethiopia.
- 324 3. Ethiopian Public Health Institute. National Public Health Emergency Operation Center.
325 COVID-19 Pandemic Preparedness and Response. Bulletin No. 26, October 26, 2020. Addis
326 Ababa, Ethiopia.
- 327 4. Ethiopian Public Health Institute. National Public Health Emergency Operation Center.
328 COVID-19 Pandemic Preparedness and Response. Bulletin No. 27, Nov 5, 2020. Addis
329 Ababa, Ethiopia.
- 330 5. World Health Organization. Coronavirus Diseases (COVID-19) Situation Reports-World,
331 October 2020
- 332 6. Taquechel K, Diwadkar AR, Sayed S, Dudley JW, Grundmeier RW, Kenyon CC, et al.
333 Pediatric Asthma Health Care Utilization, Viral Testing, and Air Pollution Changes During
334 the COVID-19 Pandemic. *J Allergy Clin Immunol Pract.* 2020; 8(10):3378-3387.e11. Doi:
335 10.1016/j.jaip.2020.07.057.
- 336 7. Dopfer C, Wetzke M, Scharff AZ, Mueller F, Dressler F, Baumann U, et al. COVID-19
337 related reduction in pediatric emergency healthcare utilization—a concerning trend. *BMC*
338 *Pediatrics.* 2020 Dec; 20(1):1-0. <https://doi.org/10.1186/s12887-020-02303-6>

- 339 8. Emanuel EJ, Persad G, Upshur R, Thome B, Parker M, Glickman A, et al. Fair allocation of
340 scarce medical resources in the time of Covid-19. *N Engl J Med* 2020; 382:2049-2055. DOI:
341 10.1056/NEJMs2005114
- 342 9. Kraemer MU, Yang CH, Gutierrez B, Wu CH, Klein B, Pigott DM, et al. The effect of human
343 mobility and control measures on the COVID-19 epidemic in China. *Science*. Science 2020
344 May 1;368(6490):493-7.
345 DOI: 10.1126/science.abb4218
- 346 10. Wamsley CE, Kramer A, Kenkel JM, Amirlak B. Trends and challenges of telehealth in an
347 academic institution: the unforeseen benefits of the COVID-19 global pandemic. *Aesthetic*
348 *Surgery Journal*. 2020 Jul 22. <https://doi.org/10.1093/asj/sjaa212>
- 349 11. Ahn, SangNam and Kim, Seonghoon and Koh, Kanghyock, Changes in Healthcare
350 Utilization, Spending, and Perceived Health during COVID-19: A Longitudinal Study from
351 Singapore (August 7, 2020). Available at
352 SSRN: <https://ssrn.com/abstract=3669090> or <http://dx.doi.org/10.2139/ssrn.3669090>
- 353 12. Lazzerini M, Barbi E, Apicella A, Marchetti F, Cardinale F, Trobia G. Delayed access or
354 provision of care in Italy resulting from fear of COVID-19. *Lancet Child Adolesc Health*.
355 2020 May 1;4(5):e10-1DOI:[https://doi.org/10.1016/S2352-4642\(20\)30108-5](https://doi.org/10.1016/S2352-4642(20)30108-5)
- 356 13. Jeffery MM, D'Onofrio G, Paek H, Platts-Mills TF, Soares WE, Hoppe JA, et al. Trends in
357 emergency department visits and hospital admissions in health care systems in 5 states in the
358 first months of the COVID-19 pandemic in the US. *JAMA Intern Med*. 2020;180(10):1328-
359 1333. doi:10.1001/jamainternmed.2020.3288
- 360 14. Elmore JG, Wang PC, Kerr KF, Schriger DL, Morrison DE, Brookmeyer R, et al. Excess
361 Patient Visits for Cough and Pulmonary Disease at a Large US Health System in the Months

- 362 Prior to the COVID-19 Pandemic: Time-Series Analysis. *J Med Internet Res*
363 2020;22(9):e21562 [doi:10.2196/21562](https://doi.org/10.2196/21562)
- 364 15. Marijon E, Karam N, Jost D, Perrot D, Frattini B, Derkenne C, et al. Out-of-hospital cardiac
365 arrest during the COVID-19 pandemic in Paris, France: a population-based, observational
366 study. *Lancet Public Health*. 2020 May 27. [https://doi.org/10.1016/S2468-2667\(20\)30117-1](https://doi.org/10.1016/S2468-2667(20)30117-1)
- 367 16. Lynn RM, Avis JL, Lenton S, Amin-Chowdhury Z, Ladhani SN. Delayed access to care and
368 late presentations in children during the COVID-19 pandemic: a snapshot survey of 4075
369 paediatricians in the UK and Ireland. *Archives of Disease in Childhood*. 2020 Jun 24.
370 <http://dx.doi.org/10.1136/archdischild-2020-319848>
- 371 17. Tam CC, Cheung KS, Lam S, Wong A, Yung A, Sze M, et al. Impact of coronavirus disease
372 2019 (COVID-19) outbreak on ST-segment–elevation myocardial infarction care in Hong
373 Kong, China. *Circ Cardiovasc Qual Outcomes*. 2020 Apr;13(4):e006631.
374 <https://doi.org/10.1161/CIRCOUTCOMES.120.006631>
- 375 18. Khairat S, Meng C, Xu Y, Edson B, Gianforcaro R. Interpreting COVID-19 and virtual care
376 trends: cohort study. *JMIR Public Health Surveill* 2020;6(2):e18811 [doi:10.2196/18811](https://doi.org/10.2196/18811)
- 377 19. Lazzerini M, Barbi E, Apicella A, Marchetti F, Cardinale F, Trobia G. Delayed access or
378 provision of care in Italy resulting from fear of COVID-19. *Lancet Child Adolesc Health*.
379 2020 May 1;4(5):e10-DOI:[https://doi.org/10.1016/S2352-4642\(20\)30108-5](https://doi.org/10.1016/S2352-4642(20)30108-5)
- 380 20. Birkmeyer JD, Barnato A, Birkmeyer N, Bessler R, Skinner J. The Impact Of The COVID-19
381 Pandemic On Hospital Admissions In The United States: Study examines trends in US
382 hospital admissions during the COVID-19 pandemic. *Health Affairs*. 2020 Nov 1:10-377.
383 <https://doi.org/10.1377/hlthaff.2020.00980>
- 384 21. Mulholland RH, Wood R, Stagg HR, Fischbacher C, Villacampa J, Simpson CR, et al. Impact
385 of COVID-19 on accident and emergency attendances and emergency and planned hospital

386 admissions in Scotland: an interrupted time-series analysis. *J RSocMed*. 2020 Oct
387 4:0141076820962447. <https://doi.org/10.1177/0141076820962447>

388 22. Feral-Pierssens AL, Claret PG, Chouihed T. Collateral damage of the COVID-19 outbreak:
389 expression of concern. *Eur J Emerg Med* 2020;27(4):233. [Eur J Emerg
390 Med10.1097/MEJ.0000000000000717](https://doi.org/10.1097/MEJ.0000000000000717)

391 23. Chaiyachati BH, Agawu A, Zorc JJ, Balamuth F. Trends in Pediatric Emergency Department
392 Utilization after Institution of Coronavirus Disease-19 Mandatory Social Distancing. *J Pediat*.
393 2020 Jul 20. doi: [10.1016/j.jpeds.2020.07.048](https://doi.org/10.1016/j.jpeds.2020.07.048)

394 24. Gonçalves-Pinho M, Mota P, Ribeiro J, Macedo S, Freitas A. The Impact of COVID-19
395 Pandemic on Psychiatric Emergency Department Visits—A Descriptive Study. *Psychiatric
396 Quarterly*. 2020 Aug 25:1-1. <https://doi.org/10.1007/s11126-020-09837-z>

397 25. Kenyon CC, Hill DA, Henrickson SE, Bryant-Stephens TC, Zorc JJ. Initial effects of the
398 COVID-19 pandemic on pediatric asthma emergency department utilization. *J Allergy Clin
399 ImmunolPract*. 2020 Sep 1;8(8): 2774-6. <https://doi.org/10.1016/j.jaip.2020.05.045>

400

Figures




Figure 1

Trends in the number of daily follow-up visits from selected clinics by gender from December 2019 to May 2020 Figure 2 shows the total number of follow-up visits and admissions for six months of Dec 2019- May 2020 compared with the same period of last year (2018/19). The number of visits at the follow up clinics decreased when compared to the same time period in 2018/19. Ward admissions at TASH decreased after the month of February, however the trend is not similar with the follow up clinics.




Figure 2

Trends in the number of daily follow-up visits from selected clinics by gender from December 2019 to May 2020 Figure 3 depicted follow up clinics visits separately. In 2019/20, the number of visits at the follow up clinics decreased when compared to the same time period in 2018/19. The relative reduction in monthly follow-up visits was observed in all clinics, except endocrine and hematology- oncology clinics.




Figure 3

Monthly follow-up visits, December 2019 to May 2020 relative to equivalent period in 2018/19. The trend of admissions in the pediatric emergency between December 1st, 2019, and May 31st, 2020 is illustrated in Figure 4. The monthly report of the number of children who were admitted to the pediatrics Emergency room (ER) showed a significant drop in the month of March and there was no rebound in the month of May ($p=0.04$).




Figure 4

Pediatric ER Admission, December 2019 to May 2020 relative to equivalent period in 2018/19. Figure 5a and 5b illustrate the monthly pediatrics and adult ward admissions and deaths respectively. Pediatrics ward admissions at TASH decreased since the month of February, with the highest drop observed in the month of May 2020. However the difference was not statistically significant ($p=0.34$). Adult medical admissions did not show any difference in 2020 compared to the previous year ($P=0.12$).





Figure 5

Figure 5a and 5b: Pediatrics and adult ward admissions, December 2019 to May 2020 relative to equivalent period in 2018/19.