A Canadian Survey of Medical Students and Undergraduate Deans on Managing Patients With Obesity

Nathan J. Katz

Queen's University

Olivia Lovrics

McMaster University

Boris Zevin (✉ bz15@queensu.ca)

Queen's University

Research Article

Keywords: obesity, medical education, medical student, curriculum, survey

Posted Date: March 7th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1363401/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: With over 26% of Canadian adults living with obesity, undergraduate medical education (UGME) should prepare medical students to manage this chronic disease. It is currently unknown how management of obesity is taught within UGME curricula in Canada. This study (1) examined the knowledge and self-reported competence of final-year medical students in managing patients with obesity, and (2) explored how this topic is taught within UGME curricula in Canada.

Methods: We distributed two online surveys: one to final-year medical students, and another to UGME deans at 9 English-speaking medical schools in Canada. The medical student survey assessed students' knowledge and self-reported competence in managing patients with obesity. The dean's survey assessed how management of patients with obesity is taught within the UGME curriculum.

Results: 133 (6.9%) and 180 (9.3%) out of 1936 eligible students completed the knowledge and self-reported competence parts of the survey, respectively. Mean knowledge score was 10.5 (2.1) out of 18. Students had greatest knowledge about etiology of obesity and goals of treatment, and poorest knowledge about physiology and maintenance of weight loss. Mean self-reported competence score was 2.5 (0.86) out of 4. Students felt most competent assessing diet for unhealthy behaviors and calculating body mass index. Five (56%) out of 9 deans completed the survey. A mean of 14.6 (5.0) curricular hours were spent on teaching management of obesity.

Conclusions: Canadian medical students lack adequate knowledge and feel inadequately prepared to manage patients with obesity. Changes to UGME curricula may help address this gap in education.

1. Background

Obesity, defined as a body mass index (BMI) $\geq 30\text{kg/m}^2$, is a public health concern in Canada.^{1,2} Common obesity-related comorbidities such as hypertension, type II diabetes, and cancer, decrease quality of life and increase mortality.^{3,4} There is emerging evidence linking obesity and poor outcomes for patients with COVID-19,^{5,6} although the association between obesity and mortality from COVID-19 remains unclear.⁷ Despite rising rates of obesity in Canada, physicians in primary-care feel poorly prepared to manage patients with obesity.^{8,9} This problem, however, is not unique to primary care; physicians in various specialties also report the same challenges.^{10–14} As such, it is plausible that educational content regarding management of patients with obesity may be best delivered during undergraduate medical education (UGME), rather than being reserved for postgraduate medical education. In fact, there is consensus in the literature that medical students should be taught approaches to management of patients with obesity;^{14–21} however, there is a lack of understanding on how this topic is currently being taught at the UGME level in Canada and whether graduating students are competent in managing patients with obesity.^{22–24} A recent survey of American medical schools demonstrated a paucity of formal education around management of patients with obesity,²⁵ while a Norwegian study found obesity-related knowledge to be inadequate in Norwegian medical students.²⁶

Our objectives with this study were to: (1) examine the knowledge and self-reported competence of final-year Canadian medical students in managing patients with obesity, and (2) explore how management of obesity is currently taught within the UGME curricula in Canada.

2. Methods

2.1. Subjects

We recruited final-year medical students and UGME deans from English-speaking medical schools in Canada. We defined final-year medical students as students who were graduating at the end of the academic year when the survey was distributed. We defined the UGME deans as the dean and/or their delegate.

2.2. Study design

We administered a cross-sectional survey between August 2019 and November 2020. The study was approved by our institution's Health Sciences & Affiliated Teaching Hospitals Research Ethics Board, and all methods were performed in accordance with the relevant guidelines and regulations [see Additional File 1 for complete study protocol, and Additional File 2 for the Checklist for Reporting Results of Internet E-Surveys (CHERRIES)].

2.3. Survey design

2.3.1. Medical students' survey

We designed this survey in two parts. Part one examined medical students' knowledge and part two examined self-reported competence in managing patients living with obesity. We adapted the knowledge part of the survey from the 2018 study by Martins and Norsett-Carr,²⁶ which consisted of 18 multiple-choice style questions. Questions were grouped into 8 domains: Physiology, Etiology, Diagnosis, Goals for Obesity treatment, Conservative Treatment of Obesity, Surgery, Consequences of Obesity, and Weight-Loss Maintenance (see Additional File 3). We adapted the 15-item questionnaire from Jay et al¹³ to explore the self-reported competence of students in managing patients living with obesity. This questionnaire used a 5As construct to explore students' ability to Assess, Advise, Agree, Assist, and Arrange management of patients living with obesity (see Additional File 4). Each self-reported competency was measured on a 4-point scale: 1) "Know very little about and not able to perform"; 2) "Know something about and somewhat able to perform"; 3) "Able to perform well"; 4) "Able to teach others how to perform".

We offered medical students, who participated in the study, the option to request personalized feedback on their answers for the knowledge part of the survey. This was done after survey completion to provide students with an opportunity for self-reflection and learning. The personalized feedback included evidence-based explanations for each correct and incorrect answer (see Additional File 5).

We piloted the survey with five final-year medical students at our institution in July of 2019. We did not collect demographic data, such as the name of medical school or the length of their medical program, to ensure anonymity of participants' responses.

2.3.2. UGME deans' survey

We designed the UGME dean's survey to explore how management of patients living with obesity is currently taught within the UGME curriculum of each school. The survey included 7-items that explored (1) number of course hours dedicated to the teaching of management of patients with obesity; (2) modalities used to teach this content (e.g., lecture, group learning, community-based learning); and (3) presence of any programs with a

specific focus on the management of patients with obesity (see Additional File 6). We piloted the dean's survey at our institution with the local UGME deans' office.

We created both the medical students' survey and the UGME deans' survey in Qualtrics (Qualtrics, Provo, UT).

2.4. Survey distribution and data collection

We contacted the UGME offices of the 14 English-based medical schools in Canada inviting them to participate in our study. Nine out of 14 medical schools agreed to proceed with the local research ethics board (REB) application to allow medical students from that school to participate in our study. One school rejected our request for participation, and 4 schools did not respond to our multiple requests for participation. Once the REB approval at each medical school was granted, we distributed the survey electronically to all final-year medical students in that school and to their UGME dean's office. We embedded the letter of information and consent material at the start of the survey. Participation was voluntary; however, we offered one \$200 CAD lottery prize to medical students as incentive to participate.

We also sent an invitation to participate in the study directly to medical students in Ontario, Canada, via an email from the Ontario Medical Students Association on October 10, 2020. The Ontario Medical Students Association has student representatives from 6 English-speaking schools in Canada. This invitation aimed to recruit students who had not had a chance to participate.

2.5. Data analysis

We scored the medical students' survey out of 18 points for the questions assessing knowledge (minimum score 0, maximum score 18), and out of 60 points for the questions assessing self-reported competence (15 questions, each scored 1, 2, 3, or 4; minimum score 15, maximum score 60). We performed a narrative analysis of the results of the dean's survey. We performed statistical analysis in Excel 2007. We conducted a Spearman's rank-order correlation to explore whether there was an association between students' knowledge scores and self-reported competence scores. We report descriptive statistics as number and percentage (categorical variables) or as mean and standard deviation (continuous variables).

3. Results

3.1. Medical students' survey

A total of 207 out of 1936 (10.7%) medical students across 9 English-speaking Canadian medical schools agreed to participate in the study. Of these, 180 students (9.3%) completed the competence part of the survey, and 133 students (6.9%) completed both knowledge and competence parts of the survey. Twenty-seven out of 207 students started the survey but did not answer any questions.

3.1.1. Knowledge regarding management of patients living with obesity

Mean score for knowledge was 10.5 out of 18 (58%, SD = 2.1, n = 133). The number of students with correct responses for each question is reported in Fig. 1. Average correct answers by knowledge domain are reported in Fig. 2. A total of 106 out of 133 (80%) students opted to receive the personalized feedback for their answers.

3.1.2. Self-reported competence in managing patients living with obesity

The overall mean competence score was 2.5 (0.9) out of 4 for the 180 students who responded to this part of the survey. Mean scores for each question are listed in Table 1. Students rated themselves as 3 or above (“able to perform well” or “able to teach others how to perform”) only on two competencies: “Determining the body mass index (BMI) from weight and height measurements” ($M = 3.59$, $SD = 0.63$) and “Assessing diet for common unhealthy behaviors associated with obesity” ($M = 3.00$, $SD = 0.74$). For the remaining 13 out of 15 competencies, students rated themselves as “knowing something about” or “knowing very little about.” Notably, for the competency of “Responding to a patient’s questions regarding treatment options including behavior change, medications, and surgery” only 4 students (2%) rated themselves as “able to teach others how to perform” and only 44 students (24%) rated themselves as “able to perform well” (see Fig. 3).

Table 1

Medical Student self-perceived competence questionnaire

Questions. I can...	M (SD)
Determine body mass index (BMI) from weight and height measurements	3.59 (0.63)
Assess diet for common unhealthy behaviours associated with obesity (e.g. sweetened beverages, nutritional quality of snacks, frequent meals from fast food restaurants, etc.)	3.00 (0.74)
Assess current level of physical activity and provide guidance for setting physical activity goals for optimal health	2.90 (0.79)
Prescribe plan for exercise / physical activity	2.60 (0.87)
Ascertain each patient's readiness and ability to work on weight loss according to health beliefs and stage of change	2.52 (0.76)
Take a targeted history and conduct a physical examination to identify common co-morbidities (e.g. arthritis, diabetes, PCOS...)	2.44 (0.76)
Discuss the effect of obesity on present and future health and personalize risk to each patient	2.40 (0.74)
Assist patient in setting realistic goals for weight loss based on making permanent lifestyle changes	2.38 (0.82)
Recognize and refer patients with eating disorders	2.34 (0.73)
Recognize and screen for common psychosocial problems in obese patients including depression, emotional eating, and binge eating	2.34 (0.77)
Collaborate with registered dietitians and refer to community nutrition resources when appropriate	2.34 (0.84)
Use motivational interviewing to change behaviour	2.32 (0.83)
Use 24-hour recall, food record, or food frequency to obtain diet history	2.24 (0.83)
Provide brief counseling intervention to help patient lose weight	2.18 (0.81)
Respond to a patient's questions regarding treatment options including behaviour change, medications, and surgery	2.11 (0.71)
Legend: Scores show Mean (M) and Standard Deviation (SD) for all responses (n = 180). Questions were scored on a 1–4 likert scale: 1 – “Know very little about and not able to perform”; 2 – “Know something about and somewhat able to perform”; 3 - “Able to perform well”; 4 – “Able to teach others how to perform”. Questionnaire adapted from Do Internists, Pediatricians, and Psychiatrists Feel Competent in Obesity Care?, by Jay et al. (2008).	

Two questions focused on examining student's self-reported competence in assessing a patient's diet. Question 1 referred to using a standardized 24-hour food frequency questionnaire (FFQ), and Question 3 explored the

topic of diet more broadly by asking students about their ability to assess for common unhealthy behaviors associated with obesity. Thirty two percent of students reported being “able to perform well” and/or “teach others how to perform” the FFQ, while 76% of students reported being “able to perform well” and/or “teach others how to perform” an assessment of diet for unhealthy behaviors.

There was no significant correlation between students’ knowledge and competence in managing patients living with obesity ($r_s(131) = 0.027$, $p = 0.754$).

3.2. UGME dean’s survey

A total of 8 of 14 UGME deans’ offices agreed to participate in the survey; however, only 5 (36%) completed the survey (see Table 2). Four of 5 schools provided an estimate for the average number of curricular hours in undergraduate medical education dedicated to teaching students about the physiology, diagnosis, and management of patients with obesity; the average of these values was 14.6 (SD = 5.0) hours over 3–4 years of medical school. All schools reported using multiple instructional modalities to deliver this content across at least two school years. Only one school used a non-traditional teaching method, which they described as a half-day nutrition and obesity program conducted at a grocery store by a chef and Registered Dietitian (School 1).

Table 2
Results of the Dean's survey (n = 5)

	Total estimated hours teaching management of patients with obesity	Instruction Modalities Used	Domains of Obesity Addressed	Use of Non-Traditional Teaching Methods	Offer Dedicated Rotation in Clerkship focused on the Management of Patients with Obesity	Years in Which Management of Patients with Obesity is Taught
School 1	21.5	LGS, SGS, PBL, Clinical Simulation, Clinical Activity	Physical Activity, Nutrition, Behavioural Therapy, Pharmacotherapy, Surgery	Yes*	No	Years 1,2,3
School 2	12	LGS, SGS, Clinical Activity, Community-Based	Physical Activity, Nutrition, Pharmacotherapy, Surgery	None	No	Years 1,2
School 3	15	LGS, SGS, Clinical Simulation, Clinical Activity	Physical Activity, Nutrition, Behavioural Therapy, Pharmacotherapy, Surgery	None	No	Years 1,2,3,4
School 4	10	LGS, SGS, Clinical Activity, Community-Based	Nutrition, Surgery	None	No	Years 2,3,4
School 5	-	LGS, Clinical Activity	Physical Activity, Nutrition, Behavioural Therapy, Pharmacotherapy, Surgery	None	No	Years 1,2,3
Legend: LGS: large-group session, SGS: small-group session, PBL: problem-based learning. *In year 1, students participate in a hands-on cooking class, scavenger hunt-style grocery store tour, and self-directed in-store budgeting activity; facilitated by Registered Dietitians and Chefs. Goal: increase skills and confidence in managing eating habits, nutrition.						

4. Discussion

4.1. Summary of results

Our survey of English-speaking final-year medical students in Canada demonstrated limited knowledge and inadequate self-reported competence in managing patients with obesity. Our survey of UGME deans in Canada demonstrated an average of 14.6 curricular hours dedicated to teaching management of patients with obesity across all years of medical school, with multiple instructional modalities used to deliver this content. Medical students were most knowledgeable on the etiology of obesity and goals of obesity treatment, and least knowledgeable on the physiology of obesity and long-term weight loss maintenance. We found no significant correlation between medical students' knowledge scores and their perceived competence in managing patients with obesity. Together, these findings suggest a gap in UGME curricula regarding management of patients with obesity.

4.2. Explanation of findings

Our findings for the knowledge scores of Canadian students are consistent with those of Martins and Norsett-Carr in Norwegian students.²⁶ Canadian and Norwegian students demonstrated similar strengths and weaknesses, including greatest knowledge about the etiology of obesity and poorest knowledge about weight loss maintenance. It is encouraging that knowledge of the etiology of obesity was a relative strength among students, since this is associated with decreased weight bias and improved clinical interactions among practicing physicians.^{18,27,28} The gaps in knowledge domains identified in Fig. 2 can be used as a guide by medical educators to develop future educational interventions aimed at management of patients with obesity. It is promising that most students (80%) opted to receive personalized feedback, which suggests that survey research like ours on underrepresented topics may offer a good opportunity both to gather data, and to provide an educational intervention that is popular with medical students.

Our finding that graduating medical students report limited competence in managing patients with obesity is novel; although consistent with those of Jay et al who found similar results among staff physicians and residents in internal medicine, pediatrics, and psychiatry.¹³ It is concerning that despite further medical training, staff and residents' scores in Jay et al's study were similar to those of medical students in our study. This finding reinforces the need for additional education on management of patients with obesity at the UGME, post-graduate medical education, and continuing professional development levels. Future competency-based medical educational interventions should aim to address the areas of weakness identified in Table 1.

We identified a discrepancy between students' self-reported competence in assessing patients' diet using tools like the FFQ and assessing diet for common unhealthy behaviors associated with obesity. This discrepancy highlights a lack of familiarity among medical students with standardized evidence-based tools available to assess a patient's diet. Familiarity with such tools should become a formal part of competency-based education on the topic of obesity for Canadian medical students.

Our results from the UGME dean's survey demonstrate that, in medical schools that responded, relatively few curricular hours are dedicated to management of patients with obesity, with variability in the topics covered. All schools discussed nutrition and surgery in their curriculum, but topics of physical activity, behavioral therapy and pharmacotherapy were less frequently addressed. By comparison, medical students spend more time learning about obesity-related comorbidities – for example, averaging 15.4 hours for diabetes alone²⁹ – which stands in contrast to the 14.5 hours spent on the chronic disease of obesity itself. These findings are consistent

with recent studies which demonstrate a persistently low prioritization of obesity-related education in contemporary medical school curricula despite ongoing calls for improvement.^{25,30–32}

4.3. Future directions

A number of educational interventions are effective at improving medical students' competence in managing patients with obesity.¹⁹ Such interventions should be incorporated and evaluated within the medical school curricula to address the competencies future physicians will require to meet the changing needs of their patients. Importantly, educational interventions should meet identified gaps. For example, the Association of American Medical Colleges in 2007 and the UK Royal College of Physicians in 2010 both described recommendations for obesity education reform at the UGME level.^{30,33} Similar initiatives should be adopted in Canada to meet the needs of Canadian medical students and patients. The recently published Clinical Practice Guideline for adults with obesity²⁸ can serve as a guide for developing competencies (entrustable professional activities) within the competency-based UGME curriculum.³⁴ In addition, formal assessment of obesity-related competencies should be conducted throughout medical training and during formal licensing exams.^{20,33}

4.4. Limitations

First, participation in our study was voluntary and the response rate was 10.7%, making our results subject to response bias and selection bias. Some medical schools declined to participate, and we only surveyed English-speaking schools, limiting the generalizability of our findings. The 36% response rate by UGME deans also limits the generalizability of our findings. Second, our study relied on students' self-assessment of their competencies in managing patients with obesity, which is susceptible to self-report bias; however, objective knowledge scores were used to complement self-report data, increasing overall reliability. Lastly, we were unable to match responses from the deans' and the students' surveys, due to confidentiality constraints, precluding us from drawing connections between program characteristics and students' performance on the knowledge and competence questionnaires.

5. Conclusions

Graduating Canadian medical students demonstrate limited knowledge and competencies in managing patients with obesity. Current UGME curricula devote limited time to teaching such competencies. Further research should focus on defining, implementing, and assessing specific competencies focused on management of patients with obesity within the undergraduate competency-based medical curriculum. Instituting such changes will ensure that future physicians are better prepared to serve members of Canadian society living with obesity.

6. List Of Abbreviations

Body Mass Index, BMI; Coronavirus Disease 2019, COVID-19; Undergraduate Medical Education, UGME; Checklist for Reporting Results of Internet E-Surveys, CHERRIES; Research Ethics Board, REB; Canadian Dollars, CAD; Mean, M; Standard Deviation, SD; Food Frequency Questionnaire, FFQ.

7. Declarations

7.1. Ethics approval and consent to participate

The study was approved by Queen's University Health Sciences & Affiliated Teaching Hospitals Research Ethics Board (#6026988). All methods were performed in accordance with the relevant guidelines and regulations. Letter of information and consent material was embedded at the start of all surveys in the Qualtrics platform online. Informed consent was obtained from all participants in the study.

7.2. Consent for publication

Not applicable.

7.3. Availability of data and materials

The datasets supporting the conclusions of this article are included within the article (and its additional files; see "Additional_File_7").

7.4. Competing interests

Regarding financial and non-financial competing interests: NJK has nothing to disclose. OL has nothing to disclose. BZ reports grants from Ethicon, outside the submitted work.

7.5. Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

7.6. Authors' contributions

NJK wrote the protocol, created the surveys, completed local and regional ethics applications, disseminated the surveys, collected the data, and prepared the manuscript for submission. OL performed data analysis and wrote the first draft of the manuscript. BZ designed the study and supervised every stage of the project. All authors participated in editing the manuscript and have read and approved the final manuscript.

7.7. Acknowledgements

The authors wish to thank the following for assistance in disseminating the survey at individual institutions: Office of Assessment and Evaluation, MD Program, Temerty Faculty of Medicine, University of Toronto; others, pending permission to be listed in Acknowledgements.

References

1. Twells LK, Gregory DM, Reddigan J, Midodzi WK. Current and predicted prevalence of obesity in Canada: a trend analysis. *CMAJ Open*. 2014;2(1):E18-26. <https://doi.org/10.9778/cmajo.20130016>

2. Statistics Canada. Overweight and obese adults, 2018. Catalogue no. 82-625-X. Health Fact Sheets. 2018. <https://www150.statcan.gc.ca/n1/pub/82-625-x/2019001/article/00005-eng.htm>. Accessed 5 September 2021.
3. Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health*. 2009;9:88. <https://doi.org/10.1186/1471-2458-9-88>
4. Luo W, Morrison H, de Groh M, et al. The burden of adult obesity in Canada. *Chronic Dis Can*. 2007;27(4):135-144.
5. Popkin BM, Du S, Green WD, et al. Individuals with obesity and COVID-19: A global perspective on the epidemiology and biological relationships. *Obesity Reviews*. 2020;21(11). <https://doi.org/10.1111/obr.13128>
6. Yang J, Ma Z, Lei Y. A meta-analysis of the association between obesity and COVID-19. *Epidemiology and Infection*. 2021;149:1-25. <https://doi.org/10.1017/s0950268820003027>
7. Zhang X, Lewis AM, Moley JR, Brestoff JR. A systematic review and meta-analysis of obesity and COVID-19 outcomes. *Scientific Reports*. 2021;11(1). <https://doi.org/10.1038/s41598-021-86694-1>
8. Barber D, Morkem R, Dalgarno N, et al. Patients eligible and referred for bariatric surgery in southeastern Ontario: Retrospective cohort study. *Can Fam Physician*. 2021;67(1):e31-e40. <https://doi.org/10.46747/cfp.6701e31>
9. Zevin B, Martin M, Dalgarno N, et al. Survey of perceptions and educational needs of primary care providers regarding management of patients with class II and III obesity in Ontario, Canada. *BMC Family Practice*. 2021;22(1):14. <https://doi.org/10.1186/s12875-020-01356-x>
10. Bucher Della Torre S, Courvoisier DS, Saldarriaga A, Martin XE, Farpour-Lambert NJ. Knowledge, attitudes, representations and declared practices of nurses and physicians about obesity in a university hospital: training is essential. *Clinical Obesity*. 2018;8(2):122-130. <https://doi.org/10.1111/cob.12238>
11. Butsch WS, Robison K, Sharma R, Knecht J, Smolarz BG. Medicine Residents are Unprepared to Effectively Treat Patients with Obesity: Results from a U.S. Internal Medicine Residency Survey. *Journal of Medical Education and Curricular Development*. 2020;7:238212052097320. <https://doi.org/10.1177/2382120520973206>
12. Glauser TA, Roepke N, Stevenin B, Dubois AM, Ahn SM. Physician knowledge about and perceptions of obesity management. *Obesity Research & Clinical Practice*. 2015;9(6):573-583. <https://doi.org/10.1016/j.orcp.2015.02.011>
13. Jay M, Gillespie C, Ark T, et al. Do Internists, Pediatricians, and Psychiatrists Feel Competent in Obesity Care? *Journal of General Internal Medicine*. 2008;23(7):1066-1070. <https://doi.org/10.1007/s11606-008-0519-y>
14. Tsai AG, Histon T, Kyle TK, Rubenstein N, Donahoo WT. Evidence of a gap in understanding obesity among physicians. *Obes Sci Pract*. 2018;4(1):46-51. <https://doi.org/10.1002/osp4.146>
15. Banasiak M, Murr MM. Medical school curricula do not address obesity as a disease. *Obes Surg*. 2001;11(6):677-679. <https://doi.org/10.1381/09608920160558597>
16. Bessesen DH. It Is Time to Put Obesity Management into the Medical Curriculum. *Obesity Management*. 2006;2(5):169-172. <https://doi.org/10.1089/obe.2006.2.169>

17. Bleich SN, Bennett WL, Gudzone KA, Cooper LA. National survey of US primary care physicians' perspectives about causes of obesity and solutions to improve care. *Bmj Open*. 2012;2(6).
<http://dx.doi.org/10.1136/bmjopen-2012-001871>
18. Dietz WH, Baur LA, Hall K, et al. Management of obesity: improvement of health-care training and systems for prevention and care. *Lancet*. 2015;385(9986):2521-2533. [https://doi.org/10.1016/s0140-6736\(14\)61748-7](https://doi.org/10.1016/s0140-6736(14)61748-7)
19. Katz NJ, Neary E, Tang N, Braund H, Cofie N, Zevin B. Undergraduate medical education interventions aimed at managing patients with obesity: A systematic review of educational effectiveness. *Obesity Reviews*. 2021. <https://doi.org/10.1111/obr.13329>
20. Kushner RF, Butsch WS, Kahan S, Machineni S, Cook S, Aronne LJ. Obesity Coverage on Medical Licensing Examinations in the United States. What Is Being Tested? *Teach Learn Med*. 2017;29(2):123-128.
<https://doi.org/10.1080/10401334.2016.1250641>
21. Roberts DH, Kane EM, Jones DB, et al. Teaching medical students about obesity: a pilot program to address an unmet need through longitudinal relationships with bariatric surgery patients. *Surg Innov*. 2011;18(2):176-183. <https://doi.org/10.1177/1553350611399298>
22. Funk LM, Jolles S, Fischer LE, Voils CI. Patient and Referring Practitioner Characteristics Associated With the Likelihood of Undergoing Bariatric Surgery: A Systematic Review. *JAMA Surg*. 2015;150(10):999-1005.
<https://doi.org/10.1001/jamasurg.2015.1250>
23. O'Brien MW, Shields CA, Oh PI, Fowles JR. Health care provider confidence and exercise prescription practices of Exercise is Medicine Canada workshop attendees. *Appl Physiol Nutr Me*. 2017;42(4):384-390.
<https://doi.org/10.1139/apnm-2016-0413>
24. Tork S, Meister KM, Uebele AL, et al. Factors Influencing Primary Care Physicians' Referral for Bariatric Surgery. *JLS*. 2015;19(3). <https://doi.org/10.4293/JLS.2015.00046>
25. Butsch WS, Kushner RF, Alford S, Smolarz BG. Low priority of obesity education leads to lack of medical students' preparedness to effectively treat patients with obesity: results from the U.S. medical school obesity education curriculum benchmark study. *BMC Med Educ*. 2020;20(1):23.
<https://doi.org/10.1186/s12909-020-1925-z>
26. Martins C, Norsett-Carr A. Obesity Knowledge among Final-Year Medical Students in Norway. *Obes Facts*. 2017;10(6):545-558. <https://doi.org/10.1159/000481351>
27. Kushner RF, Zeiss DM, Feinglass JM, Yelen M. An obesity educational intervention for medical students addressing weight bias and communication skills using standardized patients. *BMC medical education*. 2014;14(101088679):53. <https://doi.org/https://dx.doi.org/10.1186/1472-6920-14-53>
28. Wharton S, Lau DCW, Vallis M, et al. Obesity in adults: a clinical practice guideline. *CMAJ*. 2020;192(31):E875-E891. <https://doi.org/10.1503/cmaj.191707>
29. Ho JD, Woo VC. A Study of Diabetes Teaching in Canadian Medical Schools. *Canadian Journal of Diabetes*. 2016;40(2):149-151. <https://doi.org/10.1016/j.jcjd.2015.08.017>
30. RCP Foresight Report. 2010. The training of health professionals for the prevention and treatment of obesity. Report prepared for Foresight by the Royal College of Physicians. ISBN: 987-1-86016-391-3. 2010.
<https://www.rcplondon.ac.uk/file/268/download>. Accessed 5 September 2021.

31. Metcalf M, Rossie K, Stokes K, Tanner B. The Perceptions of Medical School Students and Faculty Toward Obesity Medicine Education: Survey and Needs Analysis. *JMIR Medical Education*. 2017;3(2):e22. <https://doi.org/10.2196/mededu.7361>
32. Vitolins MZ, Crandall S, Miller D, Ip E, Marion G, Spangler JG. Obesity educational interventions in U.S. medical schools: a systematic review and identified gaps. *Teach Learn Med*. 2012;24(3):267-272. <https://doi.org/10.1080/10401334.2012.692286>
33. AAMC. Report VIII: Contemporary Issues in Medicine: The Prevention and Treatment of Overweight and Obesity. Medical School Objectives Project. 2007. https://store.aamc.org/downloadable/download/sample/sample_id/57/. Accessed 5 September 2021.
34. AFMC EPA working group. AFMC Entrustable Professional Activities for the Transition from Medical School to Residency. 2021. <https://www.afmc.ca/en/faculties/entrustable-professional-activities>. Accessed 5 September 2021.

Figures

Figure 1

Medical student knowledge of obesity management scores by individual question, grouped by knowledge domain (n=133). Questions adapted from Obesity Knowledge among Final-Year medical Students in Norway, by Martins and Norsett-Carr (2017).

Average of % Correct by Knowledge Domain

Figure 2

Medical student knowledge of obesity management scores averaged by knowledge domain (n=133). Questions adapted from Obesity Knowledge among Final-Year medical Students in Norway, by Martins and Norsett-Carr (2017).

Figure 3

Medical Student Self-Assessed Competencies (n=180). Question begins with “I can...” followed by the stem below. Some stems have been condensed for presentation here. The full survey may be found in Additional File 4. Q1: Use 24-hour recall, food record, or food frequency to obtain diet history; Q2: Determine body mass index (BMI); Q3: Assess diet for unhealthy behaviours associated with obesity; Q4: Ascertain each patient’s readiness and ability to work on weight loss; Q5: Recognize and screen for common psychosocial comorbidities in those with obesity; Q6: Perform targeted history and physical examination to identify common obesity-related comorbidities; Q7: Discuss risks associated with obesity; Q8: Respond to a patient’s questions regarding treatment options; Q9: Assess level of physical activity and provide guidance; Q10: Assist in goal setting related to permanent lifestyle changes; Q11: Prescribe plan for exercise / physical activity; Q12: Use motivational interviewing to change behaviour; Q13: Provide counselling intervention regarding weight loss; Q14: Recognize and refer patients with eating disorders; Q15: Collaborate with and refer to allied health.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile1.docx](#)
- [AdditionalFile2.docx](#)
- [AdditionalFile3.docx](#)
- [AdditionalFile4.docx](#)
- [AdditionalFile5.docx](#)

- [AdditionalFile6.docx](#)
- [AdditionalFile7.xlsx](#)