

# Association Between Dietary Phytochemical Index and Breast Cancer: a Case-control Study

**Seyed Mojtaba Ghoreishy**

Tehran University of Medical Sciences <https://orcid.org/0000-0003-4215-6567>

**Azadeh Aminianfar**

Tehran University of Medical Sciences

**Sanaz Benisi-Kohansal**

Tehran university of medical sciences

**Leila Azadbakht**

Tehran University of Medical Sciences

**Ahmad Esmailzadeh** (✉ [a-esmailzadeh@sina.tums.ac.ir](mailto:a-esmailzadeh@sina.tums.ac.ir))

School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences

---

## Research article

**Keywords:** Phytochemicals, Antioxidants, Breast cancer, Diet, Case-control

**Posted Date:** December 31st, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-136491/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Association between dietary phytochemical index and breast cancer: a case-control study

Seyed Mojtaba Ghoreishy<sup>1,2</sup>, Azadeh Aminianfar<sup>3</sup>, Sanaz Benisi-Kohansal<sup>4</sup>,

Leila Azadbakht<sup>4</sup> and Ahmad Esmailzadeh<sup>4,5,6</sup>

<sup>1</sup>*Students' Scientific Research Center, Tehran University of Medical Sciences, Tehran, Iran*

<sup>2</sup>*Department of Clinical Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran*

<sup>3</sup>*Research Center for Biochemistry and Nutrition in Metabolic Diseases, Kashan University of Medical Sciences, Kashan, Iran*

<sup>4</sup>*Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran*

<sup>5</sup>*Obesity and Eating Habits Research Center, Endocrinology and Metabolism Molecular - Cellular Sciences Institute, Tehran University of Medical Sciences, Tehran, Iran*

<sup>6</sup>*Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran*

**Running Title:** Dietary phytochemical index and breast cancer

**Correspondence to:**

Ahmad Esmailzadeh, PhD  
Department of Community Nutrition,  
School of Nutritional Sciences and Dietetics,  
Tehran University of Medical Sciences,  
Tehran, P.O. Box 14155-6117  
Iran  
Tel:+98-21-88955805  
Fax:+98-21-88984861  
Email: [a-esmaillzadeh@sina.tums.ac.ir](mailto:a-esmaillzadeh@sina.tums.ac.ir)

## ABSTRACT

**Background:** Dietary intake of isoflavones has been positively associated with risk of breast cancer (BC) in some earlier studies. In addition, most studies on diet-disease associations came from western countries and limited data are available in the Middle-East.

**Methods:** This case-control study was performed on 350 women with breast cancer aged over 30 years old who were recruited from hospitals or private clinics in Isfahan, Iran. All patients were diagnosed with BC during the maximum of the last 6 months using physical examination and mammography findings. Using cluster method sampling, 700 apparently healthy age- and socioeconomic status-matched controls were randomly selected from healthy women who had no relationship with BC patients and had no familial history of BC. Data on dietary intakes were collected using a validated food frequency questionnaire. The DPI was calculated based on dietary energy derived from foods rich in phytochemicals (kcal) divided by total daily energy intake (kcal) of each participant.

**Results:** Mean  $\pm$  SD age and BMI in the study participants were  $62.4 \pm 10.8$  y and  $24.3 \pm 5.2$  kg/m<sup>2</sup>, respectively. In the crude model, participants in the highest quartile of DPI had 63% lower odds of breast cancer compared to those in the lowest quartile (95% CI: 0.26, 0.54; P-trend <0.001). After adjustment for potential confounders, this inverse association became strengthened (95% CI: 0.22, 0.49; P-trend <0.001). Further adjustment for BMI did not change the association (OR for the highest quartile vs. the lowest quartile = 0.40, 95% CI: 0.26, 0.60; P-trend <0.001).

**Conclusion:** In conclusion, a protective association was observed between DPI and BC in this case-control study. Therefore, high consumption of foods rich in phytochemicals such as fruits, vegetables and whole grains might help reducing the odds of BC among women.

**KEYWORDS:** Phytochemicals, Antioxidants, Breast cancer, Diet, Case-control

## INTRODUCTION

Breast cancer (BC) is one of the most common types of cancer in the world. It is the leading cause of cancer deaths among women in both developing and developed countries [1-4]. In most western countries, mortality rate from BC has risen rapidly to about 40% in the past 30 years [5]. In 2019, 30% of new cancer cases in the US were patients with breast cancer [5]. In Iran, breast cancer is the most prevalent cancer among women [1].

In addition to well-known risk factors for BC including age, family history, early menstruation [1], smoking [6], high adiposity [7], hormone replacement therapy and not breast feeding [8], dietary intakes play an important role [1]. In terms of dietary intakes, fruit and vegetables consumption has been inversely linked with the risk [9-12]. The protective association of these food groups with BC might be mediated through their high content of phytochemicals, which are a wide part of dietary components linked to reduced risk of several chronic disease including cardiovascular disease [13], diabetes [14], inflammatory bowel disease [15] and Alzheimer's disease [16]. These components are non-nutritive bioactive compounds including polyphenols (phenolic acids, flavonoids, isoflavones, lignans, stilbenes, curcuminoids, and calcones), Terpenoids, Organosulfurs, and phytosterols [17]. Diets containing a variety of fresh fruits and vegetables, whole grains, nuts, legumes are rich in phytochemicals. As quantification of phytochemicals in food sources is expensive and impractical for large epidemiological studies, McCarty et al developed a simple and practical tool, named dietary phytochemical index (DPI), for representing the phytochemical content of a whole diet [17]. This index is defined as the percentage of energy intake derived from phytochemical-rich foods [17-19]. Although, the association between DPI and a variety of cancers such as skin cancer [20] and gastrointestinal cancer [21] have been previously examined, limited studies are available linking DPI to the risk of BC [9, 11, 22-24]. In a case-

control study, individuals with higher DPI had a decreased risk for BC than those with the lower DPI [22]. On the other hand, dietary intake of isoflavones has been positively associated with the risk of BC in Asian women [23]. Given these controversies, it seems that additional data are required to further investigate the association between DPI and risk of breast cancer, in particular in women residing in the Middle East, where dietary intakes are different from other parts of the world. Most studies on diet-disease association came from western countries and limited data are available in Middle-East. Nutritional transition in this region has been resulted in a decreased intake of nutrient-dense foods, which are originally high in phytochemicals. Such differences might result in a different overall pattern of phytochemicals in the diet than those in western nations [25]. Therefore, we aimed to investigate the association between DPI and risk of BC among Iranian adult women.

## **METHODS**

**Study Population:** In this population-based case-control study, women aged older than 30 years were recruited from hospitals or private clinics from July 2013 to July 2015 in Isfahan, Iran. All patients were diagnosed with BC during the maximum of the last 6 months using physical examination and mammography findings. They all had primary tumors with invasive behaviors in the breast. Patients were enrolled from those who underwent BC surgery, chemotherapy, radiotherapy, or any of these. Individuals with any history of neoplastic lesions or cysts (except of current BC) or any type of hormone replacement therapy were not included in our study. Using cluster method sampling, age and socioeconomic status- adjusted controls were randomly selected from healthy women who had no relationship with BC patients and had no familial history of BC. Our inclusion criteria for control group were: being woman, having Iranian ethnicity and not having any history of malignancy, cysts and medical disorders and hormone replacement therapy

and not following a special diet. The required sample size was calculated by considering the type I error of 5% and the type II error of 20%. Using a common ratio of 25% and ratio of controls to cases as 2, and assuming a 1.5fold increase in the risk of BC following an unhealthy diet, we eventually reached 350 BC patients and 700 individuals in the control group. A written consent form was completed by all individuals. The whole project was ethically approved by the ethics committee of Isfahan University of Medical Sciences.

**Assessment of Dietary Intakes:** Using a 106-item food-based semi-quantitative Willett-format food frequency questionnaire, dietary data were collected from all participants. Details about the design and validity of the FFQ has been reported elsewhere [26]. The questionnaire included five categories of foods: (1) mixed dishes (cooked or canned, 29 items); (2) carbohydrate-based foods (different types of bread, cakes, biscuits and potato, 10 items); (3) dairy products (dairies, butter and cream, 9 items); (4) fruit and vegetables (22 items); and (5) miscellaneous food items and beverages (including sweets, fast foods, nuts, desserts and beverages, 36 items). People in the study were asked to report their food intake based on 9 options ranging from "never or less than once a month" to "12 times or more per day." We calculated the daily consumption for different foods and reported them as grams per day. Consumption of nutrients is calculated using the Nutritionist IV software, which was modified for Iranian food. The amount of nutrients consumed is obtained by adding the amount of nutrients in different foods. Our previous study showed that the validity and reliability of this FFQ in order to obtain the average long-term dietary intake was reasonable [26].

**Calculation of phytochemical index:** Dietary PI was calculated using the method developed by McCarthy [19] as follow:

DPI= (Dietary energy derived from foods rich in phytochemicals (kcal)/total daily energy intake (kcal) ×100.

The following food items were considered as phytochemical-rich foods in the present analysis: whole grains (toast, oat, bulgur and traditional Iranian breads including Sangak and Barbari), fruits (yellow, red and orange fruits), vegetables (red/orange vegetables, starchy vegetables, dark green vegetables and other vegetables), natural fruit and vegetable juices (orange juice, lemon juice, cantaloupe juice, apple juice, grapefruit juice), tomato sauces, soy products (soy bean), nuts (almond, walnut, peanut, pistachio and hazelnut), legumes (split bean, lentil, chickpea, beans, mung bean and vicia faba), seeds, olive and olive oil. Except for potatoes, which are low in phytochemicals, other vegetables are considered foods rich in phytochemicals.

### **Assessment of breast cancer**

All female patients were Iranian with a recent diagnosis of stage I-IV BC, for whom in-situ or invasive status of BC was confirmed by physical examination and mammography.

### **Assessment of other variables**

The face-to-face interview questionnaire was used to collect data on age, region (urban/rural), education (educated/non-educated), family history of breast cancer (yes/no), alcohol consumption (yes/no), smoking (non-smoker/smoker), marital status (single/ married), menopausal status (premenopausal/postmenopausal) and disease history (yes/no). Weight was measured using a digital scale without shoes and with the least possible clothing. Height was measured with an accuracy of 0.5 cm, when the person was standing without shoes in a normal position. Body mass index was obtained by dividing weight in kilograms by the square of height in meters. In order to evaluate the level of physical activity of participants, we used the short form of the International



Physical Activity Questionnaire (IPAQ). Earlier studies have shown that the information from this questionnaire is accurate and valid [27-29]. Participants were classified based on <1 hour per week (physically inactive) and  $\geq 1$  hour per week (physically active).

**Statistical Analysis:** To examine the association between dietary phytochemical index and odds of breast cancer, first we classified participants based on the quartiles of DPI. Then, ANOVA and Chi Square tests were used to evaluate the differences between continuous and categorical variables across different quartiles of DPI. In order to determine the odds ratios (ORs) and 95% confidence intervals (95% CI) for BC among different quartiles of DPI, a regression logistic was applied in different models, in which we controlled for age (continuous), residence place (urban/rural), marital status (single/married), SES (poor, rich), education (educated/non educated), family history of BC (yes/no), menopausal status (post-menopause/ pre-menopause), breast feeding (yes/no), history of disease (yes/no), supplement use (yes/no), smoking (yes/no) and alcohol (yes/no) consumption in the first model. BMI was additionally adjusted for in the second model to identify obesity-independent association. P value < 0.05 was considered as statistically significant. All analyses were done using SPSS software (version 26).

## RESULTS

Mean  $\pm$  SD of BMI and age in this study participants were  $24.3 \pm 5.2$  kg/m<sup>2</sup> and  $62.4 \pm 10.8$  y, respectively. The mean  $\pm$  SD of the DPI in our study was  $54.76 \pm 12.96$ . General characteristics of study participants across quartiles of DPI are presented in **Table 1**. Compared to people in the bottom quartile, those in the top quartile of DPI had higher weight and BMI and were more likely to use supplements, be post-menopause, married and educated. They were also less likely to reside in urban areas, have family history of BC, use alcohol, and to be smoker, physically active, breast

fed the child and were of poor socioeconomic status compared to individuals in the first quartile of DPI. No other significant association was seen across quartiles of DPI.

**Table 2** indicates dietary intakes of study participants across quartiles of DPI. Compared to the bottom quartile, individuals in the top quartile of DPI had higher intakes of protein, total fiber, folate, calcium, zinc, vegetables, grains, legumes and lower intakes of energy, carbohydrates, fats, vitamin E, meat, egg, trans FA and dairy. No other significant differences were seen in terms of dietary intakes across quartiles of DPI.

Multivariable-adjusted ORs and 95% CI for breast cancer across quartiles of DPI are presented in **Figure 1**. In the crude model, participants in the highest quartile of DPI had 63% lower odds of breast cancer compared to those in the lowest quartile (95% CI: 0.26, 0.54; P-trend <0.001). After adjustment for potential confounders, this inverse association became strengthened (95% CI: 0.22, 0.49; P-trend <0.001). Further adjustment for BMI did not change the association (OR for the highest quartile vs. the lowest quartile = 0.40, 95% CI: 0.26, 0.60; P-trend <0.001).

## DISCUSSION

In this case-control study, high intake of dietary phytochemicals was inversely associated with the odds of BC. This association remained significant after adjustment for several confounding variables. To the best of our knowledge, this study was among the first investigations on the association between DPI and odds of BC in the world.

BC is the leading cause of cancer deaths among women in both developing and developed countries [30]. In the present study, we found that people in the top quartile of DPI were 67% less likely to have breast cancer compared to those in the bottom quartile. The association between DPI and some chronic diseases including cardiovascular disease [13], diabetes [14], psychological

distress [31], Alzheimer's [16], and inflammatory bowel disease [15] have been previously investigated. However, we are aware of very limited and controversial data on the linkage between dietary phytochemicals content and risk of BC. Despite an inverse association between DPI and breast cancer in some studies [32], others failed to find any significant relationship between DPI and odds of breast cancer [24]. In agreement with our findings, another case-control study reported that women in the highest quartile of dietary phytochemical index (DPI) had 92% decreased odds of BC compared to women in the lowest quartile [22]. Post-menopausal women with higher intake of leafy vegetables, any fruits and fruit juices, had approximately 30% lower risk of BC compared to those with a lowest intake in another investigation [9]. However, in a large prospective cohort study, dietary intake of whole grain products was not associated with risk of breast cancer in postmenopausal women [24]. On the other hand, consumption of some phytochemical-rich food groups was even associated with a greater odds of BC in some other studies [33]. Moreover, some ingredients of phytochemical-rich foods were also associated with greater odds of BC in some other studies [32]. Due to the differences in the studies, such as study design, sample size, characteristics of the subjects and inclusion and exclusion criteria, there are inconsistency in the results of the studies. Therefore, more investigation are needed to explore the association between DPI and BC in the future.

There are several mechanisms which might explain the linkage between DPI and BC. Bioactive compounds in phytochemical-rich foods including soluble and insoluble fiber, sterols and stanols, lignans, chlorophyll, flavonoids, indoles, isothiocyanates, phytoestrogens, polyphenolic compounds, protease inhibitors, sulfides and other biologically active metabolites might reduce the risk of various cancers through different cellular pathway including: inhibiting phase I enzymes, inducing phase II enzymes, scavenging DNA reactive agents [34], interruption of cell

proliferation, inhibiting angiogenesis and stimulating apoptosis [35, 36]. In addition, a diet with a high phytochemical content contains high amount of antioxidants, vitamins E and C, carotenoids and various fibers [19]. These components have been shown to have positive effects on the prevention of various cancers [10, 12, 37].

Our study had several strengths. This study was among the first investigations that examined the association between DPI and odds of BC in a Middle-East country. A validated FFQ was used to obtain data on usual dietary intakes of participants. In addition, several confounders were adjusted for in the final analysis to reach an independent association between DPI and BC. Moreover, this study has applied the holistic approach of investigating total dietary phytochemicals rather than focusing on a single dietary component. However, our study had some limitations. First, the design of our case-control study is subject to selection and bias. Measurement errors may lead to misclassification of individuals based on their consumption of phytochemical-rich foods. Second, DPI may include different components in different regions and our findings may not be generalizable to all regions. Third, the status of the hormone receptor is very important in the study of BC-related factors, but there was no information about it in the current study.

## **Conclusion**

In conclusion, a protective association was observed between DPI and BC in this case-control study. Therefore, high consumption of foods rich in phytochemicals such as fruits, vegetables and whole grains might help reducing the odds of BC among women. Therefore, the efforts of the authorities to create comprehensive nutrition policies and facilitate access to foods rich in phytochemicals, in order to encourage people in the community to consume these substances is very important. Further studies with prospective design are needed to confirm this finding.

**Acknowledgements:** Not applicable

**Authors' contributions:** SMG, AA and AE contributed to study concept, search, data analysis and drafting of the manuscript. SBK and LA contributed to data processing, data analysis and drafting of the manuscript. AE supervised the research. All authors read and approved the final manuscript.

**Funding:** This study was financially supported by a grant from School of Nutrition Sciences and Dietetics, Tehran University of Medical Sciences (TUMS), Tehran, Iran

**Availability of data and materials:** The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

**Ethics approval and consent to participate:** The research project was approved by Isfahan University of Medical Sciences, Isfahan, Iran Ethics Committee. The purpose of the study was fully explained

**Consent for publication:** Participants were provided a study overview and verbal consent was attained.

**Competing interests:** The authors have no competing interests to declare

## REFERENCES

1. Marzbani B, Nazari J, Najafi F, Marzbani B, Shahabadi S, Amini M, Moradinazar M, Pashdar Y, Shakiba E, Amini S. Dietary patterns, nutrition, and risk of breast cancer: a case-control study in the west of Iran. *Epidemiol Health*. 2019;41.
2. Yang YL, Xiang ZJ, Yang JH, Wang WJ, Xiang RL. The incidence and relative risk of adverse events in patients treated with bisphosphonate therapy for breast cancer: a systematic review and meta-analysis. *Ther Adv Med Oncol*. 2019;11.
3. Carayol M, Ninot G, Senesse P, Bleuse JP, Gourgou S, Sancho-Garnier H, Sari C, Romieu I, Romieu G, Jacot W. Short- and long-term impact of adapted physical activity and diet counseling during adjuvant breast cancer therapy: the “APAD1” randomized controlled trial. *BMC Cancer*. 2019;19.
4. Bab S, Abdifard E, Elyasianfar S, Mohammadi P, Heidari M. Time trend analysis of breast cancer in Iran and its six topographical regions: a population-based study. *J Med Life*. 2019;12(2):140-149.
5. Bae K, Song SY. Comparison of the clinical effectiveness of treatments for aromatase inhibitor-induced arthralgia in breast cancer patients: a protocol for a systematic review and network meta-analysis. *BMJ Open*. 2020;10(5).
6. Momenimovahed Z, Salehiniya H. Epidemiological characteristics of and risk factors for breast cancer in the world. *Breast Cancer (Dove Med Press)*. 2019;11:151-164.
7. Agurs-Collins T, Ross SA, Dunn BK. The Many Faces of Obesity and Its Influence on Breast Cancer Risk. *Front Oncol*. 2019;9.
8. Gerber B, Müller H, Reimer T, Krause A, Friese K. Nutrition and lifestyle factors on the risk of developing breast cancer. *Breast cancer research and treatment*. 2003;79(2):265-276.
9. Fink BN, Gaudet MM, Britton JA, Abrahamson PE, Teitelbaum SL, Jacobson J, Bell P, Thomas JA, Kabat GC, Neugut AI. Fruits, vegetables, and micronutrient intake in relation to breast cancer survival. *Breast cancer research and treatment*. 2006;98(2):199-208.
10. Jaiswal McEligot A, Largent J, Ziogas A, Peel D, Anton-Culver H. Dietary fat, fiber, vegetable, and micronutrients are associated with overall survival in postmenopausal women diagnosed with breast cancer. *Nutrition and cancer*. 2006;55(2):132-140.
11. Fung TT, Chiuve SE, Willett WC, Hankinson SE, Hu FB, Holmes MD. Intake of specific fruits and vegetables in relation to risk of estrogen receptor-negative breast cancer among postmenopausal women. *Breast cancer research and treatment*. 2013;138(3):925-930.

12. Pantavos A, Ruiter R, Feskens EF, de Keyser CE, Hofman A, Stricker BH, Franco OH, Kiefte-de Jong JC. Total dietary antioxidant capacity, individual antioxidant intake and breast cancer risk: the Rotterdam study. *International journal of cancer*. 2015;136(9):2178-2186.
13. McCullough ML, Peterson JJ, Patel R, Jacques PF, Shah R, Dwyer JT. Flavonoid intake and cardiovascular disease mortality in a prospective cohort of US adults. *The American journal of clinical nutrition*. 2012;95(2):454-464.
14. Belobrajdic DP, Bird AR. The potential role of phytochemicals in wholegrain cereals for the prevention of type-2 diabetes. *Nutrition journal*. 2013;12(1):62.
15. Sung M-K, Park M-Y. Nutritional modulators of ulcerative colitis: clinical efficacies and mechanistic view. *World journal of gastroenterology: WJG*. 2013;19(7):994.
16. Beking K, Vieira A. Flavonoid intake and disability-adjusted life years due to Alzheimer's and related dementias: a population-based study involving twenty-three developed countries. *Public health nutrition*. 2010;13(9):1403-1409.
17. Eslami O, Khoshgoo M, Shidfar F. Dietary phytochemical index and overweight/obesity in children: a cross-sectional study. *BMC Res Notes*. 2020;13.
18. Vincent HK, Bourguignon CM, Taylor AG. Relationship of the Dietary Phytochemical Index to Weight Gain, Oxidative Stress and Inflammation in Overweight Young Adults. *J Hum Nutr Diet*. 2010;23(1):20-29.
19. Mirmiran P, Bahadoran Z, Golzarand M, Shiva N, Azizi F. Association between dietary phytochemical index and 3-year changes in weight, waist circumference and body adiposity index in adults: Tehran Lipid and Glucose study. *Nutr Metab (Lond)*. 2012;9:108.
20. Ng CY, Yen H, Hsiao HY, Su SC. Phytochemicals in Skin Cancer Prevention and Treatment: An Updated Review. *Int J Mol Sci*. 2018;19(4).
21. RK AL-I, Overy AJ, Büsselberg D. Phytochemicals and Gastrointestinal Cancer: Cellular Mechanisms and Effects to Change Cancer Progression. *Biomolecules*. 2020;10(1).
22. Bahadoran Z, Karimi Z, Houshiar-rad A, Mirzayi HR, Rashidkhani B. Dietary phytochemical index and the risk of breast cancer: a case control study in a population of Iranian women. *Asian Pac J Cancer Prev*. 2013;14(5):2747-2751.
23. Chen M, Rao Y, Zheng Y, Wei S, Li Y, Guo T, Yin P. Association between soy isoflavone intake and breast cancer risk for pre-and post-menopausal women: a meta-analysis of epidemiological studies. *PloS one*. 2014;9(2).

24. Egeberg R, Olsen A, Loft S, Christensen J, Johnsen NF, Overvad K, Tjønneland A. Intake of whole grain products and risk of breast cancer by hormone receptor status and histology among postmenopausal women. *International journal of cancer*. 2009;124(3):745-750.
25. Esmailzadeh A, Azadbakht L. Major dietary patterns in relation to general obesity and central adiposity among Iranian women. *The Journal of nutrition*. 2008;138(2):358-363.
26. Keshteli AH, Esmailzadeh A, Rajaie S, Askari G, Feinle-Bisset C, Adibi P. A dish-based semi-quantitative food frequency questionnaire for assessment of dietary intakes in epidemiologic studies in Iran: design and development. *International journal of preventive medicine*. 2014;5(1):29.
27. Dabbagh-Moghadam A, Mozaffari-Khosravi H, Nasiri M, Miri A, Rahdar M, Sadeghi O. Association of white and red meat consumption with general and abdominal obesity: a cross-sectional study among a population of Iranian military families in 2016. *Eating and Weight Disorders-Studies on Anorexia, Bulimia and Obesity*. 2017;22(4):717-724.
28. Anjom-Shoae J, Sadeghi O, Keshteli AH, Afshar H, Esmailzadeh A, Adibi P. The association between dietary intake of magnesium and psychiatric disorders among Iranian adults: a cross-sectional study. *British Journal of Nutrition*. 2018;120(6):693-702.
29. Miri A, Nasiri M, Zonoori S, Yarahmad F, Dabbagh-Moghadam A, Askari G, Sadeghi O, Asadi M. The association between obesity and migraine in a population of Iranian adults: a case-control study. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2018;12(5):733-736.
30. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA: a cancer journal for clinicians*. 2015;65(2):87-108.
31. Mofrad MD, Siassi F, Guilani B, Bellissimo N, Azadbakht L. Association of dietary phytochemical index and mental health in women: a cross-sectional study. *The British journal of nutrition*. 2019;121(9):1049-1056.
32. Ward HA, Kuhnle GG, Mulligan AA, Lentjes MA, Luben RN, Khaw K-T. Breast, colorectal, and prostate cancer risk in the European Prospective Investigation into Cancer and Nutrition–Norfolk in relation to phytoestrogen intake derived from an improved database. *The American journal of clinical nutrition*. 2010;91(2):440-448.
33. Hilakivi-Clarke L, Andrade JE, Helferich W. Is soy consumption good or bad for the breast? *The Journal of nutrition*. 2010;140(12):2326S-2334S.
34. Waladkhani A, Clemens MR. Effect of dietary phytochemicals on cancer development. *International journal of molecular medicine*. 1998;1(4):747-800.



35. Miller PE, Snyder DC. Phytochemicals and cancer risk: a review of the epidemiological evidence. *Nutrition in Clinical Practice*. 2012;27(5):599-612.
36. Bishayee A, Petit D, Samtani K. Angioprevention is implicated in resveratrol chemoprevention of experimental hepatocarcinogenesis. *J Carcinog Mutagen*. 2010;1:102.
37. Hui C, Qi X, Qianrong Z, Xiaoli P, Jundong Z, Mantian M. Flavonoids, flavonoid subclasses and breast cancer risk: a meta-analysis of epidemiologic studies. *PloS one*. 2013;8(1):e54318.

## Legend to figures

**Figure 1:** Multivariable-adjusted odds ratios and 95% CIs for breast cancer across quartiles of dietary phytochemical index. **A:** Crude model; **B:** Adjusted for age, residence, marital status, SES, education, family history of BC, menopausal status, breast feeding, history of disease, supplement use, smoking, alcohol; **C:** Further controlled for BMI.

# Figures




Figure 1

Multivariable-adjusted odds ratios and 95% CIs for breast cancer across quartiles of dietary phytochemical index. A: Crude model; B: Adjusted for age, residence, marital status, SES, education, family history of BC, menopausal status, breast feeding, history of disease, supplement use, smoking, alcohol; C: Further controlled for BMI.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tables.pdf](#)