

MicroRNA expression profile in the N2 phenotype neutrophils of colorectal cancer and screen of putative key mircRNAs

Liang Wang

the first affiliated hospital of kunming medical university

Jun Yang

the first affiliated hospital of kunming medical university

Jian Huang

the first affiliated hospital of kunming medical university

Zheng-Qi Wen

the first affiliated hospital of kunming medical universuty

Ning Xu

the first affiliated hospital of kunming medical university

Xuan Liu

the first affiliated hospital of kunming medical university

Jian-Hua Zhang

honghe prefecture third people's hospital

Wen-Liang Li (✉ liangwyfdiy@163.com)

the first affiliateal hospital of kunming medical university

Primary research

Keywords: MicroRNA, N2 phenotype neutrophils, colorectal cancer

Posted Date: February 10th, 2020

DOI: <https://doi.org/10.21203/rs.2.23091/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective: Colorectal cancer (CRC) is one of the most common malignant tumors in the digestive tract, which accounts for 10% of all the malignant tumors in the world. The aim of this study was to identify key genes and miRNAs in CRC diagnosis, prognosis, and therapy and to further explore the potential molecular mechanisms of CRC.

Methods: The infiltration and metastasis of neutrophils in Primary colorectal cancer tissue and Paracancerous tissue were observed by immunohistochemical staining. After inducing N2 neutrophils with TGF- β 1 in vitro, exosomes were extracted and sequenced, and then the expression differences of microRNAs were screened by using Agilent microRNA microarrays. The data were imported to the Web CARMA for differential expression analysis. The GO and KEGG enrichment analysis were performed using DIANA-MirPath v3.0 using TargetsCan database. And the corresponding targets were imported into Gephi for network analysis. The expression level of differentially expressed microRNA using quantitative Realtime polymerase chain reaction (RT-PCR) was validated.

Results: A total of 2 miRNAs were found to be associated with N2 neutrophils, in which the expression of hsa-miR-4780 was upregulated and the expression of hsa-miR-3938 was downregulated in N2 neutrophils, compared with the neutrophils. In addition, the results of miRNA-targets networks showed that the hsa-mir-3938 and hsa-mir-4780 could regulate TUSC1 and ZNF197. The expression level of hsa-miR-4780 and hsa-miR-3938 were validated in accordance with the results of RT-PCR.

Conclusion: The hsa-mir-3938 and hsa-mir-4780 were differentially expressed between N2 neutrophils and neutrophils. Moreover, the regulation of TUSC1 and ZNF197 by these DEmiRNA established the theoretical basis for the mechanism of N2 type neutrophils regulating the invasion and metastasis of CRC cells, and provided the potential biomarker for prognosis for clinical treatment of CRC

Introduction

Colorectal cancer (CRC) is one of the most common malignant tumors in the digestive tract, which accounts for 10% of all the malignant tumors in the world [1]. With the rapid development of social and economic life and the change of diet lifestyle and diet structure, the incidence and prevalence of colorectal cancer are increasing gradually [2]. Although the treatment of CRC has improved significantly in recent decades, the five year survival rate is only 8%-10% [3]. Therefore, it is more important to study the pathogenesis mechanism of CRC, and to find the right drug target to promote the effect of treatment and prolong the survival time.

Tumor microenvironment includes tumor cells themselves, various immune cells and interstitial cells. Immune cells mainly include neutrophils, macrophages, NK cells, lymphocytes and dendritic cells [4]. Many recent researches have denoted that the neutrophils play important roles in the formation of the tumor microenvironment [5]. Tumor-associated neutrophils (TANs) secrete various proteases, reactive oxygen species and cytokines, which play an important role in tumor growth, metastasis, angiogenesis

and immune regulation [6]. TANs can be further divided into two subtypes as N1 TANs and N2 TANs. N1 TANs has antitumor effect, while N2 TANs can promote the occurrence, development and metastasis of tumors [7]. In addition, A number of studies revealed that the increase of neutrophil count in peripheral blood was related to poor clinical outcomes in CRC, but the mechanism of this process was unclear [8–11].

MircRNAs (miRNAs) are small molecule RNA with a length of 20–25 nucleotides, which are involved in the regulation of various physiological and biochemical processes in organisms [12]. It has been found that miRNAs played an important regulatory role in the occurrence and development of diseases, especially tumors. If the expression of miRNAs which can regulate cell proliferation and differentiation were abnormal, it might lead to the occurrence of tumor [13]. In addition, previous studies showed that miRNAs expression were different in various tissues of CRC, and it were related to the proliferation and metastasis of CRC tumor cells. For example, studies showed that long chain noncoding RNA MIR17HG promoted metastasis of CRC via miR-17 [14], and the increased expression of miR-17 also promotes metastasis of CRC [15]. Another research showed that the miR-200 families were key factor in the occurrence of epithelial to mesenchymal transition (EMT) in tumor cells and abnormal expression of miR-200 might promote EMT in tumor cells and lead to metastasis [16]. However, the relationship between miRNA in type N2 neutrophils and CRC is still unclear and A considerable amount of research is needed in this area.

Based on the above background, we used transforming growth factor- β 1 (TGF- β 1) to induce N2 phenotype neutrophils in vitro, extracted exosomes for sequenced, and screened miRNAs with different expression. We analyzed the mechanism of N2 neutrophil miRNA regulating the invasion and metastasis of CRC cells, which provided new therapeutic target for CRC, and provided the theoretical basis for individualized treatment of CRC.

Materials And Methods

Immunohistochemical staining

Tissue samples were obtained from patients diagnosed with colorectal cancer between July 2019 and Dec 2019 at the First Affiliated Hospital of Kunming Medical University. Formalin-fixed, paraffin-embedded colon biopsy sections were deparaffinized in xylene for 20 min and rehydrated in a graded ethanol series (100-70%) for 8 min. Endogenous peroxidase were deactivated with 3% hydrogen peroxide for 5-510 min at room temperature after deparaffinization. Antigen retrieval was performed with a sodium citrate buffer (10 mM, pH = 6) for 10 min. A mouse IgM antihuman monoclonal antibody specific for CD11b, CD66b (ab197678) and matrix metalloproteinase-9 (MMP9) (BD Biosciences, San Diego, California) were used to identify neutrophils. Negative controls were stained with a rabbit IgG negative control prediluted in phosphate-buffered saline (Life Technologies). Slides were counterstained with hematoxylin. The staining protocol was the same for all samples with regard to processing, incubation times, and temperature.

TGF- β 1 induced neutrophils to N2 polarization in vitro

The acute promyelocytic leukemia (APL) cell line (NB-4) were cultured in RPMI-1640 culture medium containing 10% fetal bovine serum at 37 °C in the presence of 5% CO₂. Then, using all-trans retinoic acid (ATRA) induced the differentiation of NB-4 cells into neutrophils. According to different polarity disposal conditions, neutrophils were divided into 2 groups, normal control group and N2 polarization group (neutrophil group and neutrophil+TGF- β 1 group). Neutrophils in the normal control group were intact. Neutrophils in the N2 polarization group were incubated with 10ng/ml TGF- β 1 for 24h. Then extracted exosomes for sequenced, and screened miRNAs with different expression.

Total RNA Extraction

Total RNA was extracted from samples using a TRIzol® Reagent (Invitrogen life technologies) according to the manufacturer's introductions. RNA quantity and quality were measured by NanoDrop ND-1000. The integrity of RNA was assessed by standard denaturing agarose gel electrophoresis.

MircRNA microarrays

The Whole Human MiRNA Microarray was a broad view that represents all known miRNAs in the human transcriptome. Sequences were compiled from a broad source survey, and then verified and optimized by alignment to the assembled human transcriptome.

RNA labeling and array hybridization

Sample labeling and array hybridization were performed according to the Agilent miRNA Microarray System with miRNA Complete Labeling and Hyb Kit protocol (Agilent Technology). Briefly, total miRNA from each sample was labeled with Cyanine 3-pCp under the action of T4 RNA ligase. The labeled cRNA over the procession of inspissation and desiccation and then redissolved with water. 1 μ g of each labeled cRNA was fragmented by adding 11 μ l 10 \times Blocking Agent and 2.2 μ l of 25 \times Fragmentation Buffer, then heated at 60°C for 30 min, and finally 55 μ l 2 \times GE Hybridization buffer was added to dilute the labeled cRNA. 100 μ l of hybridization solution was dispensed into the gasket slide and assembled to the gene expression microarray slide. The slides were incubated for 17 hours at 65°C in an Agilent Hybridization Oven. The hybridized arrays were washed, fixed and scanned with using the Agilent Microarray Scanner (part number G2505C).

Data analysis

The normalized circRNA and mRNA expression data were imported to the WebCARMA for differential expression analysis. Moderated t-test (from limma package in R) was used to detect the difference between neutrophil + TGF- β 1 group (2-1, 2-2, 2-3) and neutrophil group (1-1, 1-2, 1-3). The results were stored in "miRNA_DE.csv" in the DE_analysis fold, where the column 'rawp' is the raw p value, the column 'BH' is BH adjusted FDR and 'meanM' is the mead difference between TEST group and CONTROL group.

Differentially expressed miRNAs with statistical significance between the two groups were identified through Volcano Plot filtering. Differentially expressed miRNAs between the two samples were identified through Fold Change filtering. Hierarchical Clustering was performed using the R scripts.

GO and KEGG pathway analysis

The Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis were performed using DIANA-miRPath v3.0 using Targetscan database. The target prediction was also conducted using DIANA-miRPath v3.0 (Targetscan database) and the top 10 miRNAs (based on raw p-value). GO results were mainly classified into three subgroups namely biological process (BP), cellular component (CC), and molecular function (MF).

Construction of the miRNA-targets network analysis

The corresponding targets were imported into Gephi for network analysis. The network was showed in Figure 2 (red nodes represent the miRNA and green nodes represent targets).

qRT-PCR assay

The content of the differentially expressed miRNAs in neutrophil group and neutrophil + TGF- β 1 group were detected by utilizing real-time PCR. RT-PCR was performed using the Applied Biosystems 7300 Sequence Detection System (Ambion, Austin TX, USA). Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) were quantified as an internal control for data normalization. The sequences of targets primers were as follows: Tumor suppressor candidate 1 (TUSC1) forward primer was 5'-GACTTGAGAAGCTGGAA-3', and reverse primer was 5'-TCGGGTTCTGTAGAG-3'. Zinc finger protein 197 (ZNF197) forward primer was 5'-CTTCACTTCAGAGGAATG-3', and reverse primer was 5'-ATAATTCTCCAGCATCAC-3'.

Statistical analysis

All data were presented as mean \pm SD. The comparison between two groups was conducted using unpaired Student's t-test. P values less than 0.05 were considered statistically significant.

Results

Comparison of the neutrophil infiltration among the different groups

As shown in Figure 1, the expression of CD11b (CD11b is the most important adhesion molecule on neutrophil membrane) and CD66b (CD66b is a surface specific molecular marker of tumor infiltrating neutrophils (TINs) were significantly increased in primary colorectal cancer. The MMP9 is a tumor angiogenesis protein produced by tumor-associated neutrophils, regarding the result of immunohistochemical staining, the expression of mmp9 was significantly increased in primary colorectal cancer (Figure 1). These results suggested that neutrophils were associated with tumor cell adhesion, invasion and metastasis in colorectal cancer.

Expression profiles of miRNAs in N2 neutrophils

All DEmircRNAs were displayed in the hierarchical clustering (Figure 2), with red color representing high read counts and green color representing low read counts of miRNAs. We identified a total of 799 up-regulated miRNAs and 902 down-regulated miRNAs in all samples. There were 33 miRNAs with raw p-value lower than 0.05 (only 2 have FDR lower than 0.05). These 33 miRNAs were used for downstream analysis. All the significantly different miRNAs were listed in Table 1. Based on raw p value (neutrophil + TGF- β 1 group VS. neutrophil group), the top 5 up regulated miRNA were hsa-miR-4780, hsa-miR-513b-3p, hsa-miR-5703, hsa-miR-630, hsa-miR-4684-3p, and the top 5 down regulated miRNA were hsa-miR-3938, hsa-miR-6089, hsa-miR-2114-3p, hsa-miR-1301-5p, hsa-miR-6736-3p. Moreover, there were 6/33 miRNA containing chromosomal distribution information: hsa-miR-2114-3p (chX), hsa-miR-1307-3p (ch10), hsa-miR-1273g-5p (ch1), hsa-miR-3151-5p (ch8), hsa-miR-3136-5p (ch3), hsa-miR-627-5p (ch15).

In addition, as shown in Figure 3, enrichment of total miRNAs in the 3 neutrophils and 3 N2 neutrophils was estimated in the Box Plot. In the volcano plot and Scatter Plot, the up-regulated miRNAs were represented as the red dots, while the down-regulated miRNAs were represented as the green dots (Figure 4a, Figure 4b).

GO and KEGG analysis of miRNAs

The top 10 enriched GO-BP terms, the top 9 enriched GO-MF and the top 5 enriched GO-CC terms that may be associated with the mechanism of N2 neutrophils in tumor tissue infiltration were shown in Figure 5. According to the results, the most enriched and meaningful GO-BP terms were related to “cellular nitrogen compound metabolic process,” “biosynthetic process,” “small molecule metabolic process,” “symbiosis encompassing mutualism through parasitism,” “viral process,” “neurotrophic TRK receptor signaling pathway,” “gene expression,” “cellular protein modification process,” “cellular protein metabolic process” and “catabolic process.” Within the GO-CC terms, the largest number of predicted target genes were involved in organelle, cellular component, protein complex, cytosol and nucleoplasm. Regarding the GO-MF terms, the predicted target genes were mainly associated with ion binding, enzyme binding, nucleic acid binding transcription factor activity, molecular function, cytoskeletal protein binding, enzyme regulator activity, transmembrane transporter activity, protein binding transcription factor activity and small conjugating protein binding.

Regarding the KEGG analyses, the most significantly enriched pathways of the miRNA host genes were metabolism of xenobiotics by cytochrome P450, Glycosphingolipid biosynthesis-lacto and neolacto series, Chagas disease (American trypanosomiasis), sphingolipid signaling pathway and cell adhesion molecules (CAMs) (Figure 5).

Construction of miRNA-targets networks

These 10 miRNAs in total were used for prediction of their targets and a network containing these miRNAs and genes was constructed (Figure 6). In addition, the target genes regulated by DEmRNAs were

listed, as shown in Table 2, which showed that 80 target genes were regulated by hsa-miR-3938, and 101 target genes were regulated by hsa-miR-4780. The results showed that 2 target genes, including TUSC1 and ZNF197, were regulated by hsa-mir-3938 and hsa-mir-4780 simultaneously.

The expression levels of DEmircRNA, TUSC1 and ZNF197

Figure 7a and Figure 7b showed the expression levels of DEmircRNA in neutrophil group and neutrophil + TGF- β 1 group. As shown in Figure 7, in the neutrophil + TGF- β 1 group, the expression levels of hsa-miR-3938 was significantly decreased ($P = 0.0130$), while the hsa-miR-4780 expression increased significantly ($P = 0.0188$). In addition, as shown in Figure 7c, the expression level of DEmircRNA targets also showed significant difference in the two groups. Regarding the neutrophil + TGF- β 1 group, the expression levels of TUSC1 was significantly decreased ($P = 0.0024$), while the ZNF197 expression increased significantly ($P = 0.0058$).

Discussion

CRC is the most common malignant tumor of digestive tract in the world, which seriously endangers people's health and quality of life. However, its pathogenesis has not been completely elucidated, and progression of CRC is a process that involves multiple genetic changes, multi-factor, multi-step process [17, 18]. Previous studies indicate that one third of CRC patients may develop liver metastases, and most of CRC-related death is usually attributed to distant metastasis. More and more evidences showed that neutrophils played an important role in tumor tissue infiltration to promoting the growth, invasion, angiogenesis and metastasis of various types of cancer [19, 20]. In addition, studies showed that exosomes as a novel mechanism of intercellular communication, could lead to the exchange of genetic information and reprogramming of recipient cells and transport various biomolecules, such as proteins, messenger RNAs (mRNAs), microRNAs (miRNAs), and long non-coding RNAs (lncRNAs) [21, 22]. Recent studies showed that exosomes play multiple roles in promoting tumor growth, metastasis and drug resistance. However, the function of neutrophil-derived exosomes in development, invasion and metastasis of CRC has not been well characterized. In this study, we reported mircRNA profiles in the neutrophils of CRC for the first time. Additionally, a network between miRNAs in neutrophils and targets, which may participate in tumor tissue infiltration in CRC, was constructed.

In this study, a total of 33 differentially expressed miRNAs were identified in TGF- β 1-infected neutrophils. KEGG pathway and GO category analysis demonstrated that the target miRNAs were associated with many physiological processes. GO analysis implied that the DEmircRNAs were presumably involved in the biosynthetic and small molecules metabolic process of organelles, nuclei and cytoplasm in cells. The KEGG pathway analysis showed enrichment in the biological processes of cytochrome P450 metabolism xenobiotics, Glycosphingolipid biosynthesis-lacto and neolacto series, sphingolipid signaling pathway and CAMs. The results of miRNA-targets networks showed that the hsa-mir-3938 and hsa-mir-4780 could regulate TUSC1 and ZNF197. These results suggested that the TUSC1 and ZNF197 gene regulated by

DEmirRNAs in N2 neutrophils might be related to biosynthesis, metabolism and signal transduction in colorectal cancer cells.

TUSC1 gene is located within the region of chromosome 9p that harbors tumor suppressor genes critical in carcinogenesis. It is an intronless gene which is downregulated in non-small-cell lung cancer and small-cell lung cancer cell lines, suggesting that it may play a role in lung tumorigenesis (provided by RefSeq, Jul 2008) [23]. Furthermore, A previous study provided further evidence that TUSC1 expression was downregulated in lung cancer cell lines and a trend towards higher expression of TUSC1 was correlated with longer survival times for lung cancer patients [24]. However, there is no report about biological functions and regulatory mechanisms of the TUSC1 gene in the CRC. In the current study, the results showed that TUSC1 protein decreased significantly in N2 type neutrophils compared with neutrophils. Another result showed that in N2 type neutrophils, hsa-mir-3938 expression decreased significantly, while hsa-mir-4780 expression increased significantly. In N2 neutrophils, TUSC1 gene might inhibit the adhesion, invasion and metastasis of colorectal cancer cells, which might be related to the downregulation of hsa-mir-3938 expression and the upregulation of hsa-mir-4780 expression in the tumor microenvironment.

ZNF197 gene product belongs to the zinc finger protein superfamily, members of which are regulatory proteins characterized by nucleic acid-binding zinc finger domains. The encoded protein contains 20 tandemly arrayed C2H2-type zinc fingers, a Kruppel-associated box (KRAB) domain, and a SCAN box. This gene is located in a cluster of zinc finger genes at 3p21. It is overexpressed in some thyroid papillary carcinomas (provided by RefSeq, May 2017) [25]. However, there is no report about biological functions and regulatory mechanisms of the ZNF197 gene in the CRC. In our study, the results showed that ZNF197 protein increased significantly in N2 type neutrophils group. According to the results of miRNA-targets networks, also suggesting that the ZNF197 gene regulated by hsa-mir-3938 and hsa-mir-4780 in N2 neutrophils might be related to the tumor cell growth, invasion, metastasis and occurrence/development of CRC. The current research on this area is relatively less, there is much research work needs to be done in future.

Conclusion

The hsa-mir-3938 and hsa-mir-4780 were differentially expressed between N2 neutrophils and neutrophils. Moreover, our study indicated that the regulation of TUSC1 and ZNF197 by hsa-mir-3938 and hsa-mir-4780 established the theoretical basis for the mechanism of N2 type neutrophils regulating the invasion and metastasis of CRC cells, and provided the potential biomarker for prognosis for clinical treatment of CRC, but more studies should be conducted to support this new hypothesis.

Abbreviations

A list of abbreviations was provided in Table 3.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by Yunnan Fundamental Research Projects (NO. 2019FA039), National Natural Science Foundation of China (No.31660312), and leading medical talents in Yunnan (NO. L-2017001).

Authors' contributions

Conceptualization: WLL, LW; Data curation: LW, XL; Investigation: LW, JY, JH; Methodology: LW, WLL, ZQW, NX; Resources: WLL, JHZ; Software: LW, JY, XL; Validation: LW, JY, JH; Writing - original draft: LW, JY, JH; Writing- review& editing: LW, WLL. All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Acknowledgements

Not applicable

Authors' information (optional)

Not applicable

References

1. Swiderska, M., Choromanska, B., Dabrowska, E., Konarzewska-Duchnowska, E., Choromanska, K., Szczurko, G., Mysliwiec, P., Dadan, J., Ladny, J. R., Zwierz, K., The diagnostics of colorectal cancer. Contemporary oncology (Poznan, Poland) 2014, 18, 1-6.

2. Hu, J. M., Shen, C. J., Chou, Y. C., Hung, C. F., Tian, Y. F., You, S. L., Chen, C. Y., Hsu, C. H., Hsiao, C. W., Lin, C. Y., Sun, C. A., Risk of colorectal cancer in patients with periodontal disease severity: a nationwide, population-based cohort study. *International journal of colorectal disease* 2018, 33, 349-352.
3. Hagggar, F. A., Boushey, R. P., Colorectal cancer epidemiology: incidence, mortality, survival, and risk factors. *Clinics in colon and rectal surgery* 2009, 22, 191-7.
4. Shaul, M. E., Fridlender, Z. G., Neutrophils as active regulators of the immune system in the tumor microenvironment. *Journal of leukocyte biology* 2017, 102, 343-349.
5. Joyce, J. A., Pollard, J. W., Microenvironmental regulation of metastasis. *Nature reviews. Cancer* 2009, 9, 239-52.
6. Masucci, M. T., Minopoli, M., Carriero, M. V., Tumor Associated Neutrophils. Their Role in Tumorigenesis, Metastasis, Prognosis and Therapy. *Frontiers in oncology* 2019, 9, 1146.
7. Allavena, P., Sica, A., Garlanda, C., Mantovani, A., The Yin-Yang of tumor-associated macrophages in neoplastic progression and immune surveillance. *Immunological reviews* 2008, 222, 155-61.
8. Stojkovic Lalosevic, M., Pavlovic Markovic, A., Stankovic, S., Stojkovic, M., Dimitrijevic, I., Radoman Vujacic, I., Lalic, D., Milovanovic, T., Dumic, I., Combined Diagnostic Efficacy of Neutrophil-to-Lymphocyte Ratio (NLR), Platelet-to-Lymphocyte Ratio (PLR), and Mean Platelet Volume (MPV) as Biomarkers of Systemic Inflammation in the Diagnosis of Colorectal Cancer. 2019, 2019, 6036979.
9. Pedrazzani, C., Mantovani, G., Fernandes, E., Bagante, F., Assessment of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio and platelet count as predictors of long-term outcome after R0 resection for colorectal cancer. 2017, 7, 1494.
10. Li, H., Zhao, Y., Zheng, F., Prognostic significance of elevated preoperative neutrophil-to-lymphocyte ratio for patients with colorectal cancer undergoing curative surgery: A meta-analysis. *Medicine* 2019, 98, e14126.
11. Tanio, A., Saito, H., Uejima, C., Takaya, S., Yamamoto, M., Tokuyasu, N., Sakamoto, T., Honjo, S., Ashida, K., Fujiwara, Y., A prognostic index for colorectal cancer based on preoperative absolute lymphocyte, monocyte, and neutrophil counts. *Surgery today* 2019, 49, 245-253.
12. Ma, X., Becker Buscaglia, L. E., Barker, J. R., Li, Y., MicroRNAs in NF-kappaB signaling. *Journal of molecular cell biology* 2011, 3, 159-66.
13. Hao Lai, Jie Zhang, Hong-qun Zuo et al, Advances in research on the correlation between microRNA and colorectal cancer metastasis. *Chinese Journal of New Clinical Medicine* 2019, 12, 1168-1171.
14. Xu, J., Meng, Q., Li, X., Yang, H., Xu, J., Gao, N., Sun, H., Wu, S., Familiari, G., Relucenti, M., Long Noncoding RNA MIR17HG Promotes Colorectal Cancer Progression via miR-17-5p. 2019, 79, 4882-4895.
15. Fu, F., Jiang, W., Zhou, L., Chen, Z., Circulating Exosomal miR-17-5p and miR-92a-3p Predict Pathologic Stage and Grade of Colorectal Cancer. *Translational oncology* 2018, 11, 221-232.
16. Diaz-Riascos, Z. V., Ginesta, M. M., Fabregat, J., Serrano, T., Busquets, J., Buscail, L., Cordelier, P., Capella, G., Expression and Role of MicroRNAs from the miR-200 Family in the Tumor Formation and

- Metastatic Propensity of Pancreatic Cancer. *Molecular therapy. Nucleic acids* 2019, 17, 491-503.
17. Caraglia, M., Correale, P., Giannicola, R., Staropoli, N., Botta, C., Pastina, P., Nesci, A., Caporlingua, N., Francini, E., Ridolfi, L., Mini, E., Roviello, G., Ciliberto, D., Agostino, R. M., Strangio, A., Azzarello, D., Nardone, V., Falzea, A., Cappabianca, S., Bocchetti, M., D'Arrigo, G., Tripepi, G., Tassone, P., Addeo, R., Giordano, A., Pirtoli, L., Francini, G., Tagliaferri, P., GOLFIG Chemo-Immunotherapy in Metastatic Colorectal Cancer Patients. A Critical Review on a Long-Lasting Follow-Up. *Frontiers in oncology* 2019, 9, 1102.
 18. Arai, H., Battaglin, F., Wang, J., Lo, J. H., Soni, S., Zhang, W., Lenz, H. J., Molecular insight of regorafenib treatment for colorectal cancer. *Cancer treatment reviews* 2019, 81, 101912.
 19. Mizuno, R., Kawada, K., The Role of Tumor-Associated Neutrophils in Colorectal Cancer. 2019, 20.
 20. Fridlender, Z. G., Albelda, S. M., Tumor-associated neutrophils: friend or foe? *Carcinogenesis* 2012, 33, 949-55.
 21. Zhang, X., Shi, H., Yuan, X., Jiang, P., Qian, H., Xu, W., Tumor-derived exosomes induce N2 polarization of neutrophils to promote gastric cancer cell migration. *Molecular cancer* 2018, 17, 146.
 22. Han, Q., Zhao, H., Jiang, Y., Yin, C., Zhang, J., HCC-Derived Exosomes: Critical Player and Target for Cancer Immune Escape. *Cells* 2019, 8.
 23. Shan, Z., Parker, T., Wiest, J. S., Identifying novel homozygous deletions by microsatellite analysis and characterization of tumor suppressor candidate 1 gene, TUSC1, on chromosome 9p in human lung cancer. *Oncogene* 2004, 23, 6612-20.
 24. Shan, Z., Shakoori, A., Bodaghi, S., Goldsmith, P., Jin, J., Wiest, J. S., TUSC1, a putative tumor suppressor gene, reduces tumor cell growth in vitro and tumor growth in vivo. *PloS one* 2013, 8, e66114.
 25. Li, Z., Wang, D., Na, X., Schoen, S. R., Messing, E. M., Wu, G., The VHL protein recruits a novel KRAB-A domain protein to repress HIF-1alpha transcriptional activity. *The EMBO journal* 2003, 22, 1857-67.

Tables

Table 1 Deferential expression miRNA (based on raw p-value)

Systematic_name	Mirbase accession	Raw p	BH-FDR	meanM
hsa-miR-3938	MIMAT0018353	0	0	-5.312
hsa-miR-4780	MIMAT0019939	0	0.001	5.171
hsa-miR-513b-3p	MIMAT0026749	0.002	0.965	0.672
hsa-miR-5703	MIMAT0022496	0.008	0.965	0.437
hsa-miR-630	MIMAT0003299	0.012	0.965	0.413
hsa-miR-6089	MIMAT0023714	0.015	0.965	-0.306
hsa-miR-4684-3p	MIMAT0019770	0.018	0.965	0.552
hsa-miR-2114-3p	MIMAT0011157	0.021	0.965	-0.492
hsa-miR-1301-5p	MIMAT0026639	0.022	0.965	-0.281
hsa-miR-6736-3p	MIMAT0027374	0.022	0.965	0.379
hsa-miR-1307-3p	MIMAT0005951	0.024	0.965	-0.606
hsa-miR-5010-5p	MIMAT0021043	0.027	0.965	-0.392
hsa-miR-4439	MIMAT0018957	0.028	0.965	-0.411
hsa-miR-3196	MIMAT0015080	0.029	0.965	-0.4
hsa-miR-5694	MIMAT0022487	0.035	0.965	-0.621
hsa-miR-1273g-5p	MIMAT0020602	0.035	0.965	-3.713
hsa-miR-5587-5p	MIMAT0022289	0.035	0.965	-3.713
hsa-miR-7159-5p	MIMAT0028228	0.036	0.965	-0.38
hsa-miR-4738-3p	MIMAT0019867	0.036	0.965	0.266
hsa-miR-6754-5p	MIMAT0027408	0.037	0.965	0.278
hsa-miR-3928-5p	MIMAT0027037	0.037	0.965	-0.453
hsa-miR-4754	MIMAT0019894	0.037	0.965	0.381
hsa-miR-3151-5p	MIMAT0015024	0.038	0.965	-0.544
hsa-miR-6125	MIMAT0024598	0.039	0.965	-0.265
hsa-miR-6818-5p	MIMAT0027536	0.04	0.965	-0.675
hsa-miR-615-3p	MIMAT0003283	0.041	0.965	0.564
hsa-miR-4730	MIMAT0019852	0.042	0.965	-0.905
hsa-miR-8060	MIMAT0030987	0.042	0.965	0.288
hsa-miR-5588-3p	MIMAT0022296	0.045	0.965	0.348
hsa-miR-4421	MIMAT0018934	0.045	0.965	0.561
hsa-miR-3136-5p	MIMAT0015003	0.046	0.965	0.318
hsa-miR-627-5p	MIMAT0003296	0.048	0.965	0.338
hsa-miR-6836-3p	MIMAT0027575	0.048	0.965	-0.225

Raw p, raw p value; BH-FDR, adjusted False Discovery Rate (FDR) using Benjamin & Hochberg method; meanM, the different normalized value between Test group and Control group (neutrophil+TGF- β 1 group minus neutrophil group).

Table 2 DEmRNAs target genes

miRNA	Target	N
hsa-miR-3938	AMOT,ANGPTL5,ANKRD54,ARCNI,ASAH2,ATP6V0A2,BCL10,BORA,BRWD1,CCDC141,CCDC169,CCDC73,CCSAP,CDC73,CDRT4,CPOX,CTPS2,DAZL,DDTL,DECR1,DL D,EIF2AK2,FAM216B,FARP1,FAT3,FBXO38,FGF18,GCNT2,GCSAM,GPR155,IL17RB,ISM1,KAT2A,KDEL2,KIAA1755,LAX1,LPIN3,LRP2,MIB1,MITD1,MRPS34,MTMR12,MTMR7,MYH15,NANP,NUDT15,PDHX,PNISR,PWWP2A,RELL1,RNF121,RP11-362K2.2,SLC16A1,SLC16A14,SLC16A7,SLC17A4,SLC35E4,SNX4,SS18,ST8SIA4,SYTL5,TLDC1,TMEM206,TMEM237,TMEM75,TOMM22,TROVE2,TSPAN31, TUSC1 ,TXNDC12,TXNL1,UBR5,WDFY2,WDR3,WIF1,ZMYND12, ZNF197 ,ZNF354C,ZNF629,ZNF793	80
hsa-miR-4780	AMIGO1,ANGEL2,AP3M2,ARHGAP31,ARHGAP6,ATG13,ATXN7,BAIAP2L1,BBIP1,BBS7,C6orf120,CD163,CFHR3,CIPC,CLNK,COBLL1,COL21A1,COL24A1,COL6A6,COQ3,CRHBP,DCCLK1,DDX3Y,DEPDC1B,DLEU2L,DLGAP5,ENOX2,EPM2AIP1,ERBB2IP,FUT10,GFOD1,GPR126,HELZ,HEXIM1,HNMT,IGSF23,IL6ST,ILDR2,ITGAV,KCNJ14,KLF3,LCN10,LL0XNC01-19D8.1,LSM11,MCCEMP1,MED14,MEGT1,MFAP5,MFSD11,MMP13,MMRN2,MPP1,MTERF,MTHFD2L,MVK,NAPEPLD,NCALD,NR1I3,OR5AU1,PARL,PGBD5,PID1,PKM,POLD3,PRDM7,PRIMA1,PRR34,PSMD14,PTBP3,PTGDR,PTPRD,ROCK1,RP11-296A16.1,SCML1,SERINC4,SLC25A29,SMIM3,SREK1,SRPX2,SRSF12,SUV39H2,SVIL,TCP11X2,TMEM132C,TMEM156,TMEM246,TMTC3,TOMM40L,TRMT13,TUBB, TUSC1 ,UBQLN1,VRK2,XPNPEP3,XRCC5,ZADH2,ZBTB80S, ZNF197 ,ZNF620,ZNF75D,ZSCAN23	101

The genes were regulated by hsa-mir-3938 and hsa-mir-4780 were listed in the table and were bold.

Table 3 List of abbreviations

Abbreviations	
Acute promyelocytic leukemia	APL
All-trans retinoic acid	ATRA
Biological process	BP
Cellular component	CC
Colorectal cancer	CRC
Epithelial to mesenchymal transition	EMT
Gene Ontology	GO
Glyceraldehyde 3-phosphate dehydrogenase	GAPDH
Kruppel-associated box	KRAB
Kyoto Encyclopedia of Genes and Genomes	KEGG
Long non-coding RNAs	lncRNAs
Matrix metalloproteinase-9	MMP9
Messenger RNAs	mRNAs
MircRNAs	miRNAs
Molecular function	MF
Realtime polymerase chain reaction	RT-PCR
Transforming growth factor- β 1	TGF- β 1
Tumor-associated neutrophils	TANs
Tumor infiltrating neutrophils	TINs
Tumor suppressor candidate 1	TUSC1
Zinc finger protein 197	ZNF197

Figures



Figure 1

Representative immunohistochemical images of neutrophil infiltration in primary colorectal cancer and paracancerous tissues




Figure 2

Heat map generated by hierarchical clustering of differentially expressed mircRNAs in neutrophil and N2 neutrophil samples Group 1: neutrophil samples, Group 2: N2 neutrophil samples, the highly-expressed and lowly-expressed mircRNAs are represented in red and green respectively




Figure 3

The box plot shows the enrichment of total miRNAs in each sample




Figure 4

The volcano plot and Scatter Plot a. The volcano plot represented the differentially expressed miRNAs with statistical significance; b. Scatter plot illustrated the normalized circRNA expression in both groups. The x-axis represented the miRNA level in neutrophil group, the y-axis represented the miRNA level in N2 neutrophil group. In the volcano plot and Scatter Plot, the up-regulated miRNAs were represented as the red dots, while the down-regulated miRNAs were represented as the green dots.




Figure 5

Gene Ontology and KEGG Pathway enrichment for the host genes encoding DE miRNAs




Figure 6

The predicted downstream miRNA-targets network of the validated DEMircRNAs




Figure 7

The expression levels of DEmircRNA, TUSC1 and ZNF197 a. The expression levels of hsa-miR-3938 in neutrophil group and neutrophil+TGF-β1 group; b. The expression levels of hsa-miR-4780 in neutrophil group and neutrophil+TGF-β1 group; c. The expression levels of TUSC1 and ZNF197 in neutrophil group and neutrophil+TGF-β1 group