MiR-106b-5p attenuates neuropathic pain by regulating P2X4 receptor in the spinal cord in mice

Huiying Du

Guangzhou Medical University Second Affiliated Hospital

Danlei Wu

Guangzhou Medical University Second Affiliated Hospital

Shuotao Zhong

Guangzhou Medical University Second Affiliated Hospital

Xuhong Wei

Sun Yat-sen University Zhongshan School of Medicine

Zhongmin Yuan

Guangzhou Medical University Second Affiliated Hospital

Qingjuan Gong (✉ gqj01@sohu.com)

Guangzhou Medical University Second Affiliated Hospital <https://orcid.org/0000-0002-4427-8317>

Research Article

Keywords: P2X4R, miR-106b-5p, neuropathic pain, spinal dorsal horn, spared nerve injury

Posted Date: March 1st, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1366276/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

P2X4 receptor (P2X4R) can be up-regulated after nerve injury and its mediated spinal microglial activation makes a critical contribution to pathologically enhanced pain processing in the dorsal horn. Although some studies have partly clarified the mechanism underlying altered P2X4R expression, the specific mechanism is not well understood. MicroRNAs (miRNAs) are small noncoding RNAs, which control gene expression by binding with their target mRNAs. Thus, in the present study, whether miRNA was involved in pathogenesis of neuropathic pain by regulating the P2X4R was investigated. Our results showed that P2X4R was up-regulated in the spinal dorsal horn of mice following spared nerve injury (SNI), and 69 miRNAs (46 up-regulated and 23 down-regulated miRNAs) were differently expressed (fold change > 2.0, $P < 0.05$). P2X4R was a major target of miR-106b-5p (one of down-regulated miRNAs) with bioinformatics technology and quantitative real time PCR analysis validated the expressed change of miR-106b-5p, dual-luciferase reporter assays confirmed the correlation between them. Fluorescence in situ hybridization was used to show cell co-localization of P2X4R and miR-106b-5p in the spinal dorsal horn. Transfection with miR-106b-5p mimic into BV2 cells reversed the up-regulation of P2X4R induced by LPS. Moreover, miR-106b-5p overexpression significantly attenuated neuropathic pain induced by SNI with decreased expression of P2X4R mRNA and protein in the spinal dorsal horn, intrathecal miR-106b-5p antagomir induced pain behaviors and increased the expression of P2X4R in the spinal dorsal horn of naïve mice. These data suggest that miR-106b-5p can serve as an important regulator of neuropathic pain development by targeting P2X4R.

Introduction

Pain is defined as an unpleasant sensory and emotional experience associated with actual or potential tissue damage by the International Association for the Study of pain (Merskey H 1979), and neuropathic pain is defined as “pain caused by a lesion or disease of the somatosensory nervous system” (Yan XT et al. 2018). Neuropathic pain treatment is a great challenge and the outcomes remain unsatisfactory, mainly due to barely understood pathogenesis and a lack of well- defined molecular targets (Su S et al. 2017; van Hecke O et al. 2014).

P2X receptors are non-selective cation channels that open in response to adenosine triphosphate (ATP) binding, allowing the rapid flow of ions (K^+ , Na^+ , Ca^{2+}) across the membrane (Stokes L et al. 2017). Among the seven P2X receptors, P2X4 receptor (P2X4R) displays the most widespread tissue distribution and expression in neurons and the immune system (Suurväli J et al. 2017). Studies have found that P2X4R is closely related to the immune system and plays vital functions through monocytes, macrophages, dendritic cells, neutrophils, and T lymphocytes (Antonioli L et al. 2019). A great deal of evidence has shown that P2X4R-mediated spinal microglial functions make a critical role in the process of neuropathic pain (Yamashita T et al. 2016; Jurga AM et al. 2017). Our previous study has found that ATP activates p38 mitogen-activated protein kinase (p38 MAPK) by binding to endogenous P2X4R in microglia and subsequently increases the expression of P2X4R, thus promoting long-term potentiation (Gong QJ et al. 2009), which is the basis of the hypersensitization of nociceptive neurons in the spinal

dorsal horn. Moreover, it has been demonstrated that the blockade of P2X4R inhibits neuropathic pain-related behavior by preventing MMP-9 activation, then pronociceptive interleukins (IL-1 β , IL-18, IL-6) release in chronic constriction injury (CCI)-induced rat model, suggesting that P2X4R may indeed play a significant role in neuropathic pain development by modulating neuroimmune interactions in the spinal dorsal horn and dorsal root ganglion (DRG) (Jurga AM et al. 2017). Although these studies have partly elucidated the mechanism of altered P2X4R expression, the specific mechanism is not completely clear.

MicroRNAs (miRNAs) are highly deregulated in neuropathic pain, and might be critical molecules for pain treatment (Gong Q et al. 2015; Yang CH et al. 2016). MiRNAs can negatively regulate gene expression by translationally binding with the targeted mRNAs (Zheng Q et al. 2017; Liu S et al. 2017; Guo GC et al. 2017), and can serve as indicators in neuropathic pain pathophysiology (Zhang Y et al. 2019; Xu L et al. 2019; Wen J et al. 2019). It has been found that miR-98 might depress neuropathic pain development through modulating high mobility group AT-hook 2 (HMGA2) (Zhang Y et al. 2019) and that miR-30b is involved in the development of neuropathic pain, probably by regulating the expression of Nav1.3 (Su S et al. 2017). Nevertheless, whether a miRNA is involved in the pathogenesis of neuropathic pain by regulating the P2X4R has not been discovered.

In the present study, P2X4R was up-regulated and a microarray analysis was constructed to identify differential expressed (DE) miRNAs in the spinal dorsal horn following spared nerve injury (SNI). Moreover, the interaction of potential miRNA and P2X4R was evaluated to clarify the underlying mechanism involved in the development of neuropathic pain.

Animals And Methods

Animals

Male Balb/c mice aged 8 weeks were purchased from Animal Experimental Center, Sun Yat-Sen University, China (Guangzhou, China). Mice were housed in separated cages. The room temperature was kept at 24 ± 1 °C (297 ± 1 K) and humidity 50-60% under a 12:12 light-dark cycle. The animals were allowed to access food and water ad libitum. All experimental procedures were approved by the Institutional Animal Care and Use Committee of Guangzhou Medical University and were carried out in accordance with the guideline of the National Institutes of Health on animal care and the ethical guidelines for investigation of experimental pain in conscious animal.

Cell Culture and Transfection

Culture and transfection of BV2 cells were carried out as described elsewhere (Hongxia L et al. 2019). All cells were collected in high glucose Medium (Gibco) with 10% FBS (JRSscientific, Woodland, CA, USA), 100 mg/mL streptomycin, and 100 units/mL penicillin (Quality Biological, Gaithersburg, MD, USA). The cells were incubated at 37 °C, 95% O₂, and 5% CO₂ with a seeding density of 1×10^5 BV2 cells/mL. MiRNAs mimic or inhibitor (RiboBio, Guangzhou, China) were transfected into the cells one day later. 120 μ L Opti-Mem was used to dilute miR-106b-5p mimic (50 nM)/inhibitor (100 nM) or negative control (NC). The

sequences for miR-106b-5p mimic are as follow: Forward: 5'- UAAAGUGCUGACAGUGCAGAU-3', Reverse: 5'-AUUUCACGACUGUCACGU CUA-3'; the sequence for miR-106b-5p inhibitor is 5'-AUUUCACGACUGUCACG UCUA-3'. The final concentrations of mimic and inhibitor were 50 nM and 100 nM. 7.5 μ L RNAiMAX (Invitrogen, Carlsbad, CA) was simultaneously diluted by 120 μ L Opti-Mem and mixed. After 5 min, the mixture was placed into each well and high glucose medium was added. 200 ng LPS (Sigma) diluted in water was added to each 2 mL well 6 h before the cells were collected 48 h later for qPCR and Western-blot examinations.

Cell Proliferation Assay

Cell proliferation was evaluated using a cell counting kit-8 (CCK-8) assay (Beyotime, China). BV2 cells (5×10^3) were plated onto 96-well plates, then the LPS or miRNA mimic or miRNA inhibitor was added in the BV2 cells (Xu JH et al. 2020). Each well contained 100 μ L of medium supplemented with 10 μ L of CCK8 solution. The absorbance at 450 nm was measured to assess the degree of cell proliferation after incubating the cells at 37 °C for 1 h.

Luciferase Assay

A dual luciferase reporter assay was performed as outlined for a previous procedure (Huang Y et al. 2016). The pmirGLO dual-luciferase vector (pmirGLO vector), which contained both the firefly luciferase gene and the renilla luciferase gene, was purchased from Promega (Madison, WI, USA). P2X4R 3'UTR, including the predicted binding sites of miR-106b-5p, was inserted into the 3'UTR region downstream of the firefly luciferase gene of the pmirGLO vector (pmir-GLO-UTR). HEK293 cells were cultivated in high glucose medium (Gibco) with 10% FBS (JRSscientific, Woodland, CA, USA), 100 mg/mL streptomycin, and 100 units/mL penicillin (Quality Biological, Gaithersburg, MD, USA). The cells were incubated in a humidified incubator with 5% CO₂ at 37 °C. When the HEK293 cells had a confluency of 70-80%, a P2X4R-luciferase-reporter construct (pmirGLO-P2X4R 3'-UTR vector), containing the miR-106b-5p binding motif, was co-transfected with miR-106b-5p mimic at different doses of 10, 20, and 50 nM (20 μ mol/L) or NC mimic into HEK293 cells by using RNAiMAX (Invitrogen, Carlsbad, CA). After another 48 h of culture, we used 1 $\times$ passive lysis buffer to lyse the transfected cells, and 20 μ L supernatant was achieved for luciferase activity using the Dual-Luciferase Reporter Assay System (Promega). The ratio of firefly activity to renilla activity was recognized as relative reporter activity. Experiments were performed in triplicate and repeated three times.

Surgical Procedures and Drug Infusion

SNI surgeries were performed to mice according to our previous work (Liu Z et al. 2017). Briefly, surgery and experimental procedures on mice were performed under 0.4% sodium pentobarbital anesthesia (40 mg/kg body weight, i.p.). The skin of the lateral thigh was incised and the biceps femoris muscle was dissected bluntly to expose the left sciatic nerve and its three terminal branches (the sural, common peroneal and tibial nerves). The common peroneal and the tibial nerves were tightly ligated with 5.0 silk. Then, the nerve was transected distal to the ligation, and 2-4 mm length of nerve fiber was removed. Great

care was taken to avoid any contact with or stretching of the intact sural nerve. A complete hemostasis was confirmed and the wound was sutured in two layers. The sham group was given all procedures except ligation and transection. The method of intrathecal injection was described previously (Morioka N et al. 2019) Briefly, mice were restrained with the left hand and the injection was performed with the right hand. The vertebral landmarks for L5 and L6 vertebrae were identified by palpation. An injection into the subarachnoid space between the L5 and the L6 vertebrae was done via a 27-gage needle. Entry of the needle was confirmed with the presence of a tail flick. The injection volume of all other compounds was 5 μ L. In one part, the mice were divided into three groups: sham + scramble (saline), SNI + scramble (saline) and SNI + miR-106b-5p agomir (RiboBio, Guangzhou, China). In another part, the mice were divided into two groups, naïve + scramble and naïve + miR-106b-5p antagomir (RiboBio, Guangzhou, China). Beginning 1 day after naïve or 7 days after SNI surgery, intrathecal infusion was delivered once a day for 3 days, from day 1 to 3 for naïve mice and from day 7 to 9 for SNI mice. Mice with neurological deficits were excluded from the experiment.

Behavior Tests for Allodynia

Behavioral test was conducted between 8 and 10 AM in the morning. The 50% paw withdrawal thresholds (PWMTs) were measured at postoperative days 0, 5 and 7 for the evaluation of SNI model and at postoperative days 0, 5, 7, 8, 9, 10 and 14 during continuous miR-106b-5p agomir injection in SNI mice. And PWMTs were measured at postoperative days 0, 1, 2 and 3 during continuous miR-106b-5p antagomir injection in naïve mice. PWMT was assessed with method described previously (Yang L et al. 2017), using a set of von Frey hairs with logarithmically incremental stiffness from 0.04 to 2.04 g (0.04, 0.07, 0.16, 0.4, 0.60, 1.0, 1.4, 2.04 g). The filament force evoking paw withdrawals for more than 3 times in a round of testing was defined as the mechanical threshold. The cutoff force was 4 g. The observation of a positive response (paw lifting, shaking, or licking) within 5 s.

MiRNA Microarray Analysis

To functionally investigate a possible association between the changes in miRNA expression pattern and neuropathic pain development, we analyzed the relative expression of miRNAs in the spinal dorsal horn of mice in the sham and SNI groups by performing microarray analysis. The method was described in our previous study (Gong Q et al. 2015). MiRNA expression profiling was performed using the RiboArray platform (RiboBio, Guangzhou, China). Briefly, 1 μ g of total RNA was labeled with a Cy3 using a ULS™ microRNA Labeling Kit (Kreatech, Amsterdam, Netherlands) and hybridized on the microarray. Based on Sanger miRBase version 19.0 database, RiboBio designed 1263 specific oligos for 1281 mouse miRNA, where 1263 are non-redundant sequence. In addition, 54 RiboArray™ internal controls were used. We also put some probes for location identify function. T-test *P*-value of < 0.05 and fold-change (> 2.0) were applied to determine two DE sets of genes of six experimental samples.

qRT-PCR

For the quantitative real-time PCR (qPCR), the spinal dorsal cord was pooled to obtain sufficient RNA. Total RNA was extracted with the use of Trizol (Invitrogen, Carlsbad, CA), according to the method described previously (Du H et al. 2019; Liu Z et al. 2017). The quality and quantity of RNA were measured using a Nanodrop Spectrophotometer (Thermo Scientific) and samples with an absorbance ratio at 260/280 between 1.80-2.20 were considered acceptable. RNA degradation was not assessed. RNA dilutions were made up in nuclease-free water. For analysis of mRNA, the method was described previously (Du H et al. 2019; Liu Z et al. 2017). Reverse transcription was generated using the SuperScript™ III Reverse Transcriptase (Invitrogen, Carlsbad, CA) with a Gene Amp PCR System 9700 (Applied Biosystems). For miRNA analysis, 1 µg RNA was used for reverse transcription by miRNA 1st Strand cDNA Synthesis Kit (by stem-loop) (Vazyme, #MR101) according to the instructions. Quantitative Real-time PCR (qPCR) was carried out on a real-time detection instrument ViiA 7 Real-time PCR System (Applied Biosystems) using 2 × PCR master mix (Arraystar). All experiments were replicated three times. Relative expression of genes was calculated based on the $2^{-\Delta\Delta C_t}$ method using the mouse housekeeping GAPDH/U6 gene as an endogenous control.

Western Blotting

Based on established protocol (Wu Y et al. 2017), the spinal dorsal horn was homogenized in a RIPA buffer (Sangon, Shanghai, China). After centrifugation at $12\,000 \times g$ for 15 min at 4 °C, the supernatant was collected to analyze cytosolic proteins, and protein concentrations were determined with a BCA Kit (Thermo Fisher Scientific, Rockford, IL, USA). The contents of the proteins in the samples were measured using the Bio-Rad protein assay (Bio-Rad) and were then heated at 99 °C for 5 min. Samples of protein were separated by gel electrophoresis (SDS-PAGE) and transferred onto a polyvinylidene difluoride membrane. After blocking nonspecific binding sites with 5% Milk in Tris-buffered PBS containing 0.1% Tween-20 for 1 h at room temperature (RT), the membrane was incubated with primary antibody to rabbit anti-P2X4R antibody (1:1000, Alomone Labs, Israel), rabbit anti-tubulin antibody (1:10000, Sigma, USA), and rabbit anti-GAPDH antibody (1:5000, Cell Signaling Technology, USA) overnight at 4°C. After washing the proteins were detected using horseradish peroxidase-conjugated anti-rabbit secondary antibody (1:1000, Jackson Immuno-Research, USA) and visualized using Western peroxide reagent and luminol/enhancer reagent (Clarity Western ECL Substrate, Bio-Rad). The intensity of the blots was quantified via densitometry using Image J software. All cytosol protein bands were normalized to tubulin or GAPDH.

Fluorescence In Situ Hybridization

Fluorescence in situ hybridization (FISH) was performed with FISH frozen section kit (Genepharma, Shanghai, China). The locked nucleic acid probe complementary to miR-106b-5p (5'-TAAAGTGCTGACAGTGCAGAT-3') modified with Cy3 was synthesized by Genepharma (Genepharma, Shanghai, China). The animals were terminally anesthetized and perfused with saline followed by 4% paraformaldehyde in 0.1 M phosphate buffer (PB). After perfusion, the spinal dorsal horn of the lumbar enlargement was removed and postfixed in paraformaldehyde for 4 h and then replaced with 30%

sucrose in succession until sinking. Spinal sections (20 μm) were cut in a cryostat (Leica CM1900, Leica Biosystems, Germany). Each slide was added with citrate buffer solution for 15 min, the slides were washed twice with 0.1 M PBS for 5 min. Then the slides were incubated with protease K diluent (1:1000, Genepharm) for 20 min at 37 °C. The slides were washed 3 times at RT with 2 $\times$ SSC for 1 min. The slides were dehydrated in series of ethanol washes (70%, 80%, 90% and 100%) and air-dried. Slides were incubated with denaturing solution preheated at 78 °C at RT for 8 min. Then, repeat the dehydration step with ethanol washes. 4 $\mu\text{mol/L}$ of probe in hybridization mixture were heated at 73 °C for 5 min before added to each slide and cover the cover glass and seal with FixoGum Rubber Cement (YEASEN, USA). Then, slides were incubated 12-16 h at 37 °C in dark. The slides were washed with 43 °C preheated hybridization solution for 15 min, twice with 2 $\times$ SSC for 10 min and once with 0.1 M PBS for 10 min. After washing, the slides were perforated in 0.1 M PBS containing 0.3% Triton X-100 for 30 min at RT, blocked with blocking buffer (Beyotime, China) for 10 min, then washed with 0.1 M PBS once and incubated overnight with primary antibody to rabbit anti-P2X4R antibody (1:100, Solarbio, China) at 4 °C. The sections were then treated by Alexa 488-conjugated anti-Rabbit secondary antibody (1:500, Jackson Immuno-Research, USA) for 1 h at RT. The slides were imaged under a fluorescence microscope and the images were adjusted by Image J Version.

Statistics

All data are presented as means $\pm$ SEM. For comparisons between two groups, the *P* value was evaluated and calculated using a two-tailed unpaired t-test. When there were multiple groups involved, a one-way analysis of variance (ANOVA) was used; multiple factors were compared using two-way analysis of variance (ANOVA). Values of *P* < 0.05 were considered statistically significant.

Results

Up-regulation of P2X4R in the spinal dorsal horn after SNI

Compared to baseline PWMTs observed on day -1d, SNI induced a conspicuous reduction in PWMTs (*P* < 0.001, Fig. 1a). In contrast, no changes in the PWMTs were observed in the sham group during the observation period. To determine the expression of P2X4R in the spinal dorsal horn of SNI mice, qPCR and Western blot analysis were performed. Compared to the sham group, the SNI group showed an obvious up-regulation of P2X4R mRNA expression in the spinal dorsal horn (*P* < 0.05, Fig. 2b). The Western blot results showed that the P2X4R protein strongly increased after nerve injury (*P* < 0.01, Fig. 2c), which was consistent with the data from the behavioral test. The increased expression of P2X4R mRNA and protein in the spinal dorsal horn of SNI mice confirmed the potential ability of P2X4R to aggravate SNI-induced neuropathic pain.

The differentially expressed miRNAs in neuropathic pain mice

To investigate the expression of miRNAs in neuropathic pain, the expression levels of miRNAs in the spinal dorsal horn of neuropathic pain mice with microarray were analyzed. Results showed 69 miRNAs

were differently expressed (fold change > 2.0, $P < 0.05$), including 46 up-regulated miRNAs (Fig. 3a) and 23 down-regulated miRNAs (Fig. 3b). Focus on the down-regulated miRNAs, we found that P2X4R was the target of miR-106b-5p using TargetScan software. The matched seed sequences between miR-106b-5p and P2X4R 3'UTR were highly matched between human and mice (Fig. 3c). The TargetScan software demonstrated that the seed sequence for the miR-106b-5p position (1-7) was paired with P2X4R 3'UTR from 363 bps to 388 bps in humans and from 1165 bps to 1190 bps in the mice. Then, the expression of miR-106b-5p in SNI mice was verified by the qPCR method, and the results were consistent with those of the microarray. MiR-106b-5p showed a down-regulation trend in the spinal dorsal horn of the SNI mice ($P < 0.01$, Fig. 3d). To verify whether miR-106b-5p targets P2X4R 3'UTR, a dual-luciferase reporter vector containing the sequence of P2X4R 3'UTR was designed (pmirGLO P2X4R 3'UTR). When transfecting the P2X4R 3'UTR vector with three different doses of miR-106b-5p mimic (10, 20 and 50 nM) into HEK293 cells with RNAiMax, the miR-106b-5p mimic reduced relative luciferase activity in a dose-dependent manner ($P < 0.05$, Fig. 3e).

MiR-106b-5p mimic transfection inhibits the expression of P2X4R mRNA in BV2 cells

The result was consistent with previous study (Hongxia Let al. 2019), BV2 cells were activated after stimulating by LPS (Fig. 4a-b). Further, the cell viability was verified by using CCK-8 assay. The results showed that BV2 cells viability increased significantly in a dose-dependent relationship with LPS ($P < 0.05$, Fig. 4c). Besides, miR-106b-5p mimic or inhibitor did not affect BV2 cells viability ($P > 0.05$, Fig. 4d), which indicated miR-106b-5p mimic or inhibitor had no toxic effect on BV2 cells viability. To determine whether miR-106b-5p regulated the expression of P2X4R, miR-106b-5p mimic was transfected in BV2 cells. LPS was used to stimulate the BV2 cells 42 h later. The levels of miR-106b-5p and P2X4R mRNA were measured by qPCR, and the changes in P2X4R protein expression were determined by Western blot after transfection at 48 h. Compared to the naïve group, the transfection of miR-106b-5p mimic increased the expression of miR-106b-5p ($P < 0.0001$, Fig. 5a) and decreased the expression of P2X4R mRNA ($P < 0.0001$, Fig. 5b) and protein ($P < 0.05$, Fig. 6a) in BV2 cells. Compared to the naïve group, the LPS stimulation induced a reduction of miR-106b-5p ($P < 0.05$, Fig. 5a) and a significant increase in P2X4R at the mRNA ($P < 0.0001$, Fig. 5b) and protein level ($P < 0.05$, Fig. 6a). However, miR-106b-5p overexpression, by transfecting miR-106b-5p mimic, reversed the up-regulation of P2X4R (Fig. 5b and 6a) and down-regulation of miR-106b-5p ($P < 0.0001$, Fig. 5a). In addition, we found that miR-106b-5p inhibitor transfection down-regulated miR-106b-5p ($P < 0.0001$, Fig. 5c), while it up-regulated P2X4R mRNA ($P < 0.001$, Fig. 5d) and P2X4R protein ($P < 0.01$, Fig. 6b). Taken together, it has been demonstrated that miR-106b-5p may suppress the expression of P2X4R mRNA through its binding with P2X4R 3'UTR.

P2X4R is co-localized with miR-106b-5p in spinal dorsal horn of mice

In order to determine the localization of P2X4R and miR-106b-5p, the fluorescence in situ hybridization was performed in the spinal dorsal horn. As shown in Figure 7, fluorescence in situ hybridization results showed that miR-106b-5p was co-localized with P2X4R, indicating a potential interaction between miR-106b-5p and P2X4R.

Intrathecal miR-106b-5p agomir attenuates neuropathic pain and inhibits the expression of P2X4R in the spinal dorsal horn in SNI mice

To assess the exact impact of miR-106b-5p on neuropathic pain, miR-106b-5p agomir was intrathecally delivered to SNI mice on day 7, consecutively for 3 days. As shown in Fig. 8b, qPCR results confirmed that miR-106b-5p agomir up-regulated the expression of miR-106b-5p ($P < 0.01$, Fig. 8b) in SNI mice. At day 7 after SNI, 50% paw withdrawal threshold significantly decreased, indicating neuropathic pain was established ($P < 0.0001$, Fig. 8a). From day 3 after intrathecal injection, the decreased 50% paw withdrawal threshold in SNI increased by intrathecal injection with miR-106b-5p agomir ($P < 0.05$, Fig. 8a), not scrambled miRNA ($P > 0.05$, Fig. 8a). We further tested whether miR-106b-5p agomir could repress the expression of P2X4R. QPCR and Western blot analysis showed that miR-106b-5p agomir reversed the up-regulation of P2X4R both mRNA ($P < 0.05$, Fig. 8c) and protein levels ($P < 0.001$, Fig. 8d) in SNI mice.

Intrathecal miR-106b-5p antagomir induces pain behaviors and increases the expression of P2X4R in the spinal dorsal horn of naïve mice

To further explore the regulation of P2X4R by miR-106b-5p, miR-106b-5p was down-regulated in naïve mice by intrathecal injection with miR-106b-5p antagomir ($P < 0.05$, Fig. 9b). MiR-106b-5p antagomir was applied to naïve mice for 3 days and the sensitivity to the mechanical threshold was observed. We found that the 50% paw withdrawal threshold was significantly lower in miR-106b-5p antagomir treated mice than that of scrambled miRNAs treated mice ($P < 0.01$, Fig. 9a), demonstrating that miR-106b-5p antagomir produced pain behaviors in naïve mice. Furthermore, the spinal dorsal horn tissue was harvested at day 3 to assess the expression of P2X4R at the mRNA and protein levels. As shown in Fig. 9, intrathecal miR-106b-5p antagomir increases the expression of P2X4R mRNA ($P < 0.01$, Fig. 9c), as well as P2X4R protein ($P < 0.05$, Fig. 9d) in naïve mice.

Discussion

Nervous system damage results in neuropathic pain, and miRNAs are reported to regulate neuropathic pain. In our study, P2X4R mRNA and protein were increased in the SNI mouse model. Then, 69 DE miRNAs were identified in SNI mice, in which we also found that miR-106b-5p was decreased and that it could repress neuropathic pain development by targeting P2X4R. Our findings indicated that miR-106b-5p could be used as a novel therapeutic target of neuropathic pain through regulating P2X4R.

As is already known, purinergic receptor-mediated spinal microglial functions contribute significantly to pathologically enhanced pain management (Trang T et al. 2012; Inoue K et al. 2012). In recent decades, a growing body of evidence shows the expression of spinal P2X4R was up-regulated exclusively in spinal microglia after peripheral nerve injury (PNI) or nerve damage by various reasons (Tsuda M et al. 2003; Inoue K 2019; Wang Z et al. 2019; Zhou TT et al. 2014; Teixeira JM et al. 2019). Some studies have reported that the drugs (Gabapentin, Duloxetine, Paroxetine) for treating neuropathic pain can alleviate mechanical allodynia by decreasing P2X4R expression (Lu HF et al. 2018; Yamashita T et al. 2016; Nagata K et al. 2009). In the present study, we observed that the spinal level of P2X4R was elevated when

strong tactile hypersensitivity was developed, which supports the hypothesis that P2X4R is crucial as a mediator in the development of neuropathic pain. It has been demonstrated that several signaling molecules contribute to the upregulation of P2X4R in neuropathic pain (Tsuda M et al. 2003; Biber K et al. 2011; Yuan H et al. 2010; Toyomitsu E et al. 2012; Chen XM et al. 2015; Masuda T et al. 2014; MENG X W et al. 2017). Fibronectin is vital for the expression of P2X4R via interacting with other signaling molecules. Following PNI, an IRF8-IRF5 transcriptional axis and CCL21 up-regulated expression of P2X4R (Biber K et al. 2011; Masuda T et al. 2014), while P2X4R trafficking to the cell surface from microglial lysosomes is stimulated by CCL2 (Toyomitsu E et al. 2012). Besides, PI3K-Akt pathway plays a vital role in enhancing P2X4R expression at transcriptional levels, while MEK-ERK pathway at post-transcriptional levels (Tsuda M et al. 2009). Despite recent advances, understanding the transcriptional or translational regulatory mechanisms underlying changes in the expression and function of P2X4R under neuropathic pain conditions remains a major challenge.

MiRNAs are commonly recognized as mediators of gene expression and can serve as indicators in neuropathic pain pathophysiology (Zhang Y et al. 2019; Xu L, et al. 2019; Wen J et al. 2019). There has been a focus on studies associated miRNAs with chronic neuropathic pain states. MiRNAs are highly deregulated in neuropathic pain (Gong Q et al. 2015; Yang CH et al. 2016) and might be critical molecules for treatment. For example, miR-98 may depress neuropathic pain development through modulating HMGA2 (Zhang Y et al. 2019). MiR-134-5p can inhibit neuropathic pain progression through targeting Twist1 (Ji LJ et al. 2018). In addition, miR-190a-5p can contribute to DNP via targeting SLC17A6 (Yang D et al. 2017). These findings have paid our attention to mediators regulating P2X4R expression under neuropathic pain conditions including miRNA. Herein, microarray analysis was employed and 69 miRNAs showing differential expression in the spinal dorsal horn after SNI were identified (fold change > 2, $P < 0.05$, Fig. 3a and b). Based on common mechanism, miRNA plays a regulatory role mainly by silencing its target gene (El Bezawy R et al. 2019; Zhou Y et al. 2017), thus we focused on the down-regulated miRNAs in our present study. Among the down-regulated miRNAs, miR-106b-5p was found to target P2X4R through TargetScan software (Fig. 3c). Consistent with the results of the microarray, miR-106b-5p was decreased in SNI mice by the qPCR method (Fig. 3d). Further, the dual-luciferase assay verified that miR-106b-5p negatively regulated P2X4R through combining with P2X4R 3'UTR (Fig. 3e). Moreover, Fluorescence in situ hybridization showed that miR-106b-5p was co-localized with P2X4R (Fig. 7). Our results provide evidence for the interaction between miR-106b-5p and P2X4R.

In our study, BV2 cells were activated after stimulating by LPS (Fig. 4a-b), which are consistent with classical theory (Hongxia L et al. 2019). And the viability of BV2 cells increased in a dose-dependent manner following LPS treatment (Fig. 4c). BV2 cells are typically microglia cell lines, and P2X4R is almost exclusively expressed in microglia (Yamashita T et al. 2016). Further, P2X4R is increased at both mRNA and protein levels under LPS-induced inflammatory conditions in BV2 cells, and the up-regulation of P2X4R induced by LPS was reversed by BV2 cells transfected with miR-106b-5p mimic (Fig. 5b-c). These data indicated that miR-106b-5p regulates P2X4R at the transcription level in BV2 cells.

Moreover, we demonstrated that miR-106b-5p alleviated pain by inhibiting P2X4R and the increased expression of P2X4R was reversed (Fig. 8c-d) with intrathecal administration of miR-106b-5p agomir in SNI mice. Meanwhile, intrathecal administration of miR-106b-5p antagomir contributed to pain behaviors (Fig. 9a) and the up-regulation of P2X4R (Fig. 9c-d) in naïve mice. These results revealed that miR-106b-5p may function as an effective therapeutic target for neuropathic pain through regulating P2X4R.

It has been reported that miR-106b-5p in the blood is dysregulated and may be related with neuronal impairment in acute stroke (Wang W et al. 2014), epilepsy (Wang J et al. 2015), Alzheimer's disease (AD) (Yılmaz ŞG et al. 2016), so it has been proposed as a novel potential biomarker for these diseases. Moreover, miR-106b level in the AD brain decreases (Hébert SS et al. 2009). In the rat middle cerebral artery occlusion (MCAO) model, miR-106b-5p expression is significantly increased in cerebral cortex and its antagomir has the neuroprotective effects (Li P t et al. 2017). It has been indicated that miR-106b-5p was involved in microglial activation, microglial polarization, and neuroinflammation (Sağır F et al. 2021). In addition, miR-106b-5p can promote M2 polarization of microglia through inhibiting IRF1 and disturbing the functional binding among IRF1, IFN- β and IRF5 (Shi Y et al. 2020). As mentioned previously, an IRF8-IRF5 transcriptional axis can enhance the expression of P2X4R (Masuda T et al. 2014). Thus, we conclude that miR-106b-5p might regulate the expression of P2X4R by a complex interaction among IRF1, IRF5 and IRF8, accelerate microglia M2 polarization and alleviate neuropathic pain. A recent study found that miR-106b-5p has a suppressive role in the production of brain-derived neurotrophic factor (BDNF) (Sağır F et al. 2021). Interestingly, stimulation of microglial P2X4R induces the release of BDNF (Ulmann et al. 2008), which leads to allodynia after nerve injury (Coull JA et al. 2005). These show that miR-106b-5p may target P2X4R to reduce BDNF production. Briefly, the above evidence confirms the interaction between miR-106b-5p and P2X4R. A new study has found that miR-133b-5p is proved to reverse the allodynia established in central poststroke pain rats by regulation of P2X4R gene expression in the ventral posterolateral nucleus of the thalamus (Guo X et al. 2022). Whether miR-133b-5p targets P2X4R in the spinal dorsal horn to regulate neuropathic pain remains to be further studied.

Conclusion

The regulation of miR-106b-5p on P2X4R expression in SNI-induced neuropathic pain was the focus of our present research. We observed that miR-106b-5p alleviated neuropathic pain development by down-regulating P2X4R. P2X4R can enable neuropathic pain development, which can be reversed by miR-106b-5p. Our data revealed that miR-106b-5p may function as an effective therapeutic target for neuropathic pain through regulating P2X4R.

Declarations

Ethics Approval

This study was performed in line with the principles of the National Institutes of Health on animal care and the ethical guidelines for investigation of experimental pain in conscious animal. Approval was

granted by the Institutional Animal Care and Use Committee of Guangzhou Medical University.

Consent for Publication

All authors have approved this manuscript and concur in its publication.

Competing Interests

The authors report no conflicts of interest in this work.

Availability of data and materials

The datasets generated during and analysed during the current study are available from the corresponding author on reasonable request.

Data availability statement

All relevant raw data in this research will be freely available to any researcher wishing to use them for non-commercial purposes, without breaching participant confidentiality.

Funding

This work was supported by grant from the Science & Technology Planning Project of Guangzhou (NO. 202102010051).

Author Contributions

H.D. (Huiying Du) and D.W. (Danlei Wu) performed the qPCR, Western blot, FISH, cell culture and transfection, luciferase assay, animal experiments and behavioral tests. S.Z. (Shuotao Zhong) performed the cell proliferation assay. X.W. (Xuhong Wei) and Z.Y. (Zhongmin Yuan) analyzed the data. Q.G. (Qingjuan Gong) designed and supervised the study. H.D. and Q.G. wrote the manuscript. All authors reviewed the manuscript.

Acknowledgements

We thank Yun Ling (Department of Breast Surgery, the Second Affiliated Hospital of Guangzhou Medical University, Guangzhou 510260, Guangdong, China) for guiding the revision of manuscripts.

References

1. Merskey H (1979 Jun) Pain terms: a list with definitions and notes on usage. Recommended by the IASP Subcommittee on Taxonomy. *Pain* 6(3):249. <https://pubmed.ncbi.nlm.nih.gov/460932/>
2. Yan XT, Zhao Y, Cheng XL, He XH, Wang Y, Zheng WZ, Chen H, Wang YL (2018 Jun) Inhibition of miR-200b/miR-429 contributes to neuropathic pain development through targeting zinc finger E box binding protein-1. *J Cell Physiol* 233(6):4815–4824. <https://doi.org/10.1002/jcp.26284>

3. Su S, Shao J, Zhao Q, Ren X, Cai W, Li L, Bai Q, Chen X, Xu B, Wang J, Cao J, Zang W MiR-30b Attenuates Neuropathic Pain by Regulating Voltage-Gated Sodium Channel Nav1.3 in Rats. *Front Mol Neurosci*. 2017 May5;10:126. <https://doi.org/10.3389/fnmol.2017.00126>
4. van Hecke O, Austin SK, Khan RA, Smith BH, Torrance N (2014 Apr) Neuropathic pain in the general population: a systematic review of epidemiological studies. *Pain* 155(4):654–662. <http://dx.doi.org/10.1016/j.pain.2013.11.013>
5. Stokes L, Layhadi JA, Bibic L, Dhuna K, Fountain SJ P2X4 Receptor Function in the Nervous System and Current Breakthroughs in Pharmacology. *Front Pharmacol*. 2017 May23; 8:291. <https://doi.org/10.3389/fphar.2017.00291>
6. Suurväli J, Boudinot P, Kanellopoulos J, Rüütel Boudinot S P2X4: A fast and sensitive purinergic receptor. *Biomed J*. 2017 Oct; 40 (5):245–56. <https://doi.org/10.1016/j.bj.2017.06.010>
7. Antonioli L, Blandizzi C, Fornai M, Pacher P, Lee HT, Haskó G (2019 Aug) P2X4 receptors, immunity, and sepsis. *Curr Opin Pharmacol* 47:65–74. <https://doi.org/10.1016/j.coph.2019.02.011>
8. Yamashita T, Yamamoto S, Zhang J, Kometani M, Tomiyama D, Kohno K, Tozaki-Saitoh H, Inoue K, Tsuda M Duloxetine Inhibits Microglial P2X4 Receptor Function and Alleviates Neuropathic Pain after Peripheral Nerve Injury. *PLoS One*. 2016 Oct21; 11(10):e0165189. <https://doi.org/10.1371/journal.pone.0165189>
9. Jurga AM, Piotrowska A, Makuch W, Przewlocka B, Mika J Blockade of P2X4 Receptors Inhibits Neuropathic Pain-Related Behavior by Preventing MMP-9 Activation and, Consequently, Pronociceptive Interleukin Release in a Rat Model. *Front Pharmacol*. 2017 Feb16;8:48. <https://doi.org/10.3389/fphar.2017.00048>
10. Gong QJ, Li YY, Xin WJ, Zang Y, Ren WJ, Wei XH, Li YY, Zhang T, Liu XG ATP induces long-term potentiation of C-fiber-evoked field potentials in spinal dorsal horn: the roles of P2X4 receptors and p38 MAPK in microglia. *Glia*. 2009 Apr15; 57 (6):583–91. <https://doi.org/10.1002/glia.20786>
11. Gong Q, Lu Z, Huang Q, Ruan L, Chen J, Liang Y, Wang H, Yue Y, Feng S Altered microRNAs expression profiling in mice with diabetic neuropathic pain. *Biochem Biophys Res Commun*. 2015 Jan9; 456 (2):615–20. <http://dx.doi.org/10.1016/j.bbrc.2014.12.004>
12. Yang CH, Wang Y, Sims M, Cai C, He P, Yue J, Cheng J, Boop FA, Pfeffer SR, Pfeffer LM MiRNA203 suppresses the expression of protumorigenic STAT1 in glioblastoma to inhibit tumorigenesis. *Oncotarget*. 2016 Dec20; 7 (51):84017–29. <https://doi.org/10.18632/oncotarget.12401>
13. Zheng Q, Cui X, Zhang D, Yang Y, Yan X, Liu M, Niang B, Aziz F, Liu S, Yan Q, Liu J miR-200b inhibits proliferation and metastasis of breast cancer by targeting fucosyltransferase IV and α 1,3-fucosylated glycans. *Oncogenesis*. 2017 Jul10; 6 (7):e358. <https://doi.org/10.1038/oncsis.2017.58>
14. Liu S, Zhang F, Shugart YY, Yang L, Li X, Liu Z, Sun N, Yang C, Guo X, Shi J, Wang L, Cheng L, Zhang K, Yang T, Xu Y The early growth response protein 1-miR-30a-5p-neurogenic differentiation factor 1 axis as a novel biomarker for schizophrenia diagnosis and treatment monitoring. *Transl Psychiatry*. 2017 Jan10; 7 (1):e998. <https://doi.org/10.1038/tp.2016.268>

15. Guo GC, Wang JX, Han ML, Zhang LP, Li L (2017 Apr) microRNA-761 induces aggressive phenotypes in triple-negative breast cancer cells by repressing TRIM29 expression. *Cell Oncol (Dordr)* 40(2):157–166. <https://doi.org/10.1007/s13402-016-0312-6>
16. Zhang Y, Liu HL, An LJ, Li L, Wei M, Ge DJ, Su Z miR-124-3p attenuates neuropathic pain induced by chronic sciatic nerve injury in rats via targeting EZH2. *J Cell Biochem.* 2019Apr; 120 (4):5747–55. <https://doi.org/10.1002/jcb.27861>
17. Xu L, Wang Q, Jiang W, Yu S, Zhang S (2018) MiR-34c Ameliorates Neuropathic Pain by Targeting NLRP3 in a Mouse Model of Chronic Constriction Injury. *Neuroscience.* 2019 Feb 10; 399:125 – 34. <https://doi.org/10.1016/j.neuroscience.12.030>
18. Wen J, He T, Qi F, Chen H MiR-206-3p alleviates chronic constriction injury-induced neuropathic pain through targeting HDAC4. *Exp Anim.* 2019 May8; 68 (2):213–20. <https://doi.org/10.1538/expanim.18-0091>
19. Zhang Y, Su Z, An LJ, Li L, Wei M, Ge DJ, Liu HL miR-98 acts as an inhibitor in chronic constriction injury-induced neuropathic pain via downregulation of high-mobility group AT-hook 2. *J Cell Biochem.* 2019Jun; 120 (6):10363–9. <https://doi.org/10.1002/jcb.28320>
20. Hongxia L, Yuxiao T, Zhilei S, Yan S, Yicui Q, Jiamin S, Xin X, Jianxin Y, Fengfeng M, Hui S Zinc inhibited LPS-induced inflammatory responses by upregulating A20 expression in microglia BV2 cells. *J Affect Disord.* 2019 Apr15; 249:136–42. <https://doi.org/10.1016/j.jad.2019.02.041>
21. Trang T, Beggs S, Salter MW (2012 Apr) ATP receptors gate microglia signaling in neuropathic pain. *Exp Neurol* 234(2):354–361 PMID: 22116040. <http://dx.doi.org/10.1016/j.expneurol.2011.11.012>
22. Inoue K, Tsuda M (2012 Apr) Purinergic systems, neuropathic pain and the role of microglia. *Exp Neurol* 234(2):293–301. <https://doi.org/10.1016/j.expneurol.2011.09.016>
23. Xu JH, Zhao JX, Jiang MY, Yang LP, Sun ML, Wang HW (2020 Mar) MiR-193 promotes cell proliferation and invasion by ING5/PI3K/AKT pathway of triple-negative breast cancer. *Eur Rev Med Pharmacol Sci* 24(6):3122–3129. https://doi.org/10.26355/eurev_202003_20679
24. Huang Y, Li Y, Wang FF, Lv W, Xie X, Cheng X, Over-Expressed miR-224 Promotes the Progression of Cervical Cancer via Targeting RASSF8. *PLoS One.* 2016 Sep14; 11 (9):e0162378. <https://doi.org/10.1371/journal.pone.0162378>
25. Liu Z, Liang Y, Wang H, Lu Z, Chen J, Huang Q, Sheng L, Ma Y, Du H, Gong Q LncRNA expression in the spinal cord modulated by minocycline in a mouse model of spared nerve injury. *J Pain Res.* 2017 Oct24; 10:2503–14. <https://doi.org/10.2147/jpr.s147055>
26. Morioka N, Kodama K, Tomori M, Yoshikawa K, Saeki M, Nakamura Y, Zhang FF, Hisaoka-Nakashima K, Nakata Y (2019) May Stimulation of nuclear receptor REV-ERBs suppresses production of pronociceptive molecules in cultured spinal astrocytes and ameliorates mechanical hypersensitivity of inflammatory and neuropathic pain of mice. *Brain Behav Immun.* ; 78:116 – 30. <https://doi.org/10.1016/j.bbi.2019.01.014>
27. Yang L, Dong F, Yang Q, Yang PF, Wu R, Wu QF, Wu D, Li CL, Zhong YQ, Lu YJ, Cheng X, Xu FQ, Chen L, Bao L, Zhang X (2017) FGF13 Selectively Regulates Heat Nociception by Interacting with Nav1.7.

- Neuron. 22:806–821 e9. <http://dx.doi.org/10.1016/j.neuron.2017.01.009>. 4
28. Du H, Liu Z, Tan X, Ma Y, Gong Q Identification of the Genome-wide Expression Patterns of Long Non-coding RNAs and mRNAs in Mice with Streptozotocin-induced Diabetic Neuropathic Pain. *Neuroscience*. 2019 Mar15; 402:90–103. <https://doi.org/10.1016/j.neuroscience.2018.12.040>
 29. Wu Y, Ma S, Xia Y, Lu Y, Xiao S, Cao Y, Zhuang S, Tan X, Fu Q, Xie L, Li Z, Yuan Z Loss of GCN5 leads to increased neuronal apoptosis by upregulating E2F1- and Egr-1- dependent BH3-only protein Bim. *Cell Death Dis*. 2017 Jan26; 8 (1):e2570. <https://doi.org/10.1038/cddis.2016.465>
 30. Tsuda M, Shigemoto-Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S, Salter MW, Inoue K P2X4 receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature*. 2003 Aug 14;424(6950):778 – 83. doi:10.1038/nature01786. <https://doi.org/10.1038/nature01786>
 31. Inoue K (2019 Aug) Role of the P2X4 receptor in neuropathic pain. *Curr Opin Pharmacol* 47:33–39. doi: 10.1016/j.coph.2019.02.001.. <https://doi.org/10.1016/j.coph.2019.02.001>Epub 2019 Mar 14
 32. Wang Z, Mei W, Wang Q, Guo R, Liu P, Wang Y, Zhang Z, Wang L Role of Dehydrocorybulbine in Neuropathic Pain After Spinal Cord Injury Mediated by P2X4 Receptor. *Mol Cells*. 2019 Feb28;42(2):143–150. <https://doi.org/10.14348/molcells.2018.0028>
 33. Zhou TT, Wu JR, Chen ZY, Liu ZX, Miao B Effects of dexmedetomidine on P2X4Rs, p38-MAPK and BDNF in spinal microglia in rats with spared nerve injury. *Brain Res*. 2014 Jun3; 1568:21–30. <https://doi.org/10.1016/j.brainres.2014.04.025>
 34. Teixeira JM, Dos Santos GG, Neves AF, Athie MCP, Bonet IJM, Nishijima CM, Farias FH, Figueiredo JG, Hernandez-Olmos V, Alshaibani S, Tambeli CH, Müller CE, Parada CA (2019 Feb) Diabetes-induced Neuropathic Mechanical Hyperalgesia Depends on P2X4 Receptor Activation in Dorsal Root Ganglia. *Neuroscience* 1:398158–398170. <https://doi.org/10.1016/j.neuroscience.2018.12.003>
 35. Lu HF, Xu CY, Zhang L, Gan L, Chen C, Yan MY, Guo XN, Fang Q, Xu GY, Zhang YB, Ni JQ, Zhao HR (doi: 10.1016/j.neulet.2018.09.023. Epub 2018) A new central post-stroke pain rat model: autologous blood injected thalamic hemorrhage involved increased expression of P2X4 receptor. *Neurosci Lett* 2018 Nov 20;687:124–130. <https://doi.org/10.1016/j.neulet.2018.09.023>
 36. Nagata K, Imai T, Yamashita T, Tsuda M, Tozaki-Saitoh H, Inoue K Antidepressants inhibit P2X4 receptor function: a possible involvement in neuropathic pain relief. *Mol Pain*. 2009 Apr23; 5:20. doi: 10.1186/1744-8069-5-20. <https://doi.org/10.1186/1744-8069-5-20>
 37. Biber K, Tsuda M, Tozaki-Saitoh H, Tsukamoto K, Toyomitsu E, Masuda T, Boddeke H, Inoue K Neuronal CCL21 up-regulates microglia P2X4 expression and initiates neuropathic pain development. *EMBO J*. 2011 May 4;30(9):1864-73. doi: 10.1038/emboj.2011.89. Epub 2011 Mar 25. <https://doi.org/10.1038/emboj.2011.89>
 38. \Yuan H, Zhu X, Zhou S, Chen Q, Zhu X, Ma X, He X, Tian M, Shi X Role of mast cell activation in inducing microglial cells to release neurotrophin. *J Neurosci Res*. 2010 May 1;88(6):1348-54. doi: 10.1002/jnr.22304. <https://doi.org/10.1002/jnr.22304>
 39. Toyomitsu E, Tsuda M, Yamashita T, Tozaki-Saitoh H, Tanaka Y, Inoue K (2012) Jun;8(2):301 – 10 CCL2 promotes P2X4 receptor trafficking to the cell surface of microglia. *Purinergic Signal*. doi:

- 10.1007/s11302-011-9288-x. Epub 2012 Jan 6. <https://doi.org/10.1007/s11302-011-9288-x>
40. Chen XM, Xu J, Song JG, Zheng BJ, Wang XR (2015) Jan;114(1):150-7 Electroacupuncture inhibits excessive interferon- γ evoked up-regulation of P2X4 receptor in spinal microglia in a CCI rat model for neuropathic pain. *Br J Anaesth*. doi: 10.1093/bja/aeu199. Epub 2014 Jul 29. <https://doi.org/10.1093/bja/aeu199>
 41. Masuda T, Iwamoto S, Yoshinaga R, Tozaki-Saitoh H, Nishiyama A, Mak TW, Tamura T, Tsuda M, Inoue K Transcription factor IRF5 drives P2X4R+reactive microglia gating neuropathic pain. *Nat Commun*. 2014 May13; 5:3771. doi: 10.1038/ncomms4771. <https://doi.org/10.1038/ncomms4771>
 42. Meng XW, Gao JL, Zuo JL, Wang LN, Liu SL, Jin XH, Yao M, Namaka M Toll-like receptor-4/p38 MAPK signaling in the dorsal horn contributes to P2X4 receptor activation and BDNF over-secretion in cancer induced bone pain. *Neurosci Res*. 2017 Dec; 125:37–45. doi: 10.1016/j.neures.2017.06.006. Epub 2017 Jun 28. <https://doi.org/10.1016/j.neures.2017.06.006>
 43. Tsuda M, Toyomitsu E, Kometani M, Tozaki-Saitoh H, Inoue K (2009) Sep;13(9B):3251-9 Mechanisms underlying fibronectin-induced up-regulation of P2X4R expression in microglia: distinct roles of PI3K-Akt and MEK-ERK signalling pathways. *J Cell Mol Med*. doi: 10.1111/j.1582-4934.2009.00719.x. Epub 2009 Feb 27. <https://doi.org/10.1111/j.1582-4934.2009.00719.x>
 44. Ji LJ, Su J, Xu AL, Pang B, Huang QM (2018) MiR-134-5p attenuates neuropathic pain progression through targeting Twist1. *J Cell Biochem*. Sep 6. doi: 10.1002/jcb.27486. Epub ahead of print. <https://doi.org/10.1002/jcb.27486>
 45. Yang D, Yang Q, Wei X, Liu Y, Ma D, Li J, Wan Y, Luo Y The role of miR-190a-5p contributes to diabetic neuropathic pain via targeting SLC17A6. *J Pain Res*. 2017 Oct4; 10:2395–2403. doi: 10.2147/JPR.S133755. <https://doi.org/10.2147/jpr.s133755>
 46. El Bezawy R, Tinelli S, Tortoreto M, Doldi V, Zuco V, Folini M, Stucchi C, Rancati T, Valdagni R, Gandellini P, Zaffaroni N miR-205 enhances radiation sensitivity of prostate cancer cells by impairing DNA damage repair through PKC ϵ and ZEB1 inhibition. *J Exp Clin Cancer Res*. 2019 Feb4; 38 (1):51. <https://doi.org/10.1186/s13046-019-1060-z>
 47. Zhou Y, Huang T, Siu HL, Wong CC, Dong Y, Wu F, Zhang B, Wu WK, Cheng AS, Yu J, To KF, Kang W IGF2BP3 functions as a potential oncogene and is a crucial target of miR-34a in gastric carcinogenesis. *Mol Cancer*. 2017 Apr11; 16 (1):77. <https://doi.org/10.1186/s12943-017-0647-2>
 48. Wang W, Sun G, Zhang L, Shi L, Zeng Y (2014) Nov-Dec;23(10):2607–2613 Circulating microRNAs as novel potential biomarkers for early diagnosis of acute stroke in humans. *J Stroke Cerebrovasc Dis*. doi: 10.1016/j.jstrokecerebrovasdis.2014.06.002. Epub 2014 Oct 5. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2014.06.002>
 49. Wang J, Yu JT, Tan L, Tian Y, Ma J, Tan CC, Wang HF, Liu Y, Tan MS, Jiang T, Tan L Genome-wide circulating microRNA expression profiling indicates biomarkers for epilepsy. *Sci Rep*. 2015 Mar31; 5:9522. doi: 10.1038/srep09522. <https://doi.org/10.1038/srep09522>
 50. Yilmaz ŞG, Erdal ME, Özge AA, Sungur MA (2016 Aug) Biomarkers of Alzheimer's Disease? *OMICS*. 20:456–461. doi: 10.1089/omi.2016.0099. Can Peripheral MicroRNA Expression Data Serve as

51. Hébert SS, Horré K, Nicolaï L, Bergmans B, Papadopoulou AS, Delacourte A, De Strooper B (2009 Mar) MicroRNA regulation of Alzheimer's Amyloid precursor protein expression. *Neurobiol Dis* 33(3):422–428. doi: 10.1016/j.nbd.2008.11.009.. <https://doi.org/10.1016/j.nbd.2008.11.009>Epub 2008 Dec 9
52. Li P, Shen M, Gao F, Wu J, Zhang J, Teng F, Zhang C (2017 May) An Antagomir to MicroRNA-106b-5p Ameliorates Cerebral Ischemia and Reperfusion Injury in Rats Via Inhibiting Apoptosis and Oxidative Stress. *Mol Neurobiol* 54(4):2901–2921. doi: 10.1007/s12035-016-9842-1.. <https://doi.org/10.1007/s12035-016-9842-1>Epub 2016 Mar 29
53. Sağır F, Ersoy Tunalı N, Tombul T, Koral G, Çırak S, Yılmaz V, Türkoğlu R, Tüzün E miR-132-3p, miR-106b-5p, and miR-19b-3p Are Associated with Brain-Derived Neurotrophic Factor Production and Clinical Activity in Multiple Sclerosis: A Pilot Study.*Genet Test Mol Biomarkers*. 2021Nov; 25(11):720–726. doi: 10.1089/gtmb.2021.0183. <https://doi.org/10.1089/gtmb.2021.0183>
54. Shi Y, Zhang B, Zhu J, Huang W, Han B, Wang Q, Qi C, Wang M, Liu F miR-106b-5p Inhibits IRF1/IFN- β Signaling to Promote M2 Macrophage Polarization of Glioblastoma.*Onco Targets Ther*. 2020 Jul30; 13:7479–7492. doi: 10.2147/OTT.S238975. <https://doi.org/10.2147/ott.s238975>
55. Ulmann L, Hatcher JP, Hughes JP, Chaumont S, Green PJ, Conquet F, Buell GN, Reeve AJ, Chessell IP, Rassendren F Up-regulation of P2X4 receptors in spinal microglia after peripheral nerve injury mediates BDNF release and neuropathic pain.*J Neurosci*. 2008 Oct29;28(44):11263–8. doi: 10.1523/JNEUROSCI.2308-08.2008. <https://doi.org/10.1523/jneurosci.2308-08.2008>
56. Coull JA, Beggs S, Boudreau D, Boivin D, Tsuda M, Inoue K, Gravel C, Salter MW, De Koninck Y BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain.*Nature*. 2005 Dec15;438(7070):1017–21. doi: 10.1038/nature04223. <https://doi.org/10.1038/nature04223>
57. Guo X, Lu J, Yan M, Wang Y, Yang Y, Li H, Shen H, Diao S, Ni J, Lu H, Zhao H, Chen G MicroRNA-133b-3p Targets Purinergic P2X4 Receptor to Regulate Central Poststroke Pain in Rats.*Neuroscience*. 2022 Jan15;481:60–72. doi: 10.1016/j.neuroscience.2021.10.015. Epub 2021 Oct 21. <https://doi.org/10.1016/j.neuroscience.2021.10.015>

Figures

Figure 1

The description of animal and cytology experimental process. **a** PWMTs were measured at postoperative days and after the drug administration. **b** MiRNAs mimic or inhibitor were transfected into BV2 cells.

Figure 2

The pain behavior of neuropathic pain induced by SNI and the change in expression of P2X4R mRNA and protein. **a** Responses of the paw to mechanical stimulus. *** $P < 0.001$, **** $P < 0.0001$ vs. sham. Two-way ANOVA, $n = 6$ mice. **b** The increased expression of P2X4R mRNA in the spinal dorsal horn of SNI mice, $t_4 = 3.539$, * $P < 0.05$, $n = 3$ mice. **c** The increased expression of P2X4R protein in the spinal dorsal horn of SNI mice, $t_4 = 4.670$, ** $P < 0.01$. Two-tailed unpaired t-test, $n = 3$ mice.

Figure 3

MiR-106b-5p targets P2X4R 3'UTR. **a-b** The heat map showed the differential expression of miRNA in SNI mice (fold change > 2.0 , $P < 0.05$). MiR-106b-5p was downregulated in neuropathic pain induced by SNI. **c** miR-106b-5p directly targets P2X4R 3'UTR. The matched seed region between miR-106b-5p and P2X4R 3'UTR predicted by software in red. **d** The decreased expression of miR-106b-5p in the spinal dorsal horn of SNI mice, $t_4 = 4.862$, ** $P < 0.01$. Two-tailed unpaired t-test, $n = 3$ mice. **e** Transfection of miR-106b-5p mimic with wild-type P2X4R 3'UTR plasmid vector reduced relative luciferase activity. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. vehicle; ### $P < 0.001$ vs. scramble. One-way ANOVA, $n = 3$.

Figure 4

The morphology of BV2 cells and the effect of LPS or miR-106b-5p mimic or miR-106b-5p inhibitor on BV2 cell viability. **a** The picture showed normal morphology of BV2 cells. **b** The picture presents the morphology of LPS-stimulated BV2 cells. **c** Cell viability was evaluated by CCK-8 assay. The affection of LPS on cell growth was denoted as a percentage of normal control. * $P < 0.05$, **** $P < 0.0001$ vs. naïve. One-way ANOVA, $n = 8$. **d** Compared with naïve group, the affection of those compounds on cell growth showed no different. $P > 0.05$. One-way ANOVA, $n = 5$.

Figure 5

MiR-106b-5p regulated the expression of P2X4R mRNA in BV2 cells. **a** MiR-106b-5p expression increased in the group of MiRNAs mimic. **b** P2X4R mRNA expression decreased in the group of MiRNAs mimic. **c** MiR-106b-5p expression decreased in the group of inhibitor. **d** P2X4R mRNA expression increased in the

group of inhibitor. * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$ vs. naïve; ## $P < 0.01$, #### $P < 0.0001$ vs. LPS (100 ng/mL). One-way ANOVA, $n = 3$ mice.

Figure 6

MiR-106b-5p regulated the expression of P2X4R protein in BV2 cells. **a** P2X4R protein decreased in the group of MiRNAs mimic. **b** P2X4R protein increased in the group of inhibitor. ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$ vs. naïve. One-way ANOVA, $n = 3$ mice.

Figure 7

Fluorescence in situ hybridization was used to show co-localized immunoreactivity for P2X4R and miR-106b-5p. **a** A RNA probe of miR-106b-5p to perform fluorescence in situ hybridization on frozen sections of spinal dorsal horn. **b** Immunofluorescent staining of P2X4R performed on those frozen sections of spinal dorsal horn from naïve mice. **c** P2X4R and miR-106b-5p were merged in picture. Scale bars: 200 μm .

Figure 8

MiR-106b-5p agomir down-regulated P2X4R and alleviated neuropathic pain. **a** The effect of the up-regulated of miR-106b-5p on mechanical allodynia was assessed by PWMTs at postinduction days 0, 5, 7, 8, 9, 10 and 14. **** $P < 0.0001$ vs. sham + scramble; # $P < 0.05$, ## $P < 0.01$ vs. SNI + scramble. Two-way ANOVA, $n = 6$ mice. **b** The relative expression of miR-106b-5p with intrathecal miR-106b-5p agomir/scramble in the spinal dorsal horn of SNI mice was determined by qPCR. **c** The relative expression of P2X4R mRNA with intrathecal miR-106b-5p agomir/scramble in the spinal dorsal horn of SNI mice was determined by qPCR. * $P < 0.05$, ** $P < 0.01$ vs. sham + scramble; # $P < 0.05$, ## $P < 0.01$ vs. SNI + scramble. One-way ANOVA, $n = 3$ mice. **d** P2X4R protein expression in the spinal dorsal horn of SNI mice at 14 days injected with intrathecal miR-106b-5p agomir, $t_4 = 9.357$, *** $P < 0.001$. Two-tailed unpaired t-test, $n = 3$ mice.

Figure 9

The down-regulated of miR-106b-5p and up-regulated of P2X4R intensified neuropathic pain in naïve mice. **a** The effect of the down-regulated of miR-106b-5p on mechanical allodynia was assessed by PWMTs at days -1, 1, 2 and 3. ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$ vs. naïve + scramble. Two-way ANOVA, $n = 6$ mice. **b** The relative expression of miR-106b-5p at 3 days in the spinal dorsal horn of mice after intrathecal injection, $t_4 = 3.213$, * $P < 0.05$. **c** The relative expression of P2X4R mRNA at 3 days in the spinal dorsal horn of mice after intrathecal injection, $t_4 = 6.151$, ** $P < 0.01$. **d** The relative expression of P2X4R protein at 3 days in the spinal dorsal horn of mice, $t_4 = 4.330$, * $P < 0.05$. Two-tailed unpaired t-test, $n = 3$ mice.